# Evaluating the potential of biochar in mitigating greenhouse gases emission and nitrogen retention in dairy manure based silage corn cropping systems

By

# Waqar Ashiq

A thesis submitted to the School of Graduate Studies In partial fulfillment of the requirements for the degree of

Master of Science

Boreal Ecosystems and Agricultural Sciences

School of Science and the Environment

Grenfell Campus

Memorial University of Newfoundland

August, 2018

St. John's, Newfoundland and Labrador

Evaluating the potential of biochar in mitigating greenhouse gases emission and nitrogen

retention in dairy manure based silage corn cropping systems

By

Waqar Ashiq

A Thesis submitted to the School of Graduate Studies

In partial fulfillment of the requirements for the degree of

Master of Science

Boreal Ecosystems and Agricultural Sciences

Approved:

Dean of the Graduate School

Supervisor

Date

Committee members:

Dr. Jianghua Wu

Dr. Lakshman Galagedara

#### Abstract

Greenhouse gas (GHGs) emissions from the agriculture sector have been accelerating global warming potential (GWP) and greenhouse gas intensities (GHGI). About 8 % of GHG emissions in Canada are contributed by the agriculture sector mainly through methane (CH<sub>4</sub>) and nitrous oxide (N<sub>2</sub>O). Out of these emissions, 50 % is contributed by manure and fertilizer application to land. Biochar (BC), a stable carbon-rich product has been observed to reduce GHG emissions from soil, increase soil pH, improve soil moisture, enhance nutrient retention in soil and increase biomass production in many crop plants. However, these effects are not constant across all soil types, environmental and climatic conditions, and cropping systems. This study aimed to evaluate the effect of BC on GHGs emissions, soil nitrate and ammonium retention, soil pH, plant nitrogen concentration and dry matter production in dairy manure (DM) based silage corn cropping system in western Newfoundland, Canada. Two sources of dairy manure (DM<sub>1</sub>,  $DM_2$ ), inorganic N (IN), their combination with BC ( $DM_1+B$ ,  $DM_2+B$ , and IN+B), and control ( $N_0$ ) were used as experimental treatments. Results showed that BC application to DM<sub>1</sub>, DM<sub>2</sub> and IN reduced cumulative CO<sub>2</sub> emission by 16, 25.5 and 26.5 %, CH<sub>4</sub> emission 184, 200 and 293 %, and N<sub>2</sub>O emission by 95, 86 and 93 %, respectively. BC treatments exhibited significantly higher soil moisture (SM) contents at all sampling points than non-BC treatments. It also reduced the GWP by 24.9, 34.5, and 37 %, and GHGI by 30, 37.5, 43.4 %, respectively. Furthermore, BC enhanced the  $NO_3^-$  and  $NH_4^+$ retention in topsoil (decreased their leaching to deep soil) which improved plant N concentration and dry matter yield of silage corn crop. Conclusively, BC application to

soil exhibited to be a promising tool for the mitigation of GHGs emissions, GWP, GHGI and to enhance soil fertility and crop dry matter yield simultaneously.

### Acknowledgment

My sincere gratitude to Almighty God for the grace and strength He gave me to successfully complete my master's course. First, I would like to thank my kind and respected supervisor, Dr. Mumtaz Akhtar Cheema for inviting me to work on this topic with his research team and my co-supervisor, Dr. Jianghua Wu for his purposeful advice and encouragement throughout my program, Dr. Lakshman Galagedara as the advisory committee member and Dr. Raymond Thomas for his support and comments. Special thanks to Dr. Vanessa Kavanagh and her team for providing the logistic support for execution of this research project.

I am particularly thankful to Dr. Muhammad Nadeem for his support in all phases of my research. Special thanks to Ms. Sanaz Rajabi, Ms. Crystal McCall, Dr. Tao Yuan, Mr. Waqas Ali and Mr. Muhammad Zaeem for their support during lab and field work. Special Thanks to Grenfell Campus Research office and School of graduate studies Memorial University of Newfoundland for providing tuition fee and stipend. Financial support for execution of this research project from Research and Development Corporation (RDC), NL, and department of fisheries and land resources is greatly acknowledged.

I especially thank my parents, siblings and my family members for their support and encouragement.

Waqar Ashiq

# Table of contents

Abstract	iii
Acknowledgment	. v
Table of contents	vi
List of tables	. X
List of figures	xi
List of Abbreviations	xii
Chapter 1	.1
1. General introduction and Overview	. 1
1.1. Introduction	.1
1.1.1. Overall scenario of greenhouse gas emissions	. 1
1.1.2. Agricultural greenhouse gas emissions	.2
1.1.3. Nitrogen loss through nitrate leaching and N <sub>2</sub> O	.3
1.1.4. Mitigation options	.6
1.1.5. Role of biochar role in mitigating gaseous and N losses	.7
1.2. Purpose of the thesis	.8
1.3. Thesis organization	10
1.4. References	11
1.5. Co-authorship statement	24
Chapter 2	25
2. Biochar amendment reduces greenhouse gases emission in silage corn cropping	
system following dairy manure and inorganic nitrogen application	25
2.1. Abstract	25
2.2. Introduction	26

2.3. Ma	terials and Methods	31
2.3.1.	Study site	31
2.3.2.	Experimental setup and treatments	32
2.3.3.	Crop husbandry	34
2.3.4.	Greenhouse gas sampling and analysis	37
2.3.5.	Global warming potential and greenhouse gas intensity calculation.	39
2.3.6.	Dry matter production	39
2.3.7.	Statistical analysis	40
2.4. Res	sults	41
2.4.1.	CO <sub>2</sub> flux	41
2.4.2.	CH <sub>4</sub> flux	44
2.4.3.	N <sub>2</sub> O flux	45
2.4.4.	Global warming potential, dry matter yield and greenhouse gas inter 45	nsity
2.5. Dis	cussion	49
2.5.1.	Meteorological conditions:	49
2.5.2.	CO <sub>2</sub> emission	53
2.5.3.	CH <sub>4</sub> flux	54
2.5.4.	N <sub>2</sub> O flux	55
2.6. Co	nclusion	56
2.7. Ret	erences	57
Chapter 3		73
3. Biomas	s production, and nitrogen dynamics of silage corn as influenced by	
	s production, and introgen dynamics of shage corn as influenced by	

	3.1.	Abstract	73
	3.2.	Introduction	74
	3.3.	Material and Methods:	78
	3.3.	Study site	78
	3.3.2	2. Experimental setup	78
	3.3.	B. Crop husbandry	80
	3.3.4	4. Soil sampling and analysis	81
	3.3.	5. Soil pH determination	
	3.3.	5. Dry matter production	83
	3.3.	7. Plant tissue nitrogen concentration	83
	3.3.	3. Statistical analysis	83
	3.4.	Results	84
	3.4.	Nitrate and ammonium dynamics	84
	3.4.2	2. Soil pH	90
	3.4.	B. Plant tissue N concentration	92
	3.4.4	4. Dry matter yield	94
	3.5.	Discussion	96
	3.6.	Conclusion	98
	3.7.	References	99
Cha	apter 4		114
4	. Gen	eral discussion and conclusion	114
	4.1.	Effect of biochar application on GHGs emission, GWP and GHGI	115
	4.2.	Biochar effects $NO_3^-$ and $NH_4^+$ retention, soil pH, plant N concentrat	ion and
	dry ma	tter production	

4.3.	Conclusion and Recommendations	.119
4 4		101
4.4.	References	.121

# List of tables

Table 2.1: Basic soil properties of the experimental site at Pynn's Brook Research S	
	31
Table 2.2: Chemical analysis of dairy manures used in the study	33
Table 2.3: Physio-chemical properties of biochar used in the study	36
Table 2.4: Cumulative greenhouse gas emission/absorption of CO <sub>2</sub> , CH <sub>4</sub> and N <sub>2</sub> O (Kg	g
ha <sup>-1</sup> season <sup>-1</sup> ), global warming potential (kg CO <sub>2</sub> equivalent), silage corn dry matter yi	eld
(kg ha <sup>-1</sup> ), and greenhouse gas intensity (kg CO <sub>2</sub> equivalent per kg dry matter yield),	
during growing season 2016	42
Table 2.5: Cumulative greenhouse gas emission/absorption of CO <sub>2</sub> , CH <sub>4</sub> and N <sub>2</sub> O (Kg	g
ha <sup>-1</sup> season <sup>-1</sup> ), global warming potential (kg CO <sub>2</sub> equivalent), silage corn dry matter yi	eld
(kg ha <sup>-1</sup> ), and greenhouse gas intensity (kg $CO_2$ equivalent per kg dry matter yield),	
during growing season 2017	43
Table 2.6: Weather conditions (biweekly average) during silage corn growing season i	in
2016 and 2017 at Pynn's Brook Research Station	50
Table 3.1: Silage corn genotypes used in the experiment	80
Table 3.2: Plant tissues nitrogen concentration (%) in silage corn genotypes under	
different experimental treatments during growing season 2016	93
Table 3.3: Plant tissues nitrogen concentration (%) in silage corn genotypes under	
different experimental treatments during growing season 2017	93
Table 3.4: Dry matter yield (Mg ha <sup>-1</sup> ) of three silage corn genotypes during growing	
season 2016	95
Table 3.5: Dry matter yield (Mg ha <sup>-1</sup> ) of three silage corn genotypes during growing	
season 2017	95

# List of figures

Figure 2.1: Silage corn seeding with SAMCO 2200	35
Figure 2.2: Recording soil temperature, moisture and EC (a), sampling chambers fixed	
over sampling spots in a crop row (b), GHGs sample collection with (c)	38
Figure 2.3: Temporal greenhouse gas emission during growing season 2016	17
Figure 2.4: Temporal greenhouse gas emission during growing season 2017	18
Figure 2.5: (a) Air temperature and rainfall at Pynn's Brook Research Station (b) soil	
temperature at 5cm depth, and (c) soil moisture at different greenhouse gas sampling	
dates during growing season 2016.	51
Figure 2.6: (a) Air temperature and rainfall at Pynn's Brook Research Station (b) soil	
temperature at 5cm depth, and (c) soil moisture at different greenhouse gas sampling	
dates during growing season 2017	52
Figure 3.1: Soil nitrate (NO <sub>3</sub> <sup>-</sup> ) and ammonium (NH <sub>4</sub> <sup>+</sup> ) concentrations (mg $g^{-1}$ dry soil) in	L
experimental treatments at four crop growth stages during 2016	38
Figure 3.2: Soil nitrate (NO <sub>3</sub> <sup>-</sup> ) and ammonium (NH <sub>4</sub> <sup>+</sup> ) concentrations (mg g <sup>-1</sup> dry soil) in	L
experimental treatments at four crop growth stages during 2017	39
Figure 3.3: Soil pH at 20 cm depth at different crop growth stages during 2016 and 2017	,
	<del>)</del> 1

# List of Abbreviations

BC - Biochar

- CEC Cation exchange capacity
- CFIA Canadian food inspection agency
- CH<sub>4</sub> Methane
- $CO_2$  Carbon dioxide
- DAMA Days after manure application
- DM Dairy manure
- DMY Dry matter yield
- DM<sub>1</sub> Dairy manure with high nitrogen
- DM1+B Dairy manure with high nitrogen + biochar
- DM<sub>2</sub> Dairy manure with low nitrogen
- $DM_2+B-Dairy$  manure with low nitrogen + biochar
- DOC Dissolved organic carbon
- DON Dissolved organic nitrogen
- EC Electrical conductivity
- ECD Electrical conductivity detector
- FID Flame ionization detector
- GHG Greenhouse gas
- GHGI greenhouse gas intensity
- GWP Global warming potential
- IN Inorganic nitrogen
- IN+B Inorganic nitrogen + biochar

IPCC - Intergovernmental panel on climate change

- KCL Potassium chloride
- MOP Murate of potash
- N<sub>2</sub>O Nitrous oxide
- $NH_4^+$  Ammonium
- $(NH_2)_2 CO Urea$
- $(NH_4)_2SO_4 Ammonium sulfate$
- NH<sub>4</sub>NO<sub>3</sub> Ammonium nitrate
- NO Nitric oxide
- $NO_2^- Nitrite$
- $NO_3 Nitrate$
- N<sub>2</sub>O Nitrous oxide
- $N_2 Dinitrogen$
- $Pg Petagram (10^{15})$
- PVC Polyvinyl chloride
- RCBD Randomized complete block design
- SM Soil moisture
- SOC Soil organic carbon
- SOM Soil organic matter
- SRF Slow release fertilizer
- ST Soil temperature
- TCD Thermal conductivity detector
- TSP Triple superphosphate

# **Chapter 1**

#### 1. General introduction and Overview

#### 1.1. Introduction

#### 1.1.1. Overall scenario of greenhouse gas emissions

The emission of greenhouse gases (GHGs) into the atmosphere is the greatest environmental issue of the current time. The unprecedented increase in GHG emissions lead to significant changes on the face of world climate. The main GHGs of consternation include carbon dioxide  $(CO_2)$ , methane  $(CH_4)$  and nitrous oxide  $(N_2O)$ . These gases gather in the atmosphere and cause greenhouse effect which leads to global warming. The global warming effect of GHGs can be reported using a global warming potential (GWP) value which quantifies the warming effect of GHGs relative to CO2 over a set time period (20 years, 100 years or 500 years). The GWP of CH<sub>4</sub> and N<sub>2</sub>O over a time horizon of 100 years (GWP<sub>100</sub>) is 25 and 298 times greater than CO<sub>2</sub> respectively (IPCC, 2007). The total annual anthropogenic GHGs emissions had reached to  $49\pm4.5$  (90 % confidence interval) gigatons of carbon dioxide equivalent (GtCO<sub>2</sub>eq) in 2010. Industrialized countries emit 2.5 times more GHGs than developing countries. Different economic sectors contribute towards global GHGs emissions. Energy supply sector contributed 35 % (17 GtCO<sub>2</sub>eq), agriculture, forestry and other land use contributed 24 % (12 GtCO<sub>2</sub>eq), 21 % (10 GtCO2eq) by industry, transport 14 % (7 GtCO2eq), and construction sector 6.4 % (3.2 GtCO<sub>2</sub>eq) towards a total of 49 GtCO<sub>2</sub>eq GHGs emission in 2010 (IPCC, 2014a). The main controlling forces of GHGs emissions include financial structure, the flow of income, choices of investment, policies, people behavior, consumption patterns, energy resources and land use change (IPCC, 2014a).

## 1.1.2. Agricultural greenhouse gas emissions

The agriculture industry is the largest contributor to the global anthropogenic non-CO<sub>2</sub> GHGs emission, accounting for 54 % of the global non-CO<sub>2</sub> emissions in 2005 (U.S. EPA, 2011). Individual GHGs from the agriculture sector out of total anthropogenic emissions from all sources constitute CO<sub>2</sub> (15 %), CH<sub>4</sub> (50 %) and N<sub>2</sub>O (66 %). Between 1990-2010, non-CO<sub>2</sub> emissions from the agriculture sector grew by 0.9 % per year, totaled to be 5.2-5.8 GtCO<sub>2</sub>eq per year and comprised about 10-12 % of total anthropogenic emissions (IPCC, 2014a; Tubiello et al., 2013). Agricultural N<sub>2</sub>O emissions are anticipated to increase by 35–60 % by 2030 due to increase in the use of N fertilizers and manures (FAO, 2003). Agricultural activities and practices that emit GHGs include enteric fermentation, dairy manure (DM) storages, and inorganic fertilizer application to soil, rice cultivation, manure management, crop residues and biomass burning. The emissions from enteric fermentation and soils represent about 70 %, paddy rice 9-11 %, biomass burning 6-12 % and manure management accounts for 7-8 % of total agricultural emissions. Global emissions from manure management grew by 1.1 % per year between 1961-2010 from 0.57 to 0.99 GtCO<sub>2</sub>eq per year (Herrero et al., 2013) and by 3.9 % per year from inorganic fertilizers (0.07 to 0.68  $GtCO_2$  eq per year) (Tubiello et al., 2013). Following this trend the inorganic fertilizers will become the single largest source of non-CO<sub>2</sub> GHGs after enteric fermentation in less than 10 years (IPCC, 2014a).

#### **1.1.3.** Nitrogen loss through nitrate leaching and N<sub>2</sub>O

The atmosphere consists of 70 % of inert nitrogen  $(N_2)$  by volume although there is an enormous amount of N in the atmosphere; it is the most absorbed and most limiting nutrient in the soil. Nitrogen (N) is essential plant nutrient and inevitable for plant growth and development as it is the constituent of all proteins, chlorophyll, coenzymes and nucleic acids. Therefore, appropriate C:N ratio is vital for successful crop production. The use of inorganic nitrogen (IN) is one of the major contributors towards the increased agricultural production in past decades. The total amount of N fertilizers applied to cropland increased from 11.3 Tg N per year in 1961 to 107.6 Tg N per year in 2013 (Lu and Tian, 2017). The N applied to crops or mineralized N from organic sources in soil is taken up by plants, lost in gaseous form or leached in the form of nitrates ( $NO_3^{-}$ ). Soil ammonium (NH<sub>4</sub><sup>+</sup>) concentration is usually low as most of the NH<sub>4</sub><sup>+</sup> is readily converted to  $NO_3^-$  which is not retained in the soil due to the negative charge on soil clay particles (Di and Cameron, 2002). Excessive use of IN and DM application enhance the risk of  $NO_3$  leaching and consequently increase cost of production, pollute water bodies and pose a serious threat to human health (Fan et al., 2017; Forge et al., 2016; Jokela et al., 2014; Long and Sun, 2012; Masaka et al., 2015). The  $NO_3^-$  leaching losses of 55-59 kg per hectare per year with DM application, 30-35 kg per hectare per year from compost application and 25-33 kg per hectare per year from IN application has been reported in a six-year maize-alfalfa crop rotation (Basso and Ritchie, 2005). The annual estimated loss of NO<sub>3</sub>-N from a cornfield in Manitoba was 160 kg per hectare per year (Hargrave and Shavkewich, 1997), whereas, 39-55 kg per hectare per year  $NO_3^{-1}$  load has been reported

in wheat-maize cropping system in Southern Turkey (Ibrikci et al., 2015). IN and DM applications in agricultural fields increase the concentration of  $NH_4^+$  and  $NO_3^-$  which cause a temporary surplus of these ions in the soil. These ions then undergo nitrification and denitrification processes in soil and release N<sub>2</sub>O as a byproduct (Chapuis-lardy et al., 2007; Inselsbacher et al., 2011). During nitrification,  $NH_4^+$  is oxidized to  $NO_3^-$  via nitrite  $(NO_2^-)$  releasing N<sub>2</sub>O as a byproduct (Wrage et al., 2001). This is a two-step autotrophic process. The first step is mostly carried out by ammonia oxidizing bacteria (AOB) and ammonia-oxidizing archaea (AOA) whereas, the second step is facilitated by *Nitrobacter sp.*, and *Nitrospira sp.*, under aerobic conditions (Clough et al., 2001). Nitrification process explained by Hossini et al. (2015) in the Equations (1.1) and (1.2).

$$2NH_4^+ + 3O_2 \rightarrow 2NO_2^- + 2H_2O + 4H^+$$
(1.1)

$$2NO_2^- + O_2^- \to 2NO_3^- \tag{1.2}$$

During denitrification,  $NO_3^-$  is reduced to dinitrogen (N<sub>2</sub>) via N<sub>2</sub>O. This heterotrophic process takes place under oxygen deficit conditions (Flechard et al., 2007; Toyoda et al., 2011). The denitrification process is shown in the Equations (1.3), (1.4), and (1.5) as described by (Hossini et al., 2015).

$$2NO_3^- + 2H_2O \to 2NO_2^- + 2H_2O$$
(1.3)

$$2NO_2^- + 2H_2 \rightarrow N_2O^- + H_2O + 2OH^-$$
(1.4)

$$N_2 0 + H_2 \rightarrow N_2 \uparrow + H_2 0 \tag{1.5}$$

An important factor which explains the dominance of nitrification and denitrification in the soil is "soil compaction". Soil compaction reduces the soil porosity which leads to increased water-filled pore spaces and reduced oxygen diffusion resulting in the anaerobic conditions which favor denitrification and N<sub>2</sub>O production in the soil (Bessou et al., 2010). Increase in soil temperature results in the development of anaerobic microsites in the soil causing denitrification and N<sub>2</sub>O emissions (Smith et al., 2003). Generally, soils are source of N<sub>2</sub>O, but under certain conditions like high soil moisture (SM) or low N availability, the soil may act as sink of N<sub>2</sub>O either caused by the consumption of  $N_2O$  by nitrifier during nitrifier denitrification or reduction of  $N_2O$  to  $N_2$ during denitrification (Chapuis-lardy et al., 2007). Nitrogen is applied frequently to crops and is one of the priciest inputs in corn production. The N application averages 12-15 % of the variable costs in a corn - silage corn and 18-21 % in corn - corn cropping systems (Plastina, 2018). In spite of that, its cost-benefit ratio generally surpasses that of other fertilizer inputs, but farmers must use N efficiently to maximize its value. At the same time, farmers must ensure that adequate supply of N is available to crop and its yield is not limited by the shortfall of N throughout the growing season. This means minimizing N loss through the application of the right amount of N at right time and developing strategies which decrease N losses. There are different strategies adopted to reduce GHGs emissions and  $NO_3^-$  leaching losses including nitrification inhibitors (NI), adaptation of fertilizer management practices to increase efficiency and reducing excess N application to the soil (Burney et al., 2010), Sustainable agriculture intensification (Garnett et al., 2013; Thomson et al., 2012), land drainage and biological N fixation (Rees et al., 2013).

# 1.1.4. Mitigation options

Different nutrient management strategies and crop management practices have been reported to reduce GHGs emissions and N losses. Carbon (C) sequestration has great potential to reduce global warming and climate change. Decreasing the atmospheric C by 3.5-4 Gt per year could limit the temperature increase to 2 °C by 2050 (Meinshausen et al., 2009; Minasny et al., 2017), a threshold level beyond which climate change would have a significant impact (IPCC, 2014b). This annual reduction in atmospheric  $CO_2$ concentration could be enhanced by increasing soil C sequestration in agricultural soils globally by 0.4 % per year, producing a C sink of 1.2 petagrams (Pg) per year (Paustian et al., 2016). The soils of agroecosystems have a technical potential of 1.2-3.1 billion ton C sequestration per year (Lal, 2011). Different crop management practices can improve soil C sequestration including crop residues incorporation (Coppens et al., 2006), burial of crop residues and crop rotation (Hirel et al., 2007), addition of perennial crops in rotation, no-tillage (West and Post, 2002), legume-based cropping systems (Drinkwater et al., 1998), cover cropping (Mazzoncini et al., 2011), organic amendments like manure (Maillard and Angers, 2014) and biochar (Bera et al., 2016). Nutrient management strategies have also been practiced to reduce N losses and improve N use efficiency (NUE) that includes the use of slow-release fertilizers (Ye et al., 2013), judicious

fertilizer application (Francis, 1992), variable rate technologies (VRT) (Gatti et al., 2018), use of nitrification inhibitors (NI) (Zhang et al., 2015), and plant trait selection (Ju et al., 2015). However, the integration of these practices is complicated to get dual benefits of GHGs reduction and mitigation of  $NO_3^-$  losses.

# 1.1.5. Role of biochar role in mitigating gaseous and N losses

Biochar (BC), is a recalcitrant black C material produced by the baking of organic matter under low oxygen conditions and relatively low temperature (<700 °C) (Dong et al., 2017; Lehmann and Joseph, 2009; Shackley et al., 2010). It has multifaceted benefits including reduction of GHGs emissions (Woolf et al. 2010) and soil compaction, improvement of soil pH, aggregate stability (Wang et al., 2017), soil, permeability, porosity, water holding capacity (WHC), (Basso et al., 2013; Ulyett et al., 2014; Randolph et al., 2017), soil nutrients retention (Uzoma et al., 2011), nutrient availability (Subedi et al., 2016), carbon sequestration (Atkinson et al., 2010; Khare and Goyal, 2013; Laird, 2008; Matovic, 2011), soil organic matter (SOM), cation exchange capacity (CEC), microbial growth and shelter, microbial activity and pollutant degradation (Amendola et al., 2017; Reed et al., 2017; Tan et al., 2017; Upadhyay et al., 2014; Wang et al., 2014). Organic C is mainly stored in the form of stable aromatic compounds in BC and is not decomposed easily even in suitable environmental conditions (Sohi et al., 2010). It has the ability to sequester C for thousands of years due to its recalcitrant chemical composition (Fowles, 2007). The application of BC has historic importance in some parts of the world in order to sequester C. The Terra Preta soils in the Amazon Basin have a large amount of sequestered C as a consequence of the application of BC by American

Indian people thousands of years ago (Lehmann et al., 2006). It also possesses the nutrient holding ability and improves soil physiochemical and biological life thus improves soil structure, aeration, WHC and provides microsites that act as shelter for soil microbes (Johnson et al., 2007; Lehmann et al., 2006) resulting in increased soil fertility (Koide et al., 2011). It improves the plants nutrient use efficiency which reduces nutrients leaching to watercourses thereby reduces environmental pollution (International Biochar Initiative, 2012). In addition to C sequestration, recently it has shown a great potential to mitigate GHGs emissions from agricultural soils, enhance  $NO_3^-$  retention, improve nutrient use efficiency and increase plant yield (Felber et al., 2014; Liu et al., 2012a; Taghizadeh-Toosi et al., 2011; Zhang et al., 2011). The nitrate (NO<sub>3</sub><sup>-</sup>) and ammonium  $(NH_4^+)$  retention in BC amended soils is linked with the reduction of N<sub>2</sub>O emission. The BC has high CEC due to the negative charge on its surface which allows it to retain cations such as  $NH_4^+$  (Cheng et al., 2006; Yao et al., 2012). The  $NO_3^-$  and  $NH_4^+$  retention in BC amended soils can decrease leaching of these nutrients from these soils (Winning, 2014). A significant decrease in  $NH_4^+$  loss was observed by Lehmann et al. (2002) and Angst et al. (2013). Few studies have reported no effect or increase in NO<sub>3</sub><sup>-</sup> leaching after BC application which could be due to weak adsorption and subsequent desorption of NO<sub>3</sub><sup>-</sup> by BC due to its low anion exchange capacity (Kameyama et al., 2012; Singh et al., 2010).

## **1.2.** Purpose of the thesis

The principal aim of this thesis was to investigate the potential role of BC in mitigating GHG losses, global warming potential (GWP), greenhouse gas intensity (GHGI), soil

 $NO_3^-$  and  $NH_4^+$  retention, soil pH, N uptake, and dry matter yield of silage corn following DM and inorganic N fertilizer application. Studies were carried out with the following specific objectives:

- i- To assess the GHGs emissions from organic and inorganic sources of nitrogen application in silage corn cropping systems
- ii- To determine the role of BC application in the reduction of GHG emission in silage corn cropping systems in western Newfoundland
- iii- To estimate GWP and GHGI of silage corn cropping systems
- iv- To determine the role of BC application on soil  $NO_3^-$  and  $NH_4^+$  retention in silage corn amended with DM and IN fertilizer application
- v- To compare the effects of dairy manure and IN alone and co-application of BC on soil pH, N uptake and biomass production of silage corn.

## **1.3.** Thesis organization

This thesis is divided into four chapters with the relevant literature reviewed at the start of each chapter.

Chapter one: Provides a brief overview of global GHG emissions, N losses in the form of  $NO_3^-$  and  $N_2O$  into the atmosphere, different mitigation strategies to lessen GHGs emissions and  $NO_3^-$  leaching and the potential role of BC as a mitigation strategy.

Chapter two: This chapter describes a comparative study about the effect of different organic and inorganic N sources ( $DM_1$ ,  $DM_2$ , IN and BC) on GHGs emissions, GWP, and GHGI of silage corn grown under field conditions in western Newfoundland, Canada.

Chapter three: It covers the potential role of BC on the  $NO_3^-$  and  $NH_4^+$  retention, soil pH, plant N concentration and dry matter production of different silage corn genotypes.

Chapter four: This chapter comprised of general discussion, conclusion and recommendations of the study.

#### 1.4. References

- Amendola, C., Montagnoli, A., Terzaghi, M., Trupiano, D., Oliva, F., Baronti, S.,
  Miglietta, F., Chiatante, D., Scippa, G.S., 2017. Short-term effects of biochar on
  grapevine fine root dynamics and arbuscular mycorrhizae production. Agric.
  Ecosyst. Environ. 239, 236–245.
- Angst, T.E., Patterson, C.J., Reay, D.S., Anderson, P., Peshkur, T.A., Sohi, S.P., 2013.
  Biochar Diminishes Nitrous Oxide and Nitrate Leaching from Diverse Nutrient
  Sources. J. Environ. Qual. 42, 672–682.
- Atkinson, C.J., Fitzgerald, J.D., Hipps, N.A., 2010. Potential mechanisms for achieving agricultural benefits from biochar application to temperate soils: A review. Plant Soil 337, 1–18.
- Basso, A.S., Miguez, F.E., Laird, D.A., Horton, R., Westgate, M., 2013. Assessing potential of biochar for increasing water-holding capacity of sandy soils. GCB Bioenergy 5, 132–143.
- Basso, B., Ritchie, J.T., 2005. Impact of compost, manure and inorganic fertilizer on nitrate leaching and yield for a 6-year maize-alfalfa rotation in Michigan. Agric. Ecosyst. Environ. 108, 329–341.
- Bera, T., Collins, H.P., Alva, A.K., Purakayastha, T.J., Patra, A.K., 2016. Biochar and manure effluent effects on soil biochemical properties under corn production. Appl. Soil Ecol. 107, 360–367.

- Bessou, C., Mary, B., Léonard, J., Roussel, M., Gréhan, E., Gabrielle, B., 2010.Modelling soil compaction impacts on nitrous oxide emissions in arable fields. Eur.J. Soil Sci. 61, 348–363.
- Bruinsma, J., 2003. World agriculture : towards 2015/2030, An FAO Study.
- Burney, J.A., Davis, S.J., Lobell, D.B., 2010. Greenhouse gas mitigation by agricultural intensification. Proc. Natl. Acad. Sci. 107, 12052–12057.
- Chapuis-lardy, L., Wrage, N., Metay, A., Chotte, J.L., Bernoux, M., 2007. Soils, a sink for N2O? A review. Glob. Chang. Biol. 13, 1–17.
- Cheng, C.H., Lehmann, J., Thies, J.E., Burton, S.D., Engelhard, M.H., 2006. Oxidation of black carbon by biotic and abiotic processes. Org. Geochem. 37, 1477–1488.
- Clough, T.J., Di, H.J., Cameron, K.C., Sherlock, R.R., Metherell, A.K., Clark, H., Rys,G., 2001. Accounting for the utilization of a N2O mitigation tool in the IPCCinventory methodology for agricultural soils. Nutr. Cycl. Agroecosystems 78, 1–14.
- Coppens, F., Garnier, P., De Gryze, S., Merckx, R., Recous, S., 2006. Soil moisture, carbon and nitrogen dynamics following incorporation and surface application of labelled crop residues in soil columns. Eur. J. Soil Sci. 57, 894–905.
- Di, H.J., Cameron, K.C., 2002. Nitrate leaching in temperate agroecosystems: Sources, factors and mitigating strategies. Nutr. Cycl. Agroecosystems 64, 237–256.
- Dong, X., Li, G., Lin, Q., Zhao, X., 2017. Quantity and quality changes of biochar aged for 5 years in soil under field conditions. Catena 159, 136–143.

- Drinkwater, L.E., Wagoner, P., Sarrantonio, M., 1998. Legume-based cropping systems have reuced carbon and nitrogen losses. Nature 396, 262–265.
- Fan, J., Xiao, J., Liu, D., Ye, G., Luo, J., Houlbrooke, D., Laurenson, S., Yan, J., Chen,
  L., Tian, J., Ding, W., 2017. Effect of application of dairy manure, effluent and
  inorganic fertilizer on nitrogen leaching in clayey fluvo-aquic soil: A lysimeter
  study. Sci. Total Environ. 592, 206–214.
- Felber, R., Leifeld, J., Horák, J., Neftel, A., 2014. Nitrous oxide emission reduction with greenwaste biochar: Comparison of laboratory and field experiments. Eur. J. Soil Sci. 65, 128–138.
- Flechard, C.R., Ambus, P., Skiba, U., Rees, R.M., Hensen, A., van Amstel, A., Dasselaar, A. van den P. van, Soussana, J.F., Jones, M., Clifton-Brown, J., Raschi, A., Horvath, L., Neftel, A., Jocher, M., Ammann, C., Leifeld, J., Fuhrer, J., Calanca, P., Thalman, E., Pilegaard, K., Di Marco, C., Campbell, C., Nemitz, E., Hargreaves, K.J., Levy, P.E., Ball, B.C., Jones, S.K., van de Bulk, W.C.M., Groot, T., Blom, M., Domingues, R., Kasper, G., Allard, V., Ceschia, E., Cellier, P., Laville, P., Henault, C., Bizouard, F., Abdalla, M., Williams, M., Baronti, S., Berretti, F., Grosz, B., 2007. Effects of climate and management intensity on nitrous oxide emissions in grassland systems across Europe. Agric. Ecosyst. Environ. 121, 135–152.
- Forge, T., Kenney, E., Hashimoto, N., Neilsen, D., Zebarth, B., 2016. Compost and poultry manure as preplant soil amendments for red raspberry: Comparative effects on root lesion nematodes, soil quality and risk of nitrate leaching. Agric. Ecosyst.

Environ. 223, 48–58.

- Fowles, M., 2007. Black carbon sequestration as an alternative to bioenergy. Biomass and Bioenergy 31, 426–432.
- Francis, D.D., 1992. Control mechanisms to reduce fertilizer nitrogen movement into groundwater. J. Soil Water Conserv. 47, 444–448.
- Garnett, T., Appleby, M.C., Balmford, A., Bateman, I.J., Benton, T.G., Bloomer, P.,
  Burlingame, B., Dawkins, M., Dolan, L., Fraser, D., Herrero, M., Hoffmann, I.,
  Smith, P., Thornton, P.K., Toulmin, C., Vermeulen, S.J., Godfray, H.C.J., 2013.
  Sustainable Intensifi cation in Agriculture: Premises and Policies. Philos. Trans. R.
  Soc. 341, 33–34.
- Gatti, M., Squeri, C., Garavani, A., Vercesi, A., Dosso, P., Diti, I., Poni, S., 2018. Effects of Variable Rate Nitrogen Application on cv. Barbera Performance: I. Vegetative Growth and Leaf Nutritional Status. Am. J. Enol. Vitic. ajev.2018.
- Hargrave, A.P., Shaykewich, C.F., 1997. Rainfall induced nitrogen and phosphorus losses from Manitoba soils. Can. J. Soil Sci. 77, 59–65.
- Herrero, M., Havlík, P., Valin, H., Notenbaert, A., Rufino, M.C., Thornton, P.K.,
  Blümmel, M., Weiss, F., Grace, D., Obersteiner, M., Herreroa, M., Havlikb-C, P.,
  Valinc, H., Notenbaertb, A., Rufinob, M.C., Thorntond, P.K., Bliimmelb, M.,
  Weissc, F., Grace6, D., Obersteinerc, M., 2013. Biomass use, production, feed
  efficiencies, and greenhouse gas emissions from global livestock systems. Proc.

Natl. Acad. Sci. U. S. A. 110, 20888–20893.

- Hirel, B., Le Gouis, J., Ney, B., Gallais, A., 2007. The challenge of improving nitrogen use efficiency in crop plants: Towards a more central role for genetic variability and quantitative genetics within integrated approaches. J. Exp. Bot. 58, 2369–2387.
- Hossini, H., Rezaee, A., Ayati, B., Mahvi, A.H., 2015. Simultaneous nitrification and denitrification using a polypyrrole/microbial cellulose electrode in a membraneless bio-electrochemical system. RSC Adv. 5, 72699–72708.
- Ibrikci, H., Cetin, M., Karnez, E., Flügel, W.A., Tilkici, B., Bulbul, Y., Ryan, J., 2015. Irrigation-induced nitrate losses assessed in a Mediterranean irrigation district. Agric. Water Manag. 148, 223–231.
- Inselsbacher, E., Wanek, W., Ripka, K., Hackl, E., Sessitsch, A., Strauss, J., Zechmeister-Boltenstern, S., 2011. Greenhouse gas fluxes respond to different N fertilizer types due to altered plant-soil-microbe interactions. Plant Soil 343, 17–35.
- International Biochar Initiative, 2012. Standardized Product Definition and Product Testing Guidelines for Biochar That Is Used in Soil.
- IPCC, 2014a. Climate change 2014 : mitigation : contribution of Working Group III to the third assessment report of the Intergovernmental Panel on Climate Change. Cambridge University Press.
- IPCC, 2014b. Climate change 2014: synthesis report. Contribution of Working Groups I, II and III to the fifth assessment report of the Intergovernmental Panel on Climate.

- IPCC, 2007. Climate Change 2007: The Physical Science Basis: Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change.
- Johnson, J.M.F., Franzluebbers, A.J., Weyers, S.L., Reicosky, D.C., 2007. Agricultural opportunities to mitigate greenhouse gas emissions. Environ. Pollut. 150, 107–124.
- Jokela, W.E., Bosworth, S.C., Rankin, J.J., 2014. Sidedressed dairy manure effects on corn yield and residual soil nitrate. Soil Sci. 179, 37–41.
- Ju, C., Buresh, R.J., Wang, Z., Zhang, H., Liu, L., Yang, J., Zhang, J., 2015. Root and shoot traits for rice varieties with higher grain yield and higher nitrogen use efficiency at lower nitrogen rates application. F. Crop. Res. 175, 47–55.
- Kameyama, K., Miyamoto, T., Shiono, T., Shinogi, Y., 2012. Influence of SugarcaneBagasse-derived Biochar Application on Nitrate Leaching in Calcaric Dark Red Soil.J. Environ. Qual. 41, 1131–1137.
- Khare, P., Goyal, D.K., 2013. Effect of high and low rank char on soil quality and carbon sequestration. Ecol. Eng. 52, 161–166.
- Koide, R.T., Petprakob, K., Peoples, M., 2011. Quantitative analysis of biochar in field soil. Soil Biol. Biochem. 43, 1563–1568.
- Laird, D.A., 2008. The charcoal vision: A win-win-win scenario for simultaneously producing bioenergy, permanently sequestering carbon, while improving soil and water quality. Agron. J. 100, 178–181.

Lal, R., 2011. Sequestering carbon in soils of agro-ecosystems. Food Policy 36, S33–S39.

- Lehmann, C.J., da Silva Jr, J.P., Rondon, M., C.M., D.S., Greenwood, J., Nehls, T., Steiner, C., Glaser, B., 2002. Slash-and-char - a feasible alternative for soil fertility management in the central Amazon?, in: 17th World Congress of Soil Science. Bangkok.
- Lehmann, J., Gaunt, J., Rondon, M., 2006. Bio-char sequestration in terrestrial ecosystems A review. Mitig. Adapt. Strateg. Glob. Chang. 11, 403–427.
- Lehmann, J., Joseph, S., 2009. Biochar for environmental management : An introduction, Earthscan. Routledge, USA.
- Liu, X. yu, Qu, J. jing, Li, L. qing, Zhang, A. feng, Jufeng, Z., Zheng, J. wei, Pan, G. xing, 2012. Can biochar amendment be an ecological engineering technology to depress N2O emission in rice paddies?-A cross site field experiment from South China. Ecol. Eng. 42, 168–173.
- Long, G.Q., Sun, B., 2012. Nitrogen leaching under corn cultivation stabilized after four years application of pig manure to red soil in subtropical China. Agric. Ecosyst. Environ. 146, 73–80.
- Lu, C., Tian, H., 2017. Global nitrogen and phosphorus fertilizer use for agriculture production in the past half century: shifted hot spots and nutrient imbalance. Earth Syst. Sci. Data 9, 181–192.

Maillard, É., Angers, D.A., 2014. Animal manure application and soil organic carbon

stocks: A meta-analysis. Glob. Chang. Biol. 20, 666–679.

- Masaka, J., Nyamangara, J., Wuta, M., 2015. Effect of Inorganic and Organic Fertilizer Application on Nitrate Leaching in Wetland Soil Under Field Tomato (Lycopersicon esculentum) and Leaf Rape (Brassica napus). Agric. Res. 4, 63–75.
- Matovic, D., 2011. Biochar as a viable carbon sequestration option: Global and Canadian perspective. Energy 36, 2011–2016.
- Mazzoncini, M., Sapkota, T.B., Bàrberi, P., Antichi, D., Risaliti, R., 2011. Long-term effect of tillage, nitrogen fertilization and cover crops on soil organic carbon and total nitrogen content. Soil Tillage Res. 114, 165–174.
- Meinshausen, M., Meinshausen, N., Hare, W., Raper, S.C.B., Frieler, K., Knutti, R., Frame, D.J., Allen, M.R., 2009. Greenhouse-gas emission targets for limiting global warming to 2 °C. Nature 458, 1158–1162.
- Minasny, B., Malone, B.P., McBratney, A.B., Angers, D.A., Arrouays, D., Chambers, A., Chaplot, V., Chen, Z.-S., Cheng, K., Das, B.S., Field, D.J., Gimona, A., Hedley,
  C.B., Hong, S.Y., Mandal, B., Marchant, B.P., Martin, M., McConkey, B.G.,
  Mulder, V.L., O'Rourke, S., Richer-de-Forges, A.C., Odeh, I., Padarian, J., Paustian,
  K., Pan, G., Poggio, L., Savin, I., Stolbovoy, V., Stockmann, U., Sulaeman, Y., Tsui,
  C.-C., Vågen, T.-G., van Wesemael, B., Winowiecki, L., 2017. Soil carbon 4 per
  mille. Geoderma 292, 59–86.
- Paustian, K., Lehmann, J., Ogle, S., Reay, D., Robertson, G.P., Smith, P., 2016. Climate-

smart soils. Nature 532, 49–57.

- Plastina, A., 2018. Estimated Costs of Crop Production in Iowa, Iowa State University Extension and Outreach.
- Randolph, P., Bansode, R.R., Hassan, O.A., Rehrah, D., Ravella, R., Reddy, M.R., Watts, D.W., Novak, J.M., Ahmedna, M., 2017. Effect of biochars produced from solid organic municipal waste on soil quality parameters. J. Environ. Manage. 192, 271–280.
- Reed, E.Y., Chadwick, D.R., Hill, P.W., Jones, D.L., 2017. Critical comparison of the impact of biochar and wood ash on soil organic matter cycling and grassland productivity. Soil Biol. Biochem. 110, 134–142.
- Rees, R.M., Baddeley, J.A., Bhogal, A., Ball, B.C., Chadwick, D.R., Macleod, M., Lilly,
  A., Pappa, V.A., Thorman, R.E., Watson, C.A., Williams, J.R., 2013. Nitrous oxide mitigation in UK agriculture. Soil Sci. Plant Nutr. 59, 3–15.
- Shackley, S., Sohi, S., Brownsort, P., Carter, S., 2010. An assessment of the benefits and issues associated with the application of biochar to soil-A report commissioned by the United Kingdom Department for Environment, Food and Rural Affairs, and Department of Energy and Climate Change. Edinburgh, UK.
- Singh, B.P., Hatton, B.J., Singh, B., Cowie, A.L., Kathuria, A., 2010. Influence of Biochars on Nitrous Oxide Emission and Nitrogen Leaching from Two Contrasting Soils. J. Environ. Qual. 39, 1224–1235.

- Smith, K. a., Ball, T., Conen, F., Dobbie, K.E., Massheder, J., Rey, A., 2003. Exchange of greenhousegases between soil and atmosphere: interactions of soil physical factors and biological processes. Eur. J. Soil Sci. 54, 779–791.
- Sohi, S.P., Krull, E., Lopez-Capel, E., Bol, R., 2010. A review of biochar and its use and function in soil. Adv. Agron. 105, 47–82.
- Subedi, R., Taupe, N., Pelissetti, S., Petruzzelli, L., Bertora, C., Leahy, J.J., Grignani, C., 2016. Greenhouse gas emissions and soil properties following amendment with manure-derived biochars: Influence of pyrolysis temperature and feedstock type. J. Environ. Manage. 166, 73–83.
- Taghizadeh-Toosi, A., Clough, T.J., Condron, L.M., Sherlock, R.R., Anderson, C.R.,
  Craigie, R.A., 2011. Biochar Incorporation into Pasture Soil Suppresses in situ
  Nitrous Oxide Emissions from Ruminant Urine Patches. J. Environ. Qual. 40, 468–478.
- Tan, Z., Lin, C.S.K., Ji, X., Rainey, T.J., 2017. Returning biochar to fields: A review. Appl. Soil Ecol. 116, 1–11.
- Thomson, A.J., Giannopoulos, G., Pretty, J., Baggs, E.M., Richardson, D.J., 2012.Biological sources and sinks of nitrous oxide and strategies to mitigate emissions.Philos. Trans. R. Soc. B Biol. Sci. 367, 1157–1168.
- Toyoda, S., Yano, M., Nishimura, S., Akiyama, H., Hayakawa, A., Koba, K., Sudo, S., Yagi, K., Makabe, A., Tobari, Y., Ogawa, N.O., Ohkouchi, N., Yamada, K.,

Yoshida, N., 2011. Characterization and production and consumption processes of N2 O emitted from temperate agricultural soils determined via isotopomer ratioanalysis. Global Biogeochem. Cycles 25.

- Tubiello, F.N., Salvatore, M., Rossi, S., Ferrara, A., Fitton, N., Smith, P., 2013. The FAOSTAT database of greenhouse gas emissions from agriculture. Environ. Res. Lett. 8, 015009.
- U.S. EPA, 2011. Global Anthropogenic Non-CO2 Greenhouse Gas Emissions : 1990 2030. Washington, DC. doi:EPA 430-R-12-006
- Ulyett, J., Sakrabani, R., Kibblewhite, M., Hann, M., 2014. Impact of biochar addition on water retention, nitrification and carbon dioxide evolution from two sandy loam soils. Eur. J. Soil Sci. 65, 96–104.
- Upadhyay, K.P., George, D., Swift, R.S., Galea, V., 2014. The influence of biochar on growth of lettuce and potato. J. Integr. Agric. 13, 541–546.
- Uzoma, K.C., Inoue, M., Andry, H., Zahoor, A., Nishihara, E., 2011. Influence of biochar application on sandy soil hydraulic properties and nutrient retention. J. Food, Agric. Environ. 9, 1137–1143.
- Wang, D., Fonte, S.J., Parikh, S.J., Six, J., Scow, K.M., 2017. Biochar additions can enhance soil structure and the physical stabilization of C in aggregates. Geoderma 303, 110–117.

Wang, L., Butterly, C.R., Wang, Y., Herath, H.M.S.K., Xi, Y.G., Xiao, X.J., 2014. Effect

of crop residue biochar on soil acidity amelioration in strongly acidic tea garden soils. Soil Use Manag. 30, 119–128.

- West, T.O., Post, W., 2002. Soil organic carbon sequestration rates by tillage and crop rotation: A global data analysis. Soil Sci. Soc. Am. J. 66, 1930–1946.
- Winning, N.J., 2014. Greenhouse gas emissions from Scottish arable agriculture and the potential for biochar to be used as an agricultural greenhouse gas mitigation option.University of Edinburgh.
- Woolf, D., Amonette, J.E., Street-Perrott, F.A., Lehmann, J., Joseph, S., 2010. Sustainable biochar to mitigate global climate change. Nat. Commun. 1, 56.
- Wrage, N., Velthof, G.L., Van Beusichem, M.L., Oenema, O., 2001. Role of nitrifier denitrification in the production of nitrous oxide. Soil Biol. Biochem. 33, 1723– 1732.
- Yao, Y., Gao, B., Zhang, M., Inyang, M., Zimmerman, A.R., 2012. Effect of biochar amendment on sorption and leaching of nitrate, ammonium, and phosphate in a sandy soil. Chemosphere 89, 1467–1471.
- Ye, Y., Liang, X., Chen, Y., Liu, J., Gu, J., Guo, R., Li, L., 2013. Alternate wetting and drying irrigation and controlled-release nitrogen fertilizer in late-season rice. Effects on dry matter accumulation, yield, water and nitrogen use. F. Crop. Res. 144, 212– 224.

Zhang, A., Liu, Y., Pan, G., Hussain, Q., Li, L., Zheng, J., Zhang, X., 2011. Effect of

biochar amendment on maize yield and greenhouse gas emissions from a soil organic carbon poor calcareous loamy soil from Central China Plain. Plant Soil 351, 263–275.

Zhang, M., Fan, C.H., Li, Q.L., Li, B., Zhu, Y.Y., Xiong, Z.Q., 2015. A 2-yr field assessment of the effects of chemical and biological nitrification inhibitors on nitrous oxide emissions and nitrogen use efficiency in an intensively managed vegetable cropping system. Agric. Ecosyst. Environ. 201, 43–50.
## **1.5.** Co-authorship statement

Manuscripts based on the chapter 2, entitled "Biochar amendment reduces greenhouse gases emission in silage corn cropping system following dairy manure and inorganic nitrogen application" and chapter 3 "Biomass production, and nitrogen dynamics of silage corn as influenced by organic and inorganic nitrogen sources and biochar amendment" will be submitted to Agriculture, Ecosystems and Environment (Ashiq, W., Nadeem, M., Ali, W., Zaeem, M., Wu, J., Galagedara, L., Kavanagh, V., Cheema, M. 2018). Wagar Ashiq, the thesis author will be the primary author and Dr. Cheema (supervisor), will be the corresponding and the last author. Dr. Wu (co-supervisor) and Dr. Galagedara (committee member) will be sixth and seventh authors, respectively. Dr. Kavanagh, research collaborator, Department of Fisheries and Land Resources will be the eighth author. For the work in Chapter 2 and chapter 3, Dr. Cheema wrote the research grants, developed the layout of this research field trial and helped in results interpretation. Mr. Wagar was responsible for the data collection, analysis, and writing of the manuscript. Dr. Nadeem, Mr. Ali, Mr. Zaeem, contributed in all steps of field and lab work. Dr. Wu provided specific guidance on GHG sample analysis and Dr. Kavanagh helped in experimental layout and seeding of the experiment. Drs. Galagedara, and Nadeem helped in statistical analyses and manuscript editing.

#### Chapter 2

# 2. Biochar amendment reduces greenhouse gases emission in silage corn cropping system following dairy manure and inorganic nitrogen application

# 2.1. Abstract

Biochar (BC) is considered as a mitigation tool for agricultural greenhouse gas (GHG) emissions. To access the effect of biochar application on the GHG emissions from organic and inorganic nitrogen sources applied to silage corn field, a two-year field experiment was conducted in Newfoundland, Canada. The treatments comprised of 1) dairy manure with high N (DM<sub>1</sub>:0.37 % N), 2) dairy manure with low N (DM<sub>2</sub>:0.13 % N), 3) Inorganic nitrogen (IN), 4) DM<sub>1</sub> + BC, 5) DM<sub>2</sub> + BC, 6) IN + BC, and 7) Control (N<sub>0</sub>). Overall, BC application to DM<sub>1</sub>, DM<sub>2</sub> and IN reduced cumulative CO<sub>2</sub> emission by 16, 25.5 and 26.5 %, lessened cumulative CH<sub>4</sub> emission by 184, 200 and 293 %, lowered cumulative N<sub>2</sub>O emission by 95, 86 and 93 % respectively. It also enhanced the silage corn dry matter yield by 6.8, 4.8, and 11 %, decreased global warming potential (GWP) by 25, 34.6 and 37 %, and contracted greenhouse gas intensity (GHGI) by 29.8, 37.6, and 43 % respectively. In conclusion, BC application demonstrated a great potential to decrease GHG emissions an increase crop yield simultaneously.

**Keywords:** greenhouse gas emissions, global warming potential, greenhouse gas intensity, silage corn, dairy manure, biochar

## 2.2. Introduction

Greenhouse gases (GHGs) emitted from different sources gather into the atmosphere and cause higher atmospheric temperature leading to climate change. Anthropogenic greenhouse gas (GHG) emissions have a major contribution towards global warming and climate change and have reached to 49.5 Gt CO<sub>2</sub> equivalent per year in 2010 (IPCC, 2014a). According to Intergovernmental Panel on Climate Change (IPCC, 2014a) "global GHGs emissions have increased to unprecedented levels and must be pulled down by 40-70 % compared to 2010 values by mid-century and near to zero by the end of the century to limit the increase in global mean temperature to 2 °C". Agriculture sector contributes 11-14 % towards global anthropogenic GHG emissions (Conway, 2012; Smith et al., 2007; Tubiello et al., 2015) and these agricultural emissions are increasing at around 1 % per year (Lamb et al., 2016). In 2011, about 8 % of the total GHG emissions in Canada were contributed by the agriculture sector largely through methane CH<sub>4</sub>) and nitrous oxide (N<sub>2</sub>O) (Agriculture and Agri-Food Canada). From 1981-2011, N<sub>2</sub>O emissions had increased by 31 % and CH<sub>4</sub> emission by 2 % from agricultural soils in Canada which were attributed to increased use of nitrogen fertilizers and dairy industry (Agriculture and Agri-Food Canada). Livestock sector produces approximately seven billion tons  $(7x10^9)$ Mg) of animal manure per year worldwide (Thangarajan et al., 2013). Canadian dairy and livestock sector produces about half a million tons of manure daily which equals to 180 million tons per year (Statistics Canada, 2006). Animal manure is historically known as a rich source of macro and micronutrients, when applied to agricultural soil it improves nutrient availability, soil organic matter (SOM) contents, soil organic carbon (SOC), cation exchange capacity (CEC), and water holding capacity (WHC) etc. (Bolan et al., 2004; Diacono and Montemurro, 2010; Maillard and Angers, 2014). Dairy manure (DM) application to maize increased C sequestration, soil N, P, K concentrations, N, and K uptake, maize biomass and harvest index (Khan et al., 2007). DM application to maize crop improved soil water use, increased SOM, N, P, K, Cu, Zn, Mn, improved biomass allocation into shoot and grains, increased water productivity by 3-5 % and maize yield by 5-10 % (Matsi et al., 2015; X. Wang et al., 2017). However, application of DM and IN to agricultural soils cause emission of a significant amount of GHGs including  $CO_2$ ,  $CH_4$ and N<sub>2</sub>O (Amon et al., 2006; HUANG et al., 2017). Whereas, DM application to soil emits 32.7 % more GHGs than IN alone and these emissions may offset the benefits of improving SOC by DM application (Barneze et al., 2014; M. Zhou et al., 2017). Shortchain volatile fatty acids in the DM are easily available to methanogenic archaea and cause CH<sub>4</sub> outburst into the atmosphere immediately after application (Hrapovic and Rowe, 2002; Sherlock et al., 2002). The total amount of IN applied to agricultural soils is 107.6 Tg N per year worldwide (Lu and Tian, 2017), of which 17 Tg nitrogen is lost every year in the form of N<sub>2</sub>O into the atmosphere and the loss is expected to increase four times by 2100 due to increased application of IN (Galloway et al., 2008; Schlesinger, 2009). Reduction of GHG emission and C sequestration has great potential to reduce global warming and climate change. Decreasing the atmospheric carbon by 3.5-4 Gt per year would limit the temperature increase to 2 °C by 2050 (Meinshausen et al., 2009; Minasny et al., 2017), a threshold level beyond which climate change would have a momentous impact (IPCC, 2014b). This annual reduction in atmospheric C concentration could be enhanced by increasing soil C sequestration in agricultural soils globally by 0.4

% per year, producing a C sink of 1.2 petagrams (Pg) per year (Paustian et al., 2016). The soils of agroecosystems have a technical potential of 1.2-3.1 billion ton C sequestration per year (Lal, 2011). A variety of management practices promote soil C sequestration including crop residues incorporation (Coppens et al., 2006), addition of perennial crops in rotation, no-tillage (West and Post, 2002), cover cropping (Mazzoncini et al., 2011), organic amendments like manure (Maillard and Angers, 2014) and biochar (Bera et al., 2016). Some of these practices may be disadvantageous as they increase GHG emissions into the atmosphere like manure application and crop residues incorporation increase  $N_2O$ emission (Li et al., 2005; M. Zhou et al., 2017). However, it has been reported that BC application to soil increase C sequestration (Bruun et al., 2012; Hernandez-Soriano et al., 2016; D. A. Laird et al., 2010; D. Wang et al., 2017), increase soil microbial biomass (H. Zhou et al., 2017), enhance WHC and water use efficiency (Ippolito et al., 2016), improve nutrient holding capacity (Hagemann et al., 2017; Laird et al., 2010; Laird et al., 2010), increase crop yield (Liu et al., 2017; Usman et al., 2016; Zhang et al., 2011) and decrease GHGs emission (Ahmed et al., 2016; Cayuela et al., 2013; Chang et al., 2016; Jia et al., 2012; Lan et al., 2017; Liu et al., 2012b; Sun et al., 2014; Wang et al., 2013). Integration of BC in agricultural systems has been proposed as an effective management option to mitigate GHG emissions from soils (Hawthorne et al., 2017; Lehmann, 2007; Thomazini et al., 2015; Van Zwieten et al., 2010a). However, before using BC as a C sequestration tool, it must be verified that its addition does not create adverse effects, e.g. increased GHG emission (Schimmelpfennig et al., 2014). In a two-year field experiment, application of BC increased SOC, pH, total N, and crop productivity however there was no effect on the GWP and GHGI during the first year, but during the second year it

decreased GWP and GHGI by 7-18 % and 12-38 %, respectively, in rice cropping system (Zhang et al., 2012). Wheat straw derived BC application at the rate of 24 ton per hectare and 48 ton per hectare decreased GWP by 30.7 and 35.6 %, respectively in double rice cropping system in China (Liu et al., 2014). BC application in intensive vegetable cropping systems with four consecutive vegetable crops had no influence on CH<sub>4</sub> emission while decreased N<sub>2</sub>O emission by 1.7-25.4 %, net GWP by 89-700 % and GHGI by 89-644 %, respectively (Li et al., 2015). BC application to soil can reduce GHG emission, GWP and GHGI of maize crop along with improving soil physiochemical and biological properties (Sun et al., 2017; Tan et al., 2017; Yang et al., 2017). It modifies the nutrient transformations in the soil and reduces the emission of GHGs (Castaldi et al., 2011; Liu et al., 2012; Laufer and Tomlinson, 2012; Liu et al., 2017a). BC application to soil amended with slurry reduced the cumulative N<sub>2</sub>O and CO<sub>2</sub> emission by 63, and 84 %, respectively while had no effect on CH<sub>4</sub> emissions during first 15 days of slurry application (Brennan et al., 2015). Application of BC to maize crop under field conditions could reduce N<sub>2</sub>O emissions by 41.8-52 % (Hüppi et al., 2015; Zhang et al., 2011). In a meta-analysis it was found that BC has the potential to reduce  $N_2O$  emissions by 49±5 % and this reduction depends on the degree of polymerization and aromaticity of biochar i.e. biochar with low H :  $C_{org}$  ratio reduces N<sub>2</sub>O emission more than BC having high H :  $C_{org}$ ratio (Cayuela et al., 2015). Soil aeration is significantly increased after BC application which decreases denitrification and reduces N<sub>2</sub>O emission (Case et al., 2012; Suddick and Six, 2013). Biochar surface absorbs soil  $NO_3^-$  and reduces the substrate for nitrification thus help to reduce N<sub>2</sub>O emissions (Mizuta et al., 2004; Taghizadeh-Toosi et al., 2011). Biochar application to soil could also reduce CH<sub>4</sub> emission resulting from manure application. After a series of laboratory and field studies it was found that slurry application increased  $N_2O$  and  $CH_4$  emissions from the soil while BC application with slurry increased plant biomass, increased C sequestration, decreased  $N_2O$  and  $CO_2$  emission and increased  $CH_4$  oxidation (Schimmelpfennig et al., 2014). Application of bamboo char and straw char reduced  $CH_4$  emissions from waterlogged paddy soil by 51 % and 91 % respectively, which was attributed to inhibition of methanogenic activity and increased  $CH_4$  oxidation (Liu et al., 2011).

There have been numerous studies documenting the GHGs emissions from agricultural soils in tropical areas; however, there is a lack of information concerning the GHG emissions under field conditions in the cool climatic region of Newfoundland. To address this issue and deficit in information, the current study was designed with the following objectives:

- i. To assess the GHGs emission from organic and inorganic sources of nitrogen application in silage corn cropping systems.
- ii. To determine the role of BC application in enhancing biomass production and reduction in GHG emission in silage corn cropping systems in NL.
- iii. To estimate GWP and GHGI of silage corn cropping systems under different dairy manure and biochar treatments

# 2.3. Materials and Methods

# 2.3.1. Study site

A field experiment was carried out at Pynn's Brook Research Station, Pasadena (49°04'21.9"N, 57°33'37.4"W) in Newfoundland, Canada, during 2016 and 2017 growing seasons. The soil of this area is classified as rapidly drained, Orthic Humo-Ferric Podzol with reddish brown to brown color. The soil has developed on gravely sandy fluvial deposit of mixed lithology. Due to high coarse fragments and rapid drainage it has a limited agricultural use. The best-suited crops for this soil are hey and forage crops (Kirby, 1988). Basic physio-chemical properties of soil can be seen in Table 2.1.

Site characteristics	Description
Soil class	Orthic Humo-Ferric Podzol
Soil texture (10-15 cm depth)	Gravelly loamy sand: sand (82±3.4 %), silt
	(11.6±2.4 %), clay (6.4±1.2 %)
Soil parental material	Channery, gravely sandy stratified fluvial deposit
Elevation	45 m
Soil drainage class	Well to rapidly drained
Soil pH	6.3 (2016), 6.8 (2017)
Average bulk density	$1.31\pm0.07 \text{ g cm}^{-3}$
Average porosity	51±0.03 %
Gravel	20 % in top 5 cm layer
Average soil organic matter	3.10 %
CEC	12 cmol/kg
Previous Crop (2015)	Silage corn

Table 2.1: Basic soil properties of the experimental site at Pynn's Brook Research Station

#### **2.3.2.** Experimental setup and treatments

The experiment comprised of three nitrogen sources amended with BC. Treatments included were; 1) DM with high N conc. (0.37 %) designated as DM<sub>1</sub>, 2) DM with low N conc. (0.13 %), designated as  $DM_2$ , 3) Inorganic nitrogen (IN), 4)  $DM_1$ +Biochar, 5)  $DM_2$ +Biochar, 6) IN + Biochar, and 7) control (N<sub>0</sub>). DM was collected from two dairy farms (Larch Grove and Rideout Junior) located in Cormack area near Deer Lake, Newfoundland and Labrador (NL). Soil and DM samples were sent for detailed nutrient analyses to Soil, Plant and Feed Laboratory, Department of Fisheries and Land Resources, St. John's, NL. DM from Larch Grove farm (designated as DM<sub>1</sub>) exhibited high concentration of N, P, K, Ca, Mg, Fe, Mn, Zn, B whereas, DM sourced from Rideout Junior farm (designated as DM<sub>2</sub>) had a low concentration of N, P, K, Ca, Mg, Fe, Mn, Zn, B (Table 2.2). DM was applied before seeding in respective plots according to local farmers practice i.e. 30,000 liters per hectare. Fertilizers were applied to fulfill the required nutrients based on DM and soil analyses reports and regional recommendations of the crop. Ammonium nitrate (AN), triple superphosphate (TSP) and murate of potash (MOP) were used as nitrogen (N), phosphorus (P), and potash (K) sources, respectively and were applied @ 215,110, 225 kg per hectare.  $DM_1$ ,  $DM_2$  and the entire IN were applied before crop seeding during 2016 while in 2017  $DM_1$  and  $DM_2$  were applied to all respective treatments (DM<sub>1</sub>, DM<sub>2</sub>, DM<sub>1</sub>+B, DM<sub>2</sub>+B) before seeding but the IN fertilizer was applied in two splits (first dose: 6 leaf stage, second dose: 12 leaf stage). The experimental design was a randomized complete block (RCBD) with three replications and net plot size was 4.8 meters x 1.5 meters.

Characteristic	Larch Grove farm (DM <sub>1</sub> )		Rideout Junio	<u>or farm (DM<sub>2</sub>)</u>
(as received basis)	2016	2017	2016	2017
Dry matter (%)	9.33	10.9	3.57	1.70
рН	6.80	6.80	7.00	7.10
Total Nitrogen (%)	0.37	0.44	0.14	0.12
Total Phosphorus (%)	0.06	0.08	0.02	0.01
Total Potassium (%)	0.38	0.37	0.12	0.12
Total Calcium (%)	0.16	0.19	0.059	0.04
Total Magnesium (%)	0.07	0.07	0.02	0.01
Total Iron (ppm)	49.0	68	19.0	7.00
Total Manganese (ppm)	23.0	21.0	9.00	5.00
Total Copper (ppm)	4.70	4.50	33.0	20.0
Total Zinc (ppm)	17.0	21.0	8.00	5.00
Total Boron (ppm)	3.00	3.40	1.00	0.50
Total Sodium (ppm)	911	904	275	241

Table 2.2: Chemical analysis of dairy manures used in the study

## 2.3.3. Crop husbandry

DM and BC were applied one day before seeding and mixed in the soil to 15 cm depth. Seeding of silage corn hybrid (Yukon R) was done with the SAMCO 2200 system (SAMCO Agricultural Manufacturing Ltd) on May 24 and May 23 during 2016 and 2017, respectively. This system has an advantage that it can cover the seed rows with degradable polythene sheet while seeding, which allows accumulating maximum heat during the cold season for seed germination (Figure 2.1). This sheet had several pin holes which allow the trapped air under the sheet to escape and keep it tight to the soil. These pin holes weaken the sheet allowing the plants easy access through the sheet while maintaining soil temperature. Seeding rate for the crop was 90,900 seeds per hectare. Weeds were controlled with the spray of Roundup WeatherMax on July 09, 2016 and July 08, 2017 at the rate of 2 L per hectare. BC used in the study was purchased from AirTerra Inc. located in Calgary, Alberta, and is a registered BC product with the Canadian Food Inspection Agency (CFIA), which is the first in Canada. It was produced from yellow pine wood pyrolyzed at 500 °C for 30 min in oxygen-limited conditions. BC was applied @ 20 tons ha<sup>-1</sup> (Liu et al., 2012b). The detailed BC analyses report conducted by Gabilan laboratory, Salinas, California, USA can be seen in Table 2.3.



Figure 2.1: Silage corn seeding with SAMCO 2200

Property	Wet basis	Dry weight basis	
pH	9	-	
ECe (mmhos/cm	0.43	-	
Moisture (%)	15.2		
WHC (mL water per 100g dry char)	74.9	74.9	
Volatile matter (%)		8.5	
Ash (%)		6.7	
Fixed carbon (%)		84.5	
Н (%)		0.68	
O (%)		7.84	
N (%)		0.22	
S (%)		0	
H/C		0.1	
O/C		0.07	
Total ash (%)	6	7.1	
Recalcitrant carbon (%)	64.6	76.2	
Neutralizing value (% as CaCO <sub>3</sub> )	4.2	4.9	
Carbonate value (% as CaCO <sub>3</sub> )	0.5	0.6	
Butane activity (g/100g dry char)		5.1	
Bulk density $(Mg/m^3)$	0.23	0.19	
Particle density (acetone) (g/cc)		1.57	
Solid space (% v/v)		12.5	
Void space (% $v/v$ )		87.5	

Table 2.3: Physio-chemical properties of biochar used in the study

#### 2.3.4. Greenhouse gas sampling and analysis

GHG samples were collected weekly in the first month and then fortnightly for the whole growing seasons using static chamber method (Holland et al., 1999). The Polyvinyl chloride (PVC) collars with an inner diameter of 26 cm were inserted permanently to a depth of 10 cm in each plot one week before the start of 1<sup>st</sup> GHGs sampling to mitigate any placement disturbance. A 50 cm high PVC chamber with 26 cm diameter and covering lid was fixed on the top of each collar during GHGs sampling. Chamber top lid had tubing outlets connected with three-way stopcocks with Luer-lock tip. For each measurement, four gas samples were taken from the chamber using a 30 mL non-sterile syringe fitted with a three-way stopcock (BD Luer-lock tip) at 10 min intervals (0, 10, 20 and 30 min after lid closure) (Wang et al., 2012; Chen et al., 2015). To minimize any effect of diurnal variation in emissions, the samples were taken at the same time of the day (9 am - 3 pm) on each sampling occasion. During each GHG sampling event, soil moisture (SM) content (volume basis) and EC (5 cm depth), and soil temperature (ST) (5 cm and 20 cm depth) were also monitored by SM and temperature probes (EC-TM model, Decagon Devices Inc.) from each treatment (Figure 2.2).

GHG samples were transferred to evacuated clear Labco Exetainer® glass vials (Vial type 3-soda glass, height 101 mm, diameter 15.5 mm, capacity 12 mL) sealed with gastight neoprene septum. Quantification of GHGs i.e.  $CO_2$ ,  $CH_4$  and  $N_2O$  was carried out by gas chromatography (SICON GC-456 Bruker) equipped with thermal conductivity detector (TCD), flame ionization detector (FID), and electron capture detector (ECD) (Collier et al., 2014). All the fluxes were adjusted for headspace volume and chamber area as explained by (Holland et al., 1999), and calculated by linear regression using all time points sampled:  $F = (dC/dt) \times V/A$  (where, V is volume of the chamber, A is the area covered by chamber, and dC/dt is the rate of concentration change). Cumulative GHG fluxes during the experimental period were calculated by multiplying the mean fluxes of two successive determinations by the length of the period between samplings and adding that amount to the previous cumulative total as described in Equation (2.1) (Cai et al., 2013; Menéndez et al., 2006).

Cumulative flux = 
$$\sum_{i=1}^{n} (F_i + F_{i+1}) / 2 \times (t_{i+1} - t_i) \times 24$$
 (2.1)

Where *F* is the GHGs flux (mg m<sup>-2</sup> h<sup>-1</sup>), *i* is the i<sup>th</sup> measurement, the term of  $(t_{i+1} - t_i)$  is the days between two adjacent sampling events, and *n* is the total number of sampling events.



Figure 2.2: Recording soil temperature, moisture and EC (a), sampling chambers fixed over sampling spots in a crop row (b), GHGs sample collection with (c)

#### 2.3.5. Global warming potential and greenhouse gas intensity calculation

Global warming potential is the relative measure of how much warming is caused by a certain gas as compared to same mass of  $CO_2$ . Whereas greenhouse gas intensity is the measurement of the total emissions from a system per unit of the produce. GWP of  $CO_2$ ,  $CH_4$ , and  $N_2O$  and GHGI were calculated by Equation (2.2) and (2.3), respectively (Yang et al., 2017; Zhang et al., 2013, 2012; Z. S. Zhang et al., 2014).

$$GWP (kg CO_{2 eq}) = CO_2 + CH_4 \times 25 + N_2 O \times 298$$
(2.2)

GHGI (kg 
$$CO_{2 eq}$$
 per kg dry matter yield) =  $\frac{GWP}{dry matter yield}$  (2.3)

## 2.3.6. Dry matter production

Plants were harvested from a 1 m<sup>2</sup> area at black layer stage from each plot and their fresh weight was recorded. Plants were oven dried at 70  $^{\circ}$ C for 48 h and dry matter yield (DMY) was calculated from each treatment plot using Equations(2.4) and (2.5). Percent dry matter was calculated by;

Percent dry matter (%) = 
$$\frac{\text{oven dry weight}}{\text{fresh weight}} \times 100$$
 (2.4)

DMY was calculated by multiplying dry matter percentage with the fresh weight of plants

DMY per hectare = Percent dry matter × fresh weight  $m^{-2} \times 10,000$  (2.5)

# 2.3.7. Statistical analysis

The analysis of variance (ANOVA) was used to determine the effect of different treatments on the emission of CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O, DMY, GWP and GHGI. Where treatment effects were significant, the means were compared with LSD ( $\alpha = 0.05$ ). The data were analyzed using the Statistix 10 software package (Analytical software, FL, USA) and figures were prepared using SigmaPlot 12.0 software program (Systat Software Inc., San Jose, CA).

#### 2.4. Results

## 2.4.1. CO<sub>2</sub> flux

Data presented in Tables 2.4 and 2.5 show that DM<sub>1</sub>, DM<sub>2</sub>, and IN alone and in combination with BC had significantly (p < 0.05) affected CO<sub>2</sub> emission during both growing seasons. Cumulative CO<sub>2</sub> emission was maximum (7,834 kg ha<sup>-1</sup> season<sup>-1</sup>) in  $DM_1$  treatment and minimum cumulative  $CO_2$  emission (5,576 kg ha<sup>-1</sup> season<sup>-1</sup>) was observed in IN+B treatment during the 2016 growing season (Table 2.4). In the 2017 growing season,  $CO_2$  emission pattern in treatments was the same as 2016, but cumulative emission was lower. The  $DM_1$  treatment emitted more cumulative  $CO_2$  (7,078) kg ha<sup>-1</sup> season<sup>-1</sup>) while minimum (3,800 kg ha<sup>-1</sup> season<sup>-1</sup>) was noted in the IN+B treatment (Table 2.5). BC application to  $DM_1$ ,  $DM_2$  and IN significantly (p < 0.05) reduced cumulative CO<sub>2</sub> emission by 17, 25 and 26 % in 2016 (Table 1), while it was reduced by 15, 26 and 27 % in 2017, respectively. Significant temporal variation in CO<sub>2</sub> emission was noted in both years. Maximum  $CO_2$  emission (439 mg m<sup>-2</sup> h<sup>-1</sup>) was observed 90 days after manure application (DAMA) in DM<sub>2</sub> treatment and minimum emission (77 mg m<sup>-2</sup>  $h^{\text{-1}}$  ) was noticed in  $N_0$  (control) treatment at 146 DAMA during 2016 (Figure 2.3a). In 2017, CO<sub>2</sub> emission at 60 DAMA was greatest (462 mg m<sup>-2</sup> h<sup>-1</sup>) in DM<sub>1</sub> and least emission (44 mg m<sup>-2</sup> h<sup>-1</sup>) was recorded in N<sub>0</sub> treatment at 29 DAMA (Figure 2.4a).

Table 2.4: Cumulative greenhouse gas emission/absorption of  $CO_2$ ,  $CH_4$  and  $N_2O$  (Kg ha<sup>-1</sup> season<sup>-1</sup>), global warming potential (kg  $CO_2$  equivalent), silage corn dry matter yield (kg ha<sup>-1</sup>), and greenhouse gas intensity (kg  $CO_2$  equivalent per kg dry matter yield), during growing season 2016.

Treatment	CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O	GWP	DMY	GHGI
DM1	7834±476 <sup>a</sup>	1.26±0.9 <sup>a</sup>	1.69±0.2 <sup>a</sup>	8372±433 <sup>a</sup>	19797±173°	$0.42{\pm}0.02^{a}$
DM1+B	6430±169 <sup>b</sup>	-1.42±0.1 <sup>b</sup>	-0.15±0.1 <sup>d</sup>	6350±203 <sup>b</sup>	21050±125 <sup>a</sup>	0.30±0.01 <sup>b</sup>
DM2	7652±31 <sup>a</sup>	0.83±0.5 <sup>a</sup>	2.17±0.1 <sup>a</sup>	8319±293 <sup>a</sup>	19567±240 <sup>c</sup>	0.42±0.01 <sup>a</sup>
DM2+B	5666±16 <sup>b</sup>	-1.01±0.4 <sup>b</sup>	$0.59{\pm}0.1^{bc}$	5819±202 <sup>cd</sup>	$20433 \pm 176^{b}$	0.28±0.01 <sup>bc</sup>
IN	7566±37 <sup>a</sup>	$0.86{\pm}0.5^{a}$	1.80±0.3 <sup>a</sup>	8126±332 <sup>a</sup>	18813±135 <sup>d</sup>	$0.43 \pm 0.02^{a}$
IN+B	$5576 \pm 29^{b}$	-1.69±0.6 <sup>b</sup>	$0.24{\pm}0.1^{bc}$	$5607 \pm 268^{d}$	20533±176 <sup>ab</sup>	0.27±0.01 <sup>c</sup>
N0	5961±11 <sup>b</sup>	-0.10±0.2 <sup>ab</sup>	$0.89{\pm}0.0^{b}$	6224±105 <sup>bc</sup>	15300±152 <sup>e</sup>	$0.40 \pm 0.00^{a}$

Means sharing common letters in each column are not significantly different (at 0.05 probability level).

Table 2.5: Cumulative greenhouse gas emission/absorption of  $CO_2$ ,  $CH_4$  and  $N_2O$  (Kg ha<sup>-1</sup> season<sup>-1</sup>), global warming potential (kg  $CO_2$  equivalent), silage corn dry matter yield (kg ha<sup>-1</sup>), and greenhouse gas intensity (kg  $CO_2$  equivalent per kg dry matter yield), during growing season 2017

Treatment	CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O	GWP	DMY	GHGI
DM1	7078±639 <sup>a</sup>	11.6±3 <sup>a</sup>	1.95±0.27 <sup>a</sup>	7953±660 <sup>a</sup>	15983±258 <sup>bc</sup>	0.49±0.04 <sup>a</sup>
DM1+B	5957±714 <sup>ab</sup>	-6.5±3 <sup>bc</sup>	0.33±0.13 <sup>b</sup>	5894±653 <sup>b</sup>	17160±105 <sup>a</sup>	$0.34{\pm}0.04^{c}$
DM2	$5601{\pm}806^{abc}$	11.5±2 <sup>a</sup>	1.63±0.10 <sup>a</sup>	$6377 \pm 770^{b}$	15667±218 <sup>c</sup>	$0.40{\pm}0.04^{b}$
DM2+B	4100±754 <sup>bc</sup>	-9.1±5 <sup>bc</sup>	-0.01±0.51 <sup>b</sup>	3868±999 <sup>cd</sup>	16483±44 <sup>ab</sup>	$0.23{\pm}0.06^d$
IN	$5248 \pm 740^{abc}$	9.9±3 <sup>a</sup>	1.47±0.20 <sup>a</sup>	5936±659 <sup>b</sup>	$14580 {\pm} 408^{d}$	$0.40{\pm}0.05^{b}$
IN+B	3800±465 <sup>c</sup>	-19±5°	0.19±0.16 <sup>b</sup>	3382±539 <sup>d</sup>	16483±130 <sup>ab</sup>	$0.20{\pm}0.03^{d}$
N0	3997±561 <sup>bc</sup>	-0.28±5 <sup>ab</sup>	$0.96 {\pm} 0.56^{ab}$	4277±745 <sup>c</sup>	11200±152 <sup>e</sup>	$0.38 \pm 0.06^{bc}$

Means sharing common letters in each column are not significantly different (at 0.05 probability level)

#### 2.4.2. CH<sub>4</sub> flux

Experimental treatments (DM<sub>1</sub>, DM<sub>2</sub>, and IN alone and in combination with BC) had significantly (p < 0.05) affected CH<sub>4</sub> emission during 2016 and 2017 growing seasons (Table 2.4 & 2.5). Comparison of treatment's means showed that cumulative CH<sub>4</sub> emission was greatest (1.26 kg ha<sup>-1</sup> season<sup>-1</sup>) in DM<sub>1</sub> treatment, whereas, IN+B treatment exhibited maximum (1.69 kg ha<sup>-1</sup> season<sup>-1</sup>) cumulative CH<sub>4</sub> absorption during 2016 growing season (Table 2.4). However, DM<sub>1</sub>, DM<sub>2</sub>, and IN treatments were statistically non-significant with each other. In the 2017 growing season, cumulative CH<sub>4</sub> emission was higher as compared to 2016 and DM<sub>1</sub> treatment emitted more cumulative CH<sub>4</sub> (11.6 kg ha<sup>-1</sup> season<sup>-1</sup>), while maximum absorption (19.5 kg ha<sup>-1</sup> season<sup>-1</sup>) was noted in the IN+B treatment (Table 2.5). BC application to DM<sub>1</sub>, DM<sub>2</sub> and IN treatments significantly (p < 0.05) reduced cumulative CH<sub>4</sub> emission/increased absorption by 213, 221 and 295 % in 2016 (Table 1), while it was 156, 179 and 291 % in the 2017 growing season, respectively. There was a significant temporal variation in CH<sub>4</sub> emission/absorption during both years, which may be attributed to great variation in soil temperature during both growing seasons (Helbig et al., 2017). Maximum CH<sub>4</sub> emission (0.13 mg m<sup>-2</sup> h<sup>-1</sup>) was noted in DM<sub>1</sub> treatment after 103 DAMA and maximum absorption (0.17 mg m<sup>-2</sup> h<sup>-1</sup>) was observed in IN+B treatment at 20 DAMA during 2016 (Figure 2.3b). In 2017 growing season, DM<sub>2</sub> treatment emitted more CH<sub>4</sub> emission (1.78 mg m<sup>-2</sup> h<sup>-1</sup>) at 127 DAMA and IN+B treatment showed the highest absorption (1.2 mg m<sup>-2</sup> h<sup>-1</sup>) at 127 DAMA (Figure 2.4b).

#### 2.4.3. N<sub>2</sub>O flux

Data presented in Tables 2.4 and 2.5 show that DM<sub>1</sub>, DM<sub>2</sub>, and IN treatments alone and in combination with BC had significantly (p < 0.05) affected N<sub>2</sub>O emission during both growing seasons. Cumulative N<sub>2</sub>O emission was maximum (2.17 kg ha<sup>-1</sup> season<sup>-1</sup>) in DM<sub>2</sub> treatment compared to minimum cumulative N<sub>2</sub>O emission (-0.15 kg ha<sup>-1</sup> season<sup>-1</sup>) that was observed in DM<sub>1</sub>+B treatment during 2016 growing season (Table 2.4). Whereas, in 2017 DM1 treatment produced more cumulative N<sub>2</sub>O (1.95 kg ha<sup>-1</sup> season<sup>-1</sup>) while DM<sub>2</sub>+B treatment emitted minimum (-0.01 kg ha<sup>-1</sup> season<sup>-1</sup>) N<sub>2</sub>O (Table 2.5). BC application to  $DM_1$ ,  $DM_2$  and IN significantly (p < 0.05) reduced cumulative N<sub>2</sub>O emission by 108, 72 and 86 % in 2016 (Table 2.4), while in 2017 it was reduced by 82, 100 and 86 %, respectively (Table 2.5) Significant temporal variation in N<sub>2</sub>O emission was also noted in both years. N<sub>2</sub>O emission was greatest (0.25 mg m<sup>-2</sup> h<sup>-1</sup>) in DM<sub>2</sub> treatment at 34 DAMA and minimum emission (-0.07 mg m<sup>-2</sup> h<sup>-1</sup>) was noted in the DM<sub>1</sub>+B treatment at 117 DAMA during 2016 (Figure 2.3c). In 2017, N<sub>2</sub>O emission at 127 DAMA was greatest (0.15 mg m<sup>-2</sup> h<sup>-1</sup>) in DM<sub>1</sub> and least emission (-0.05 mg m<sup>-2</sup> h<sup>-1</sup>) was recorded in the  $DM_1$ +B treatment at 46 DAMA (Figure 2.4c).

## 2.4.4. Global warming potential, dry matter yield and greenhouse gas intensity

GWP and GHGI of all experimental treatments are shown in Table 2.4 for the 2016 growing season and in Table 2.5 for the 2017 growing season. There was a significant (p<0.05) effect of treatments on the GWP and GHGI during both years. During 2016, the highest GWP (8,372 kg CO<sub>2</sub> equivalent) and the lowest GWP (5,607 kg CO<sub>2</sub> equivalent)

were observed in DM<sub>1</sub> and IN+B, respectively. DMY ranged from 15,300 kg ha<sup>-1</sup> in the control to 21,050 kg ha<sup>-1</sup> in DM<sub>1</sub>+B. Maximum GHGI (0.43 kg CO<sub>2</sub> equivalent kg<sup>-1</sup> dry matter) was in IN whereas minimum (0.27 kg CO<sub>2</sub> equivalent kg<sup>-1</sup> dry matter) was observed in IN+B. BC application to DM<sub>1</sub>, DM<sub>2</sub>, and IN reduced GWP by 24, 30, and 31 % respectively, increased dry matter production by 6.3, 4.4, and 9 %, and decreased GHGI by 28.9, 33 and 36.8 % respectively. **During 2017**, the maximum GWP (7,954 kg CO<sub>2</sub> equivalent) and the lowest (3,382 kg CO<sub>2</sub> equivalent) were observed in DM<sub>1</sub> and IN+B respectively. DMY ranged from 11,200 kg ha<sup>-1</sup> in control to 17,160 kg ha<sup>-1</sup> in DM<sub>1</sub>+B. GHGI was highest in DM<sub>1</sub> (0.49 kg CO<sub>2</sub> equivalent kg<sup>-1</sup> dry matter) whereas minimum (0.20 kg CO<sub>2</sub> equivalent kg<sup>-1</sup> dry matter) was observed in IN+B. BC application to DM<sub>1</sub>, DM<sub>2</sub>, and IN reduced GWP by 26, 39, and 43 %, increased DMY by 7, 5, and 13 %, and decreased GHGI by 31, 42 and 50 %, respectively.



Greenhouse gas sampling dates-2016 (DAMA)

Figure 2.3: Temporal greenhouse gas emission during growing season 2016

(a)  $CO_2$  (b)  $CH_4$  (c)  $N_2O$ , Solid circle ( $DM_1$ ), empty circle ( $DM_1+B$ ), solid triangle ( $DM_2$ ), empty triangle ( $DM_2+B$ ), Solid square (IN), empty square (IN+B), solid diamond ( $N_0$ )



Greenhouse gas sampling dates-2017 (DAMA)

Figure 2.4: Temporal greenhouse gas emission during growing season 2017

(a)  $CO_2$  (b)  $CH_4$  (c)  $N_2O$ , Solid circle ( $DM_1$ ), empty circle ( $DM_1+B$ ), solid triangle ( $DM_2$ ), empty triangle ( $DM_2+B$ ), Solid square (IN), empty square (IN+B), solid diamond ( $N_0$ )

# 2.5. Discussion

# 2.5.1. Meteorological conditions:

Newfoundland has a unique climate and characterized with relatively short and cool growing period from May to October. Relatively low temperature was recorded during crop seeding (May) and harvesting (October) periods during both study years. However, mean maximum, mean minimum and mean average temperature was slightly lower during 2017 than 2016 with few exemptions. The average seasonal rainfall was significantly lower (30 % less) in 2017 compared to 2016 (Table 2.6). Well distributed rainfall was recorded during 2016 whereas the second growing season was characterized not only by low rainfall but also a completely dry period during June where the crop was at the active growth stage.

Growth period	Mean max. Temp. (°C)	Mean min. temp. (°C)	Average temp. (°C)	Rain (mm)	Mean Max. Temp. (°C)	Mean Min. temp. (°C)	Average temp. (°C)	Rain (mm)
-	2016			2017				
May 01-15	12.06	0.26	6.16	37	12.8	-1.46	5.66	11
May 16-31	16.06	2.68	9.37	47	13.06	0.0	6.53	47
June 01-15	14.86	4.40	9.63	107	15.33	3.53	9.43	35
June 16-30	23.26	7.73	15.5	41	21.46	8.40	14.93	45
July 01-15	21.0	7.93	14.46	42	23.33	8.8	16.06	29
July 16-31	24.81	10.93	17.87	40	24.18	8.31	16.25	12
August 01-15	23.06	10.06	16.56	27	23.8	9.46	16.63	58
August 16-31	21.75	10.0	15.87	112	22.12	6.87	14.5	32
September 01-15	18.4	7.46	12.93	98	19.06	7.53	13.3	80
September 16-30	14.2	3.26	8.73	35	15.33	2.86	9.1	76
October 01-20	14.05	0.4	7.22	118	12.8	2.4	7.6	76
Total				704				501

Table 2.6: Weather conditions (biweekly average) during silage corn growing season in2016 and 2017 at Pynn's Brook Research Station



Greenhouse gas sampling date-2016 (DAMA)

Figure 2.5: (a) Air temperature and rainfall at Pynn's Brook Research Station (b) soil temperature at 5cm depth, and (c) soil moisture at different greenhouse gas sampling dates during growing season 2016.

Solid circle  $(DM_1)$ , empty circle  $(DM_1+B)$ , solid triangle  $(DM_2)$ , empty triangle  $(DM_2+B)$ , Solid square (IN), empty square (IN+B), solid diamond (N<sub>0</sub>)



Greenhouse gas sampling dates-2017 (DAMA)

Figure 2.6: (a) Air temperature and rainfall at Pynn's Brook Research Station (b) soil temperature at 5cm depth, and (c) soil moisture at different greenhouse gas sampling dates during growing season 2017.

Solid circle  $(DM_1)$ , empty circle  $(DM_1+B)$ , solid triangle  $(DM_2)$ , empty triangle  $(DM_2+B)$ , Solid square (IN), empty square (IN+B), solid diamond (N<sub>0</sub>)

#### **2.5.2.** $CO_2$ emission

Maximum CO<sub>2</sub> emission reached the peak of 439 mg  $m^{-2}$   $h^{-1}$  at 90 DAMA in the 2016 growing season. This peak could be directly related to highest ST and lowest SM, during the period between 79-90 DAMA (Figure 2.5b). In 2017, the maximum CO<sub>2</sub> emission peak occurred at 60 DAMA when ST was the highest (25 °C) and SM was the lowest of the whole growing season (Figure 2.6). Similarly, the minimum  $CO_2$  emission events in both growing seasons were related to ST and SM. During 2016, minimum CO<sub>2</sub> emission was observed at 146 DAMA when ST was the lowest (9.8 °C) and SM was the maximum (35.7 %) (Figure 2.5), while in 2017, minimum emission was recorded at 29 DAMA when ST was relatively low (16.5 °C) and SM was relatively high (31 %). There was a significant reduction in CO<sub>2</sub> emission in BC treatments (DM<sub>1</sub>+B, DM<sub>2</sub>+B, IN+B) as compared to non-BC treatments (DM<sub>1</sub>, DM<sub>2</sub>, IN) (Figure 2.6). DM incorporation to soil increased cumulative  $CO_2$  emission over the season. DM application increases the soil CO<sub>2</sub> emission directly from C compounds in the DM and also by inducing a priming effect on native soil C (Bol et al., 2003). It had been reported in several studies that BC application reduces CO<sub>2</sub> emission. BC application to DM amended soil reduced cumulative CO<sub>2</sub> emission by 84 % most probably due to sorption of CO<sub>2</sub> on BC surface or a reduction in the availability of labile C (Brennan et al., 2015). Both positive and negative response of BC have been reported on CO<sub>2</sub> emission, for example, (Spokas and Reicosky, 2009) of the sixteen BC types evaluated, three have reduced, five have increased and eight have no impact on CO<sub>2</sub> emissions from agricultural soils (Spokas and Reicosky, 2009). Manure treatments  $(DM_1, DM_2)$  have high CO<sub>2</sub> emission rates than IN

and BC treatments, which can also be supported with previous studies (Agegnehu et al., 2016; Lentz et al., 2014; Schimmelpfennig et al., 2014). The decline in dissolved organic carbon (DOC) from native SOC after BC addition reduced the decomposition of organic C which reduced CO<sub>2</sub> emission from the soil by 64-68 % (Lu et al., 2014). BC induced negative priming effect and slowed the breakdown of SOM by different mechanisms including; (1) sorption of enzymes responsible for SOM breakdown, (2) shift in microbial metabolism, (3) enhanced stability of soil aggregates and microbial community shift towards low C turnover bacteria taxa (Zheng et al., 2018), and (4) decreased bioavailability of SOC via adsorption on BC large surface area (Sheng and Zhu, 2018).

# 2.5.3. CH<sub>4</sub> flux

The studied site acted both as a source and a sink of CH<sub>4</sub>. During the 2016 growing season, the highest CH<sub>4</sub> emission (0.13 mg m<sup>-2</sup> h<sup>-1</sup>) occurred at 103 DAMA when the SM was the highest (36.5 %) (Figure 2.5), whereas in 2017, the highest emission (1.78 mg m<sup>-2</sup> h<sup>-1</sup>) was recorded at 127 DAMA when the SM was the highest (40 %) (Figure 2.6). Minimum CH<sub>4</sub> emission of -0.17 mg m<sup>-2</sup> h<sup>-1</sup> (from IN+B in 2016 and -1.2 mg m<sup>-2</sup> h<sup>-1</sup> (from IN+B) in 2017 was not related to SM or ST. Increase in CH<sub>4</sub> production after DM application as in this experiment has been reported in previous studies. Short-chain fatty acids present in DM become available to methanogenic archaea after application to land and cause CH<sub>4</sub> outbursts (Hrapovic and Rowe, 2002; Sherlock et al., 2002). Significant reduction in CH<sub>4</sub> emission was observed in BC treatments (DM<sub>1</sub>+B, DM<sub>2</sub>+B, IN+B) as compared to non-BC treatments (DM<sub>1</sub>, DM<sub>2</sub>, IN). The decrease in CH<sub>4</sub> emission after BC application might be due to the stimulation of methanotrophic activity or the increased

abundance of the methanotrophic proteobacterial community (Feng et al., 2012; Liu et al., 2011). Kim et al (2017) studied the effect of BC and slow release fertilizer (SRF) on rice yield and  $CH_4$  emission and concluded that BC suppressed methanogenesis by increasing the oxygen supply in the soil through increased aeration.

# 2.5.4. N<sub>2</sub>O flux

The experimental site was a source as well as a sink of N<sub>2</sub>O. The highest N<sub>2</sub>O emission peak (0.25 mg m<sup>-2</sup> h<sup>-1</sup> from DM<sub>2</sub>) at 34 DAMA was correlated to the highest ST (19.9 °C) during 2016 (Figure 2.5). Whereas, the high emission peak (0.15 mg m<sup>-2</sup> h<sup>-1</sup> from DM<sub>1</sub>) at 127 DAMA in 2017 growing season was related to the highest SM (33.5 %) at that day (Figure 2.6). BC amendment suppressed N<sub>2</sub>O emissions from DM<sub>1</sub>, DM<sub>2</sub> and IN during both years. The decrease in  $N_2O$  emissions with BC incorporation was observed by several researchers previously (Augustenborg et al., 2012; Singh et al., 2010; Spokas and Reicosky, 2009; Taghizadeh-Toosi et al., 2011; Van Zwieten et al., 2010b; Yanai et al., 2007). There are several mechanisms by which BC could reduce N<sub>2</sub>O emissions. Application of BC improves soil aeration by reducing the soil bulk density resulting in a decrease in the activity of denitrifiers in paddy fields (Zhang et al., 2010). Reduction of N<sub>2</sub>O emissions after BC amendment had been explained due to different mechanisms including modification of SM, increased aeration, inhibition of nitrifier and denitrifier communities (Laird et al., 2009; Yanai et al., 2007). BC application to soil accelerates; (1) the growth of soil microbes (e.g. Bradyrhizobiaceae and Hyphomicrobiaceae families) that can decrease  $N_2O$  emission by supporting denitrification of  $NO_3^-$  to  $N_2$ , (2) the mycobacterial reduction of  $NO_3^-$  to  $NH_4^+$ , and (3) adsorption of  $NH_4^+$  on BC surface

decreasing the abundance of microorganisms involved in nitrification of  $NH_4^+$  to nitrite  $(NO_2^-)$  (Anderson et al., 2011). BC particles adsorb  $NH_4^+$  on their surface and, reduce its availability for nitrification, as a result, N<sub>2</sub>O emission is declined (Berglund et al., 2004; Lehmann et al., 2006).

# 2.6. Conclusion

BC application reduced the cumulative GHG emission during both growing seasons from silage corn cropping system in western Newfoundland. Overall, BC application to DM<sub>1</sub>, DM<sub>2</sub> and IN decreased cumulative CO<sub>2</sub> emission by 16, 25.5 and 26.5 %, cumulative CH<sub>4</sub> emission by 184, 200 and 293 %, and cumulative N<sub>2</sub>O emission by 95, 86 and 93 %, respectively. BC treatments exhibited significantly higher SM contents at all sampling points than non-BC treatments. BC incorporation also reduced the GWP by 24.9, 34.5, and 37 %, and GHGI by 30, 37.5, and 43.4 % in DM<sub>1</sub>, DM<sub>2</sub> and IN treatment, respectively. Based on the results of this study, it is concluded that BC application to soil with DM and IN fertilizer have a great potential to reduce GHGs emissions, global warming and climate change without compromising the dry matter yield of silage corn crop in western Newfoundland.

#### 2.7. References

- Agegnehu, G., Bass, A.M., Nelson, P.N., Bird, M.I., 2016. Benefits of biochar, compost and biochar-compost for soil quality, maize yield and greenhouse gas emissions in a tropical agricultural soil. Sci. Total Environ. 543, 295–306.
- Agriculture and Agri-Food Canada, n.d. Agricultural Greenhouse Gas Indicator -Agriculture and Agri-Food Canada (AAFC) [WWW Document].
- Ahmed, A., Kurian, J., Raghavan, V., 2016. Biochar influences on agricultural soils, crop production, and the environment: A review. Environ. Rev. 24, 495–502.
- Amon, B., Kryvoruchko, V., Amon, T., Zechmeister-Boltenstern, S., 2006. Methane, nitrous oxide and ammonia emissions during storage and after application of dairy cattle slurry and influence of slurry treatment. Agric. Ecosyst. Environ. 112, 153– 162.
- Anderson, C.R., Condron, L.M., Clough, T.J., Fiers, M., Stewart, A., Hill, R.A., Sherlock, R.R., 2011. Biochar induced soil microbial community change: Implications for biogeochemical cycling of carbon, nitrogen and phosphorus. Pedobiologia Int. J. Soil Biol. 54, 309–320.
- Augustenborg, C.A., Hepp, S., Kammann, C., Hagan, D., Schmidt, O., Müller, C., 2012.
  Biochar and Earthworm Effects on Soil Nitrous Oxide and Carbon Dioxide
  Emissions. J. Environ. Qual. 41, 1203–1209.

Barneze, A.S., Mazzetto, A.M., Zani, C.F., Misselbrook, T., Cerri, C.C., 2014. Nitrous

oxide emissions from soil due to urine deposition by grazing cattle in Brazil. Atmos. Environ. 92, 394–397.

- Bera, T., Collins, H.P., Alva, A.K., Purakayastha, T.J., Patra, A.K., 2016. Biochar and manure effluent effects on soil biochemical properties under corn production. Appl. Soil Ecol. 107, 360–367.
- Berglund, L.M., Deluca, T.H., Zackrisson, O., 2004. Activated carbon amendments to soil alters nitrification rates in Scots pine forests. Soil Biol. Biochem. 36, 2067– 2073.
- Bol, R., Moering, J., Kuzyakov, Y., Amelung, W., 2003. Quantification of priming and CO2 respiration sources following slurry-C incorporation into two grassland soils with different C content. Rapid Commun. Mass Spectrom. 17, 2585–2590.
- Bolan, N.S., Adriano, D.C., Mahimairaja, S., 2004. Distribution and bioavailability of trace elements in livestock and poultry manure by-products. Crit. Rev. Environ. Sci. Technol. 34, 291–338.
- Brennan, R.B., Healy, M.G., Fenton, O., Lanigan, G.J., 2015. The effect of chemical amendments used for phosphorus abatement on greenhouse gas and ammonia emissions from dairy cattle slurry: Synergies and pollution swapping. PLoS One 10, e0111965.
- Bruun, E.W., Ambus, P., Egsgaard, H., Hauggaard-Nielsen, H., 2012. Effects of slow and fast pyrolysis biochar on soil C and N turnover dynamics. Soil Biol. Biochem. 46,

- Cai, Y., Ding, W., Luo, J., 2013. Nitrous oxide emissions from Chinese maize-wheat rotation systems: A 3-year field measurement. Atmos. Environ. 65, 112–122.
- Case, S.D.C., McNamara, N.P., Reay, D.S., Whitaker, J., 2012. The effect of biochar addition on N2O and CO2 emissions from a sandy loam soil - The role of soil aeration. Soil Biol. Biochem. 51, 125–134.
- Castaldi, S., Riondino, M., Baronti, S., Esposito, F.R.R., Marzaioli, R., Rutigliano,
  F.A.A., Vaccari, F.P.P., Miglietta, F., 2011. Impact of biochar application to a
  Mediterranean wheat crop on soil microbial activity and greenhouse gas fluxes.
  Chemosphere 85, 1464–1471.
- Cayuela, M.L., Jeffery, S., van Zwieten, L., 2015. The molar H:Corg ratio of biochar is a key factor in mitigating N2O emissions from soil. Agric. Ecosyst. Environ. 202, 135–138.
- Cayuela, M.L., Sanchez-Monedero, M.A., Roig, A., Hanley, K., Enders, A., Lehmann, J., 2013. Biochar and denitrification in soils: when, how much and why does biochar reduce N2O emissions? Sci. Rep. 3, 1732.
- Chang, J., Clay, D.E., Clay, S.A., Chintala, R., Miller, J.M., Schumacher, T., 2016.
  Biochar reduced nitrous oxide and carbon dioxide emissions from soil with different water and temperature cycles. Agron. J. 108, 2214–2221.

Chen, J., Kim, H., Yoo, G., 2015. Effects of biochar addition on CO2 and N2O emissions
following fertilizer application to a cultivated grassland soil. PLoS One 10, e0126841.

- Collier, S.M., Ruark, M.D., Oates, L.G., Jokela, W.E., Dell, C.J., 2014. Measurement of greenhouse gas flux from agricultural soils using static chambers. J. Vis. Exp. e52110.
- Conway, G., 2012. How to create resilient agriculture. Appropr. Technol. 39, 12-14.
- Coppens, F., Garnier, P., De Gryze, S., Merckx, R., Recous, S., 2006. Soil moisture, carbon and nitrogen dynamics following incorporation and surface application of labelled crop residues in soil columns. Eur. J. Soil Sci. 57, 894–905.
- Diacono, M., Montemurro, F., 2010. Long-term effects of organic amendments on soil fertility. A review. Agron. Sustain. Dev. 30, 401–422.
- Feng, Y., Xu, Y., Yu, Y., Xie, Z., Lin, X., 2012. Mechanisms of biochar decreasing methane emission from Chinese paddy soils. Soil Biol. Biochem. 46, 80–88.
- Galloway, J.N., Townsend, A.R., Erisman, J.W., Bekunda, M., Cai, Z., Freney, J.R.,
  Martinelli, L.A., Seitzinger, S.P., Sutton, M.A., 2008. Transformation of the nitrogen cycle: recent trends, questions, and potential solutions. Science (80-. ). 320, 889–892.
- Hagemann, N., Kammann, C.I., Schmidt, H.P., Kappler, A., Behrens, S., 2017. Nitrate capture and slow release in biochar amended compost and soil. PLoS One 12, e0171214.

- Hawthorne, I., Johnson, M.S., Jassal, R.S., Black, T.A., Grant, N.J., Smukler, S.M., 2017. Application of biochar and nitrogen influences fluxes of CO2, CH4 and N2O in a forest soil. J. Environ. Manage. 192, 203–214.
- Helbig, M., Quinton, W.L., Sonnentag, O., 2017. Warmer spring conditions increase annual methane emissions from a boreal peat landscape with sporadic permafrost. Environ. Res. Lett. 12.
- Hernandez-Soriano, M.C., Kerré, B., Kopittke, P.M., Horemans, B., Smolders, E., 2016. Biochar affects carbon composition and stability in soil: a combined spectroscopymicroscopy study. Sci. Rep. 6, 25127.
- Holland, E.A., Robertson, phiilip G., Greenberg, J., Groffman, P.M., Boone, R.D., Gosz,J.R., 1999. Soil CO2, N2O, and CH4 exchange. Stand. Soil Methods Long-TermEcol. Res. 185–201.
- Hrapovic, L., Rowe, R.K., 2002. Intrinsic degradation of volatile fatty acids in laboratorycompacted clayey soil. J. Contam. Hydrol. 58, 221–242.
- HUANG, J., SUI, P., GAO, W., CHEN, Y., 2017. Effect of Maize-Soybean Intercropping on Soil Nitrous Oxide Emissions in Silt Loam Soil of the North China Plain. Pedosphere.
- Hüppi, R., Felber, R., Neftel, A., Six, J., Leifeld, J., 2015. Effect of biochar and liming on soil nitrous oxide emissions from a temperate maize cropping system. Soil 1, 707–717.

- IPCC, 2014a. Climate change 2014 : mitigation : contribution of Working Group III to the third assessment report of the Intergovernmental Panel on Climate Change. Cambridge University Press.
- IPCC, 2014b. Climate change 2014: synthesis report. Contribution of Working Groups I, II and III to the fifth assessment report of the Intergovernmental Panel on Climate.
- Ippolito, J.A.A., Stromberger, M.E.E., Lentz, R.D.D., Dungan, R.S.S., 2016. Hardwood biochar and manure co-application to a calcareous soil. Chemosphere 142, 84–91.
- Jia, J.X., Li, B., Chen, Z.Z., Xie, Z.B., Xiong, Z.Q., 2012. Effects of biochar application on vegetable production and emissions of N2O and CH4. Soil Sci. Plant Nutr. 58, 503–509.
- Khan, A.U.H., Iqbal, M., Islam, K.R., 2007. Dairy manure and tillage effects on soil fertility and corn yields. Bioresour. Technol. 98, 1972–1979.
- Kim, J., Yoo, G., Kim, D., Ding, W., Kang, H., 2017. Combined application of biochar and slow-release fertilizer reduces methane emission but enhances rice yield by different mechanisms. Appl. Soil Ecol. 117, 57–62.
- Kirby, G.E., 1988. Soils of the Pasadena-Deer Lake area, Newfoundland. St. John's, Newfoundland.
- Laird, D., Fleming, P., Wang, B., Horton, R., Karlen, D., 2010. Biochar impact on nutrient leaching from a Midwestern agricultural soil. Geoderma 158, 436–442.
- Laird, D.A., Brown, R.C., Amonette, J.E., Lehmann, J., 2009. Review of the pyrolysis

platform for coproducing bio-oil and biochar. Biofuels, Bioprod. Biorefining 3, 547– 562.

- Laird, D.A., Fleming, P., Davis, D.D., Horton, R., Wang, B., Karlen, D.L., 2010. Impact of biochar amendments on the quality of a typical Midwestern agricultural soil. Geoderma 158, 443–449.
- Lal, R., 2011. Sequestering carbon in soils of agro-ecosystems. Food Policy 36, S33–S39.
- Lamb, A., Green, R., Bateman, I., Broadmeadow, M., Bruce, T., Burney, J., Carey, P.,
  Chadwick, D., Crane, E., Field, R., Goulding, K., Griffiths, H., Hastings, A., Kasoar,
  T., Kindred, D., Phalan, B., Pickett, J., Smith, P., Wall, E., zu Ermgassen, E.K.H.J.,
  Balmford, A., 2016. The potential for land sparing to offset greenhouse gas
  emissions from agriculture. Nat. Clim. Chang. 6, 1–5.
- Lan, Z.M., Chen, C.R., Rashti, M.R., Yang, H., Zhang, D.K., 2017. Stoichiometric ratio of dissolved organic carbon to nitrate regulates nitrous oxide emission from the biochar-amended soils. Sci. Total Environ. 576, 559–571.
- Laufer, J., Tomlinson, T., 2012. Biochar Field Studies : An IBI Research Summary.
- Lehmann, J., 2007. Bio-energy in the black. Front. Ecol. Environ. 5, 381–387.
- Lehmann, J., Gaunt, J., Rondon, M., 2006. Bio-char sequestration in terrestrial ecosystems A review. Mitig. Adapt. Strateg. Glob. Chang. 11, 403–427.
- Lentz, R.D., Ippolito, J.A., Spokas, K.A., 2014. Biochar and Manure Effects on Net Nitrogen Mineralization and Greenhouse Gas Emissions from Calcareous Soil under

Corn. Soil Sci. Soc. Am. J. 78, 1641–1655.

- Li, B., Fan, C.H., Zhang, H., Chen, Z.Z., Sun, L.Y., Xiong, Z.Q., 2015. Combined effects of nitrogen fertilization and biochar on the net global warming potential, greenhouse gas intensity and net ecosystem economic budget in intensive vegetable agriculture in southeastern China. Atmos. Environ. 100, 10–19.
- Li, C., Frolking, S., Butterbach-Bahl, K., 2005. Carbon sequestration in arable soils is likely to increase nitrous oxide emissions, offsetting reductions in climate radiative forcing. Clim. Change 72, 321–338.
- Liu, J., Shen, J., Li, Y., Su, Y., Ge, T., Jones, D.L., Wu, J., 2014. Effects of biochar amendment on the net greenhouse gas emission and greenhouse gas intensity in a Chinese double rice cropping system. Eur. J. Soil Biol. 65, 30–39.
- Liu, L., Wang, Y., Yan, X., Li, J., Jiao, N., Hu, S., 2017. Biochar amendments increase the yield advantage of legume-based intercropping systems over monoculture. Agric. Ecosyst. Environ. 237, 16–23.
- Liu, Q., Liu, B., Zhang, Y., Lin, Z., Zhu, T., Sun, R., Wang, X., Ma, J., Bei, Q., Liu, G., Lin, X., Xie, Z., 2017. Can biochar alleviate soil compaction stress on wheat growth and mitigate soil N2O emissions? Soil Biol. Biochem. 104, 8–17.
- Liu, X. yu, Qu, J. jing, Li, L. qing, Zhang, A. feng, Jufeng, Z., Zheng, J. wei, Pan, G. xing, 2012. Can biochar amendment be an ecological engineering technology to depress N 2O emission in rice paddies?-A cross site field experiment from South

China. Ecol. Eng. 42, 168–173.

- Liu, Y., Yang, M., Wu, Y., Wang, H., Chen, Y., Wu, W., 2011. Reducing CH4 and CO2 emissions from waterlogged paddy soil with biochar. J. Soils Sediments 11, 930– 939.
- Lu, C., Tian, H., 2017. Global nitrogen and phosphorus fertilizer use for agriculture production in the past half century: shifted hot spots and nutrient imbalance. Earth Syst. Sci. Data 9, 181–192.
- Lu, W., Ding, W., Zhang, J., Li, Y., Luo, J., Bolan, N., Xie, Z., 2014. Biochar suppressed the decomposition of organic carbon in a cultivated sandy loam soil: A negative priming effect. Soil Biol. Biochem. 76, 12–21.
- Maillard, É., Angers, D.A., 2014. Animal manure application and soil organic carbon stocks: A meta-analysis. Glob. Chang. Biol. 20, 666–679.
- Matsi, T., Lithourgidis, A.S., Barbayiannis, N., 2015. Effect of liquid cattle manure on soil chemical properties and corn growth in northern greece. Exp. Agric. 51, 435–450.
- Mazzoncini, M., Sapkota, T.B., Bàrberi, P., Antichi, D., Risaliti, R., 2011. Long-term effect of tillage, nitrogen fertilization and cover crops on soil organic carbon and total nitrogen content. Soil Tillage Res. 114, 165–174.
- Meinshausen, M., Meinshausen, N., Hare, W., Raper, S.C.B., Frieler, K., Knutti, R., Frame, D.J., Allen, M.R., 2009. Greenhouse-gas emission targets for limiting global

warming to 2 °C. Nature 458, 1158–1162.

- Menéndez, S., Merino, P., Pinto, M., González-Murua, C., Estavillo, J.M., 2006. 3,4-Dimethylpyrazol phosphate effect on nitrous oxide, nitric oxide, ammonia, and carbon dioxide emissions from grasslands. J. Environ. Qual. 35, 973–981.
- Minasny, B., Malone, B.P., McBratney, A.B., Angers, D.A., Arrouays, D., Chambers, A., Chaplot, V., Chen, Z.-S., Cheng, K., Das, B.S., Field, D.J., Gimona, A., Hedley,
  C.B., Hong, S.Y., Mandal, B., Marchant, B.P., Martin, M., McConkey, B.G.,
  Mulder, V.L., O'Rourke, S., Richer-de-Forges, A.C., Odeh, I., Padarian, J., Paustian,
  K., Pan, G., Poggio, L., Savin, I., Stolbovoy, V., Stockmann, U., Sulaeman, Y., Tsui,
  C.-C., Vågen, T.-G., van Wesemael, B., Winowiecki, L., 2017. Soil carbon 4 per
  mille. Geoderma 292, 59–86.
- Mizuta, K., Matsumoto, T., Hatate, Y., Nishihara, K., Nakanishi, T., 2004. Removal of nitrate-nitrogen from drinking water using bamboo powder charcoal. Bioresour. Technol. 95, 255–257.
- Paustian, K., Lehmann, J., Ogle, S., Reay, D., Robertson, G.P., Smith, P., 2016. Climatesmart soils. Nature 532, 49–57.
- SAMCO Agricultural Manufacturing Ltd, n.d. Samco 2200.
- Schimmelpfennig, S., Müller, C., Grünhage, L., Koch, C., Kammann, C., 2014. Biochar, hydrochar and uncarbonized feedstock application to permanent grassland-Effects on greenhouse gas emissions and plant growth. Agric. Ecosyst. Environ. 191, 39–52.

- Schlesinger, W.H., 2009. On the fate of anthropogenic nitrogen. Proc. Natl. Acad. Sci. 106, 203–208.
- Sheng, Y., Zhu, L., 2018. Biochar alters microbial community and carbon sequestration potential across different soil pH. Sci. Total Environ. 622–623, 1391–1399.
- Sherlock, R.R., Sommer, S.G., Khan, R.Z., Wood, C.W., Guertal, E.A., Freney, J.R., Dawson, C.O., Cameron, K.C., 2002. Ammonia, Methane, and Nitrous Oxide Emission from Pig Slurry Applied to a Pasture in New Zealand. J. Environ. Qual. 31, 1491–1501.
- Singh, B.P., Hatton, B.J., Singh, B., Cowie, A.L., Kathuria, A., 2010. Influence of Biochars on Nitrous Oxide Emission and Nitrogen Leaching from Two Contrasting Soils. J. Environ. Qual. 39, 1224–1235.
- Smith, P., Martino, D., Cai, Z., Gwary, D., Janzen, H., Kumar, P., McCarl, B., Ogle, S.,
  O'Mara, F., Rice, C., Scholes, B., Sirotenko, O., Howden, M., McAllister, T., Pan,
  G., Romanenkov, V., Schneider, U., Towprayoon, S., 2007. Policy and technological constraints to implementation of greenhouse gas mitigation options in agriculture.
  Agric. Ecosyst. Environ. 118, 6–28.
- Spokas, K.A., Reicosky, D.C., 2009. Impacts of sixteen different biochars on soil greenhouse gas production. Ann. Environ. Sci. 3, 179–193.
- Statistics Canada, 2006. A geographical profile of livestock manure production in Canada.

- Suddick, E.C., Six, J., 2013. An estimation of annual nitrous oxide emissions and soil quality following the amendment of high temperature walnut shell biochar and compost to a small scale vegetable crop rotation. Sci. Total Environ. 465, 298–307.
- Sun, L., Li, L., Chen, Z., Wang, J., Xiong, Z., 2014. Combined effects of nitrogen deposition and biochar application on emissions of N2 O, CO2 and NH3 from agricultural and forest soils. Soil Sci. Plant Nutr. 60, 254–265.
- Sun, Z., Sänger, A., Rebensburg, P., Lentzsch, P., Wirth, S., Kaupenjohann, M., Meyer-Aurich, A., 2017. Contrasting effects of biochar on N2O emission and N uptake at different N fertilizer levels on a temperate sandy loam. Sci. Total Environ. 578, 557– 565.
- Taghizadeh-Toosi, A., Clough, T.J., Condron, L.M., Sherlock, R.R., Anderson, C.R.,
  Craigie, R.A., 2011. Biochar Incorporation into Pasture Soil Suppresses in situ
  Nitrous Oxide Emissions from Ruminant Urine Patches. J. Environ. Qual. 40, 468–478.
- Tan, Z., Lin, C.S.K., Ji, X., Rainey, T.J., 2017. Returning biochar to fields: A review. Appl. Soil Ecol. 116, 1–11.
- Thangarajan, R., Bolan, N.S., Tian, G., Naidu, R., Kunhikrishnan, A., 2013. Role of organic amendment application on greenhouse gas emission from soil. Sci. Total Environ. 465, 72–96.
- Thomazini, A., Spokas, K., Hall, K., Ippolito, J., Lentz, R., Novak, J., 2015. GHG

impacts of biochar: Predictability for the same biochar. Agric. Ecosyst. Environ. 207, 183–191.

- Tubiello, F.N., Salvatore, M., Ferrara, A.F., House, J., Federici, S., Rossi, S., Biancalani,
  R., Condor Golec, R.D., Jacobs, H., Flammini, A., Prosperi, P., Cardenas-Galindo,
  P., Schmidhuber, J., Sanz Sanchez, M.J., Srivastava, N., Smith, P., 2015. The
  Contribution of Agriculture, Forestry and other Land Use activities to Global
  Warming, 1990-2012. Glob. Chang. Biol. 21, 2655–2660.
- Usman, A.R.A., Al-Wabel, M.I., Ok, Y.S., Al-Harbi, A., Wahb-Allah, M., El-Naggar,
  A.H., Ahmad, M., Al-Faraj, A., Al-Omran, A., 2016. Conocarpus Biochar Induces
  Changes in Soil Nutrient Availability and Tomato Growth Under Saline Irrigation.
  Pedosphere 26, 27–38.
- Van Zwieten, L., Kimber, S., Morris, S., Chan, K.Y., Downie, A., Rust, J., Joseph, S., Cowie, A., 2010a. Effects of biochar from slow pyrolysis of papermill waste on agronomic performance and soil fertility. Plant Soil 327, 235–246.
- Van Zwieten, L., Kimber, S., Morris, S., Downie, A., Berger, E., Rust, J., Scheer, C., 2010b. Influence of biochars on flux of N2O and CO2from Ferrosol. Aust. J. Soil Res. 48, 555–568.
- Wang, D., Fonte, S.J., Parikh, S.J., Six, J., Scow, K.M., 2017. Biochar additions can enhance soil structure and the physical stabilization of C in aggregates. Geoderma 303, 110–117.

- Wang, J., Pan, X., Liu, Y., Zhang, X., Xiong, Z., 2012. Effects of biochar amendment in two soils on greenhouse gas emissions and crop production. Plant Soil 360, 287– 298.
- Wang, X., Ren, Y., Zhang, S., Chen, Y., Wang, N., 2017. Applications of organic manure increased maize (Zea mays L.) yield and water productivity in a semi-arid region.Agric. Water Manag. 187, 88–98.
- Wang, Z., Zheng, H., Luo, Y., Deng, X., Herbert, S., Xing, B., 2013. Characterization and influence of biochars on nitrous oxide emission from agricultural soil. Environ.Pollut. 174, 289–296.
- West, T.O., Post, W., 2002. Soil organic carbon sequestration rates by tillage and crop rotation: A global data analysis. Soil Sci. Soc. Am. J. 66, 1930–1946.
- Yanai, Y., Toyota, K., Okazaki, M., 2007. Effects of charcoal addition on N2O emissions from soil resulting from rewetting air-dried soil in short-term laboratory experiments. Soil Sci. Plant Nutr. 53, 181–188.
- Yang, X., Lan, Y., Meng, J., Chen, W., Huang, Y., Cheng, X., He, T., Cao, T., Liu, Z., Jiang, L., Gao, J., 2017. Effects of maize stover and its derived biochar on greenhouse gases emissions and C-budget of brown earth in Northeast China. Environ. Sci. Pollut. Res. 24, 8200–8209.
- Zhang, A., Bian, R., Hussain, Q., Li, L., Pan, G., Zheng, J.J., Zhang, X., Zheng, J.J., 2013. Change in net global warming potential of a rice-wheat cropping system with

biochar soil amendment in a rice paddy from China. Agric. Ecosyst. Environ. 173, 37–45.

- Zhang, A., Bian, R., Pan, G., Cui, L., Hussain, Q., Li, L., Zheng, J.J.J.J., Zheng, J.J.J.J., Zhang, X., Han, X., Yu, X., 2012. Effects of biochar amendment on soil quality, crop yield and greenhouse gas emission in a Chinese rice paddy: A field study of 2 consecutive rice growing cycles. F. Crop. Res. 127, 153–160.
- Zhang, A., Cui, L., Pan, G., Li, L., Hussain, Q., Zhang, X., Zheng, J., Crowley, D., 2010.
  Effect of biochar amendment on yield and methane and nitrous oxide emissions from a rice paddy from Tai Lake plain, China. Agric. Ecosyst. Environ. 139, 469–475.
- Zhang, A., Liu, Y., Pan, G., Hussain, Q., Li, L., Zheng, J., Zhang, X., 2011. Effect of biochar amendment on maize yield and greenhouse gas emissions from a soil organic carbon poor calcareous loamy soil from Central China Plain. Plant Soil 351, 263–275.
- Zhang, Z.S., Cao, C.G., Guo, L.J., Li, C.F., 2014. The effects of rape residue mulching on net global warming potential and greenhouse gas intensity from no-tillage paddy fields. Sci. World J. 2014.
- Zheng, H., Wang, X., Luo, X., Wang, Z., Xing, B., 2018. Biochar-induced negative carbon mineralization priming effects in a coastal wetland soil: Roles of soil aggregation and microbial modulation. Sci. Total Environ. 610–611, 951–960.

Zhou, H., Zhang, D., Wang, P., Liu, X., Cheng, K., Li, L., Zheng, J.J., Zhang, X., Zheng,

J.J., Crowley, D., van Zwieten, L., Pan, G., 2017. Changes in microbial biomass and the metabolic quotient with biochar addition to agricultural soils: A Meta-analysis. Agric. Ecosyst. Environ. 239, 80–89.

Zhou, M., Zhu, B.O., Wang, S., Zhu, X., Vereecken, H., 2017. Stimulation of N2O emission by manure application to agricultural soils may largely offset carbon benefits: a global meta-analysis. Glob. Chang. Biol. 23, 4068–4083.

## Chapter 3

# 3. Biomass production, and nitrogen dynamics of silage corn as influenced by organic and inorganic nitrogen sources and biochar amendment

# 3.1. Abstract

Most of the nitrogen (N) applied to crops is leached down in the form of nitrates (NO<sub>3</sub><sup>-</sup>) resulting in reduced N use efficiency. To assess the potential of biochar (BC) application for the mitigation of NO<sub>3</sub><sup>-</sup> leaching and retention of NO<sub>3</sub><sup>-</sup> and ammonium (NH<sub>4</sub><sup>+</sup>), a two year field study was conducted with two dairy manure sources (DM<sub>1</sub>, DM<sub>2</sub>), inorganic N (IN), their combination with BC and a control (no N). Three silage corn genotypes (A4177G3RIB, DKC26-28 RIB, Yukon R) were used. BC application significantly reduced the NO<sub>3</sub><sup>-</sup> and NH<sub>4</sub><sup>+</sup> movement to deep soil and it enhanced their retention in topsoil, increased soil pH, enhanced N concentration in plant tissues and increased dry matter yield in all silage corn genotypes. On average, BC addition to DM<sub>1</sub>, DM<sub>2</sub>, and IN enhanced N uptake by 13.5, 11.5 and 17.3 % and dry matter yield by 6, 5.5, and 8.75 %, respectively. Conclusively, BC application to soil could improve soil pH, reduce NO<sub>3</sub><sup>-</sup> and NH<sub>4</sub><sup>+</sup> loses by increasing their residence time in soil, hence N concentration and dry matter production in silage corn cropping systems in western Newfoundland.

Keywords: nitrate/ammonium retention, silage corn, dairy manure, biochar

## 3.2. Introduction

Dairy manure (DM) and inorganic nitrogen (IN) fertilizers are the major sources of plant nutrients and are being applied to crops worldwide to boost the agricultural productivity (Jokela et al. 2014; Wang et al. 201 7c; Parker et al. 2018). DM application improves physiochemical properties of soil and agronomic performance of plants e.g. soil organic carbon (SOC), bulk density, soil aggregation, nutrient status and uptake, crop growth and yield (Forge et al., 2016; Martínez et al., 2017). However, excessive IN fertilizers and DM application enhance the risk of NO<sub>3</sub><sup>-</sup> leaching in different cropping systems and consequently increase cost of production, pollute water bodies and pose a serious threat to human health (Fan et al., 2017; Forge et al., 2016; Jokela et al., 2014; Long and Sun, 2012; Masaka et al., 2015). It has been reported that 34-92 % of the N is leached from manure application to soil was in the form of  $NO_3^-$  and 14-57 % in dissolved organic nitrogen (DON) form (Fan et al., 2017). However, soil NO<sub>3</sub><sup>-</sup> concentration and leaching is site-specific and is mainly driven by rainfall, management practices (cover crop, fertilizer sources, crop rotation), and soil texture (Gaines and Gaines, 1994; Jabloun et al., 2015; Jean et al., 2000). For instance,  $NO_3^-$  leaching losses of 55-59 kg per hectare per year with DM application, 30-35 kg per hectare per year from compost application and 25-33 kg per hectare per year from IN application has been reported in a six-year corn - alfalfa crop rotation (Basso and Ritchie, 2005). Significant variation in NO<sub>3</sub><sup>-</sup> losses has been observed in different cropping systems, but corn-based cropping systems have been found to have the highest  $NO_3$  leaching (Hargrave and Shaykewich, 1997; Hernandez-Ramirez et al., 2011; van Es et al., 2006). The annual estimated loss of  $NO_3^-$  from cornfield amended with Ammonium nitrate (NH<sub>4</sub>NO<sub>3</sub>) in Manitoba was 160 kg per hectare per year (Hargrave and Shaykewich, 1997), whereas, in Southern Turkey, the annual NO<sub>3</sub><sup>-</sup> load was 39-55 kg per hectare per year in wheat-corn cropping system when a mixture of NH<sub>4</sub>NO<sub>3</sub>, ammonium sulfate (NH<sub>4</sub>)<sub>2</sub> SO<sub>4</sub>, urea (NH<sub>2</sub>)<sub>2</sub> CO, and compound fertilizer was applied (Ibrikci et al., 2015). These N losses indicate ineffectiveness in the current nutrient management strategies or management practices that result not only in environmental pollution but also an economic loss to the farmers (Güereña et al., 2013). Different nutrient management strategies and crop management practices have been practiced to reduce N losses and improve N use efficiency (NUE). These include, slow-release fertilizers (Ye et al., 2013), burial of crop residues and crop rotation (Hirel et al., 2007), judicious use of fertilizers (Francis, 1992), legume-based cropping systems (Drinkwater et al., 1998), variable rate technologies (VRT) (Gatti et al., 2018), use of nitrification inhibitors (NI) (Zhang et al., 2015), and plant trait selection (Ju et al., 2015).

Biochar (BC) is a form of black carbon (C) created by thermal degradation of organic material (e.g., wood, manure, leaves, etc.) in zero or low oxygen environments (Lehmann and Joseph, 2009). BC is recalcitrant in nature (Spokas, 2010) and its reactive surfaces are capable of sorbing and exchanging nutrients and native organic matter (Liang et al., 2006); therefore, there is a great potential and interest in utilizing BC as a soil amendment to sequester C and improve soil fertility in agricultural soils. Additionally, BC application could be one of the best approaches to improve N retention in topsoil, reduce NO<sub>3</sub><sup>-</sup> leaching and improve soil fertility in agricultural systems (Haider et al., 2015; Knowles et al., 2011; Laird et al., 2010; Lehmann, 2007). BC application decreases soil bulk density,

increases porosity, pH, nutrient use efficiency, N<sub>2</sub>-fixation, soil saturation, water holding capacity (WHC) (Busch et al., 2012; Busscher et al., 2010; Harter et al., 2014; Hussain et al., 2017; Jervin et al., 2017; Kammann et al., 2012; Karhu et al., 2011), enhances Ammonium  $(NH_4^+)$  retention and availability, increases urease activity preventing ammonia losses, and eventually reduces NO<sub>3</sub><sup>-</sup> leaching loses (Amendola et al., 2017; Cao et al., 2017; Huang et al., 2017; Sun et al., 2017). BC amendment increased hydraulic conductivity, soil water availability and infiltration (Asai et al., 2009; Baronti et al., 2014; Buss et al., 2012; Ippolito et al., 2012), and improved soil aeration (Case et al., 2012; Cayuela et al., 2013) and nutrient retention (Clough et al., 2013; Ventura et al., 2012; L. Wang et al., 2017). BC can also increase soil microbial activities, alter microbial community structure and extracellular enzymatic activities (Foster et al., 2016; Gul et al., 2015; Lu et al., 2015). There is a significant effect of feedstock source and pyrolysis process on characteristics of BC that consequently affect the physiochemical and biological properties of soil (Borchard et al., 2014; Gul et al., 2015; Lentz and Ippolito, 2012; Schmidt et al., 2014; Spokas and Reicosky, 2009; Vitkova et al., 2017). Application of BC produced at 550 °C increased soil NO<sub>3</sub><sup>-</sup> concentration, nutrient uptake, and corn dry matter yield (Haider et al., 2015; Smider and Singh, 2014). Pinewood BC amendment at the rate of 0.5, 2.5 and 10 % w/w in sandy soil reduced NO<sub>3</sub><sup>-</sup> leaching by 26, 42 and 96 %, respectively (Sika and Hardie, 2014). BC amendment to soil may increase NO<sub>3</sub><sup>-</sup> residence time in soil possibly due to unconventional H-bonding in micro and nano-pores (Kammann et al., 2015), or development of functional groups and organomineral complexes on BC surface (Joseph et al., 2013; Lin et al., 2013; Prost et al., 2013), acting as slow-release fertilizer which allows plants to absorb more NO<sub>3</sub><sup>-</sup> from the soil (Hagemann et al., 2017; Kameyama et al., 2012; Uzoma et al., 2011). A significant increase in inorganic N pool (NH<sub>4</sub><sup>+</sup>-N, NO<sub>3</sub><sup>-</sup>-N), leaf N concentration, photosynthesis rate, and pod yield have been recorded with BC application (Xu et al., 2015). Incorporation of rice straw BC to acidic soil accelerated nitrification but decreased NO<sub>3</sub><sup>-</sup> leaching and improved N adsorption (Zhao et al., 2014). BC amendment to a degraded Chernozem soil at 5 % (w/w) application rate significantly improved spinach growth, increased uptake of K in plant tissues, free amino acid contents, and proline content but limited Ca, Mg and Na concentrations (Zemanová et al., 2017). Contrary to reported above (Tammeorg et al., 2014) observed that BC application did not improve N uptake and grain yield of wheat, faba bean and turnip rape in a three-year field experiment. In Newfoundland, most of the cultivated soils are acidic and sandy and the growing season is short with frequent rainfall events. The average precipitation received in this area is 1113 mm per year with less than 410 mm as snow during last 30 years (1986-2016) recorded from the nearest weather station in Deer Lake by Environment Canada (Badewa, 2017). This is a most favorable condition for N leaching. Most of the N applied to sandy soils is lost and is not available to crops (Gaines and Gaines, 1994; Jabloun et al., 2015; Jean et al., 2000), which results in low crop productivity and economic loss. Therefore, I hypothesized that BC amendment would reduce nitrate leaching; enhance soil pH and biomass production of silage corn in the podzolic soil. This research project was planned with the following specific objectives;

i- To investigate the role of BC amendment on soil's NO<sub>3</sub><sup>-</sup> and NH<sub>4</sub><sup>+</sup> dynamics after dairy manure and IN fertilizer application in silage corn cropping systems.

ii- To compare the effects of DM, IN alone and co-application of BC on soil pH, N concentration and biomass production of silage corn.

# **3.3.** Material and Methods:

### 3.3.1. Study site

A field experiment was conducted at Pynn's Brook Research Station, Pasadena (49°04'21.9"N, 57°33'37.4"W), Newfoundland and Labrador (NL), Canada, during 2016 and 2017 growing seasons. The soil was classified as rapidly drained, Orthic Humo-Ferric Podzol with reddish brown to brown color, and developed on gravely sandy fluvial deposit of mixed lithology (Kirby, 1988). Moreover, this soil has limited agricultural use because of its high coarse fragment contents and rapid drainage (Kirby, 1988). Basic physio-chemical properties of the soil are described in Table 2.1.

#### **3.3.2.** Experimental setup

The experiment comprised of seven treatments with organic and inorganic N sources and BC. Treatments included were; 1) DM with high N conc. (0.37 %) designated as  $DM_1$ , 2) DM with low N conc. (0.13 %), designated as  $DM_2$ , 3) IN, 4)  $DM_1$  + BC, 5)  $DM_2$  + BC, 6) IN + BC, and 7) control (N<sub>0</sub>). The experimental design was a randomized complete block (RCBD) in a factorial setting with three replications. Plots were 1.5 m wide and 4.8 m long. DM procured from Larch Grove and Rideout's dairy operations located at Cormack area near Deer Lake, NL was used. DM samples were collected from these two dairy operations and were sent to Soil, Plant and Feed Laboratory, Department of

Fisheries and Land Resources, St. John's, NL for detailed analysis. Larch Grove's DM had a high concentration of macro and micronutrients and was designated as DM<sub>1</sub> while DM from Rideout's dairy had a low concentration of macro and micronutrients and was titled as  $DM_2$  (Table 2.2). Representative soil sample collected from the experimental site was also sent to the same laboratory for physiochemical analyses. DM was applied each year before seeding in corresponding plots @ 30,000 liters per hectare (local farmers practice). Inorganic fertilizers were applied to fulfill the required NPK nutrients based on DM and soil analyses reports. Ammonium nitrate ( $NH_4NO_3$ ), triple superphosphate (TSP) and murate of potash (MOP) were used as nitrogen (N), phosphorus (P), and potash (K) sources, and were applied at the rate of 215,110, 225 kg ha<sup>-1</sup>, respectively. DM<sub>1</sub>, DM<sub>2</sub> and the entire IN were applied before crop seeding during 2016 while in 2017  $DM_1$  and  $DM_2$ were applied to all respective treatments (DM<sub>1</sub>, DM<sub>2</sub>, DM<sub>1</sub>+B, DM<sub>2</sub>+B) before seeding but the IN fertilizer was applied in two splits (first dose: 6 leaf stage, second dose: 12 leaf stage). BC was applied only once in 2016 and incorporated to top 15 cm of the soil before DM application. BC is produced at AirTerra Inc. located in Calgary, Alberta, and is a registered BC product with the Canadian Food Inspection Agency (CFIA). This BC was produced from yellow pine wood pyrolyzed at 500 °C for 30 min by slow pyrolysis in oxygen-limited conditions. BC was applied @ 20 tons ha<sup>-1</sup> as reported by (Liu et al., 2012b). The detailed BC analyses report conducted by Gabilan laboratory, Salinas, California, USA can be seen in (Table 2.3).

#### **3.3.3.** Crop husbandry

Three silage corn genotypes, Yukon R, A4177G3 RIB, and DKC26-28 RIB were selected and used as test hybrids in this experiment. Selection of these hybrids was made on the basis of high biomass production performance in a previous field trial conducted during 2015 and low heating unit requirements. Detailed information about hybrids can be seen in Table 3.1. Seeding was done on May 24, 2016 and May 23, 2017 with the SAMCO 2200 system (SAMCO Agricultural Manufacturing Ltd.). SAMCO 3 in 1 machine which sows seed, sprays the soil with pre-emergence herbicide, and lays a thin layer of biodegradable plastic film over the seedbed. This operation protects the young plants from late frost, increases the soil temperature and thereby maximizes silage corn yield per hectare. Biodegradable plastic provides additional heat units which enhance seed germination during cold and frost days (Figure 2.1). Seeding rate for the crop was 90,900 seeds per hectare. Weeds were controlled with the spray of Roundup WeatherMax on July 09, 2016 and July 08, 2017 at the rate of 2 L per hectare.

Number	Genotype name	CHU	Company	Trait
1	Yukon R	2150	Brett Young	RR2
2	A4177G3 RIB	2175	Pride	VT3/RR
3	DKC26-28 RIB	2150	DEKLAB	GENVT2P

Table 3.1: Silage corn genotypes used in the experiment.

CHU = corn heat units, RIB= Refuge is in the bag, RR = Roundup Ready, VT3= VT TriplePro insect protection, RR2 = resistance gene to Roundup® and Factor 540®

#### **3.3.4.** Soil sampling and analysis

Soil sampling for  $NO_3^-$  and  $NH_4^+$  determination was done from 20 cm and 40 cm depths at four stages (seedling emergence, 6 leaves, 12 leaf and black layer stage) during 2016 and (6 leaves, 12 leaves, tasseling and black layer stage) in 2017. Samples were collected with augur and sealed in marked plastic bags and transferred to the laboratory within six hours of collection, where they were stored at -20 °C until further analysis. Soil samples were sieved through a 2 mm sieve to remove stones and other residues. Sieved soil (5 g) was weighed in a pre-weighed aluminum dish, and dried overnight in an oven at 105  $^{\circ}$ C and final dry weight was recorded. Moisture factor was calculated by weight difference. Sieved soil sample (5 g) was taken into a 125 mL Erlenmeyer flask and 50 mL, 2 molars KCl (potassium chloride) solution was added (1:10 soil to solution ratio) and shaken for 30 min on a reciprocating shaker (Cao et al., 2017; Carter et al., 2008; Heman et al., 2016; Sika and Hardie, 2014), and filtered into sterile 50 mL plastic tubes. The filtrate was stored at -20  $^{\circ}$ C until further analysis. NO<sub>3</sub><sup>-</sup> and Ammonium NH<sub>4</sub><sup>+</sup> concentrations in the soil extract were determined using AutoAnalyzer (Seal analytical continuous flow analyzer (AA3 HR) (Cao et al., 2017; Heman et al., 2016). NO<sub>3</sub><sup>-</sup> is reduced to nitrite (NO2) by a cadmium-copper reduction column at a pH of 8, NO2 ion then reacts with sulfanilamide to form a diazo compound. This compound then reacts with N-1naphthylethylenediamine dihydrochloride to form a reddish-purple azo dye. NH4<sup>+</sup> was determined using the salicylate chemistry and the results obtained were calculated using the formulas in Equations (3.1), (3.2), and (3.3).

Moisture factor was calculated by dividing weight of moist soil by dried soil weight

Moisture factor (MF) = 
$$\frac{\text{Moist soil weight (g)}}{\text{Oven dry weight (g)}}$$
 (3.1)

Soil NO<sub>3</sub><sup>-</sup> and NH<sub>4</sub><sup>+</sup> in moist soil was calculated by;

$$NO_{3}^{-} \text{ or } NH_{4}^{+} \text{ in moist soil (mg g}^{-1} \text{ of wet soil)}$$

$$= NO_{3}^{-} \text{ or } NH_{4}^{+} \text{ in soil extract (mg L}^{-1}) \times 10$$
(3.2)

Nitrate and ammonium quantity was multiplied with 10 as the soil-to-solution ratio was 1:10 in extract.

Soil NO<sub>3</sub><sup>-</sup> and NH<sub>4</sub><sup>+</sup> in dry soil was calculated by;

 $NO_3^-$  or  $NH_4^+$  in dry soil (mg g<sup>-1</sup> of dry soil) =  $NO_3^-$  or  $NH_4^+$  in moist soil × MF (3.3)

# 3.3.5. Soil pH determination

Soil pH was determined from the samples collected for NO<sub>3</sub><sup>-</sup> and NH<sub>4</sub><sup>+</sup> determination at three crop growth stages during 2016 and two stages in 2017. Air-dried soil samples (10 g) were taken in long plastic tubes and 20 mL, 0.01 M CaCl<sub>2</sub> (calcium chloride) was added to each tube. It was stirred for 30 minutes and let it stand for one hour (Carter et al., 2008). The pH was measured using benchtop pH meter (Oakton Instruments) (Zhang et al., 2014).

## 3.3.6. Dry matter production

Plants were harvested from a 1 m<sup>2</sup> area at black layer stage from each plot and their fresh weight was recorded. Plants were oven dried at 70  $^{\circ}$ C for 48 h and dry matter yield was calculated from each treatment plot using the formula given in Equation (2.4) and (2.5).

# 3.3.7. Plant tissue nitrogen concentration

Three dried plants from each treatment were grounded using Wiley Mill (Arthur H. Thomas) and then with CryoMill (Retsch, Germany) to a fine powder. Ground plant samples  $(4.5\pm0.5 \text{ mg})$  were weighed in tin capsules and analyzed with PerkinElmer CHN 2400 Series II for total nitrogen contents.

## 3.3.8. Statistical analysis

The analysis of variance (ANOVA) was used to determine the effect of different treatments on soil NO<sub>3</sub><sup>-</sup> and NH<sub>4</sub><sup>+</sup>, soil pH, plant N concentration and biomass production. Where treatment effects were significant, the means were compared with LSD ( $\alpha = 0.05$ ). The data were analyzed using the Statistix 10 software package (Analytical software, FL, USA) and figures were prepared using SigmaPlot 12.0 software program (Systat Software Inc., San Jose, CA).

#### 3.4. Results

## 3.4.1. Nitrate and ammonium dynamics

Soil  $NO_3^-$  and  $NH_4^+$  concentration in DM, IN alone and BC amended treatments varied significantly (p<0.05) during both growing seasons from IN, manure treatments (DM<sub>1</sub>, DM<sub>2</sub>, IN) and control (N0).

## Nitrate and ammonium during 2016

 $DM_1$ ,  $DM_2$  and the entire IN were applied before crop seeding. The NO<sub>3</sub><sup>-</sup> concentration in all treatments ranged from 7.3-55 (mg g<sup>-1</sup> dry soil) in top 20 cm soil layer at seedling emergence stage of the crop. Soil  $NO_3^-$  concentration in all treatments and stages at 20 cm depth is shown in the Figure 3.1a and 40 cm depth in Figure 3.1b. BC treatments  $(DM_1+B, DM_2+B, and IN+B)$  have high soil NO<sub>3</sub><sup>-</sup> concentration, but it was not significantly different from their non-BC treatments (DM<sub>1</sub>, DM<sub>2</sub> and IN) (Figure 3.1a) as N was applied at the same rate in all plots (DM treatments received some IN as well to balance the N requirement of the crop). Soil NO<sub>3</sub><sup>-</sup> concentration in deep soil (40 cm) ranged from 4.3-35.3 (mg g<sup>-1</sup> dry soil) in all treatments. BC treatments have relatively low NO<sub>3</sub><sup>-</sup> concentration in deep soil layer as compared to non-BC treatments which indicate less NO<sub>3</sub><sup>-</sup> movement to the deep soil at seedling emergence stage of the crop (Figure 3.1b). Soil NO<sub>3</sub><sup>-</sup> increased after seedling establishment and it reached to 78-120 (mg g<sup>-1</sup> dry soil) in all treatments (except control where it was 6.5 mg g<sup>-1</sup> dry soil) in topsoil layer at 6 leaf stage of the crop. Here BC treatments have significantly higher  $NO_3^-$  concentration (p<0.05) than non-BC treatments and the control (Figure 3.1a) which was due to the retention of NO<sub>3</sub><sup>-</sup> in BC amended soil as more NO<sub>3</sub><sup>-</sup> moved to deep soil

layer in non-BC treatments (Figure 3.1b). At 12 leaf stage there was a significant effect of BC treatments on NO<sub>3</sub><sup>-</sup> concentration (p < 0.05) in the topsoil as most of the NO<sub>3</sub><sup>-</sup> in DM<sub>1</sub>, DM<sub>2</sub>, and IN leached to deep soil but BC treatments have more NO<sub>3</sub><sup>-</sup> retained in the topsoil layer. At black layer stage there was very little NO<sub>3</sub><sup>-</sup> amount left in soil ranging from 1.13 (mg g<sup>-1</sup> dry soil) in control to 4.9 (mg g<sup>-1</sup> dry soil) in  $DM_1+B$  in topsoil and 1.40 (mg  $g^{-1}$  dry soil) in control to 2.5 (mg  $g^{-1}$  dry soil) in DM<sub>1</sub> treatments in deep soil layer. Soil NH<sub>4</sub><sup>+</sup> concentration in all treatments and stages at 20 cm depth is shown in the Figure 3.1c and 40 cm depth in Figure 3.1d. There was significantly higher  $NH_4^+$ concentration present in the topsoil layer at seedling emergence stage as compared to other stages and dropped to near zero at black layer stage. The NH<sub>4</sub><sup>+</sup> concentration ranged from 3.2 (mg g<sup>-1</sup> dry soil) in the control treatment to 80.1 (mg g<sup>-1</sup> dry soil) in DM<sub>1</sub>+B at seedling emergence stage. There was significantly higher NH<sub>4</sub><sup>+</sup> concentration in BC treatments as compared to non-BC treatments and control in 20 cm soil depth (Figure 3.1c). No significant difference between BC and non-BC treatments with respect to the NH4<sup>+</sup> concentration at 40 cm soil depth was observed whereas control treatment had the lowest  $NH_4^+$  at seedling emergence stage. At 6 leaf stage  $NH_4^+$  in the topsoil layer decreased as compared to seedling emergence stage, and BC treatments had significantly high  $NH_4^+$  than non-BC treatments and control. Whereas in deep soil layer  $NH_4^+$ increased as compared to the seedling stage with significantly lower  $NH_4^+$  in BC treatments than non-BC treatments as most of the NH<sub>4</sub><sup>+</sup> retained in BC in topsoil. At 12 leaf stage  $NH_4^+$  concentration further decreased and it ranged from 1.2 (mg (g<sup>-1</sup> dry soil) in control to 25.4 (mg ( $g^{-1}$  dry soil) in DM<sub>1</sub>+B at 20 cm depth whereas, in deep soil NH<sub>4</sub><sup>+</sup> ranged from 1.4 (mg (g<sup>-1</sup> dry soil) in control to 20 (mg (g<sup>-1</sup> dry soil) in DM<sub>1</sub>+B. At black

layer stage  $NH_4^+$  dropped to near zero. Maximum  $NH_4^+$  i.e. 1.1 (mg (g<sup>-1</sup> dry soil) was found in  $DM_1+B$  whereas 0 (mg (g<sup>-1</sup> dry soil) in control treatment.

### Nitrate and ammonium during 2017

DM<sub>1</sub> and DM<sub>2</sub> were applied to all respective treatments (DM<sub>1</sub>, DM<sub>2</sub>, DM<sub>1</sub>+B, DM<sub>2</sub>+B) before seeding but the IN fertilizer was applied in two splits (first dose: two days after 6 leaf stage sampling, second dose: one day before 12 leaf stage sampling). Soil NO<sub>3</sub><sup>-</sup> and  $NH_4^+$  concentrations were measured at four crop growth stages (6 leaves, 12 leaves, tasseling and black layer stage). Soil NO<sub>3</sub><sup>-</sup> concentration was low at 6 leaf stage as compared to 12 leaves and tasseling stage. Manure treatments (DM<sub>1</sub>, DM<sub>2</sub>, DM<sub>1</sub>+B,  $DM_2+B$ ) have high  $NO_3^-$  concentration in soil than IN, and IN+B as fertilizer was not applied to any treatment before seeding.  $DM_1+B$ , and  $DM_2+B$  had low  $NO_3^$ concentration at 40 cm soil depth as most of the NO3<sup>-</sup> retained in topsoil in biochar (Figure 3.2a). At 12 leaf stage, NO<sub>3</sub><sup>-</sup> concentration increased in DM+BC treatments and BC helped to reduce  $NO_3^-$  movement to the deep soil (Figure 3.2b). At the tasseling stage, as there was fertilizer (NH<sub>4</sub>NO<sub>3</sub>) application one day before sampling so there was high NO<sub>3</sub> observed in all treatments except control. There was more NO<sub>3</sub> in BC treatments at 20 cm depth and in DM<sub>1</sub>, DM<sub>2</sub>, and IN at 40 cm depth (Figure 3.2c). At black layer stage, soil NO<sub>3</sub><sup>-</sup> deceased in all treatments with significantly high NO<sub>3</sub><sup>-</sup> concentration in BC treatments than DM<sub>1</sub>, DM<sub>2</sub>, IN and control (Figure 3.2d). Soil NH<sub>4</sub><sup>+</sup> concentrations were lowest at both soil depths at 6 leaves stage of the crop. But after that an increase in NH<sub>4</sub><sup>+</sup> was observed in  $DM_1$ ,  $DM_1+B$ ,  $DM_2$ ,  $DM_2+B$  at 12 leaves the stage with the highest  $NH_4^+$  as 41.2 (mg g<sup>-1</sup> dry soil) in DM<sub>1</sub>+B at 20 cm soil layer. At tasseling stage as sampling was done one day after fertilizer application there was an increase in  $\mathrm{NH_4^+}$  concentration and it reached a maximum of 46.3 (mg g<sup>-1</sup> dry soil) in IN+B. At black layer stage the  $NH_4^+$  was significantly higher in  $DM_1+B$ ,  $DM_2+B$ , and IN+B than  $DM_1$ ,  $DM_2$ , IN and control (Figure 3.2c).



Figure 3.1: Soil nitrate (NO<sub>3</sub><sup>-</sup>) and ammonium (NH<sub>4</sub><sup>+</sup>) concentrations (mg g<sup>-1</sup> dry soil) in experimental treatments at four crop growth stages during 2016

(a)  $NO_3^-$  at 20 cm depth (b)  $NO_3^-$  at 40 cm depth (c)  $NH_4^+$  at 20 cm depth (d)  $NH_4^+$  at 40 cm depth, filled circle (seedling emergence), empty circle (6 leaves stage), filled triangle (12 leaves stage), empty triangle (Black layer stage)



Figure 3.2: Soil nitrate (NO<sub>3</sub><sup>-</sup>) and ammonium (NH<sub>4</sub><sup>+</sup>) concentrations (mg g<sup>-1</sup> dry soil) in experimental treatments at four crop growth stages during 2017

(a)  $NO_3^-$  at 20 cm depth (b)  $NO_3^-$  at 40 cm depth (c)  $NH_4^+$  at 20 cm depth (d)  $NH_4^+$  at 40 cm depth, filled circle (seedling emergence), empty circle (6 leaves stage), filled triangle (12 leaves stage), empty triangle (Black layer stage)

#### **3.4.2.** Soil pH

Soil pH in experimental treatments was measured at three crop growth stages in 2016 and at two stages during 2017. There was a significant temporal variation in soil pH in all treatments measured at different crop growth stages in both seasons (Figure 3.3). BC treatments have high soil pH at each sampling event than non-BC treatments. Generally, soil pH increased with DM application; however, a significant increase was noticed with BC amendment and declined as the season proceeded. For example, DM<sub>1</sub>, DM<sub>2</sub>, and IN treatments exhibited pH values of 6.23, 6.17 and 6.02 at the seedling stage during 2016. BC addition to these treatments significantly increased pH to 6.46, 6.34, and 6.21 respectively (Fig. 8). There was a reduction in soil pH in all treatments except IN, IN+B, and the control at 12 leaf stage compared to seedling stage. The mean values of soil pH at 12 leaf stage in all treatments were 6.20, 6.31, 6.15, 5.20, 6.15, 6.23, and 6.0 in DM<sub>1</sub>,  $DM_1+B$ ,  $DM_2$ ,  $DM_2+B$ , IN, IN+B, and the Control. At black layer stage, soil pH further decreased in all treatments with relatively high pH in BC amended treatments. At the end of the growing season of 2016 (black layer stage), the BC amended treatments have high soil pH by 0.18, 0.10, and 0.06 units than non-BC amended treatments. At six-leaf stage during 2017, soil pH was stable, however again decreased at black layer stage in 2017 but BC treatments have relatively high pH. The mean values of soil pH at black layer stage (2017) in all treatments were 5.87, 6.05, 5.50, 5.70, 5.68, 5.73, and 5.50 in DM<sub>1</sub>, DM<sub>1</sub>+B,  $DM_2$ ,  $DM_2$ +B, IN, IN+B, and control treatment with an overall increase by 0.14 unit in BC than non-BC amended treatments.



Figure 3.3: Soil pH at 20 cm depth at different crop growth stages during 2016 and 2017 Solid circle (DM<sub>1</sub>), empty circle (DM<sub>1</sub>+B), solid triangle (DM<sub>2</sub>), empty triangle (DM<sub>2</sub>+B), Solid square (IN), empty square (IN+B), solid diamond (N<sub>0</sub>)

### 3.4.3. Plant tissue N concentration

DM and IN alone and co-application with BC significantly (p < 0.05) influenced the N concentration in silage corn genotypes. BC amended treatments enhanced the N concentration in silage corn tissues, however, maximum N concentration was noted in IN+B treatment, compared to the minimum concentration in the control treatment. Among the genotypes, A4177G3 RIB exhibited higher N concentration compared to DKC26-28 RIB and Yukon R during 2016 (Table 3.2). Overall, maximum mean N concentration was noted in IN+BC treatment compared to control treatment. BC addition to DM<sub>1</sub>, DM<sub>2</sub>, and IN enhanced N concentration by 10, 13 and 27 % respectively. In 2017, A4177G3 RIB exhibited the highest tissue N concentration in IN+BC treatment, whereas lowest was noted in DKC26-28 RIB in control treatment. Overall, BC addition to DM<sub>1</sub>, DM<sub>2</sub>, and IN enhanced N concentration by 17, 8 and 7.6 % respectively. N concentration was maximum in DM<sub>1</sub>+BC treatment compared to control treatment whereas, A4177G3 RIB genotype was the most efficient in N uptake followed by Yukon R and DKC26-28 RIB.

Treatments	A4177G3 RIB	DKC26-28 RIB	Yukon R	Mean
DM1	$1.25 \pm 0.15^{abcd}$	$1.03 \pm 0.03^{defg}$	$1.07\pm0.17^{cdefg}$	$1.11 \pm 0.07^{BC}$
DM1+B	1.28±0.01 <sup>abc</sup>	$1.07\pm0.13^{cdefg}$	$1.32 \pm 0.05^{ab}$	$1.22 \pm 0.05^{AB}$
DM2	$1.09 \pm 0.06^{bcdefg}$	$1.17\pm0.12^{bcdef}$	$0.99 \pm 0.12^{efg}$	$1.08 \pm 0.06^{BC}$
DM2+B	$1.45 \pm 0.05^{a}$	$1.08 \pm 0.04^{bcdefg}$	$1.14{\pm}0.05b^{cdefg}$	$1.23 \pm 0.06^{AB}$
IN	1.18±0.03 <sup>bcde</sup>	$0.93{\pm}0.06^{fg}$	$0.96{\pm}0.04^{efg}$	$1.02{\pm}0.03^{\circ}$
IN+B	1.47±0.02 <sup>a</sup>	$1.16\pm0.11^{bcdefg}$	$1.28 \pm 0.04^{abc}$	1.30±0.05 <sup>A</sup>
N0	$1.11\pm0.01^{bcdefg}$	$0.93{\pm}0.02^{fg}$	0.92±0.09 <sup>g</sup>	$0.99 \pm 0.05^{\circ}$
	1.26±0.03 <sup>A</sup>	1.05±0.03 <sup>B</sup>	1.10±0.04 <sup>B</sup>	

Table 3.2: Plant tissues nitrogen concentration (%) in silage corn genotypes under different experimental treatments during growing season 2016

Means sharing common letters are not significantly different at 0.05 probability level. Table 3.3: Plant tissues nitrogen concentration (%) in silage corn genotypes under different experimental treatments during growing season 2017

Treatments	A4177G3 RIB	DKC26-28 RIB	Yukon R	Mean
DM1	$1.40\pm0.03^{abcd}$	$1.08 \pm 0.00^{fg}$	$1.24 \pm 0.05^{def}$	$1.24 \pm 0.05^{\circ}$
DM1+B	$1.41 \pm 0.01^{abcd}$	$1.45 \pm 0.03^{abc}$	1.50±0.06 <sup>ab</sup>	1.45±0.02 <sup>A</sup>
DM2	$1.40 \pm 0.03^{abcd}$	1.10±0.06 <sup>fg</sup>	$1.22{\pm}0.04^{defg}$	$1.24{\pm}0.05^{C}$
DM2+B	1.45±0.08 <sup>abc</sup>	$1.26 \pm 0.07^{cdef}$	$1.31 \pm 0.02b^{bcde}$	$1.34{\pm}0.04^{ABC}$
IN	$1.47 \pm 0.03^{ab}$	1.10±0.16 <sup>fg</sup>	1.32±0.01 <sup>bcde</sup>	$1.30{\pm}0.07^{BC}$
IN+B	$1.54{\pm}0.07^{a}$	$1.31 \pm 0.03^{bcde}$	1.33±0.01 <sup>bcde</sup>	$1.39{\pm}0.04^{AB}$
N0	1.14±0.03 <sup>efg</sup>	1.03±0.16 <sup>g</sup>	1.09±0.13 <sup>fg</sup>	1.08±0.06 <sup>D</sup>
	1.40±0.03 <sup>A</sup>	1.19±0.04 <sup>C</sup>	1.29±0.03 <sup>B</sup>	

Means sharing common letters are not significantly different at 0.05 probability level.

## 3.4.4. Dry matter yield

The application of DM<sub>1</sub>, DM<sub>2</sub> and IN alone or in combination with BC had significantly (p < 0.05) affected dry matter yield (DMY) of silage corn. BC amended treatments  $(DM_1+B, DM_2+B, and IN+B)$  yielded significantly (p<0.05) higher biomass than without BC and the control treatment. Genotype  $\times$  N source interaction (p < 0.05) had significantly influenced dry matter yield during both years. Yukon R produced the maximum dry matter yield of 21 Mg ha<sup>-1</sup> in DM<sub>1</sub>+BC treatment, compared to minimum dry matter production (14 Mg ha<sup>-1</sup>) by A4177G3 RIB in the control treatment (Table 3.4). N sources had a significant effect on dry matter production in all genotypes. BC addition to IN, DM<sub>1</sub>, and DM<sub>2</sub> increased dry matter yield by 5, 3, and 4.5 % in A4177G3 RIB, 3.6, 7.5 and 8 % in DKC26-28 RIB, 6, 4, and 8 % in Yukon R, respectively. Overall, BC application to IN,  $DM_1$  and  $DM_2$  increased the dry matter production by 5, 5, and 7 % respectively during 2016. Similarly, in 2017 growing season, the highest dry matter yield of 17.1 Mg ha<sup>-1</sup> was observed in Yukon R while minimum 10.1 Mg ha<sup>-1</sup> in A4177G3 RIB in the control treatment (Table 3.5). BC addition to IN, DM<sub>1</sub>, and DM<sub>2</sub> increased dry matter yield by 6.8, 3, and 7 % in A4177G3 RIB and 7.6, 9 and 12.5 % in DKC26-28 RIB and 6.8, 5, and 11.5 % in Yukon R, respectively. Overall BC application to IN, DM<sub>1</sub> and DM<sub>2</sub> increased the dry matter production by 7, 6, and 10.5 %, respectively.

Treatment/Genotype	A4177G3 RIB	DKC26-28 RIB	Yukon R	Mean
DM <sub>1</sub>	$18.0\pm0.14^{ij}$	19.3±0.28 <sup>de</sup>	19.7±0.17 <sup>b</sup>	19.0±0.28 <sup>C</sup>
DM <sub>1</sub> +B	19.0±0.12 <sup>ef</sup>	$20.0 \pm 0.12b^{c}$	21.0±0.12 <sup>a</sup>	$20.0{\pm}0.29^{A}$
$DM_2$	18.0±0.18 <sup>ij</sup>	$18.3 \pm 0.11^{hi}$	$19.5 \pm 0.24^{d}$	18.6±0.25 <sup>D</sup>
DM <sub>2</sub> +B	18.6±0.15 <sup>fgh</sup>	$19.8 \pm 0.17^{cd}$	$20.4{\pm}0.17^{b}$	19.6±0.28 <sup>B</sup>
IN	17.6±0.18 <sup>j</sup>	17.9±0.16 <sup>ij</sup>	18.8±0.13 <sup>ef</sup>	$18.1 \pm 0.19^{E}$
IN+B	18.4±0.29 <sup>ghi</sup>	19.5±0.16 <sup>cde</sup>	20.5±0.17 <sup>c</sup>	$19.5 \pm 0.32^{B}$
N0	$14.0\pm0.15^{1}$	$14.3 \pm 0.15^{1}$	$15.3 \pm 0.15^{i}$	14.5±0.21 <sup>F</sup>
	17.6±0.35 <sup>C</sup>	18.4±0.41 <sup>B</sup>	19.3±0.40 <sup>A</sup>	

Table 3.4: Dry matter yield (Mg ha<sup>-1</sup>) of three silage corn genotypes during growing season 2016

Means sharing common letters are not significantly different at 0.05 probability level

Table 3.5: Dry matter yield (Mg ha<sup>-1</sup>) of three silage corn genotypes during growing season 2017

Treatment/Genotype	A4177G3 RIB	DKC26-28 RIB	Yukon R	Mean
DM1	14.1±0.13 <sup>hijk</sup>	14.9±0.19 <sup>fg</sup>	$15.9 \pm 0.25^{bcd}$	$15.0\pm0.22^{C}$
DM1+B	15.1±0.23 <sup>ef</sup>	$16.1 \pm 0.09^{bc}$	17.1±0.10 <sup>a</sup>	16.1±0.25 <sup>A</sup>
DM2	$14.0{\pm}0.08i^{jk}$	$14.1 \pm 0.16^{hij}$	15.6±0.21 <sup>cde</sup>	14.6±0.23 <sup>D</sup>
DM2+B	14.5±0.06 <sup>ghi</sup>	$15.6 \pm 0.04^{de}$	$16.4 \pm 0.04^{b}$	15.5±0.24 <sup>B</sup>
IN	13.6±0.18 <sup>k</sup>	$13.7 \pm 0.39^{jk}$	$14.5{\pm}0.40^{\text{gh}}$	$13.9 \pm 0.18^{E}$
IN+B	14.6±0.11 <sup>fg</sup>	15.7±0.11 <sup>cd</sup>	$16.4\pm\!0.13^{b}$	$15.6 \pm 0.24^{B}$
N0	10.1±0.15 <sup>m</sup>	10.3±0.03 <sup>m</sup>	$11.2\pm0.15^{1}$	$10.5 \pm 0.14^{F}$
	$13.7 \pm 0.35^{\circ}$	14.3±0.41 <sup>B</sup>	15.3±0.42 <sup>A</sup>	

Means sharing common letters are not significantly different at 0.05 probability level
#### 3.5. Discussion

Soil NO<sub>3</sub><sup>-</sup> retention after BC addition had been reported in various studies. In this study, NO<sub>3</sub><sup>-</sup> and NH<sub>4</sub><sup>+</sup> concentration increased in BC amended treatments in the top 20 cm soil layer where BC prevented the downward movement of  $NO_3^-$  and  $NH_4^+$  to a deep soil layer. Manure and inorganic nitrogen fertilizer application increased NO<sub>3</sub><sup>-</sup> and NH<sub>4</sub><sup>+</sup> in all manure and BC amended treatments compared to the control. For example, increased soil NO<sub>3</sub> was observed after swine slurry application (Bertora et al., 2008). BC amended treatments further increased NO<sub>3</sub><sup>-</sup> and NH<sub>4</sub><sup>+</sup> concentration than manure and IN treatments alone which could be attributed to the adsorption of these ions on BC surfaces which decreased their downward movement. Acid functional groups present on BC surface i.e. carboxylic, hydroxyl, lactone, lactol, phenol and carbonyls which have a negative charge and attract NH<sub>4</sub><sup>+</sup> ions (Amonette and Joseph, 2009; Brennan et al., 2001; Montes-Morán et al., 2004; Zheng et al., 2010). Other functional groups i.e. chromenes, ketones and pyrones etc. also exist on BC surface which facilitate NO<sub>3</sub><sup>-</sup> adsorption to its surface (Amonette and Joseph, 2009; Montes-Morán et al., 2004). Another possible mechanism could be unconventional H-bonding between NO<sub>3</sub><sup>-</sup> and BC surface, which might increase the NO<sub>3</sub><sup>-</sup> adsorption on BC surface (Kammann et al., 2015; Lawrinenko, 2014; Mukherjee et al., 2011). Over the time, the adsorbed N could be desorbed and become available in soil (Kameyama et al., 2012; Taghizadeh-Toosi et al., 2012), depending on BC adsorption capacity, amount of BC applied, soil cation and anion exchange capacity, soil microbial community and crop N demand (Clough et al., 2013). Application of BC to soil increased soil N mineralization, enhanced nitrification ( $NH_4^+ \rightarrow NO_2^- \rightarrow NO_3^-$ ) by 34 %, suppressed denitrification (NO<sub>3</sub><sup>-</sup>  $\rightarrow$  NO<sub>2</sub><sup>-</sup>  $\rightarrow$  NO  $\rightarrow$  N<sub>2</sub>O  $\rightarrow$  N<sub>2</sub>) by 37 %, and reduced cumulative N<sub>2</sub>O emission by 91 % in a sandy loam soil (Case et al., 2015). BC induced reduction in denitrification can be explained by changes in soil aeration brought by low soil bulk density and high WHC which lower the denitrifier activity (Karhu et al. 2011; Basso et al. 2013). Most of the biochars decrease soil acidity due to their high pH buffering capacity and alkaline nature (pH > 7) at least 1.5 units higher than acid soils (pH < 5.5). BC used in this study was alkaline (pH = 9), and at each sampling point, BC amended treatments exhibited high soil pH compared to non-BC amended treatments and the control. The carbonates and oxides formed during pyrolysis from the cations (Ca, Mg, K, Na etc.) in feedstock react with H<sup>+</sup> and monomeric aluminum species in acidic soils and increased soil pH (Brewer et al., 2012; Enders et al., 2012; Novak et al., 2009). In addition to the carbonates and oxides,  $-COO^-$  (-COOH) and  $-O^-$  (-OH) also play important role in BC alkalinity (Yuan et al., 2011).

BC increased NH<sub>4</sub><sup>+</sup> retention in the soil and improved N uptake (Sun et al., 2017), enhanced lettuce yield and nutrient concentrations in plant tissues (Upadhyay et al., 2014; Woldetsadik et al., 2017), increased soil pH, CEC, Ca, total C, N uptake and biomass production of wheat up to 250 % (Van Zwieten et al., 2010a). It diminished the nutrient losses due to winter freeze-thaw cycle and increased N uptake in the subsequent crop (Zhou et al., 2017). Gunes et al. (2014) reported that BC application increased uptake of N, P, K and reduced Fe, Cu, Zn, and Mn in lettuce grown in alkaline soils. DM application to soil decreased the soil pH while BC amendment increased soil pH along with increasing *Lolium perenne* biomass by 29 % in a growth experiment (Schimmelpfennig et al., 2014). There was significantly low dry matter recorded during 2017 because 30 % less rain was received in this season especially during the crop active growth period (July-August). However, BC increased dry matter yield during both years. Maximum DMY was observed in Yukon R while maximum N concentration was observed in A4177G3 RIB. The N concentration of corn hybrids is influenced by their genetic characters and environment (Gautam et al., 2011). The increase in plant N concentration and DMY could be attributed to improvement in soil fertility, soil physical properties and nutrient retention after BC application. Contrasting effects of BC application on N uptake and plant yield had been reported in the literature depending upon BC feedstock, pyrolysis conditions, soil type etc. (Agegnehu et al., 2016; Borchard et al., 2014; Tammeorg et al., 2014; Upadhyay et al., 2014; Vitkova et al., 2017; Wang et al., 2016; Woldetsadik et al., 2017).

#### 3.6. Conclusion

Application of BC to  $DM_1$ ,  $DM_2$ , and IN increased the soil  $NO_3^-$ ,  $NH_4^+$  retention, soil pH, plant N concentration, and DMY. BC application to these treatments decreased leaching which improved plant N concentration and DMY of silage corn crop. BC application also decreased soil acidity. At each sampling interval, high soil pH was observed in BC treatments than non-BC treatments. On an average, BC addition to  $DM_1$ ,  $DM_2$ , and IN enhanced N concentration by 13.5, 11.5 and 17.3 %, respectively. Overall BC application to IN,  $DM_1$  and  $DM_2$  increased the dry matter production by 6, 5.5, and 8.75 %, respectively.

- Agegnehu, G., Bass, A.M., Nelson, P.N., Bird, M.I., 2016. Benefits of biochar, compost and biochar-compost for soil quality, maize yield and greenhouse gas emissions in a tropical agricultural soil. Sci. Total Environ. 543, 295–306.
- Amendola, C., Montagnoli, A., Terzaghi, M., Trupiano, D., Oliva, F., Baronti, S.,
  Miglietta, F., Chiatante, D., Scippa, G.S., 2017. Short-term effects of biochar on
  grapevine fine root dynamics and arbuscular mycorrhizae production. Agric.
  Ecosyst. Environ. 239, 236–245.
- Amonette, J.E., Joseph, S., 2009. Characteristics of biochar: microchemical properties., in: Biochar for Environmental Management; Science and Technology. p. 33.
- Asai, H., Samson, B.K., Stephan, H.M., Songyikhangsuthor, K., Homma, K., Kiyono, Y., Inoue, Y., Shiraiwa, T., Horie, T., 2009. Biochar amendment techniques for upland rice production in Northern Laos. 1. Soil physical properties, leaf SPAD and grain yield. F. Crop. Res. 111, 81–84.
- Badewa, E.A., 2017. Apparent Electrical Conductivity Mapping in Managed Podzols Using Multicoil and Multifrequency EMI Sensor Measurments. Memorial University of Newfoundland.
- Baronti, S., Vaccari, F.P., Miglietta, F., Calzolari, C., Lugato, E., Orlandini, S., Pini, R., Zulian, C., Genesio, L., 2014. Impact of biochar application on plant water relations in Vitis vinifera (L.). Eur. J. Agron. 53, 38–44.

- Basso, A.S., Miguez, F.E., Laird, D.A., Horton, R., Westgate, M., 2013. Assessing potential of biochar for increasing water-holding capacity of sandy soils. GCB Bioenergy 5, 132–143.
- Basso, B., Ritchie, J.T., 2005. Impact of compost, manure and inorganic fertilizer on nitrate leaching and yield for a 6-year maize-alfalfa rotation in Michigan. Agric. Ecosyst. Environ. 108, 329–341.
- Bertora, C., Alluvione, F., Zavattaro, L., van Groenigen, J.W., Velthof, G., Grignani, C.,
  2008. Pig slurry treatment modifies slurry composition, N2O, and CO2 emissions
  after soil incorporation. Soil Biol. Biochem. 40, 1999–2006.
- Borchard, N., Siemens, J., Ladd, B., Möller, A., Amelung, W., 2014. Application of biochars to sandy and silty soil failed to increase maize yield under common agricultural practice. Soil Tillage Res. 144, 184–194.
- Brennan, J.K., Bandosz, T.J., Thomson, K.T., Gubbins, K.E., 2001. Water in porous carbons. Colloids Surfaces A Physicochem. Eng. Asp. 187–188, 539–568.
- Brewer, C.E., Hu, Y.-Y., Schmidt-Rohr, K., Loynachan, T.E., Laird, D.A., Brown, R.C., 2012. Extent of Pyrolysis Impacts on Fast Pyrolysis Biochar Properties. J. Environ. Qual. 41, 1115–1122.
- Busch, D., Kammann, C., Grünhage, L., Müller, C., 2012. Simple Biotoxicity Tests for Evaluation of Carbonaceous Soil Additives: Establishment and Reproducibility of Four Test Procedures. J. Environ. Qual. 41, 1023–1032.

- Buss, W., Kammann, C., Koyro, H.-W., 2012. Biochar reduces copper toxicity in Chenopodium quinoa Willd. in a sandy soil. J. Environ. Qual. 41, 1157–1165.
- Busscher, W.J., Novak, J.M., Evans, D.E., Watts, D.W., Niandou, M.A.S., Ahmedna, M.,
  2010. Influence of pecan biochar on physical properties of a Norfolk loamy sand.
  Soil Sci. 175, 10–14.
- Cao, T., Meng, J., Liang, H., Yang, X., Chen, W., 2017. Can biochar provide ammonium and nitrate to poor soils ?. Soil column incubation. J. Soil Sci. Plant Nutr. 17, 253– 265.
- Carter, M.R., Gregorich, E.G., Carter, Gregorich, Carter, M.R., Gregorich, E.G., 2008. Soil Sampling and Methods of Analysis, 2nd ed. Taylor & Francis.
- Case, S.D.C., McNamara, N.P., Reay, D.S., Whitaker, J., 2012. The effect of biochar addition on N2O and CO2 emissions from a sandy loam soil - The role of soil aeration. Soil Biol. Biochem. 51, 125–134.
- Case, S.D.D.C., McNamara, N.P., Reay, D.S., Stott, A.W., Grant, H.K., Whitaker, J.,
  2015. Biochar suppresses N2O emissions while maintaining N availability in a sandy loam soil. Soil Biol. Biochem. 81, 178–185.
- Cayuela, M.L., Sanchez-Monedero, M.A., Roig, A., Hanley, K., Enders, A., Lehmann, J.,
  2013. Biochar and denitrification in soils: when, how much and why does biochar
  reduce N2O emissions? Sci. Rep. 3, 1732.
- Clough, T.J., Condron, L.M., Kammann, C., Müller, C., 2013. A Review of Biochar and

Soil Nitrogen Dynamics. Agronomy 3, 275–293.

- Drinkwater, L.E., Wagoner, P., Sarrantonio, M., 1998. Legume-based cropping systems have reuced carbon and nitrogen losses. Nature 396, 262–265.
- Enders, A., Hanley, K., Whitman, T., Joseph, S., Lehmann, J., 2012. Characterization of biochars to evaluate recalcitrance and agronomic performance. Bioresour. Technol. 114, 644–653.
- Fan, J., Xiao, J., Liu, D., Ye, G., Luo, J., Houlbrooke, D., Laurenson, S., Yan, J., Chen,
  L., Tian, J., Ding, W., 2017. Effect of application of dairy manure, effluent and
  inorganic fertilizer on nitrogen leaching in clayey fluvo-aquic soil: A lysimeter
  study. Sci. Total Environ. 592, 206–214.
- Forge, T., Kenney, E., Hashimoto, N., Neilsen, D., Zebarth, B., 2016. Compost and poultry manure as preplant soil amendments for red raspberry: Comparative effects on root lesion nematodes, soil quality and risk of nitrate leaching. Agric. Ecosyst. Environ. 223, 48–58.
- Foster, E.J., Hansen, N., Wallenstein, M., Cotrufo, M.F., 2016. Biochar and manure amendments impact soil nutrients and microbial enzymatic activities in a semi-arid irrigated maize cropping system. Agric. Ecosyst. Environ. 233, 404–414.
- Francis, D.D., 1992. Control mechanisms to reduce fertilizer nitrogen movement into groundwater. J. Soil Water Conserv. 47, 444–448.
- Gaines, T.P., Gaines, S.T., 1994. Soil texture effect on nitrate leaching in soil percolates.

Commun. Soil Sci. Plant Anal. 25, 2561–2570.

- Gatti, M., Squeri, C., Garavani, A., Vercesi, A., Dosso, P., Diti, I., Poni, S., 2018. Effects of Variable Rate Nitrogen Application on cv. Barbera Performance: I. Vegetative Growth and Leaf Nutritional Status. Am. J. Enol. Vitic. ajev.2018.
- Gautam, P., Gustafson, D., Wicks III, Z., 2011. Study on nitrogen concentration of corn hybrids. Am. J. Agric. Biol. Sci. 6, 148–154.
- Griffin, D.E., Wang, D., Parikh, S.J., Scow, K.M., 2017. Short-lived effects of walnut shell biochar on soils and crop yields in a long-term field experiment. Agric. Ecosyst. Environ. 236, 21–29.
- Güereña, D., Lehmann, J., Hanley, K., Enders, A., Hyland, C., Riha, S., 2013. Nitrogen dynamics following field application of biochar in a temperate North American maize-based production system. Plant Soil 365, 239–254.
- Gul, S., Whalen, J.K., Thomas, B.W., Sachdeva, V., Deng, H., 2015. Physico-chemical properties and microbial responses in biochar-amended soils: Mechanisms and future directions. Agric. Ecosyst. Environ. 206, 46–59.
- Gunes, A., Inal, A., Taskin, M.B., Sahin, O., Kaya, E.C., Atakol, A., 2014. Effect of phosphorus-enriched biochar and poultry manure on growth and mineral composition of lettuce (Lactuca sativa L. cv.) grown in alkaline soil. Soil Use Manag. 30, 182–188.
- Hagemann, N., Kammann, C.I., Schmidt, H.P., Kappler, A., Behrens, S., 2017. Nitrate

capture and slow release in biochar amended compost and soil. PLoS One 12, e0171214.

- Haider, G., Koyro, H.W., Azam, F., Steffens, D., Müller, C., Kammann, C., 2015.Biochar but not humic acid product amendment affected maize yields via improving plant-soil moisture relations. Plant Soil 395, 141–157.
- Haider, G., Steffens, D., Moser, G., Müller, C., Claudia, I.K., 2017. Biochar reduced nitrate leaching and improved soil moisture content without yield improvements in a four-year field study. Agric. Ecosyst. Environ. 237, 80–94.
- Hargrave, A.P., Shaykewich, C.F., 1997. Rainfall induced nitrogen and phosphorus losses from Manitoba soils. Can. J. Soil Sci. 77, 59–65.
- Harter, J., Krause, H.-M., Schuettler, S., Ruser, R., Fromme, M., Scholten, T., Kappler,A., Behrens, S., 2014. Linking N2O emissions from biochar-amended soil to the structure and function of the N-cycling microbial community. ISME J. 8, 660–674.
- Heman, L., Lihua, C., Changchang, X., Hong, Y. and, Baoguo, L., 2016. Distribution characteristics of soil organic carbon and nitrogen in farmland and adjacent natural grassland in Tibet. Int. J. Agric. Biol. Eng. 9, 135–145.
- Hernandez-Ramirez, G., Brouder, S.M., Ruark, M.D., Turco, R.F., 2011. Nitrate, Phosphate, and Ammonium Loads at Subsurface Drains: Agroecosystems and Nitrogen Management. J. Environ. Qual. 40, 1229–1240.
- Hirel, B., Le Gouis, J., Ney, B., Gallais, A., 2007. The challenge of improving nitrogen

use efficiency in crop plants: Towards a more central role for genetic variability and quantitative genetics within integrated approaches. J. Exp. Bot. 58, 2369–2387.

- Huang, M., Zhou, X., Chen, J., Cao, F., Jiang, L., Zou, Y., 2017. Interaction of Changes in pH and Urease Activity Induced by Biochar Addition Affects Ammonia Volatilization on an Acid Paddy Soil Following Application of Urea. Commun. Soil Sci. Plant Anal. 48, 107–112.
- Hussain, M., Farooq, M., Nawaz, A., Al-Sadi, A.M., Solaiman, Z.M., Alghamdi, S.S.,Ammara, U., Ok, Y.S., Siddique, K.H.M., 2017. Biochar for crop production:potential benefits and risks. J. Soils Sediments 17, 685–716.
- Ibrikci, H., Cetin, M., Karnez, E., Flügel, W.A., Tilkici, B., Bulbul, Y., Ryan, J., 2015. Irrigation-induced nitrate losses assessed in a Mediterranean irrigation district. Agric. Water Manag. 148, 223–231.
- Ippolito, J.A., Laird, D.A., Busscher, W.J., 2012. Environmental benefits of biochar. J. Environ. Qual. 41, 967–972.
- Jabloun, M., Schelde, K., Tao, F., Olesen, J.E., 2015. Effect of temperature and precipitation on nitrate leaching from organic cereal cropping systems in Denmark. Eur. J. Agron. 62, 55–64.
- Jean, M., Es, V., Harold, M., Charissa, L., Larry, D., Fred, R., 2000. Nitrate leaching and nitrogen budget as affected by maize nitrogen rate and soil type. J. Environ. Qual. 29, 1813–1820.

- Jervin, P., Menchie, V.A., Rosario, R., 2017. Soil Properties and Response of Spring Onion to Different Levels of Biochar. J. Agric. Technol. 13, 131–137.
- Jokela, W.E., Bosworth, S.C., Rankin, J.J., 2014. Sidedressed dairy manure effects on corn yield and residual soil nitrate. Soil Sci. 179, 37–41.
- Joseph, S., Graber, E., Chia, C., Munroe, P., Donne, S., Thomas, T., Nielsen, S., Marjo, C., Rutlidge, H., Pan, G., Li, L., Taylor, P., Rawal, A., Hook, J., 2013. Shifting paradigms : development of high-efficiency biochar fertilizers based on nanostructures and soluble components. Carbon Manag. 4, 323–343.
- Ju, C., Buresh, R.J., Wang, Z., Zhang, H., Liu, L., Yang, J., Zhang, J., 2015. Root and shoot traits for rice varieties with higher grain yield and higher nitrogen use efficiency at lower nitrogen rates application. F. Crop. Res. 175, 47–55.
- Kameyama, K., Miyamoto, T., Shiono, T., Shinogi, Y., 2012. Influence of SugarcaneBagasse-derived Biochar Application on Nitrate Leaching in Calcaric Dark Red Soil.J. Environ. Qual. 41, 1131–1137.
- Kammann, C., Ratering, S., Eckhard, C., Muller, C., 2012. Biochar and Hydrochar Effects on Greenhouse Gas (Carbon Dioxide, Nitrous Oxide, and Methane) Fluxes from Soils. J. Environ. Qual. 41, 1052–1066.
- Kammann, C.I., Schmidt, H.-P., Messerschmidt, N., Linsel, S., Steffens, D., Müller, C., Koyro, H.-W., Conte, P., Stephen, J., 2015. Plant growth improvement mediated by nitrate capture in co-composted biochar. Sci. Rep. 5, 11080.

- Karhu, K., Mattila, T., Bergström, I., Regina, K., 2011. Biochar addition to agricultural soil increased CH4uptake and water holding capacity - Results from a short-term pilot field study. Agric. Ecosyst. Environ. 140, 309–313.
- Kirby, G.E., 1988. Soils of the Pasadena-Deer Lake area, Newfoundland. St. John's, Newfoundland.
- Knowles, O.A., Robinson, B.H., Contangelo, A., Clucas, L., 2011. Biochar for the mitigation of nitrate leaching from soil amended with biosolids. Sci. Total Environ. 409, 3206–3210.
- Laird, D.A., Fleming, P., Davis, D.D., Horton, R., Wang, B., Karlen, D.L., 2010. Impact of biochar amendments on the quality of a typical Midwestern agricultural soil. Geoderma 158, 443–449.
- Lawrinenko, M., 2014. Anion exchange capacity of biochar. Green Chem. Lowa State University, Lowa, USA.

Lehmann, J., 2007. Bio-energy in the black. Front. Ecol. Environ. 5, 381–387.

- Lentz, R.D., Ippolito, J. a., 2012. Biochar and Manure Affect Calcareous Soil and Corn Silage Nutrient Concentrations and Uptake. J. Environ. Qual. 41, 1033–1043.
- Lentz, R.D., Ippolito, J.A., Spokas, K.A., 2014. Biochar and Manure Effects on Net Nitrogen Mineralization and Greenhouse Gas Emissions from Calcareous Soil under Corn. Soil Sci. Soc. Am. J. 78, 1641–1655.
- Lin, Y., Munroe, P., Joseph, S., Ziolkowski, A., Zwieten, L. Van, Kimber, S., Rust, J.,

2013. Chemosphere Chemical and structural analysis of enhanced biochars : Thermally treated mixtures of biochar , chicken litter , clay and minerals. Chemosphere 91, 35–40.

- Liu, X. yu, Qu, J. jing, Li, L. qing, Zhang, A. feng, Jufeng, Z., Zheng, J. wei, Pan, G. xing, 2012. Can biochar amendment be an ecological engineering technology to depress N 2O emission in rice paddies?-A cross site field experiment from South China. Ecol. Eng. 42, 168–173.
- Long, G.Q., Sun, B., 2012. Nitrogen leaching under corn cultivation stabilized after four years application of pig manure to red soil in subtropical China. Agric. Ecosyst.
   Environ. 146, 73–80.
- Lu, H., Lashari, M.S., Liu, X., Ji, H., Li, L., Zheng, J., Kibue, G.W., Joseph, S., Pan, G., 2015. Changes in soil microbial community structure and enzyme activity with amendment of biochar-manure compost and pyroligneous solution in a saline soil from Central China. Eur. J. Soil Biol. 70, 67–76.
- Martínez, E., Domingo, F., Roselló, A., Serra, J., Boixadera, J., Lloveras, J., 2017. The effects of dairy cattle manure and mineral N fertilizer on irrigated maize and soil N and organic C. Eur. J. Agron. 83, 78–85.
- Masaka, J., Nyamangara, J., Wuta, M., 2015. Effect of Inorganic and Organic Fertilizer Application on Nitrate Leaching in Wetland Soil Under Field Tomato (Lycopersicon esculentum) and Leaf Rape (Brassica napus). Agric. Res. 4, 63–75.

- Montes-Morán, M.A., Suárez, D., Menéndez, J.A., Fuente, E., 2004. On the nature of basic sites on carbon surfaces: An overview. Carbon N. Y. 42, 1219–1225.
- Mukherjee, A., Zimmerman, A.R., Harris, W., 2011. Surface chemistry variations among a series of laboratory-produced biochars. Geoderma 163, 247–255.
- Novak, J.M., Busscher, W.J., Laird, D.L., Ahmedna, M., Watts, D.W., Niandou, M.A.S., 2009. Impact of Biochar Amendment on Fertility of a Southeastern Coastal Plain Soil. Soil Sci. 174, 105–112.
- Parker, D., Venhaus, D., Robinson, C., Marek, T., Sweeten, J., 2018. Corn Yield and Soil Fertility with Combined Use of Raw or Composted Beef Manure and Inorganic Fertilizers on the Texas Northern High Plains. Compost Sci. Util. 1–12.
- Prost, K., Borchard, N., Siemens, J., Kautz, T., Séquaris, J.-M., Möller, A., Amelung, W.,
  2013. Biochar Affected by Composting with Farmyard Manure. J. Environ. Qual.
  42, 164–172.
- SAMCO Agricultural Manufacturing Ltd, n.d. Samco 2200.
- Schimmelpfennig, S., Müller, C., Grünhage, L., Koch, C., Kammann, C., 2014. Biochar, hydrochar and uncarbonized feedstock application to permanent grassland-Effects on greenhouse gas emissions and plant growth. Agric. Ecosyst. Environ. 191, 39–52.
- Schmidt, H.P., Kammann, C., Niggli, C., Evangelou, M.W.H., Mackie, K.A., Abiven, S.,2014. Biochar and biochar-compost as soil amendments to a vineyard soil:Influences on plant growth, nutrient uptake, plant health and grape quality. Agric.

Ecosyst. Environ. 191, 117–123.

- Sika, M.P., Hardie, A.G., 2014. Effect of pine wood biochar on ammonium nitrate leaching and availability in a South African sandy soil. Eur. J. Soil Sci. 65, 113–119.
- Smider, B., Singh, B., 2014. Agronomic performance of a high ash biochar in two contrasting soils. Agric. Ecosyst. Environ. 191, 99–107.
- Spokas, K.A., Reicosky, D.C., 2009. Impacts of sixteen different biochars on soil greenhouse gas production. Ann. Environ. Sci. 3, 179–193.
- Sun, H., Lu, H., Chu, L., Shao, H., Shi, W., 2017. Biochar applied with appropriate rates can reduce N leaching, keep N retention and not increase NH3 volatilization in a coastal saline soil. Sci. Total Environ. 575, 820–825.
- Sun, Z., Sänger, A., Rebensburg, P., Lentzsch, P., Wirth, S., Kaupenjohann, M., Meyer-Aurich, A., 2017. Contrasting effects of biochar on N2O emission and N uptake at different N fertilizer levels on a temperate sandy loam. Sci. Total Environ. 578, 557– 565.
- Taghizadeh-Toosi, A., Clough, T.J., Sherlock, R.R., Condron, L.M., 2012. Biochar adsorbed ammonia is bioavailable. Plant Soil 350, 57–69.
- Tammeorg, P., Simojoki, A., Mäkelä, P., Stoddard, F.L., Alakukku, L., Helenius, J.,
  2014. Biochar application to a fertile sandy clay loam in boreal conditions: Effects on soil properties and yield formation of wheat, turnip rape and faba bean. Plant Soil 374, 89–107.

- Upadhyay, K.P., George, D., Swift, R.S., Galea, V., 2014. The influence of biochar on growth of lettuce and potato. J. Integr. Agric. 13, 541–546.
- Uzoma, K.C., Inoue, M., Andry, H., Zahoor, A., Nishihara, E., 2011. Influence of biochar application on sandy soil hydraulic properties and nutrient retention. J. Food, Agric. Environ. 9, 1137–1143.
- van Es, H.M., Sogbedji, J.M., Schindelbeck, R.R., 2006. Effect of Manure Application Timing, Crop, and Soil Type on Nitrate Leaching. J. Environ. Qual. 35, 670–679.
- Van Zwieten, L., Kimber, S., Morris, S., Chan, K.Y., Downie, A., Rust, J., Joseph, S., Cowie, A., 2010. Effects of biochar from slow pyrolysis of papermill waste on agronomic performance and soil fertility. Plant Soil 327, 235–246.
- Ventura, M., Sorrenti, G., Panzacchi, P., George, E., Tonon, G., 2012. Biochar reduces short-term nitrate leaching from a horizon in an apple orchard. J. Environ. Qual. 42, 76–82.
- Vitkova, J., Kondrlova, E., Rodny, M., Surda, P., Horak, J., 2017. Analysis of soil water content and crop yield after biochar application in field conditions. platn soil Environ. 63, 569–573.
- Wang, L., Wu, J., Wang, Z., Li, W., Zhang, K., 2017. Effects of Three Biochars as Adsorbents on Soils Adsorbing Ammonium Nitrogen in Biogas Slurry. J. Chem. 2017.
- Wang, X., Ren, Y., Zhang, S., Chen, Y., Wang, N., 2017. Applications of organic manure

increased maize (Zea mays L.) yield and water productivity in a semi-arid region. Agric. Water Manag. 187, 88–98.

- Wang, Y., Zhang, L., Yang, H., Yan, G., Xu, Z., Chen, C., Zhang, D., 2016. Biochar nutrient availability rather than its water holding capacity governs the growth of both C3 and C4 plants. J. Soils Sediments 16, 801–810.
- Woldetsadik, D., Drechsel, P., Marschner, B., Itanna, F., Gebrekidan, H., 2017. Effect of biochar derived from faecal matter on yield and nutrient content of lettuce (Lactuca sativa) in two contrasting soils. Environ. Syst. Res. 6.
- Xu, C.Y., Hosseini-Bai, S., Hao, Y., Rachaputi, R.C.N., Wang, H., Xu, Z., Wallace, H.,
  2015. Effect of biochar amendment on yield and photosynthesis of peanut on two
  types of soils. Environ. Sci. Pollut. Res. 22, 6112–6125.
- Ye, Y., Liang, X., Chen, Y., Liu, J., Gu, J., Guo, R., Li, L., 2013. Alternate wetting and drying irrigation and controlled-release nitrogen fertilizer in late-season rice. Effects on dry matter accumulation, yield, water and nitrogen use. F. Crop. Res. 144, 212– 224.
- Yuan, J.H., Xu, R.K., Zhang, H., 2011. The forms of alkalis in the biochar produced from crop residues at different temperatures. Bioresour. Technol. 102, 3488–3497.
- Zemanová, V., Břendová, K., Pavlíková, D., Kubátová, P., Tlustoš, P., 2017. Effect of biochar application on the content of nutrients(Ca, Fe, K, Mg, Na, P) and amino acids in subsequently growing spinach and mustard. Plant, Soil Environ. 63, 322–

- Zhang, F., Li, J., He, Z., 2014. A new method for nutrients removal and recovery from wastewater using a bioelectrochemical system. Bioresour. Technol. 166, 630–634.
- Zhang, M., Fan, C.H., Li, Q.L., Li, B., Zhu, Y.Y., Xiong, Z.Q., 2015. A 2-yr field assessment of the effects of chemical and biological nitrification inhibitors on nitrous oxide emissions and nitrogen use efficiency in an intensively managed vegetable cropping system. Agric. Ecosyst. Environ. 201, 43–50.
- Zhao, X., Wang, S., Xing, G., 2014. Nitrification, acidification, and nitrogen leaching from subtropical cropland soils as affected by rice straw-based biochar: Laboratory incubation and column leaching studies. J. Soils Sediments 14, 471–482.
- Zheng, W., Sharma, B.K., Rajagopalan, N., 2010. Using Biochar as a Soil Amendment for Sustainable Agriculture, Illinois Department of Agriculture.
- Zhou, Y., Berruti, F., Greenhalf, C., Tian, X., Henry, H.A.L., 2017. Increased retention of soil nitrogen over winter by biochar application: Implications of biochar pyrolysis temperature for plant nitrogen availability. Agric. Ecosyst. Environ. 236, 61–68.

### Chapter 4

## 4. General discussion and conclusion

The specific objectives of this thesis were:

- i- To assess the GHGs emissions from organic and inorganic sources of nitrogen application in silage corn cropping systems
- ii- To determine the role of BC application in the reduction of GHG emission in silage corn cropping systems in western Newfoundland
- iii- To estimate GWP and GHGI of silage corn cropping systems
- iv- To determine the role of BC application on soil  $NO_3^-$  and  $NH_4^+$  retention in silage corn amended with DM and IN fertilizer application
- v- To compare the effects of dairy manure and IN alone and co-application of BC on soil pH, N uptake and biomass production of silage corn.

This thesis sought to achieve these objectives through two main experimental works as described in chapters 2 and 3. Estimation of GHG emission, GWP and GHGI under DM, and IN application and the potential role of BC in mitigating GHG emissions, GWP and GHGI have been described in chapter 2. Whereas,  $NO_3^-$  and  $NH_4^+$  retention/losses in the soil with DM, and IN application alone and combined application of BC, plant N uptake, soil pH and dry matter yield of silage corn genotypes are presented in chapter 3.

#### 4.1. Effect of biochar application on GHGs emission, GWP and GHGI

About 8 % of the total GHGs emissions in Canada are contributed by the agriculture sector largely through methane (CH<sub>4</sub>) and nitrous oxide (N<sub>2</sub>O) (Kebreab et al., 2006). N<sub>2</sub>O emissions from agricultural soils represent 3 % of anthropogenic sources in Canada (Environment and Climate Change Canada, 2017). DM and inorganic fertilizers application to agricultural soils emit significant amount of GHGs including CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O (Amon et al., 2006; Burton et al., 2008; Kebreab et al., 2006), and emissions were more pronounced with DM application than chemical fertilizers (Barneze et al., 2014; M. Zhou et al., 2017). The C compounds in the DM increase the soil  $CO_2$  emission by inducing a priming effect on native soil C (Bol et al., 2003). However, application of BC decreases the decomposition of soil organic matter (SOM) present in the soil so, it remains in the soil for a longer period (Cui et al., 2017) and reduces the cumulative  $CO_2$  emission either by the sorption of  $CO_2$  on its surface or by reducing the availability of labile C (Brennan et al., 2015). In the present study  $DM_1$  (low N dairy manure) and  $DM_2$  (high N dairy manure) treatment produced significantly higher CO<sub>2</sub> emission than IN and BC amended treatments, which is in accordance with some previous studies (Agegnehu et al., 2016; Lentz et al., 2014; Schimmelpfennig et al., 2014). BC application induces negative priming effect and decelerates the breakdown of SOM by the sorption of enzymes responsible for SOM breakdown, shifting the microbial metabolism, by increasing the stability of soil aggregates (Zheng et al., 2018) and by reducing the bioavailability of soil organic carbon (SOC) via adsorption on BC large surface area (Sheng and Zhu, 2018). It

decreases the dissolved organic carbon (DOC) from native SOC and reduces the decomposition of SOC after IN addition which reduces CO2 emission from the soil (Lu et al., 2014). High  $CH_4$  emission after DM application as observed in our experiment had been reported in other studies (Troy et al., 2013). Short-chain fatty acids present in manure are readily available to methanogenic archaea and cause  $CH_4$ emissions after application to the soil (Hrapovic and Rowe, 2002; Sherlock et al., 2002). Significant reduction in CH<sub>4</sub> emission was observed in BC amended treatments (DM<sub>1</sub>+B, DM<sub>2</sub>+B, IN+B) as compared to non-BC treatments (DM<sub>1</sub>, DM<sub>2</sub>, IN). The decrease in CH<sub>4</sub> emission after BC application may be due to the stimulation of methanotrophic activity or the increased abundance of methanotrophic proteobacterial community abundance (Feng et al., 2012; Liu et al., 2011). The BC suppresses methanogenesis by increasing the oxygen supply in the soil through increased aeration (Kim et al., 2017). There are several mechanisms by which BC could reduce N<sub>2</sub>O emissions. It improves soil aeration by reducing the soil bulk density which decreases the activity of denitrifiers in the soil (Zhang et al., 2010). The reduction in N<sub>2</sub>O emissions after BC application may be due to modification of SM, increased aeration, inhibition of nitrifier and denitrifier communities (Laird et al., 2009; Yanai et al., 2007). Application of BC (1) may accelerate the growth of soil microbes which reduce N<sub>2</sub>O emission by supporting denitrification of NO<sub>3</sub><sup>-</sup> to N<sub>2</sub> (2) facilitates the mycobacterial reduction of  $NO_3^-$  to  $NH_4^+$  (3) adsorbs  $NH_4^+$  on its surface (4) decrease the abundance of microorganisms involved in nitrification of  $NH_4^+$  to nitrite (NO<sub>2</sub>) (Anderson et al., 2011) (5) act as "electron shuttle" facilitating electron transfer to soil denitrifying microbes (Cayuela et al. 2013 ). BC adsorb  $NH_4^+$  on its surface and

reduces its availability for nitrification as a result, N<sub>2</sub>O emission is declined (Berglund et al., 2004; Lehmann et al., 2006). Application of BC to soil increased soil N mineralization, suppressed denitrification and reduced cumulative N<sub>2</sub>O emission by 91 % in a sandy loam soil (Case et al., 2015). Application of pine wood chips BC produced at 550 °C to kurosol soil (pH = 5) increased the NO<sub>3</sub><sup>-</sup> concentration in the soil and decreased the abundance of *nar*G (a gene involved in NO<sub>3</sub><sup>-</sup> reduction to NO<sub>2</sub><sup>-</sup>) (Bai et al., 2015). It also increased the abundance of nosZ (a gene involved in N<sub>2</sub>O reduction to N<sub>2</sub>) by providing suitable conditions for nosZ including increased soil pH and microbial respiration (Van Zwieten et al., 2014). Similarly, some other studies also confirmed that BC increased the abundance of nosZ, nirK, and nirS (both NO<sub>2</sub><sup>-</sup> reductase genes) and favored the last step of denitrification (converted  $N_2O$  to  $N_2$ ) which ultimately decreased N<sub>2</sub>O emission (Cayuela et al., 2013; Ducey et al., 2013; Harter et al., 2014; Van Zwieten et al., 2010b). Molar H:C ratio of BC also affects N<sub>2</sub>O emission. BCs with high H: Corg ratio is more effective in reducing N<sub>2</sub>O emission. For example, BC having a molar H: Corg ratio less than 0.3 (have high degree of aromaticity) decreased N<sub>2</sub>O emission by 73 % whereas, BCs with a molar H: Corg ratio more than 0.5 decreased N2O emission by 40 % (Cayuela et al., 2015). BC with high H:C ratio reduces the bioavailability of C for the growth of denitrifying communities (Van Zwieten et al., 2014). It has been reported in several studies that BC application improves soil aeration and increase oxygen supply, these conditions decrease the rate of denitrification in the soil. In the present study, more NO<sub>3</sub><sup>-</sup> and NH<sub>4</sub><sup>+</sup> was observed to be retained in BC amended soils as described in chapter 3. So that, the reduction in  $N_2O$  emissions was most probably due to the reduction in denitrification or due to complete denitrification (reduction of  $N_2O$  to  $N_2$ ) after BC application (Cayuela et al. 2013, 2014, 2015).

# 4.2. Biochar effects NO<sub>3</sub><sup>-</sup> and NH<sub>4</sub><sup>+</sup> retention, soil pH, plant N concentration and dry matter production

In this study, DM and IN application to soil increased NO<sub>3</sub><sup>-</sup> and NH<sub>4</sub><sup>+</sup> retention (concentration) in 20 cm soil in all treatments compared to the control treatment. The BC application reduced the movement of  $NO_3^-$  and  $NH_4^+$  to 40 cm deep soil layer which could be attributed due to the adsorption of these ions to BC surface area. Similar results have been reported in a 4-year long field experiment where BC increased the NO3 concentration in topsoil and decreased its movement to deep soil layer (Haider et al., 2017). Acid functional groups present at BC surface i.e. carboxylic, hydroxyl, lactone, lactol, phenol and carbonyls attract  $NH_4^+$  ions (Amonette and Joseph, 2009; Brennan et al., 2001; Montes-Morán et al., 2004; Zheng et al., 2010). Some other functional groups i.e. chromenes, ketones and pyrones also exist on BC surface which facilitates NO<sub>3</sub> adsorption to its surface (Amonette and Joseph, 2009; Montes-Morán et al., 2004). Unconventional H-bonding between  $NO_3^-$  and BC surface possibly increased the  $NO_3^$ adsorption on BC surface (Kammann et al., 2015; Lawrinenko, 2014; Mukherjee et al., 2011). Over the time, the adsorbed N could be desorbed and become available in soil (Kameyama et al., 2012; Taghizadeh-Toosi et al., 2012). Application of BC to soil augmented soil N mineralization, boosted nitrification and inhibited denitrification (Case et al., 2015). Most of the biochars increase soil acidity due to their high pH buffering capacity and alkaline nature. In the present study, BC amended treatments exhibited high soil pH as compared to non-BC and the control treatments at each sampling point. The carbonates and oxides formed during pyrolysis from the cations (Ca, Mg, K, Na etc.) in feedstock react with  $H^+$  and monomeric aluminum species in acidic soils and increase soil pH (Brewer et al., 2012; Enders et al., 2012; Novak et al., 2009). The BC increased  $NH_4^+$  retention in the soil and improved N uptake (Z. Sun et al., 2017), enhanced lettuce yield and nutrient concentrations in plant tissues (Upadhyay et al., 2014; Woldetsadik et al., 2017), increased soil pH, CEC, Ca, total C, N uptake and biomass production of wheat up to 250 % (Van Zwieten et al., 2010a). The increase in plant N concentration and dry matter yield (DMY) could be attributed to improvement in soil fertility, soil physical properties and nutrient retention after BC application (Mukherjee et al., 2014; Randolph et al., 2017).

## 4.3. Conclusion and Recommendations

BC application to  $DM_1$ ,  $DM_2$  and IN significantly reduced GHG emissions, decreased GWP and lowered GHGI of silage corn cropping system tested in western Newfoundland. It also improved soil pH, increased soil  $NO_3^-$  and  $NH_4^+$  retention, enhanced N concentration in plant tissues, and DMY of silage corn during the two-year field experiment. BC amended treatments reduced  $CO_2$  emission by 22 %, CH<sub>4</sub> emission by 225 %, N<sub>2</sub>O emission by 91 %, GWP by 32 %, and GHGI by 37 % compared to the control treatment. Additionally, BC amended treatments also improved Plant N concentration by 16 %, dry matter yield by 6.7 % of silage corn during two consecutive growing seasons. This study was conducted in a cool climate system. The soil of the study site was rapidly drained, Orthic Humo-Ferric Podzol which have loamy sand

texture with pH of 6.3. Based on the results of this two-year field study it was concluded that:

- i- Different sources of N have a significant impact on GHGs emissions, IN  $(NH_4^+ \text{ and } NO_3^-)$  concentrations in soil, soil pH, plant N concentration and DMY of silage corn in cool climate cropping system.
- ii- Pinewood BC could be used to mitigate GHG emissions, decrease GWP and GHGI with great success, and to increase soil pH, decrease  $NH_4^+$  and  $NO_3^-$  losses to the deep soil, increase plant N concentration and DMY.
- iii- BC maintains high soil moisture which favors crop growth during a dry spell.

Further studies exploring the role of soil bulk density, porosity, SOC contents on GHGs emissions are required as BC amendment to soil modifies soil physical properties. It also affects soil biota which affects the emission of GHGs (methanogenic, methanotrophic communities, soil nitrifying and denitrifying microbe's population) which can provide more insights into the mechanism underlying the BC role in GHGs emission reduction.

- Agegnehu, G., Bass, A.M., Nelson, P.N., Bird, M.I., 2016. Benefits of biochar, compost and biochar-compost for soil quality, maize yield and greenhouse gas emissions in a tropical agricultural soil. Sci. Total Environ. 543, 295–306.
- Amon, B., Kryvoruchko, V., Amon, T., Zechmeister-Boltenstern, S., 2006. Methane, nitrous oxide and ammonia emissions during storage and after application of dairy cattle slurry and influence of slurry treatment. Agric. Ecosyst. Environ. 112, 153– 162.
- Amonette, J.E., Joseph, S., 2009. Characteristics of biochar: microchemical properties., in: Biochar for Environmental Management; Science and Technology. p. 33.
- Anderson, C.R., Condron, L.M., Clough, T.J., Fiers, M., Stewart, A., Hill, R.A., Sherlock, R.R., 2011. Biochar induced soil microbial community change: Implications for biogeochemical cycling of carbon, nitrogen and phosphorus. Pedobiologia Int. J. Soil Biol. 54, 309–320.
- Bai, S.H., Reverchon, F., Xu, C.Y., Xu, Z., Blumfield, T.J., Zhao, H., Van Zwieten, L.,
  Wallace, H.M., 2015. Wood biochar increases nitrogen retention in field settings
  mainly through abiotic processes. Soil Biol. Biochem. 90, 232–240.
- Barneze, A.S., Mazzetto, A.M., Zani, C.F., Misselbrook, T., Cerri, C.C., 2014. Nitrous oxide emissions from soil due to urine deposition by grazing cattle in Brazil. Atmos. Environ. 92, 394–397.

- Berglund, L.M., Deluca, T.H., Zackrisson, O., 2004. Activated carbon amendments to soil alters nitrification rates in Scots pine forests. Soil Biol. Biochem. 36, 2067– 2073.
- Bol, R., Moering, J., Kuzyakov, Y., Amelung, W., 2003. Quantification of priming and CO2 respiration sources following slurry-C incorporation into two grassland soils with different C content. Rapid Commun. Mass Spectrom. 17, 2585–2590.
- Brennan, J.K., Bandosz, T.J., Thomson, K.T., Gubbins, K.E., 2001. Water in porous carbons. Colloids Surfaces A Physicochem. Eng. Asp. 187–188, 539–568.
- Brennan, R.B., Healy, M.G., Fenton, O., Lanigan, G.J., 2015. The effect of chemical amendments used for phosphorus abatement on greenhouse gas and ammonia emissions from dairy cattle slurry: Synergies and pollution swapping. PLoS One 10, e0111965.
- Brewer, C.E., Hu, Y.-Y., Schmidt-Rohr, K., Loynachan, T.E., Laird, D.A., Brown, R.C.,
  2012. Extent of Pyrolysis Impacts on Fast Pyrolysis Biochar Properties. J. Environ.
  Qual. 41, 1115–1122.
- Burton, D.L., Li, X., Grant, C.A., 2008. Influence of fertilizer nitrogen source and management practice on N2O emissions from two Black Chernozemic soils. Can. J. Soil Sci. 88, 219–227.
- Case, S.D.D.C., McNamara, N.P., Reay, D.S., Stott, A.W., Grant, H.K., Whitaker, J., 2015. Biochar suppresses N2O emissions while maintaining N availability in a sandy

loam soil. Soil Biol. Biochem. 81, 178–185.

- Cayuela, M.L., Jeffery, S., van Zwieten, L., 2015. The molar H:Corg ratio of biochar is a key factor in mitigating N2O emissions from soil. Agric. Ecosyst. Environ. 202, 135–138.
- Cayuela, M.L., Sanchez-Monedero, M.A., Roig, A., Hanley, K., Enders, A., Lehmann, J., 2013. Biochar and denitrification in soils: when, how much and why does biochar reduce N2O emissions? Sci. Rep. 3, 1732.
- Cayuela, M.L., van Zwieten, L., Singh, B.P., Jeffery, S., Roig, A., Sánchez-Monedero,
  M.A., 2014. Biochar's role in mitigating soil nitrous oxide emissions: A review and
  meta-analysis. Agric. Ecosyst. Environ. 191, 5–16.
- Cui, J., Ge, T., Kuzyakov, Y., Nie, M., Fang, C., Tang, B., Zhou, C., 2017. Interactions between biochar and litter priming: A three-source 14C and 13C partitioning study. Soil Biol. Biochem. 104, 49–58.
- Ducey, T.F., Ippolito, J.A., Cantrell, K.B., Novak, J.M., Lentz, R.D., 2013. Addition of activated switchgrass biochar to an aridic subsoil increases microbial nitrogen cycling gene abundances. Appl. Soil Ecol. 65, 65–72.
- Enders, A., Hanley, K., Whitman, T., Joseph, S., Lehmann, J., 2012. Characterization of biochars to evaluate recalcitrance and agronomic performance. Bioresour. Technol. 114, 644–653.

Environment and Climate Change Canada, 2017. National Inventory Report - The

Canadian Government's Submission to the UN Framework Convention on Climate Change 1990–2005: Greenhouse Gas Sources and Sinks in Canada.

- Feng, Y., Xu, Y., Yu, Y., Xie, Z., Lin, X., 2012. Mechanisms of biochar decreasing methane emission from Chinese paddy soils. Soil Biol. Biochem. 46, 80–88.
- Haider, G., Steffens, D., Moser, G., Müller, C., Claudia, I.K., 2017. Biochar reduced nitrate leaching and improved soil moisture content without yield improvements in a four-year field study. Agric. Ecosyst. Environ. 237, 80–94.
- Harter, J., Krause, H.-M., Schuettler, S., Ruser, R., Fromme, M., Scholten, T., Kappler,A., Behrens, S., 2014. Linking N2O emissions from biochar-amended soil to the structure and function of the N-cycling microbial community. ISME J. 8, 660–674.
- Hrapovic, L., Rowe, R.K., 2002. Intrinsic degradation of volatile fatty acids in laboratorycompacted clayey soil. J. Contam. Hydrol. 58, 221–242.
- Kameyama, K., Miyamoto, T., Shiono, T., Shinogi, Y., 2012. Influence of SugarcaneBagasse-derived Biochar Application on Nitrate Leaching in Calcaric Dark Red Soil.J. Environ. Qual. 41, 1131–1137.
- Kammann, C.I., Schmidt, H.-P., Messerschmidt, N., Linsel, S., Steffens, D., Müller, C., Koyro, H.-W., Conte, P., Stephen, J., 2015. Plant growth improvement mediated by nitrate capture in co-composted biochar. Sci. Rep. 5, 11080.
- Kebreab, E., Clark, K., Wagner, R.C., France, J., 2006. Methane and nitrous oxide emissions from Canadian animal agriculture: A review. Can. J. Anim. Sci. 86, 135–

- Kim, J., Yoo, G., Kim, D., Ding, W., Kang, H., 2017. Combined application of biochar and slow-release fertilizer reduces methane emission but enhances rice yield by different mechanisms. Appl. Soil Ecol. 117, 57–62.
- Laird, D.A., Brown, R.C., Amonette, J.E., Lehmann, J., 2009. Review of the pyrolysis platform for coproducing bio-oil and biochar. Biofuels, Bioprod. Biorefining 3, 547–562.
- Lawrinenko, M., 2014. Anion exchange capacity of biochar. Green Chem. Lowa State University, Lowa, USA.
- Lehmann, J., Gaunt, J., Rondon, M., 2006. Bio-char sequestration in terrestrial ecosystems A review. Mitig. Adapt. Strateg. Glob. Chang. 11, 403–427.
- Lentz, R.D., Ippolito, J.A., Spokas, K.A., 2014. Biochar and Manure Effects on Net Nitrogen Mineralization and Greenhouse Gas Emissions from Calcareous Soil under Corn. Soil Sci. Soc. Am. J. 78, 1641–1655.
- Liu, Y., Yang, M., Wu, Y., Wang, H., Chen, Y., Wu, W., 2011. Reducing CH4 and CO2 emissions from waterlogged paddy soil with biochar. J. Soils Sediments 11, 930– 939.
- Lu, W., Ding, W., Zhang, J., Li, Y., Luo, J., Bolan, N., Xie, Z., 2014. Biochar suppressed the decomposition of organic carbon in a cultivated sandy loam soil: A negative priming effect. Soil Biol. Biochem. 76, 12–21.

- Montes-Morán, M.A., Suárez, D., Menéndez, J.A., Fuente, E., 2004. On the nature of basic sites on carbon surfaces: An overview. Carbon N. Y. 42, 1219–1225.
- Mukherjee, A., Lal, R., Zimmerman, A.R., 2014. Effects of biochar and other amendments on the physical properties and greenhouse gas emissions of an artificially degraded soil. Sci. Total Environ. 487, 26–36. doi:10.1016/j.scitotenv.2014.03.141
- Mukherjee, A., Zimmerman, A.R., Harris, W., 2011. Surface chemistry variations among a series of laboratory-produced biochars. Geoderma 163, 247–255.
- Novak, J.M., Busscher, W.J., Laird, D.L., Ahmedna, M., Watts, D.W., Niandou, M.A.S., 2009. Impact of Biochar Amendment on Fertility of a Southeastern Coastal Plain Soil. Soil Sci. 174, 105–112.
- Randolph, P., Bansode, R.R., Hassan, O.A., Rehrah, D., Ravella, R., Reddy, M.R., Watts, D.W., Novak, J.M., Ahmedna, M., 2017. Effect of biochars produced from solid organic municipal waste on soil quality parameters. J. Environ. Manage. 192, 271–280.
- Schimmelpfennig, S., Müller, C., Grünhage, L., Koch, C., Kammann, C., 2014. Biochar, hydrochar and uncarbonized feedstock application to permanent grassland-Effects on greenhouse gas emissions and plant growth. Agric. Ecosyst. Environ. 191, 39–52.
- Sheng, Y., Zhu, L., 2018. Biochar alters microbial community and carbon sequestration potential across different soil pH. Sci. Total Environ. 622–623, 1391–1399.

- Sherlock, R.R., Sommer, S.G., Khan, R.Z., Wood, C.W., Guertal, E.A., Freney, J.R., Dawson, C.O., Cameron, K.C., 2002. Ammonia, Methane, and Nitrous Oxide Emission from Pig Slurry Applied to a Pasture in New Zealand. J. Environ. Qual. 31, 1491–1501.
- Sun, Z., Sänger, A., Rebensburg, P., Lentzsch, P., Wirth, S., Kaupenjohann, M., Meyer-Aurich, A., 2017. Contrasting effects of biochar on N2O emission and N uptake at different N fertilizer levels on a temperate sandy loam. Sci. Total Environ. 578, 557– 565.
- Taghizadeh-Toosi, A., Clough, T.J., Sherlock, R.R., Condron, L.M., 2012. Biochar adsorbed ammonia is bioavailable. Plant Soil 350, 57–69.
- Troy, S.M., Lawlor, P.G., O' Flynn, C.J., Healy, M.G., 2013. Impact of biochar addition to soil on greenhouse gas emissions following pig manure application. Soil Biol. Biochem. 60, 173–181.
- Upadhyay, K.P., George, D., Swift, R.S., Galea, V., 2014. The influence of biochar on growth of lettuce and potato. J. Integr. Agric. 13, 541–546.
- Van Zwieten, L., Kimber, S., Morris, S., Chan, K.Y., Downie, A., Rust, J., Joseph, S., Cowie, A., 2010a. Effects of biochar from slow pyrolysis of papermill waste on agronomic performance and soil fertility. Plant Soil 327, 235–246.
- Van Zwieten, L., Kimber, S., Morris, S., Downie, A., Berger, E., Rust, J., Scheer, C.,2010b. Influence of biochars on flux of N2O and CO2from Ferrosol. Aust. J. Soil

Res. 48, 555–568.

- Van Zwieten, L., Singh, B.P., Kimber, S.W.L., Murphy, D. V., Macdonald, L.M., Rust,
  J., Morris, S., 2014. An incubation study investigating the mechanisms that impact
  N2O flux from soil following biochar application. Agric. Ecosyst. Environ. 191, 53–62.
- Woldetsadik, D., Drechsel, P., Marschner, B., Itanna, F., Gebrekidan, H., 2017. Effect of biochar derived from faecal matter on yield and nutrient content of lettuce (Lactuca sativa) in two contrasting soils. Environ. Syst. Res. 6.
- Yanai, Y., Toyota, K., Okazaki, M., 2007. Effects of charcoal addition on N2O emissions from soil resulting from rewetting air-dried soil in short-term laboratory experiments. Soil Sci. Plant Nutr. 53, 181–188.
- Zhang, A., Cui, L., Pan, G., Li, L., Hussain, Q., Zhang, X., Zheng, J., Crowley, D., 2010.Effect of biochar amendment on yield and methane and nitrous oxide emissions from a rice paddy from Tai Lake plain, China. Agric. Ecosyst. Environ. 139, 469–475.
- Zheng, H., Wang, X., Luo, X., Wang, Z., Xing, B., 2018. Biochar-induced negative carbon mineralization priming effects in a coastal wetland soil: Roles of soil aggregation and microbial modulation. Sci. Total Environ. 610–611, 951–960.
- Zheng, W., Sharma, B.K., Rajagopalan, N., 2010. Using Biochar as a Soil Amendment for Sustainable Agriculture, Illinois Department of Agriculture.
- Zhou, M., Zhu, B.O., Wang, S., Zhu, X., Vereecken, H., 2017. Stimulation of N2O

emission by manure application to agricultural soils may largely offset carbon benefits: a global meta-analysis. Glob. Chang. Biol. 23, 4068–4083.