

Identification of Sex-Specific Genetic Polymorphisms Associated with Asthma in Middle-Aged and Older Canadian Adults: An Analysis of CLSA Data

Ugochukwu Odimba¹, Ambikaipakan Senthilselvan², Jamie Farrell^{1,3}, Zhiwei Gao¹

¹Clinical Epidemiology Unit, Division of Community Health and Humanities, Faculty of Medicine, Memorial University, St. John's, Newfoundland and Labrador, Canada; ²School of Public Health, University of Alberta, Edmonton, Alberta, Canada; ³Faculty of Medicine, Health Science Centre (Respirology Department), Memorial University, St John's, Newfoundland and Labrador, Canada

Correspondence: Zhiwei Gao, Clinical Epidemiology Unit, Division of Community Health and Humanities, Faculty of Medicine, Memorial University of Newfoundland, St. John's, Newfoundland and Labrador, A1B 3V6, Canada, Tel +17098646523, Email zhiwei.gao@med.mun.ca

Purpose: Asthma is a chronic heterogeneous respiratory disease resulting from a complex interplay between genetic variations and environmental exposures. There are sex disparities in the prevalence and severity of asthma in males and females. Asthma prevalence is higher in males during childhood but increases in females in adulthood. The mechanisms underlying these sex differences are not well understood; nevertheless, genetic variations, hormonal changes, and environmental influences are thought to play important roles. This study aimed to identify sex-specific genetic variants associated with asthma using CLSA genomic and questionnaire data.

Methods: First, we conducted a genome-wide SNP-by-sex interaction analysis on 23,323 individuals, examining 416,562 single nucleotide polymorphisms (SNPs) after quality control, followed by sex-stratified survey logistic regression of SNPs with interaction p-value less than 10^{-5} .

Results: Out of the 49 SNPs with interaction p-value less than 10^{-5} , a sex-stratified survey logistic regression showed that five male-specific SNPs (rs6701638, rs17071077, rs254804, rs6013213, and rs2968822) in/near KIF26B, NMBR, PEPD, RTN4, and NFATC2 loci, and three female-specific SNPs (rs2968801, rs2864052, and rs9525931) in/near RTN4, and SERP2 loci were significantly associated with asthma after Bonferroni correction. An SNP (rs36213) in the EPHB1 gene was significantly associated with an increased risk of asthma in males [OR=1.35, 95% CI (1.14, 1.60)] but with a reduced risk of asthma in females [OR=0.84, 95% CI (0.76, 0.92)] after Bonferroni correction.

Conclusion: We discovered novel sex-specific genetic markers in/near the KIF26B, RTN4, EPHB1, NMBR, SERP2, PEPD, and NFATC2 genes that could potentially shed light on the sex differences in asthma susceptibility in males and females. Future mechanistic studies are required to understand better the underlying sex-related pathways of the identified loci in asthma development.

Keywords: asthma, single nucleotide polymorphism, sex-specific, CLSA

Introduction

Asthma is a chronic inflammatory respiratory disease that affects both children and adults. It is characterized by reversible airflow obstruction, airway remodeling, hyperresponsiveness, shortness of breath, wheezing, and mucus production. In 2019, there were more than 262 million asthma cases globally, and the age-standardized point prevalence was highest in the high-income North American region.¹ In Canada, between 2011 and 2012, an estimated 3.8 million people one year and older lived with asthma.² Asthma prevalence and incidence differ in childhood and adulthood, and sex differences in prevalence, incidence, susceptibility, and severity have long been recognized.³⁻⁵ During childhood, the prevalence of asthma is higher in males than in females, while in puberty and adulthood, females are more likely to develop asthma than males.⁶

Sexual dimorphism has been reported to influence the severity, pathogenesis, pathophysiology, progression, and susceptibility of various human diseases and treatments, including asthma.^{7–10} Women with asthma tend to have higher mortality and morbidity,¹¹ greater perception of symptom bothersomeness, and poorer quality of life than men with asthma.^{12,13} Studies using antigen-challenged mouse asthma models have shown that female mice had greater levels of antigen-specific serum IgE, eosinophils, and Th2 cytokine than male mice in the lung tissue and bronchoalveolar lavage fluid.^{14,15} Furthermore, airway hyperresponsiveness has been reported to be greater in male mice than in female mice at six weeks of age and reduced at twelve weeks of age.¹⁶

Although the underlying biological mechanism of respiratory diseases and sex differences in risk are not fully understood, it is commonly believed that anatomical differences, genetic predisposition, hormonal changes, environmental exposures, and gene and environment interactions play an important role.^{4,17–19}

Genetic predisposition plays a vital role in asthma pathogenesis. More than a hundred independent genetic loci have been associated with asthma.^{20,21} Studies have shown that several gene polymorphisms and gene expressions associated with asthma are sex-specific.^{22–25} For instance, a large genome-wide association study (GWAS) using EVE Asthma Genetics Consortium discovered six sex-specific asthma risk loci, with two male-specific SNPs in/near IRF and RAB11FIP2 gene and four female-specific SNPs in/near RAP1GAP2, C6orf118, ERBB4, and AK057517.²⁵ Other sex-specific polymorphisms in many genes have been identified, eg, a male-specific SNP in the LCORL gene was significantly associated with childhood asthma,²² and a female-specific polymorphism in the COX-2₋₇₆₅ gene was significantly associated with bronchial asthma.²⁶ Most GWAS of asthma generally include limited information on social and environmental factors and do not have enough sample size to investigate sex-specific genetic effects; this study aims to identify the sex-specific genetic polymorphisms significantly associated with asthma in middle-aged and older Canadian adults using the Canadian Longitudinal Study on Aging (CLSA) data (Baseline Comprehensive and Genomic datasets).

Methods and Materials

Study Population

This study included 26,622 individuals with complete genotyping information from the Canadian Longitudinal Study for Aging (CLSA) comprehensive cohort.²⁷ The CLSA comprehensive cohort comprises over 30,000 middle-aged and older Canadian adults between the ages of 45 and 85 years. Baseline data on a wide range of variables, including socio-demographic and socio-economic factors, information on lifestyle and health behaviors, physical measurements including height, weight, pulmonary function test, and biospecimens (blood and urine), were obtained through in-person home interviews and visits to data collection sites (DCS).^{28,29}

Definition of Asthma

A positive response to the following question from the CLSA questionnaire determined the presence of self-reported physician-diagnosed asthma, “Has a doctor ever told you that you have asthma?” However, we excluded the following participants from this study: participants with missing responses, participants who self-reported a physician diagnosis of COPD, and participants who self-reported a physician diagnosis of asthma and COPD.

Covariates

The following potential confounders were considered: age and principal components of genetic ancestry were included as continuous variables. Categorical variables included were age groups (45 to 54 years, 55 to 64 years, 65 to 74 years, and over 75 years), biological sex (male vs female), smoking status (current, never, and former smokers), marital status (single/never married, married/common-law relationship, widowed/divorced/separated), education level (less than post-secondary education, post-secondary but not university education, and university education/others), total personal income and total household income (Less than \$20,000, \$20,000 to less than \$50,000, \$50,000 to less than \$100,000, \$100,000 or more), province of recruitment (Prairies, British Columbia, Eastern provinces, Ontario and Quebec), retirement status

(Retired completely vs retired/partly retired), homeownership (owned vs rented/others), and urban/rural dwelling (urban core vs rural/not urban core).

Genotyping, Sample, and Marker Quality Control

Genotyping of 794,409 genetic markers was undertaken using Affymetrix UK Biobank Axiom array.²⁷ Detailed sample and markers quality control have been described previously.²⁷ Samples and markers that failed the quality control requirements were excluded. Samples with inconsistent sex information (discrepancy between reported sex and sex determined from genetic data) and high rates of genotype missingness (> 5%). SNPs in sex chromosomes, SNPs with low genotype call rates (< 99%), SNPs with minor allele frequency < 1% (MAF < 0.01), and SNPs deviating from the Hardy Weinberg equilibrium threshold of 1e-10 were excluded. Using PLINK's *Indep-pairwise* command (*Indep-pairwise* 50, 5, 0.5), we generated a subset of SNPs in approximate linkage equilibrium. Post-quality control, we had 2,799 asthma cases, 20,524 controls, and 416,562 SNPs for genome-wide SNP-by-sex interaction analysis.

Statistical Analysis

Descriptive statistics, mean (SE), and frequency (%) for continuous and categorical variables were presented to describe the study population. Characteristics between male and female participants were compared using chi-square tests and Student's *t*-test. Trimmed inflation and analytic weights provided in the CLSA data (CLSA Sample Weights Version 1.2) were used for descriptive and regression analyses.

In order to identify sex-specific SNPs associated with asthma, first, we performed a multivariate logistic regression using PLINK 1.90b6.2^{30,31} to identify significant interaction terms between a sex variable, and each of the SNPs in the regression analysis of asthma after controlling for age, sex, smoking status, and the first four principal components of genetic ancestry. Second, we performed a sex-stratified analysis for those SNPs with an SNP-by-sex interaction p-value less than 10⁻⁵ using survey-specific logistic regression (*Proc Surveylogistic*) in SAS 9.4 version. This approach allowed us to include sampling weights, complex survey design variables and adjust for potential confounders.

A purposeful model selection method was used to identify potential confounders for inclusion in the final model of the sex-stratified analysis. The covariates with a p-value ≤ 0.20 from the univariate analysis were entered into an interim multivariate model. The least significant covariate was then removed one at a time until only covariates with significant p-values (p ≤ 0.05) and clinically important factors remained in the model. The following covariates were included in the final model of the sex-stratified analysis for asthma: age, smoking status, province of recruitment, marital status, total personal income, and retirement status. The "sex variable" was included as a domain factor in the sex-stratified multivariate survey logistic regression to examine one SNP at a time after controlling for the first four principal components, age, smoking status, province of recruitment, marital status, total personal income, and retirement status.

Three inheritance models (dominant, recessive, and additive) for each variant were evaluated. However, only the inheritance model with the smallest AIC value (the best-fitted model) was selected and presented. Bonferroni correction was applied to control for multiple comparisons. The association's strength was reported as an odds ratio with 95% confidence intervals. We created regional association plots using LocusZoom³² for the significant SNPs from the sex-stratified analysis. All statistical analysis was performed using Plink 1.90b6.2³⁰ and SAS 9.4 (SAS Institute Inc, Cary, NC).

Results

Population Characteristics

Table 1 compares the characteristics of the study population between males and females. Female subjects were more likely to have self-reported physician-diagnosed asthma than males (14.0% vs 10.6%). The mean age of females was significantly greater than that of males (60.0 vs 59.1, p ≤ 0.0001). The distribution of smoking status, marital status, urban or rural dwelling, homeownership, total household, personal income, retirement status, province of recruitment, and highest education status also differed significantly between males and females. The proportion of current and former

Table 1 Characteristics of the Study Population

Characteristics	Males (n=13,343) (%)	Females (n= 13,279) (%)	p-value*
<u>Asthma</u>			
Yes	10.6	14	<0.0001
Age, mean \pm SEM	59.1 \pm 0.10	60.0 \pm 0.10	<0.0001
<u>Lifestyle and Health Behavior</u>			
Smoking Status			<0.0001
Current Smoker	10	8.5	
Never Smoker	47.6	51.4	
Former Smoker	42.4	40.1	
<u>Socio-Demographic and Economic Factors</u>			
Marital status			<0.0001
Single/never married	8.1	8.2	
Married/Living in common law	82.3	70.2	
Widowed/Separated/Divorced	9.6	21.6	
Urban or Rural Dwelling			0.003
Urban core	86.6	85	
Rural/not Urban core	13.4	15	
Homeownership			0.0008
Owned	86.7	85	
Rented/Others	13.3	15	
Total household income			<0.0001
Less than \$20,000	3.4	5.6	
\$20,000 to less than \$50,000	15	22.2	
\$50,000 to less than \$100,000	32.5	34.2	
\$100,000 or more	49.1	38	
Total personal income			<0.0001
Less than \$20,000	7.8	22.9	
\$20,000 to less than \$50,000	27.9	39.9	
\$50,000 to less than \$100,000	40.7	28.9	
\$100,000 or more	23.6	8.2	
Retirement Status			<0.0001
Retired completely	29	38	
Not retired/partly retired	71	62	
Province of recruitment			<0.0001
Prairies	22.2	19.1	
British Columbia	28.9	30	
Eastern province ^a	5.2	6.2	
Ontario	13.5	13.6	
Quebec	30.2	31.1	
Highest Education Status			<0.0001
No post-secondary Education	7.2	8.4	
Non-university post-secondary	31.4	34.8	
University post-secondary	61.4	56.8	

Notes: *p-values for overall comparison between males and females; ^aNewfoundland and Labrador; Nova Scotia.

smokers was significantly greater in males than in females. The percentage of widowed, separated, or divorced females (21.6%) was significantly higher than that of males (9.6%).

Results of SNP-by-Sex Interaction Analysis

After quality control, 416,562 SNPs were included in the GWAS of SNP by sex interaction analysis. As indicated in Table 2, 49 variants showed a p-value less than 10^{-5} for the interaction term. The polymorphisms of rs7676077 and

Table 2 49 Signals of the SNP-by-Sex Interactions with p-value Less Than 10^{-5} for Asthma

CHR	SNP	A1/A2	MAF	Gene/Nearest Gene	OR (95% CI)	Interaction p-value	Functional Consequence
4	rs7676077	T/C	0.308	SLC34A2	0.73 (0.641, 0.821)	3.84E-07	Intergenic variant
1	rs6701638	G/A	0.493	KIF26B	1.33 (1.188, 1.492)	7.88E-07	Intron variant
2	rs2968822	T/C	0.403	RTN4	1.33 (1.184, 1.489)	1.24E-06	Intron variant
2	rs2968801	G/A	0.297	RTN4	1.35 (1.190, 1.521)	2.14E-06	Intergenic variant
14	rs10467731	G/A	0.014	SLC38A6	0.27 (0.158, 0.473)	3.52E-06	Regulatory region variant
20	rs11699160	C/T	0.353	RPS21	0.76 (0.671, 0.852)	4.40E-06	Intergenic variant
18	rs4800396	C/T	0.169	CTAGE1	0.70 (0.602, 0.817)	4.94E-06	Intergenic variant
18	rs3911730	A/C	0.079	RTTN	0.62 (0.503, 0.766)	8.95E-06	Missense variant
2	rs2006878	A/G	0.271	EML6	1.33 (1.173, 1.510)	9.06E-06	Intron variant
1	rs3737247	G/A	0.384	SMYD3	0.77 (0.687, 0.869)	1.53E-05	Intron variant
1	rs7519461	T/C	0.331	KIF26B	1.30 (1.154, 1.468)	1.84E-05	Intron variant
13	rs117190915	C/T	0.034	LINC00408	0.48 (0.338, 0.669)	1.94E-05	Intron variant
21	rs34026679	A/C	0.041	APP	1.88 (1.405, 2.506)	1.99E-05	Intergenic variant
20	rs79496646	C/T	0.038	KIF16B	1.87 (1.403, 2.504)	2.12E-05	Intergenic variant
7	rs62442915	T/C	0.039	MAD1L1	1.88 (1.406, 2.520)	2.13E-05	Intron variant
6	rs17071077	G/A	0.278	NMBR	0.76 (0.664, 0.860)	2.21E-05	Intergenic variant
5	rs75073573	A/C	0.199	FGF18	0.73 (0.631, 0.845)	2.37E-05	Intron variant
13	rs9525931	G/A	0.245	SERP2	1.33 (1.166, 1.523)	2.50E-05	Intergenic variant
13	rs628778	T/C	0.066	FARPI	0.61 (0.481, 0.767)	2.78E-05	Synonymous variant
19	rs254804	G/T	0.181	PEPD	1.37 (1.180, 1.578)	2.83E-05	Intergenic variant
10	rs72804487	T/C	0.018	BICC1	0.38 (0.238, 0.595)	2.85E-05	Intergenic variant
5	rs61027918	G/A	0.173	TENM2	0.72 (0.622, 0.842)	3.00E-05	Intron variant
3	rs35644024	A/G	0.073	TGFBR2	0.61 (0.487, 0.772)	3.07E-05	Intergenic variant
13	rs74438452	G/T	0.019	DACH1	2.67 (1.680, 4.237)	3.21E-05	Intron variant
16	rs111933452	A/G	0.049	CDH5	1.75 (1.342, 2.272)	3.31E-05	Intergenic variant
8	rs73191549	A/C	0.087	MSRA	1.52 (1.243, 1.848)	3.93E-05	Intron variant
7	rs4718733	G/A	0.317	NA	1.29 (1.141, 1.457)	4.51E-05	Intergenic variant

(Continued)

Table 2 (Continued).

CHR	SNP	A1/A2	MAF	Gene/Nearest Gene	OR (95% CI)	Interaction p-value	Functional Consequence
6	rs1416037	A/G	0.086	HACE1	1.54 (1.250, 1.887)	4.51E-05	Intergenic variant
15	rs16974494	C/G	0.231	SEC11A	1.32 (1.155, 1.509)	4.64E-05	Intron variant
21	rs382583	T/C	0.109	RBM11	0.68 (0.562, 0.818)	5.08E-05	Intron variant
3	rs1488342	T/G	0.368	LOC101927394	0.78 (0.694, 0.881)	5.27E-05	Intron variant
3	rs36213	G/C	0.487	EPHB1	1.26 (1.128, 1.414)	5.43E-05	Intron variant
5	rs6893959	A/G	0.439	CD180	1.26 (1.128, 1.416)	5.63E-05	Intergenic variant
16	rs7200747	C/G	0.396	ZNF598	0.79 (0.700, 0.884)	5.74E-05	Intron variant
20	rs6013213	A/G	0.445	NFATC2	0.79 (0.704, 0.886)	6.06E-05	Intron variant
2	rs10205651	C/T	0.131	NA	0.71 (0.605, 0.841)	6.10E-05	Regulatory region variant
2	rs2864052	A/G	0.239	RTN4	1.31 (1.147, 1.493)	6.11E-05	Intron variant
7	rs59889022	T/C	0.122	LOC100507642	0.69 (0.582, 0.831)	6.53E-05	Intergenic variant
2	rs3821261	C/A	0.075	TGFA	0.64 (0.515, 0.799)	7.35E-05	Intron variant
20	rs117107888	T/C	0.054	RALGAPA2	1.68 (1.300, 2.178)	7.67E-05	Intron variant
16	rs4513116	G/A	0.019	RBFOX1	0.37 (0.226, 0.606)	7.92E-05	Intergenic variant
2	rs1527243	C/T	0.160	TSN	0.73 (0.629, 0.856)	8.18E-05	Intergenic variant
15	rs837132	C/T	0.262	MEIS2	1.29 (1.137, 1.467)	8.30E-05	Intron variant
13	rs7998573	T/C	0.012	SLITRK6	0.29 (0.163, 0.545)	8.38E-05	Intron variant
1	rs868376	T/C	0.279	PBX1	0.77 (0.679, 0.879)	8.56E-05	Intergenic variant
3	rs73877940	T/A	0.033	ARL14	0.49 (0.348, 0.704)	8.92E-05	Regulatory region variant
16	rs72799714	G/T	0.054	NA	0.58 (0.436, 0.759)	9.11E-05	Intergenic variant
6	rs6926101	T/C	0.057	SLC18B1	0.61 (0.477, 0.782)	9.25E-05	Missense variant
20	rs6041383	T/C	0.029	LOC101929486	0.46 (0.314, 0.681)	9.47E-05	Intergenic variant

Notes: SNP-by-Sex Interaction GWAS on Asthma controlling for Age, Sex, smoking status, and top 4 principal components. Genes contain variants or are located within 500kb of variants.

Abbreviations: Chr, Chromosome; SNP, Single Nucleotide Polymorphism; MAF, Minor allele frequency; OR, Odds ratio; A1/A2, Minor/Major allele; NA, No Annotation.

rs6701638 in/near the SLC34A2 and KIF26B genes (interaction p-values = 3.84×10^{-7} and 7.88×10^{-7} respectively) were close to reaching the genome-wide significance p-value threshold (Figure 1A). There was no evidence of population stratification, as indicated by the genomic inflation factor ($\lambda = 1.025$) in the Q-Q plot shown in Figure 1B.

Results of Sex Stratified Analysis

Figure 2 presents regional plots of the significant sex-specific SNPs from the sex-stratified multivariate survey logistic regression. After adjusting for multiple comparisons using Bonferroni correction at $\alpha \leq 0.001$ (0.05/49) significance level, we identified five male-specific SNPs (rs6701638, rs2968822, rs17071077, rs254804, and rs6013213) in/near KIF26B, RTN4, NMBR, PEPD, and NFATC2 loci and three female-specific SNPs (rs2968801, rs2864052 and rs9525931) in/near RTN4, and SERP2 genes (Table 3, Figure 3A and 3B). The polymorphism rs36213 in the EPHB1 gene was significantly associated with an increased risk of asthma [OR=1.35, p= 0.0004, 95% CI (1.14, 1.60)] in males and a reduced risk of asthma [OR=0.84, p= 0.0003, 95% CI (0.76, 0.92)] in females (Table 3, Figure 3A and 3B).




Figure 1 (A) Manhattan plot for the SNP-by-sex interaction GWAS. The X-axis indicates the chromosomal positions of each variant and the Y-axis shows the negative Log p-values. The blue horizontal line indicates the suggestive significance threshold (1×10^{-5}), and the red horizontal line shows the genome-wide significance threshold (5×10^{-8}). (B) Quantile-Quantile plot showing the distribution of observed versus expected p-values. The genomic inflation factor, lambda (λ) = 1.025.

In the additive genetic model, males with homozygous GG of rs6701638 (minor allele: G) in the KIF26B gene, homozygous GG of rs254804 (minor allele: G) near the PEPD gene, and homozygous TT of rs2968822 (minor allele: T) in RTN4 gene exhibited an increased risk of asthma (ORs: 1.24, 1.27, and 1.22) than those with wild-type (AA for rs6701638, TT for rs254804 and CC for rs2968822). In addition, males with homozygous GG of rs17071077 (minor allele: G) near the NMBR gene and homozygous AA of rs6013213 (minor allele: A) in the NFATC2 gene showed reduced risk of asthma (ORs: 0.79 and 0.77) compared to males with wild-type (AA for rs17071077 and GG for rs6013213).

Under the dominant genetic model, females with homozygous GG or heterozygous GA for rs2968801 (minor allele: G) and homozygous AA or heterozygous AG for rs2864052 (minor allele: A) in the RTN4 gene had a reduced risk of asthma (ORs: 0.79 and 0.77, respectively) compared to females with the wild-type (AA for rs2968801 and GG for rs2864052). Under the additive genetic model, females carrying homozygous GG for rs9525931 (minor allele: G) near the SERP2 gene demonstrated a lower risk of asthma (OR: 0.80) compared to females with the wild-type (AA for rs9525931).

Discussion

In this study, we identified sex-specific SNPs associated with asthma among middle-aged and older Canadians. Five male-specific SNPs (rs6701638, rs17071077, rs254804, rs6013213 and rs2968822) and three female-specific SNPs (rs2968801, rs2864052 and rs9525931) were significantly associated with asthma. Three of the five male-specific SNPs were associated with an increased risk of asthma, with an odds ratio ranging from 1.22 to 1.27. The other two male-specific SNPs were associated with a reduced risk of asthma (OR = 0.79 and OR = 0.77). All three female-specific SNPs were associated with a reduced risk of asthma, with ORs ranging from 0.77 to 0.80. The polymorphism rs36213 was significantly associated with an increased risk of asthma (OR = 1.35) in males but with a reduced risk of asthma (OR = 0.84) in females. In previous studies, male and female-specific genetic markers have been associated with asthma.^{22,25} However, these studies did not consider the influence of socio-demographic, socio-economic, and lifestyle factors.

The five male-specific polymorphisms, ie, rs6701638 in the KIF26B, rs254804 near the PEPD, rs17071077 near the NMBR, rs6013213 in the NFATC2, and rs2968822 near the RTN4 genes are located at 1q44, 19q13.11, 6q24.1, 20q13.2 and 2p16.1 genomic regions, respectively. The three female-specific polymorphisms, ie, rs2968801 near the RTN4, rs2864052 in the RTN4, and rs9525931 near the SERP2 genes, are located at 2p16.1 and 13q14.11 cytogenetic regions.

The polymorphism rs36213 (minor allele: G), which was significantly associated with an increased risk of asthma in males and a decreased risk of asthma in females, is an intronic variant in the EPH receptor B1 gene (EPHB1). EPHB1 is a receptor for Ephrin-B ligands (Ephrin-B1, Ephrin-B2, Ephrin-B3). EPH receptors are the largest family members of receptor tyrosine kinase (RTK), which are expressed in various immune cells, including CD4⁺ and CD8⁺ T cells,




Figure 2 Regional plots for the nine sex-specific SNPs show their respective locations. **(A)** SNP rs6701638 in KIF26B gene. **(B)** SNP rs17071077 near NMBR gene. **(C)** SNP rs254804 near PEPD gene. **(D)** SNP rs6013213 in NFATC2 gene. **(E)** SNP rs2968822 in RTN4 gene. **(F)** SNP rs2968801 near RTN4 gene. **(G)** SNP rs2864052 in RTN4 gene. **(H)** SNP rs9525931 near SERP2 gene. **(I)** SNP rs36213 in EPHB1 gene.

lymphocytes, monocytes, and granulocytes.^{33–35} Furthermore, the EPH receptor-ephrin ligand interaction has been linked to immune cell activation, T cell differentiation, proliferation, and migration.^{34,35} Previous research has linked allergic rhinitis and asthma to EPH receptors, especially the EPHB2 gene, an important paralog of the EPHB1 gene.³⁶ Recently, animal studies have illustrated that sex hormones augmented the sex-specific effect of EPHB receptors and Ephrin-B ligands.^{37–39} Wang et al³⁷ found that deleting EPHB4 from vascular smooth muscle cells of mice resulted in hypotension in males but not females. In a different experiment, female but not male Ephrin-B3 null mice had higher blood pressure and increased vascular smooth muscle cell contractions than wild-type.³⁸ In addition, estrogen enhanced increased vascular smooth muscle contraction in female Ephrin-b3 null mice compared with wild type, whereas testosterone reduced it.³⁸ These suggest that sex hormones may modulate ephrin-Eph receptor functions in males and females and, as such, may play crucial sex-specific role in asthma susceptibility.

This study also discovered three variants in/near the RTN4 gene, SNPs rs2968822 (minor allele: T), rs2968801 (minor allele: G), and rs2864052 (minor allele: A), associated with asthma in males and females, respectively. While the

Table 3 Result of Sex Stratified Analysis for Asthma

Chr	Gene /Nearest Gene	SNP (A1/A2) [Functional Consequence]	Male		Female	
			OR (95% CI)	p-value	OR (95% CI)	p-value
1	KIF26B	rs6701638 (G/A) [Intron variant]	1.24 (1.11, 1.38) ^A	0.0001*	0.91 (0.83, 1.00) ^A	0.0503
6	NMBR	rs17071077 (G/A) [Intergenic variant]	0.79 (0.70, 0.91) ^A	0.0006*	PM: 1.05 (0.92, 1.19) ^D WT: I	0.5095
19	PEPD	rs254804 (G/T) [Intergenic variant]	1.27 (1.11, 1.45) ^A	0.0005*	PM: 0.89 (0.77, 1.03) ^D WT: I	0.1167
20	NFATC2	rs6013213 (A/G) [Intron variant]	0.77 (0.69, 0.86) ^A	<0.0001*	PM: 1.04 (0.89, 1.19) ^D WT: I	0.6293
2	RTN4	rs2968822 (T/C) [Intron variant]	1.22 (1.09, 1.36) ^A	0.0004*	PM: 0.85 (0.74, 0.97) ^D WT: I	0.0174
2	RTN4	rs2968801 (G/A) [Intergenic variant]	1.12 (0.99, 1.25) ^A	0.0587	PM: 0.79 (0.69, 0.91) ^D WT: I	0.0007*
2	RTN4	rs2864052 (A/G) [Intron Variant]	1.09 (0.96, 1.23) ^A	0.1843	PM: 0.77 (0.67, 0.89) ^D WT: I	0.0003*
13	SERP2	rs9525931 (G/A) [Intergenic variant]	1.07 (0.95, 1.21) ^A	0.2716	0.80 (0.72, 0.89) ^A	<0.0001*
3	EPHBI	rs36213 (G/C) [Intron variant]	PM: 1.35 (1.14, 1.60) ^R WT: I	0.0004*	0.84 (0.76, 0.92) ^A	0.0003*

Notes: Sex-stratified multivariate survey logistic regression of top interaction single nucleotide polymorphisms for asthma, adjusting for age, smoking status, province of recruitment, marital status, total personal income, retirement status, and first four principal components. *Statistically significant after adjusting for multiple comparisons (0.05/49 = $p \leq 0.001$). Functional consequence of each variant is presented in bold. Genes contain variants or are located within 500kb of variants.

Abbreviations: A, Additive; D, Dominant; R, Recessive; PM, Polymorphism; WT, Wild type; Chr, Chromosome; SNP, Single Nucleotide Polymorphism; A1/A2, Minor/Major Allele; OR, Odds ratio; CI, Confidence Interval.

intronic SNP rs2968822 indicated an increased risk of asthma in males, Intergenic and intronic SNPs rs2968801 and rs2864052 showed a protective effect against asthma risk in females. RTN4 (Reticulon 4), commonly known as a neurite outgrowth inhibitor (Nogo) with three major isoforms, most notably NOGO-B (RTN4B), has previously been shown to




Figure 3 The forest plots show the odds ratio distribution for the sex-specific variants associated with asthma from the sex-stratified analysis. (A) Males. (B) Females. Only significant SNPs after Bonferroni adjustment were plotted. The blue diamonds represent effect sizes (Odds ratio), and the lines extending from the blue diamonds are the 95% confidence intervals.

be expressed in the lung tissue and airway epithelium.^{40,41} Numerous animal and human studies have examined the role of RTN4 in asthma and immune responses.^{41–45} In an experimental study using mice, Wright et al⁴¹ demonstrated that RTN4 inhibits Th2-mediated inflammation in the lungs, airway epithelium, and smooth muscles by showing that eosinophils levels and Th2 cytokines (IL-13, IL-4, and IL-5) were higher in RTN-A/B knockout mice when compared to wild-type mice.⁴¹ In addition, RTN4B has also been shown to enhance the production of the inflammatory cytokine after the stimulation of various nucleic acid-sensing toll-like receptors, including TLR9, TLR3, and TLR7, in macrophages and serum.⁴³ Studies have shown that males produce higher TNF- α (pro-inflammatory cytokine) and IL-10 (anti-inflammatory cytokine) than females after the stimulation of TLR3 and TLR9 or viral infection in peripheral blood mononuclear cells (PBMCs).⁴⁶ However, a review article reported that older men or men with androgen deficiency express higher pro-inflammatory cytokines and lower anti-inflammatory cytokines than older or menopausal women.⁴⁷ These suggest that sex hormones may distinctively impact RTN4-mediated expressions of Th2 immune responses, toll-like receptors, and the production of inflammatory and suppressive cytokines in males and females.

An intergenic SNP rs9525931 near the Stress-Associated Endoplasmic Reticulum Protein 2 gene (SERP2) had a female-specific association with asthma. Variants near the SERP2 gene have previously been associated with FEV₁ and FEV₁/FVC ratio in a GWAS of lung function and COPD.⁴⁸ Furthermore, CpGs and SNPs mapped to the SERP2 gene have been reported to be associated with Alzheimer's disease in females.⁴⁹

An intronic variant rs6013213 (minor allele: A) in the NFATC2 gene was protective against the risk of asthma in males only. NFATC2, a member of the nuclear factor of activated T cells (NFAT) family, regulates Th cell immune response, differentiation, and the expression of induced cytokines such as IL-2, IL-3, IL-4, IL-10, and TNF-alpha.⁵⁰ Previous genome-wide association studies have associated several variants in the NFATC2 gene with asthma and allergic disease, including hay fever, eczema, and allergic rhinitis.^{51–55} NFATC2 mRNA levels in PBMCs of allergic asthmatics were higher than in healthy non-asthmatic controls.⁵⁶ Furthermore, NFATC2 mRNA has been demonstrated to correlate positively with IL-5-induced eosinophils in asthma.⁵⁶ Animal and human studies have shown that estrogen and cigarette smoke exposure increased NFAT mRNA expression in the airways of females.^{57,58} Furthermore, NFAT has been reported to have a regulatory function in estradiol-mediated MUC5AC mRNA and protein expression in airway epithelial cells.⁵⁸ MUC5AC is a marker for mucus production in the airway epithelial cells. Thus, it is reasonable to suggest that NFATC2 may influence sex-hormone-induced mucus secretion differently in male and female asthmatic bronchial epithelial cells.

The intergenic variant rs17071077 (minor allele: G) near the Neuromedin B Receptor (NMBR) gene exhibited a significant protective effect against the risk of asthma in males. NMBR is a G protein-coupled peptide receptor that binds with the regulatory neuropeptide Neuromedin B (NMB).⁵⁹ NMB/NMBR, widely expressed in the lungs, bronchoepithelial cells, pulmonary neuroendocrine cells, brain, and testis at the protein and mRNA levels,^{59–61} has been shown to promote fetal lung development.⁶² Several studies have shown that NMBR plays an innate immune defense role against respiratory viral infection by enhancing IFN-alpha and reducing the expression of IL-6.⁶³ Females have been shown to have greater Type I IFN responses and are less susceptible to viral infections than males.⁶⁴ We reasonably infer that any abnormality or polymorphic alterations in the NMBR gene may have a higher impact on males.

In our study, males with the intergenic variant rs254804 near the PEPD gene had a significantly higher risk of asthma susceptibility. PEPD is a gene that encodes prolydase, an enzyme involved in collagen metabolism, wound healing, inflammation, angiogenesis, and cell growth.⁶⁵ Numerous studies have shown that airway remodeling in asthma is largely due to the deposition of extracellular matrix protein, including collagen fibers, around the airway smooth muscle layers.^{66,67} This suggests that polymorphisms of the PEPD gene may contribute to airway remodeling. Patients with mutations in the PEPD gene and dysfunctional prolydase enzyme have been reported to have asthma and asthma-like airway disease.^{68,69} Studies have shown that serum prolydase activity was associated with bronchial asthma.^{70,71} Higher serum prolydase levels appear to be associated with higher oxidative stress and lower antioxidant levels.^{72,73} The homeostatic balance between the oxidant and antioxidant systems is impaired in asthma,⁷⁴ with oxidative stress increasing tissue damage, triggering the production of pro-inflammatory mediators, and exacerbating airway inflammation.⁷⁴ Studies have shown that males exhibit higher levels of oxidative stress, oxidative stress biomarkers, reactive oxygen species, and lower antioxidant capacity than females.⁷⁵

In our current study, an intron SNP rs6701638 in the Kinesin Family Member 26B gene (KIF26B) exhibited a significantly increased risk of asthma susceptibility in males but not females. Several variants of the genes in the 1q43-q44 region have been associated with asthma and atopic asthma.⁷⁶ White et al demonstrated that a variant in the KIF26B gene within the 1q43-44 region was associated with atopic asthma.⁷⁶ However, the association's p-value in that study was not significant after multiple testing adjustments. KIF26B is a target of the Wnt5a-Ror signaling pathway.⁷⁷ Studies have revealed that non-canonical Wnt5a-Ror signaling activation reduces KIF26B protein expression levels. Susman et al⁷⁷ demonstrated that the KIF26B protein expression level increased in embryonic fibroblast of Wnt5a null mice compared to the wild type. Dysregulation of Wnt5a signaling has been implicated in disrupted alveologenesis and the development of chronic lung disease, including asthma.⁷⁸ Furthermore, loss of Wnt5a in male mice resulted in abnormal reproductive organ development.⁷⁹ Taking together, we suggest that the alteration of the Wnt5a-Ror-KIF26B signaling pathway may have a deleterious impact in males and could play a crucial role in sex-specific asthma pathogenesis.

In our study, most sex-specific polymorphisms associated with asthma susceptibility may directly or indirectly interact with sex hormones in modulating immunoregulation, immune cell population, airway remodeling, oxidative stress, and lung function differently in males and females.

This study has strengths and limitations. A major strength was the size of the study population and the inclusion of lifestyle, socio-economic and socio-demographic factors in our analysis. Regarding the limitations, we used all asthma cases in our analysis without stratifying asthma by the age of onset (childhood onset and adulthood onset); this could potentially mask the identification of sex-specific loci associated with childhood and adulthood onset asthma. We reckon that misclassification of asthma is possible with self-reported physician-diagnosed asthma. Misclassification of the outcome variable may result in a bias towards the null. This may lead to an underestimation of the observed association. However, self-reported physician diagnosis of asthma has been used in genome-wide association and large population-based studies to identify sex-specific genetic and clinical characteristics of asthma.^{22,80} Our genome-wide SNP-by-sex interaction testing found no variants that met the genome-wide significance p-value criterion ($p \leq 5 \times 10^{-8}$). Genome-wide interaction testing generally requires a very large sample size to achieve adequate statistical power to detect genome-wide significant interaction p-value. There was no independent replication cohort in our study to corroborate our findings. Future replication of these findings in an independent cohort will increase their generalizability. Our work lacks functional follow-up and enrichment analysis of sex-specific genetic variants associated with asthma; hence we cannot demonstrate a causal relationship. Future experimental investigations can address the underlying biological processes for the sex differences.

In conclusion, we found evidence of sex-specific polymorphisms associated with asthma susceptibility in/near the KIF26B, RTN4, EPHB1, NMBR, SERP2, PEPD, and NFATC2 genes. Most of the identified loci may potentially play direct or indirect roles in the immune modulatory mechanism, oxidative stress, and airway remodeling. Future studies aimed at finding sex-specific expressions of these loci and the related pathways of action would give functional insights into their sex-specific involvement in asthma pathogenesis.

Data Sharing Statement

The datasets used in this study are not readily available. Data are only available from the Canadian Longitudinal Study on Aging (www.clsa-elcv.ca) for researchers who meet the criteria for access to de-identified CLSA data.

Ethics Approval

This CLSA project received ethics approval at two levels. Consent to participate was obtained for all participants under the CLSA harmonized multi-university ethics process approved by the Hamilton Integrated Research Ethics Board (HiREB), Hamilton Health Sciences/McMaster University. Simon Fraser University (SFU) was a participating institution in the CLSA data collection, and the SFU Office of Research Services Ethics Committee reviewed all consent material prior to data collection (SFU ORS #2018s0139). This study was also conducted following the principles of the Declaration of Helsinki, and ethical approval was obtained from the University of Alberta Health Research Ethics Board (Pro00091377_REN3) and Memorial University Health Research Ethics Board (HREB # 2019.072).

Acknowledgment

This research was made possible using the data/biospecimens collected by the Canadian Longitudinal Study on Aging (CLSA). The Government of Canada provides funding for the Canadian Longitudinal Study on Aging (CLSA) through the Canadian Institutes of Health Research (CIHR) under grant reference: LSA 94473 and the Canada Foundation for Innovation, as well as the following provinces, Newfoundland and Labrador, Nova Scotia, Quebec, Ontario, Manitoba, Alberta, and British Columbia. This research has been conducted using CLSA's Baseline Comprehensive Dataset version 4.0 and Genomics Dataset version 3.0, under Application Number 19CA006. The CLSA is led by Drs. Parminder Raina, Christina Wolfson, and Susan Kirkland. The opinions expressed in this manuscript are solely those of the author and do not represent those of the Canadian Longitudinal Study on Aging.

Funding

The Canadian Institutes of Health Research (CIHR) Catalyst Grant provided funding for this project (ACD 162989).

Disclosure

The authors declare no conflicts of interest in this work.

References

1. Safiri S, Carson-Chahhoud K, Karamzad N, et al. Prevalence, deaths, and disability-adjusted life-years due to asthma and its attributable risk factors in 204 countries and territories, 1990–2019. *Chest*. 2022;161(2):318–329. doi:10.1016/j.chest.2021.09.042
2. Report from the Canadian Chronic Disease Surveillance System: Asthma and Chronic Obstructive Pulmonary Disease (COPD) in Canada, 2018. Available from: <https://www.canada.ca/content/dam/phac-aspc/documents/services/publications/diseases-conditions/asthma-chronic-obstructive-pulmonary-disease-canada-2018/pub-eng.pdf>. Accessed May 5, 2023.
3. Postma DS. Gender differences in asthma development and progression. *Gen Med*. 2007;4:S133–S146. doi:10.1016/S1550-8579(07)80054-4
4. Chowdhury NU, Guntur VP, Newcomb DC, Wechsler ME. Sex and gender in asthma. *Eur Respir Rev*. 2021;30(162):210067. doi:10.1183/16000617.0067-2021
5. Ekpruke CD, Silveyra P. Sex differences in airway remodeling and inflammation: clinical and biological factors. *Front Allergy*. 2022;29(3):875295. doi:10.3389/falgy.2022.875295
6. Fuseini H, Newcomb DC. Mechanisms driving gender differences in asthma. *Curr Allergy Asthma Rep*. 2017;17(3):19. doi:10.1007/s11882-017-0686-1
7. Koleade A, Farrell J, Mugford G, Gao Z. Female-specific risk factors associated with risk of ACO (asthma COPD overlap) in aboriginal people. *J Asthma*. 2020;57(9):925–932. doi:10.1080/02770903.2019.1621890
8. Wadden D, Allwood Newhook LA, Twells L, Farrell J, Gao Z. Sex-specific association between childhood BMI trajectories and asthma phenotypes. *Int J Pediatr*. 2018;2:e9057435.
9. Gao Z, Dosman JA, Rennie DC, et al. Gender-specific associations between polymorphisms in the Toll-like receptor (TLR) genes and lung function among workers in swine operations. *J Toxicol Environ Health A*. 2018;81(22):1186–1198. doi:10.1080/15287394.2018.1544523
10. Amegadzie JE, Gamble JM, Farrell J. Gender differences in inhaled pharmacotherapy utilization in patients with obstructive airway diseases (OADs): a population-based study. *COPD*. 2020;30(15):2355–2366. doi:10.2147/COPD.S264580
11. Pignataro FS, Bonini M, Forgiione A, Melandri S, Usmani OS. Asthma and gender: the female lung. *Pharmacol Res*. 2017;1(119):384–390. doi:10.1016/j.phrs.2017.02.017
12. Chhabra SK, Chhabra P. Gender differences in perception of dyspnea, assessment of control, and quality of life in asthma. *J Asthma*. 2011;48(6):609–615. doi:10.3109/02770903.2011.587577
13. Zillmer LR, Gazzotti MR, Nascimento OA, Montealegre F, Fish J, Jardim JR. Gender differences in the perception of asthma and respiratory symptoms in a population sample of asthma patients in four Brazilian cities. *J Bras Pneumol*. 2014;40:591–598. doi:10.1590/S1806-37132014000600002
14. Blacquièrre MJ, Hylkema MN, Postma DS, Geerlings M, Timens W, Melgert BN. Airway inflammation and remodeling in two mouse models of asthma: comparison of males and females. *Int Arch Allergy Immunol*. 2010;153(2):173–181. doi:10.1159/000312635
15. Takeda M, Tanabe M, Ito W, et al. Gender difference in allergic airway remodelling and immunoglobulin production in mouse model of asthma. *Respirology*. 2013;18(5):797–806. doi:10.1111/resp.12078
16. McKenzie R, Burton MD, Royce SG, Tang MLK. Age and sex influences on airway hyperresponsiveness. *J Asthma*. 2010;47(6):651–654. doi:10.3109/02770901003692801
17. Gao Z, Dosman JA, Rennie DC, et al. Association of Toll-like receptor 2 gene polymorphisms with lung function in workers in swine operations. *Ann Allergy Asthma Immunol*. 2013;110(1):44–50.e1. doi:10.1016/j.anai.2012.11.003
18. Gao Z, Dosman JA, Rennie DC, et al. Effects of tumor necrosis factor (TNF) gene polymorphisms on the association between smoking and lung function among workers in swine operations. *J Toxicol Environ Health A*. 2021;84(13):536–552. doi:10.1080/15287394.2021.1896404
19. Gao Z, Rennie DC, Senthilselvan A. Allergic rhinitis and genetic components: focus on Toll-like receptors (TLRs) gene polymorphism. *Appl Clin Genet*. 2010;16:109–120.
20. El-Husseini ZW, Gosens R, Dekker F, Koppelman GH. The genetics of asthma and the promise of genomics-guided drug target discovery. *Lancet Respir Med*. 2020;8(10):1045–1056. doi:10.1016/S2213-2600(20)30363-5
21. Odimba U, Senthilselvan A, Farrell J, Gao Z. Current knowledge of asthma-COPD overlap (ACO) genetic risk factors, characteristics, and prognosis. *COPD*. 2021;18(5):585–595. doi:10.1080/15412555.2021.1980870

22. Espuela-Ortiz A, Herrera-Luis E, Lorenzo-Díaz F, et al. Role of sex on the genetic susceptibility to childhood asthma in latinos and African Americans. *J Pers Med*. 2021;11(11):1140. doi:10.3390/jpm11111140
23. Mersha TB, Martin LJ, Biagini Myers JM, et al. Genomic architecture of asthma differs by sex. *Genomics*. 2015;106(1):15–22. doi:10.1016/j.ygeno.2015.03.003
24. Gautam Y, Afanador Y, Abebe T, López JE, Mersha TB. Genome-wide analysis revealed sex-specific gene expression in asthmatics. *Hum Mol Genet*. 2019;28(15):2600–2614. doi:10.1093/hmg/ddz074
25. Myers RA, Scott NM, Gauderman WJ, et al. Genome-wide interaction studies reveal sex-specific asthma risk alleles. *Hum Mol Genet*. 2014;23(19):5251–5259. doi:10.1093/hmg/ddu222
26. Szczeklik W, Sanak M, Szczeklik A. Functional effects and gender association of COX-2 gene polymorphism G-765C in bronchial asthma. *J Allergy Clin Immunol*. 2004;114(2):248–253. doi:10.1016/j.jaci.2004.05.030
27. Forgetta V, Li R, Darmond-Zwaig C, et al. Cohort profile: genomic data for 26 622 individuals from the Canadian Longitudinal Study on Aging (CLSA). *BMJ Open*. 2022;12(3):e059021. doi:10.1136/bmjopen-2021-059021
28. Raina PS, Wolfson C, Kirkland SA, et al. The Canadian Longitudinal Study on Aging (CLSA). *Can J Aging*. 2009;28(3):221–229. doi:10.1017/S0714980809990055
29. Raina P, Wolfson C, Kirkland S, et al. Cohort profile: the Canadian Longitudinal Study on Aging (CLSA). *Int J Epidemiol*. 2019;48(6):1752–1753j. doi:10.1093/ije/dyz173
30. Purcell S, Chang C. General usage - PLINK 1.9 [Internet]; 2022. Available from: https://www.cog-genomics.org/plink/1.9/general_usage. Accessed May 5, 2023.
31. Chang CC, Chow CC, Tellier LC, Vattikuti S, Purcell SM, Lee JJ. Second-generation PLINK: rising to the challenge of larger and richer datasets. *GigaScience*. 2015;4:S13742. doi:10.1186/s13742-015-0047-8
32. Pruim RJ, Welch RP, Sanna S, et al. LocusZoom: regional visualization of genome-wide association scan results. *Bioinformatics*. 2010;26(18):2336–2337. doi:10.1093/bioinformatics/btq419
33. Ivanov AI, Romanovsky AA. Putative dual role of ephrin-Eph receptor interactions in inflammation. *IUBMB Life*. 2006;58(7):389–394. doi:10.1080/15216540600756004
34. Wu J, Luo H. Recent advances on T-cell regulation by receptor tyrosine kinases. *Curr Opin Hematol*. 2005;12(4):292–297. doi:10.1097/01.moh.0000166497.26397.9f
35. Darling TK, Lamb TJ. Emerging roles for Eph receptors and ephrin ligands in immunity. *Front Immunol*. 2019;10. doi:10.3389/fimmu.2019.01473
36. Wagener AH, Zwinderman AH, Luiten S, et al. The impact of allergic rhinitis and asthma on human nasal and bronchial epithelial gene expression. *PLoS One*. 2013;8(11):e80257. doi:10.1371/journal.pone.0080257
37. Wang Y, Thorin E, Luo H, et al. EPHB4 protein expression in vascular smooth muscle cells regulates their contractility, and EPHB4 deletion leads to hypotension in mice. *J Biol Chem*. 2015;290(22):14235–14244. doi:10.1074/jbc.M114.621615
38. Wang Y, Wu Z, Thorin E, et al. Estrogen and testosterone in concert with EFNB3 regulate vascular smooth muscle cell contractility and blood pressure. *Am J Physiol Heart Circ Physiol*. 2016;310(7):H861–H872. doi:10.1152/ajpheart.00873.2015
39. Luo H, Wu Z, Tremblay J, et al. Receptor tyrosine kinase Ephb6 regulates vascular smooth muscle contractility and modulates blood pressure in concert with sex hormones. *J Biol Chem*. 2012;287(9):6819–6829. doi:10.1074/jbc.M111.293365
40. Oertle T, Schwab ME. Nogo and its paRTNers. *Trends Cell Biol*. 2003;13(4):187–194. doi:10.1016/S0962-8924(03)00035-7
41. Wright PL, Yu J, Di YPP, et al. Epithelial reticulon 4B (Nogo-B) is an endogenous regulator of Th2-driven lung inflammation. *J Exp Med*. 2010;207(12):2595–2607. doi:10.1084/jem.20100786
42. Scatena R, Chupp GL, Perez MF, et al. The role of neurite outgrowth inhibitory protein-B in poorly controlled asthma. *Am Thoracic Society Int Conf Abst*. 2011:A1349–A1349. doi:10.1164/ajrccm-conference.2011.183.1_MeetingAbstracts.A1349
43. Kimura T, Endo S, Inui M, Saitoh SI, Miyake K, Takai T. Endoplasmic protein Nogo-B (RTN4-B) interacts with GRAMD4 and regulates TLR9-mediated innate immune responses. *J Immunol*. 2015;194(11):5426–5436. doi:10.4049/jimmunol.1402006
44. Xu W, Hong W, Shao Y, Ning Y, Cai Z, Li Q. Nogo-B regulates migration and contraction of airway smooth muscle cells by decreasing ARPC 2/3 and increasing MYL-9 expression. *Respir Res*. 2011;12(1):14. doi:10.1186/1465-9921-12-14
45. Ullah HMA, Elfadl AK, Park S, et al. Nogo-A is critical for pro-inflammatory gene regulation in myocytes and macrophages. *Cells*. 2021;10(2):282. doi:10.3390/cells10020282
46. Torcia MG, Nencioni L, Clemente AM, et al. Sex differences in the response to viral infections: TLR8 and TLR9 ligand stimulation induce higher IL10 production in males. *PLoS One*. 2012;7(6):e39853. doi:10.1371/journal.pone.0039853
47. Klein SL, Flanagan KL. Sex differences in immune responses. *Nat Rev Immunol*. 2016;16(10):626–638. doi:10.1038/nri.2016.90
48. Sakornsakolpat P, Prokopenko D, Lamontagne M, et al. Genetic landscape of chronic obstructive pulmonary disease identifies heterogeneous cell-type and phenotype associations. *Nat Genet*. 2019;51(3):494–505. doi:10.1038/s41588-018-0342-2
49. Zhang L, Young JI, Gomez L, et al. Sex-specific DNA methylation differences in Alzheimer's disease pathology. *Acta Neuropathol Commun*. 2021;9(1):77. doi:10.1186/s40478-021-01177-8
50. Fric J, Zelante T, Wong AYW, Mertes A, Yu HB, Ricciardi-Castagnoli P. NFAT control of innate immunity. *Blood*. 2012;120(7):1380–1389. doi:10.1182/blood-2012-02-404475
51. Han Y, Jia Q, Jahani PS, et al. Genome-wide analysis highlights contribution of immune system pathways to the genetic architecture of asthma. *Nat Commun*. 2020;11(1):1776. doi:10.1038/s41467-020-15649-3
52. Johansson Å, Rask-Andersen M, Karlsson T, Ek WE. Genome-wide association analysis of 350 000 Caucasians from the UK Biobank identifies novel loci for asthma, hay fever and eczema. *Hum Mol Genet*. 2019;28(23):4022–4041. doi:10.1093/hmg/ddz175
53. Waage J, Standl M, Curtin JA, et al. Genome-wide association and HLA fine-mapping studies identify risk loci and genetic pathways underlying allergic rhinitis. *Nat Genet*. 2018;50(8):1072–1080. doi:10.1038/s41588-018-0157-1
54. Ferreira MA, Vonk JM, Baurecht H, et al. Shared genetic origin of asthma, hay fever and eczema elucidates allergic disease biology. *Nat Genet*. 2017;49(12):1752–1757. doi:10.1038/ng.3985
55. Hinds DA, McMahon G, Kiefer AK, et al. A genome-wide association meta-analysis of self-reported allergy identifies shared and allergy-specific susceptibility loci. *Nat Genet*. 2013;45(8):907–911. doi:10.1038/ng.2686

56. Jakobi M, Kiefer A, Mirzakhani H, et al. Role of nuclear factor of activated T cells 2 (NFATc2) in allergic asthma. *Immun Inflamm Dis.* 2020;8(4):704–712. doi:10.1002/iid3.360
57. Tam A, Churg A, Wright JL, et al. Sex differences in airway remodeling in a mouse model of chronic obstructive pulmonary disease. *Am J Respir Crit Care Med.* 2016;193(8):825–834. doi:10.1164/rccm.201503-0487OC
58. Tam A, Wadsworth S, Dorscheid D, Man SFP, Sin DD. Estradiol increases mucus synthesis in bronchial epithelial cells. *PLoS One.* 2014;9(6):e100633. doi:10.1371/journal.pone.0100633
59. Jensen RT, Batten JF, Spindel ER, Benya RV. International union of pharmacology. LXVIII. Mammalian bombesin receptors: nomenclature, distribution, pharmacology, signaling, and functions in normal and disease states. *Pharmacol Rev.* 2008;60(1):1–42. doi:10.1124/pr.107.07108
60. Kaczyńska K, Zajac D, Wojciechowski P, Jampolska M. Regulatory peptides in asthma. *Int J Mol Sci.* 2021;22(24):13656. doi:10.3390/ijms222413656
61. Kane MA, Toi-Scott M, Johnson GL, Kelley KK, Boose D, Escobedo-Morse A. Bombesin-like peptide receptors in human bronchial epithelial cells. *Peptides.* 1996;17(1):111–118. doi:10.1016/0196-9781(95)02088-8
62. Shan L, Emanuel RL, Dewald D, et al. Bombesin-like peptide receptor gene expression, regulation, and function in fetal murine lung. *Am J Physiol Lung Cell Mol Physiol.* 2004;286(1):L165–L173. doi:10.1152/ajplung.00436.2002
63. Yang G, Huang H, Tang M, et al. Role of neuromedin B and its receptor in the innate immune responses against influenza A virus infection in vitro and in vivo. *Vet Res.* 2019;50(1):80. doi:10.1186/s13567-019-0695-2
64. Pujantell M, Altfeld M. Consequences of sex differences in Type I IFN responses for the regulation of antiviral immunity. *Front Immunol.* 2022;13. doi:10.3389/fimmu.2022.986840
65. Eni-Aganga I, Lanaghan ZM, Balasubramaniam M, Dash C, Pandhare J. PROLIDASE: a review from discovery to its role in health and disease. *Front Mol Biosci.* 2021;8. doi:10.3389/fmolb.2021.723003
66. Araujo BB, Dolnikoff M, Silva LFF, et al. Extracellular matrix components and regulators in the airway smooth muscle in asthma. *Eur Respir J.* 2008;32(1):61–69. doi:10.1183/09031936.00147807
67. Parameswaran K, Willems-Widyastuti A, Alagappan VKT, Radford K, Kranenburg AR, Sharma HS. Role of extracellular matrix and its regulators in human airway smooth muscle biology. *Cell Biochem Biophys.* 2006;44(1):139–146. doi:10.1385/CBB:44:1:139
68. Alrumayyan N, Slauenwhite D, McAlpine SM, et al. Prolidase deficiency, a rare inborn error of immunity, clinical phenotypes, immunological features, and proposed treatments in twins. *Allergy Asthma Clin Immunol.* 2022;18(1):17. doi:10.1186/s13223-022-00658-2
69. Wang H, Kurien BT, Lundgren D, et al. A nonsense mutation of PEPD in four Amish children with prolidase deficiency. *Am J Med Genet A.* 2006;140(6):580–585. doi:10.1002/ajmg.a.31134
70. Kaleli S, Akkaya A, Akdogan M, Gültekin F. The effects of different treatments on prolidase and antioxidant enzyme activities in patients with bronchial asthma. *Environ Toxicol Pharmacol.* 2006;22(1):35–39. doi:10.1016/j.etap.2005.11.001
71. Cakmak A, Zeyrek D, Atas A, Celik H, Aksoy N, Erel O. Serum prolidase activity and oxidative status in patients with bronchial asthma. *J Clin Lab Anal.* 2009;23(2):132–138. doi:10.1002/jcla.20303
72. Geçit İ, Aslan M, Gunes M, et al. Serum prolidase activity, oxidative stress, and nitric oxide levels in patients with bladder cancer. *J Cancer Res Clin Oncol.* 2012;138(5):739–743. doi:10.1007/s00432-011-1136-4
73. Piriñçi N, Kaba M, Geçit İ, et al. Serum prolidase activity, oxidative stress, and antioxidant enzyme levels in patients with renal cell carcinoma. *Toxicol Ind Health.* 2016;32(2):193–199. doi:10.1177/0748233713498924
74. Sahiner UM, Birben E, Erzurum S, Sackesen C, Kalayci O. Oxidative stress in asthma. *World Allergy Organ J.* 2011;4(10):151–158. doi:10.1097/WHOX.0b013e318232389e
75. Kander MC, Cui Y, Liu Z. Gender difference in oxidative stress: a new look at the mechanisms for cardiovascular diseases. *J Cell Mol Med.* 2017;21(5):1024–1032. doi:10.1111/jcmm.13038
76. White JH, Chiano M, Wigglesworth M, et al. Identification of a novel asthma susceptibility gene on chromosome 1qter and its functional evaluation. *Hum Mol Genet.* 2008;17(13):1890–1903. doi:10.1093/hmg/ddn087
77. Susman MW, Karuna EP, Kunz RC, et al. Kinesin superfamily protein Kif26b links Wnt5a-Ror signaling to the control of cell and tissue behaviors in vertebrates. *eLife.* 2017;6:e26509. doi:10.7554/eLife.26509
78. Li C, Smith SM, Peinado N, et al. WNT5a-ROR signaling is essential for alveologenesis. *Cells.* 2020;9(2):384. doi:10.3390/cells9020384
79. Chawengsaksohak K, Svingen T, Ng ET, et al. Loss of Wnt5a disrupts primordial germ cell migration and male sexual development in mice¹. *Biol Reprod.* 2012;86(1):1–12. doi:10.1095/biolreprod.111.095232
80. Veerasingam E, Gao Z, Beach J, Senthilselvan A. Sex-specific characteristics for the coexistence of asthma and COPD in the Canadian population: a cross-sectional analysis of CLSA data. *J Asthma.* 2022;4:1–22.