



UNIVERSITY OF CALIFORNIA PRESS  
JOURNALS + DIGITAL PUBLISHING



---

Tonic Immobility Responses of Herring Gull Chicks

Author(s): William Montevecchi

Source: *The Condor*, Vol. 80, No. 2 (Summer, 1978), pp. 248-249

Published by: [University of California Press](#) on behalf of the [Cooper Ornithological Society](#)

Stable URL: <http://www.jstor.org/stable/1367931>

Accessed: 25/06/2013 13:26

---

Your use of the JSTOR archive indicates your acceptance of the Terms & Conditions of Use, available at  
<http://www.jstor.org/page/info/about/policies/terms.jsp>

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact support@jstor.org.



University of California Press and Cooper Ornithological Society are collaborating with JSTOR to digitize, preserve and extend access to *The Condor*.

<http://www.jstor.org>

(Table 1), thus resulting in a stable proportion of calcium in the bone. Depletion of cortical bone during egg laying was not evident. The geese in this study did not appear to suffer from a calcium deficiency. Medullary bone seemed to be sufficient to provide the calcium for the average clutch, because the femur and tibiotarsus bone and ash weights, and their calcium contents were greater at the end of egg laying than during the non-reproductive season. However, the contribution of minerals from other bones, especially ribs, sternum, vertebrae and pelvic girdle should be examined for a definitive conclusion (Taylor and Moore 1954).

This study was financed, in part, by a University of California, Davis, Agricultural Experiment Station faculty grant to D. G. Raveling. The U.S. Fish and Wildlife Service provided transportation, food and logistic support in Alaska. We wish to express our thanks to E. Collins and C. J. Lensink, U.S. Fish and Wildlife Service, and M. R. Petersen, M. L. Wege, and M. R. McLandress for field and laboratory assistance.

#### LITERATURE CITED

- CHEATUM, E. L. 1949. Bone marrow as an index of malnutrition in deer. N.Y. Conservationist 3:19-22.
- GRAU, C. R. 1976. Ring structure of avian egg yolk. Poultry Sci. 55:1418-1422.
- HANSON, H. C. 1962. The dynamics of condition factors in Canada geese and their relation to seasonal stresses. Arct. Inst. N. Am. Tech. Pap. No. 12.
- MICKELSON, P. G. 1975. Breeding biology of cackling geese and associated species on the Yukon-Kuskokwim Delta, Alaska. Wildl. Monogr. No. 45.
- NELSON, U. C. AND H. A. HANSEN. 1959. The cackling goose—its migration and management. N. Am. Wildl. and Natural Resources Conf. Trans. 24:174-186.
- SIMKISS, K. 1967. Calcium in reproductive physiology. Reinhold Publ., N.Y.
- TAYLOR, T. G. AND J. H. MOORE. 1954. Skeletal depletion in hens laying on a low calcium diet. Br. J. Nutr. 8:112-124.
- TAYLOR, T. G. AND J. H. MOORE. 1956. The effect of calcium depletion on the chemical composition of bone minerals in laying hens. Br. J. Nutr. 10:250-263.
- TAYLOR, T. G., T. R. MORRIS AND F. HERTELENDY. 1962. The effect of pituitary hormones on ovulation in calcium-deficient pullets. Vet. Rec. 74:123-125.
- TAYLOR, T. G., K. SIMKISS AND D. A. STRINGER. 1971. The skeleton: its structure and metabolism, pp. 621-640. In D. J. Bell and B. M. Freeman [eds.], Physiology and biochemistry of the domestic fowl. Vol. 2. Academic Press, N.Y.
- URIST, M. R. 1959. The effects of calcium deprivation upon the blood, adrenal cortex, ovary, and skeleton in domestic fowl. Recent Prog. Horm. Res. 15:455-477.

Address of first and third authors: Division of Wildlife and Fisheries Biology, University of California, Davis, CA 95616. Address of second author: Department of Avian Sciences, University of California, Davis, CA 95616. Accepted for publication 23 February 1978.

Condor, 80:248-249  
© The Cooper Ornithological Society 1978

## TONIC IMMOBILITY RESPONSES OF HERRING GULL CHICKS

WILLIAM MONTEVECCHI

Following physical restraint, many birds exhibit a state of pronounced immobility (Armstrong 1947). Such tonic immobility (TI), which may be accompanied by eye closure, leg extension and muscle tremors, has often been referred to as "death feign-

ing." Animals in this state are sensitive to external stimulation. Immobility of captured prey can inhibit the attack behavior of a predator and may allow for escape, if the predator is distracted (Sargeant and Eberhardt 1975, Gallup 1977). With very few exceptions (e.g. Sargeant and Eberhardt 1975) almost all studies of tonic immobility in birds have been conducted in the laboratory with domesticated species.

During June and July 1976, 54 Herring Gull chicks (*Larus argentatus*) and eight Great Black-backed Gull chicks (*L. marinus*) on Little Bell Island, Con-

TABLE 1. Tonic immobility reactivity of Herring Gull chicks in relation to weight.

Group	Weight range (g)	N	TI duration (s) ( $\bar{x} \pm S.E. s$ )	No response (< 5 s)	Maximum response (600 s)
1	50-190	6	12.3 $\pm$ 9.6	5 (83%)	0
2	205-340	7	70.4 $\pm$ 37.7	2 (29%)	0
1+2	50-340	13	43.6 $\pm$ 21.7	7 (54%)	0
3	420-580	11	293.2 $\pm$ 69.3	1 (9%)	3 (27%)
4	600-675	8	392.3 $\pm$ 82.6	0	4 (50%)
3+4	420-675	19	387.5 $\pm$ 51.7	1 (5%)	7 (37%)
5	710-790	13	457.8 $\pm$ 63.8	0	9 (69%)
6	830-1175	9	317.7 $\pm$ 93.0	0	4 (44%)
5+6	710-1175	22	400.5 $\pm$ 54.3	0	13 (59%)
<i>L. marinus</i>	490-1200	8	410.0 $\pm$ 87.5	0	4 (50%)



FIGURE 1. Herring Gull chick immobilized following manual restraint.

ception Bay, Newfoundland ( $47^{\circ}38'N$ ,  $52^{\circ}27'W$ , see Noseworthy and Lien (1976) for a description of the island) were tested for TI reactions. Each chick was placed on its back on level ground near the point of capture and restrained (not squeezed) with both hands (thumbs on the sternum) for 15 s after which the investigator removed his hands and sat back about 1 m from the chick. Tests ended when a chick got to its feet or 10 min elapsed; chicks were then weighed (Pesola scales) and banded. Most chicks had not been handled previously.

Older (heavier) chicks showed proportionately more and longer TI responses than younger ones (Table 1). The average duration of immobility for Weight Group 1 was not significantly different from that of Group 2 ( $F = 1.92$ ,  $P = 0.19$ ) but was significantly shorter than that of Group 3 ( $F = 8.66$ ,  $P < 0.01$ ). The mean durations of Group 2 and of Groups 1 and 2 combined were both significantly shorter than that of Group 3 ( $F = 5.78$ ,  $P < 0.05$  and  $F = 13.59$ ,  $P < 0.001$ , respectively). Groups 4, 5 and 6 were not significantly different ( $F = 1.66$ ,  $P = 0.21$ ). TI duration and weight showed a significant positive correlation ( $r = +0.50$ ,  $P < 0.001$ ), and the linear regression of TI

duration on chick weight was also significant ( $Y = 54X + 432.62$ ,  $F = 17.43$ ,  $P < 0.001$ ).

As a possible result of developing fear motivation (Ratner and Thompson 1960) domestic chicks show increasing susceptibility to TI inductions with age. These reactions begin to appear around 7–10 days posthatch, an age when thermoregulatory mechanisms are developed and heart rate approaches adult levels (Braud and Ginsburg 1973). Younger gull chicks showed shorter TI reactions (righted themselves sooner) than older ones; this is interesting because young Herring Gull chicks are less mobile and slower to move than older ones following displacements from the nest (Noseworthy and Lien 1976). Weight data of the Herring Gull chicks indicate that TI reactivity begins around 15–20 days posthatch (420–580 g, Kadlec et al. 1969).

Mortality among larid chicks is greatest during the first week posthatch and appears to be largely due to pecking by adults (e.g. Kadlec et al. 1969). TI reactivity begins and increases as mortality rates are decreasing during the nestling period. Ventral TI inductions, unlike the dorsal ones used in the present tests, are effective in eliciting TI in neonatal domestic chicks (Braud and Ginsburg 1973). The potential anti-predator adaptiveness of TI responses remains to be studied. Tonic immobility may be of interest to some ornithologists who may find the dorsal induction procedure useful when banding chicks.

I am grateful to Bruce Bursey for help with the tests. Work was supported by a Faculty Research Grant from Memorial University of Newfoundland and a National Research Council of Canada Grant awarded to Dr. Jon Lien.

#### LITERATURE CITED

- ARMSTRONG, E. A. 1947. Bird display and behavior. Lindsey Press, London.
- BRAUD, W. G. AND H. J. GINSBURG. 1973. Immobility reactions in the domestic fowl (*Gallus gallus*) less than 7 days old: Resolution of a paradox. *Anim. Behav.* 21:104–107.
- GALLUP, G. G., JR. 1977. Tonic immobility: The role of fear and predation. *Psychol. Rec.* 1:41–61.
- KADLEC, J. A., W. H. DRURY, JR. AND D. K. ONION. 1969. Growth and mortality of Herring Gull chicks. *Bird-Banding* 40:22–223.
- NOSEWORTHY, C. M. AND J. LIEN. 1976. Ontogeny of nesting habitat recognition and preference in neonatal Herring Gull chicks, *Larus argentatus* Pontoppidan. *Anim. Behav.* 24:637–651.
- RATNER, S. C. AND R. W. THOMPSON. 1960. Immobility reactions (fear) of domestic fowl as a function of age and prior experience. *Anim. Behav.* 8:186–191.
- SARGEANT, A. B. AND L. E. EBERHARDT. 1975. Death feigning by ducks in response to predation by Red Foxes (*Vulpes fulva*). *Am. Midl. Nat.* 94:108–119.

Department of Psychology, Memorial University of Newfoundland, St. John's, Newfoundland A1C 5S7 Canada. Accepted for publication 6 October 1977.