


"THE METAZOAN PARASITES OF THREE SPECIES OF
SCOTERS (MELANITTA PERSPICILLATA (LINNAEUS,
1758), M. NIGRA (LINNAEUS, 1758) AND
M. FUSCA (LINNAEUS, 1758))"

CENTRE FOR NEWFOUNDLAND STUDIES

**TOTAL OF 10 PAGES ONLY
MAY BE XEROXED**

(Without Author's Permission)

CHARLES EDWARD BOURGEOIS



000237



The Metazoan parasites of
three species of scoters (*Melanitta perspicillata*
(Linnaeus, 1758), *M. nigra* (Linnaeus, 1758) and
M. fusca (Linnaeus, 1758))

by



Charles Edward Bourgeois

A thesis submitted in partial fulfillment
of the requirements for the degree of
Master of Science.

Department of Biology
Memorial University of Newfoundland
St. John's, Newfoundland, Canada

December, 1980

ABSTRACT

One hundred and seventy-five (94 surf, 36 white-winged and 45 black) scoters from four localities (British Columbia, New Brunswick, Labrador and Norway) were examined for metazoan parasites, with 91% (159 birds: 86 surf, 33 white-winged and 40 black scoters) being infected. Approximately 55 species of parasites were found (43 from the surf, 29 from the white-winged and 34 from the black), including 45 which were new host records. There were 4 new North American records, 4 new Canadian records and 1 new European record. A checklist of parasites harboured by these birds is presented.

The number and percent of birds infected, range of numbers and mean number of parasites per infected bird are given. Also for each host species the number and percent of each sex and age class infected by individual parasites is given. Parasite species are discussed individually with regard to topics such as location within host, new host records and minor variations from their original description.

The ecology of the parasites is discussed; the preferred habitat, their geographical distribution and their preference for marine or freshwater environments are presented. A short discussion of host specificity is included and the parasites of each host are categorized. Six parasites were found to occur across the geographic area sampled, with 23 helminths cycling through the marine environment and 19 through the freshwater environment. Throughout the work the parasite burdens of each host species are contrasted. In addition the number and percent of species occurring in each geographic locale is compared and contrasted. Cluster analysis revealed that

neither individual birds of a species nor species of birds could be distinguished using parasite fauna and geographical locality as criteria.

One species of parasite, *Polymorphus botulus*, was noted to cause damage to its host but was not considered to be lethal.

ACKNOWLEDGEMENTS

I wish to express my sincere thanks to Dr. W. Threlfall, for his helpful suggestions, editorial assistance and encouragement throughout this study and preparation of this manuscript. Furthermore, I am indebted to him for the New Brunswick samples.

I am grateful to Messrs. C. V. Chandra and O. Blagdon for their assistance in collecting the Labrador sample of birds, in very poor weather conditions. The British Columbia birds were provided by K. Vermeer to whom I express my sincere appreciation and to D. Cannings for packing and shipping them. R. T. Barrett kindly provided the Norwegian birds.

The author expresses his gratitude to Dr. R. D. Price for verifying the *Holomonopon* identifications, and Dr. H. L. Ching for her advice, and the specimens, which she so freely gave. The author expresses his thanks to Prof. S. DeBlock for his assistance in verifying the authors identifications of the microphallid trematodes.

Also, the author wishes to thank Mr. D. Paul for helping procure birds from the R.C.M.P. and his issuance of Scientific Kill Permit # ASK-25-78 which aided this study greatly.

The author wishes to thank Mr. R. Ficken and Mrs. C. Emerson for their assistance.

The author wishes to acknowledge Mary-Louise Porter (Dub) for her assistance in editing this manuscript.

The Author wishes to acknowledge the aid given by Doctors A. Whittick and A. Bush with regard to their advice on cluster analysis.

This study was supported by a Provincial Government Bursary to the Author, and a Northern Sciences Training Grant and NSERC Grant (A-3500) to Dr. W. Threlfall.

LIST OF CONTENTS

<u>Title:</u>	<u>Page:</u>
Abstract	i
Acknowledgements	iii
List of Tables	v
List of Figures	vii
List of Appendices	viii
Introduction	1
Methods and Materials	6
Results and Discussion	16.
Trematoda	16
Cestoda	33
Nematoda	36
Acanthocephala	44
Mallophaga	47
Acarina	53
General Discussion	54
Parasite Distribution	54
Host Specificity	67
Life-Cycles	72
The Influence of Age and Sex of the Host on Parasite Burden	74
Summary	78A
Literature Cited	79

LIST OF TABLES

<u>No.:</u>		<u>Page:</u>
Table 1.	Collection dates and localities for scoters examined.	7-9*
Table 2.	Sex and age of scoters examined.	11
Table 3.	Numbers of scoters lacking various organs, due to a variety of reasons.	12
Table 4.	Details of infection of surf scoters, white-winged scoters and black scoters with Trematoda.	17-20
Table 5.	Range of measurements of <i>Gymnophallus</i> sp. (type 1) and the gymnophallid sp. found in this study.	28
Table 6.	Comparison of measurements of <i>Cotylurus strigeoides</i> and <i>Cotylurus raquet</i> with specimens recovered in this study.	32
Table 7.	Details of infection of surf, white-winged and black scoters with Cestoda.	35
Table 8.	Details of infection of surf, white-winged and black scoters with Nematoda.	37-38
Table 9.	Details of infection of surf, white-winged and black scoters with Acanthocephala.	46
Table 10.	Details of infestation of surf scoters with Mallophaga.	48
Table 11.	Details of infestation of white-winged scoters with Mallophaga.	49
Table 12.	Details of infestation of black scoters with Mallophaga.	50
Table 13.	Details of infection of scoters and individual sample areas by major helminth groups.	55
Table 14.	Number of species of each major helminth group in the different sample areas.	56

<u>No.:</u>		<u>Page:</u>
Table 15.	Details of parasite locations and parasite communities possible in scoters examined in this study.	58
Table 16.	Geographical distribution of helminth parasites recovered from scoters in this study.	59-60
Table 17.	New host records noted in this study.	62
Table 18.	Categorization of parasites recovered in this study.	69-71
Table 19.	Environment in which parasite life-cycles are completed.	73
Table 20.	Parasites found to occur in only immature or adult birds.	75

LIST OF FIGURES

NO.:		FOLLOWING PAGE:
Figure 1.	Showing the similarity in the morphology of <i>A. T. elegans</i> and <i>B. N. attenuatus</i> .	23
Figure 2:	Showing Nodular taeniasis on the intestine of a black scoter.	45
Figure 3:	Cluster analysis comparing individual surf scoters with respect to presence of helminth species.	64
Figure 4:	Cluster analysis comparing individual white-winged scoters with respect to presence of helminth species.	65
Figure 5:	Cluster analysis comparing individual black scoters with respect to presence of helminth species.	66

LIST OF APPENDICIES

<u>NO.:</u>		<u>PAGE:</u>
Appendix 1.	Checklist, compiled from original papers and other checklists, of Metazoan parasites recovered from the Surf Scoter, White-Winged Scoter and Black Scoter (excluding Acarina.)	91
Appendix 2.	Comparisons of the helminth recovered from 6 <i>M. f. fusca</i> to those recovered from the 30 <i>M. f. deglandi</i> .	110
Appendix 3.	Details of infection of Scoters with helminths by age and sex.	112

INTRODUCTION

The genus *Melanitta* Boie, 1822 contains the diving ducks commonly known as scoters. These birds are members of the Mergal, the tribe of ducks, commonly referred to as sea ducks, which spend most of their time in the marine environment only frequenting inland waters to reproduce. Palmer (1976) notes that there are three valid species in the genus namely, *M. perspicillata* (Linnaeus, 1758) the surf scoter, *M. nigra* (Linnaeus, 1758) the black scoter and *M. fusca* (Linnaeus, 1758) the white-winged scoter, with five recognized subspecies in the genus. The scoters are Holarctic in distribution, with the surf scoter being indigenous to North America and having no recognizable subspecies. This bird breeds from Alaska through the northern boreal forest to Labrador. Its wintering areas are the eastern and western coasts of North America south to Florida and California respectively.

The black scoter has two recognizable subspecies, *M. n. americana* the Nearctic and eastern Siberian race and *M. n. nigra* the western Palearctic race. *Melanitta n. americana* breeds in Alaska, Kodiak Island and the interior of the Ungava Peninsula as well as Grand Lake in Newfoundland. The wintering range of this bird is similar to that of *M. perspicillata*. *Melanitta n. nigra* breeds in Iceland, Britain,

Ireland, Sweden, Finland and Western U.S.S.R. It appears that each breeding population of *M. n. nigra* moves south for the winter with wintering populations commencing around Norway and extending south to the Ivory Coast of Africa (Johnsgard, 1978; Palmer, 1976).

The white-winged scoter has three recognized subspecies, *M. f. deglandi* the American white-winged scoter, *M. f. fusca* the European white-winged scoter and *M. f. stejnegeri* the Asiatic white-winged scoter. *Melanitta f. deglandi* breeds in North America from western Ontario to British Columbia and up into Alaska. This subspecies winters from the Alaskan coast to California in the west and from Newfoundland to Florida in the east. *Melanitta f. fusca* breeds across northern Europe and Asia from Scandinavia to the Yenisei River in Western Siberia, and winters in the Atlantic Ocean from Norway to Spain and on the Caspian Sea. *Melanitta f. stejnegeri* breeds in eastern Asia from the Altai to Kamchatka Peninsula and winters on the coast of eastern Asia to Japan and China.

Prior to 1973, the black scoter was placed in the genus *Oidemia* (Flemming, 1822) because of a morphological difference between this species and the other two scoters. The black scoter has a simple syrinx, while the other two species both have a bulbous enlargement of the trachea. However, they have a similar biology and are very similar in appearance and are now placed in a single genus; a fact which is no longer a

source of disagreement. If indeed the three scoters are quite similar then very little difference would be expected in the parasite burdens of birds collected from the same geographical area. With regard to the taxonomy of the genus *Melanitta* some authors consider *M. f. fusca* to be a separate species (the velvet scoter) and *M. deglandi* to be the white-winged scoter thus creating four species in the genus. However, the author chooses to follow the works of Delacour (1959) and Palmer (1976) which recognize only three valid species.

This study was initiated in 1978 to compare the parasite burdens of these three closely related species of ducks. There are many reports of parasites collected from the black and white-winged scoters but relatively few from the surf scoter (Appendix 1). However, these reports of parasites from scoters tend to be qualitative in nature rather than quantitative. Buscher (1965) compared the parasite fauna of three species of ducks along a North American migration route and found that the burden changed from the northern to southern environment. To date, no workers have looked at differences in the parasite burden of three closely related bird species along an east-west gradient. Thus, the author was curious to see if the parasite burdens did differ longitudinally, as a result of geographical distribution of the scoters.

Rhode (1964, *vide* Graham, 1966) has pointed out that just listing the hosts of a helminth tends to lump frequently parasitized hosts with those of accidental origin. Hence, when a single bird is autopsied for parasites, very little insight is obtained about the host/parasite relationships. Conversely, when a large sample of birds is surveyed, it is possible to deduce more information on the host/parasite relationships e.g. host specificity of the parasites, food relations of the host, etc.. Various workers have recognized the above (e.g. Dogiel 1964; Gallimore 1964; McDonald 1969.), with the first two workers devising systems that categorized the host as main, auxiliary, rare, etc.. The system devised by McDonald (1969) differs in that he lists the parasite as being accidental, rare, infrequent, frequent or common in waterfowl; but doesn't give any criteria for his grouping. Given a large sample of birds one can place parasites into McDonald's categories utilizing stated criteria for each category, this being one of the goals of the present work.

BezuBik (1956), Cornwell and Cowan (1963) and Buscher (1965) concluded that the diet of a bird is the single most important factor in determining the bird's helminth fauna. Scoters are omnivorous birds that feed predominantly on animal matter, and thus, should be more adequately described as carnivores that consume some plant material. Cottam (1939) examined the stomachs of 168 surf, 819 white-winged and 124 black scoters and found animal material to make up 88%, 94% and 90% of their diets res-

pectively. Molluscs made up 60.8%, 75.3% and 65.2% of the animal material respectively. Moulting anatids eat-predominately vegetable matter (Cornwell and Cowan, 1963; Buscher, 1965). In addition, Buscher (1965) states, "The incidence of helminths in adult ducks ... was greatly reduced when the ducks were flightless during the moulting period."

The objectives of this work are thus, to determine and quantify the parasitic burden of three closely related species of ducks along an east-west gradient; to define the status of each parasite species found in these ducks and to relate the parasites found to the ecology of the ducks.

METHODS AND MATERIALS

A total of 175 scoters (94, *M. perspicillata*, 45 *M. nigra* and 36 *M. fusca*) were collected across large parts of their respective geographical ranges, from 1973-1978. Data on the locality and date of collection of these birds is summarized in Table 1. The birds were all collected with a 12-gauge shotgun using various sizes of shot. The majority of birds were collected during their moulting season thus accounting for the large number of males (i.e. males undertake a moult migration to the sea where they congregate in large flocks while the females remain on the breeding grounds with their young).

The New Brunswick birds were obtained from the Royal Canadian Mounted Police, after they had been confiscated from hunters who were taking them illegally. As a result, no information is available on how soon they were frozen after collection. All other birds were either deep frozen or preserved in 10% formalin within six hours of their collection. The Labrador sample of birds were the only specimens collected by the author.

Birds were aged using plumage (Carney and Geis, 1964) and secondary sexual characters, and/or the presence or absence of a Bursa of Fabricius. Two age classes were recognized in this study: (1) immature (prebreeding bird) - a bird that was collected in its first or second calendar year, that possessed

TABLE 1. COLLECTION DATES AND LOCALITIES FOR SCOTERS EXAMINED.

BIRD SPECIES	LOCAL	YEAR	MONTH								TOTAL
			JAN.	FEB.	APR./MAY	JUN.	JUL.	AUG.	SEPT.	SUMMER*	
Surf Scoter	New Brunswick -Shippegah	1973	-	-	25	-	-	-	-	-	25
	British Columbia										
	-Roberts Bank	1974	-	-	-	-	-	-	-	11	11
	-Iona	1974	14	-	-	-	-	-	-	-	14
	-Boundary Bay	1974	-	5	-	-	9	1	-	-	15
	-'?'	1974	-	-	-	-	1	-	-	4	5
	Newfoundland										
	-Porcupine Strand (Lab.)	1978	-	-	-	-	-	22	-	-	22
	-Porcupine Hbr. (Lab.)	1978	-	-	-	-	-	-	2	-	2

TABLE 1. CONTINUED.

BIRD SPECIES	LOCAL	YEAR	MONTH							SUMMER*	TOTAL
			JAN.	FEB.	APR./MAY	JUN.	JUL.	AUG.	SEPT.		
White-winged scoter	New Brunswick -Shippegan	1973	-	-	3	-	-	-	-	-	3
	British Columbia										
	-Roberts Bank	1974	-	-	-	-	-	-	-	4	4
	-Boundary Bay	1974	-	3	-	1	1	-	-	-	5
	-Salt Spring Islands	1974	-	-	-	-	-	-	-	5	5
	-Salt Spring Islands	1978	-	8	-	-	-	-	-	-	8
	Newfoundland										
	-Porcupine Strand (Lab.)	1978	-	-	-	-	-	5	-	-	5
	Norway										
	-Balsfjord Tromsø	1978	-	6	-	-	-	-	-	-	6

TABLE 1. CONTINUED.

BIRD SPECIES	LOCAL	YEAR	MONTH								TOTAL
			JAN.	FEB.	APR./MAY	JUN.	JUL.	AUG.	SEPT.	SUMMER*	
Black Scoter	New Brunswick -Shippegan	1973	-	-	27	-	-	-	-	-	27
	British Columbia -Roberts Bank	1974	-	-	-	-	-	-	-	10	10
	-Iona	1974	-	-	-	-	-	-	-	3	3
	Newfoundland -Porcupine Strand (Lab.)	1978	-	-	-	-	-	5	-	-	5

*No further information on date of collection is possible.

a Bursa of Fabricius and juvenile plumage; (2) adult - a bird that was two calendar years of age or older that lacked a Bursa of Fabricius and bore adult plumage. Birds were sexed using the appearance of their genital organs, when possible, and secondary sexual characteristics. Sexes and ages of the birds examined in this study are given in Table 2.

Necropsies were carried out as described by Turner (1974). In addition to necropsying the specimens the weight (g.), culmen length and depth, wing length, tail length and tarsus length was obtained (all measurements in mm.), whenever possible. Due to the birds being treated in a variety of ways by the various collectors not all birds were whole (Table 3). In addition, 27 of the British Columbia surf scoters had their food contents removed (25 had their proventriculus and gizzard emptied while 2 had their gizzard emptied). Of the British Columbia black scoters, 7 had the contents of their proventriculus and gizzard removed. Of the 1974 British Columbia white-winged scoters, 4 had their gizzard emptied and an additional 4 had their proventriculus and gizzard emptied.

The removal of food stuffs was carried out in such a way as to leave the lining of the proventriculus intact and while some parasites were undoubtedly lost, the author believes such losses were minimal. The helminth burdens recorded represent a minimum, especially in those birds that had various portions of their viscera removed.

TABLE 2
SEX AND AGE OF SCOTERS EXAMINED.

BIRD	LOCALITY COLLT'D	FEMALE		MALE		SEX UNKNOWN	SAMPLE SIZE
		ADULT	IMMATURE	ADULT	IMMATURE		
Surf Scoters	New Brunswick	2	0	23	0	0	25
	British Columbia	14	1	22	8	0	45
	Labrador	0	5	11	8	0	24
	Total	16	6	56	16	0	94
White-Winged Scoters	New Brunswick	0	0	3	0	0	3
	British Columbia	2	2	16	1	1	22
	Labrador	0	0	3	2	0	5
	Norway	1	0	4	1	0	6
Total	3	2	26	4	1	36	
Black scoters	New Brunswick	7	0	20	0	0	27
	British Columbia	1	2	10	0	0	13
	Labrador	0	2	2	1	0	5
	Total	8	4	32	1	0	45

NOTE: The one bird that could not be sexed was missing its head, and its genital organs were not developed.

TABLE 3. NUMBERS OF SCOTERS LACKING VARIOUS ORGANS, DUE TO A VARIETY OF REASONS.

BIRD SPECIES	LOCAL	A	B	C	D	E	F	G	H	I	J	K	L
Surf scoter	British Columbia	-	2	7	18	12	31	32	-	-	-	-	-
	Labrador Porcupine Hbr.	2	-	-	-	-	-	-	2	2	2	2	1
Black scoter	British Columbia	-	-	-	-	1	12	12	-	-	-	-	-
White-winged scoter	British Columbia	1	1	4	-	3	6	6	-	-	-	-	-
	Norway	-	-	-	-	-	6	6	6	6	-	6	-

A - Esophagus
 B - Proventriculus
 C - Gizzard
 D - Liver
 E - Gall bladder
 F - Spleen
 G - Pancreas
 H - Lungs
 I - Kidneys
 J - Gonads
 K - Cloaca
 L - Bursa

Removal of ectoparasites was in accordance with Eveleigh (1974). After this procedure was completed the birds were rinsed, with a jet of water, into a sieve and any debris collected was examined for ectoparasites. Only a selected sample of each species of bird that was examined for ectoparasites was included in the results (48 surf scoters, 17 white-winged scoters and 17 black scoters), due to the fact that some birds were poorly wrapped and had almost certainly lost some of their ectoparasites.

The black scoters necropsied in this study were all *M. n. americana* and the white-winged scoters save 6 were *M. f. deglandi*, with the former being *M. f. fusca* (the Norwegian birds). Appendix 2 compares the parasites recovered from the 6 *M. f. fusca* to those recovered from the 30 *M. f. deglandi*.

The methods of fixation, preservation, staining and mounting of helminths are those set forth in Andrews (1974), while ectoparasites were treated utilizing the method outlined by Palma (1978). A scanning electron microscope was used to aid in the identification of one species of trematode, whilst it proved necessary to section another species.

Infections are recorded as prevalence (percentage of ducks infected) and intensity (average number of parasites per infected duck).

The number of schistosomes and cestodes collected refer to minimum numbers; this is a result of only searching for schistosomes until specimens were recovered i.e. presence or absence. Cestode numbers were arrived at by counting the number of scolices present and placing that number between the nearest two numbers on a five/digit scale (i.e. if 42 scolices were recovered from a host, it was assumed that the bird had between 40-45 cestodes). However, due to their small size, destrobilization and digestion of the cestodes by the host on its death, some may have been overlooked. The number of *Fimbriaroides* present was arrived at by counting the number of fan shaped neck segments present. Parasites were categorized as immature or adult by meristic characters and/or presence of ova in females.

Due to taxonomic confusion within the genus *Hymenolepis* all tritesticular Hymenolepids are referred to as *Hymenolepis* s.l.. Specific determination of Hymenolepids collected in this study was not carried out due to the condition of the specimens (destrobilization, digestion and lack of rostellar hooks).

The classifications utilized in this work are of Emerson, 1972 (for lice), Yamaguti, 1961, 1963 and 1971 (for nematodes, acanthocephalans and trematodes respectively) and McDonald 1969c (for cestodes).

Data were analyzed using cluster analysis, G-tests
and the Mann-Whitney U-test.

RESULTS AND DISCUSSION

A total of 37 different genera (at least 50 species) of metazoan parasites were recovered from the scoters examined in this study. Necropsies yielded 29 genera (39 species), 21 genera (28 species) and 25 genera (33 species) of parasites from the surf, white-winged and black scoters respectively.

Trematoda

The 175 scoters examined yielded 18 genera and 23 species of trematodes (surf scoters 11 genera (15 species); white-winged scoters 11 genera (14 species); black scoters 11 genera (15 species)) (Table 4). A total of 55 (58.5%) surf, 28 (77.5%) white-winged and 17 (37.8%) black scoters were infected with these parasites.

Austrobilharzia terrigalensis Johnston, 1927 was noted in 8 (8.5%) surf, 2 (5.6%) white-winged and 4 (8.9%) of the black scoters examined (Table 4). The finding of this trematode in each host constitutes a new host record. In all instances, except one, the specimens were removed from the blood vessels associated with the small intestine. The exception contained schistosomes in the blood vessels associated with the liver and gall bladder:

TABLE 4. DETAILS OF INFECTION OF SURF SCOTERS, WHITE-WINGED SCOTERS AND BLACK SCOTERS WITH TREMATODA.

DIGENEA	SURF SCOTER				WHITE-WINGED SCOTER				BLACK SCOTER			
	A	B	C	D	A	B	C	D	A	B	C	D
<i>Austrobilharzia terrigalensis</i>	8.5	2.5	1-5	20 17,3,0	5.6	1.5	1-2	3 3,0,0	8.9	1.0	1	4 4,0,0
<i>Eggocotyle lunata</i>	2.1	2.0	1-3	4	-	-	-	-	-	-	-	-
<i>Notocotylus attenuatus</i>	1.1	4.0	4	4	2.8	14.0	14	14 9ad. 5imm.	2.2	17.0	17	17
<i>Catatropis verrucosa</i>	-	-	-	-	8.3	4.7	3-8	14	-	-	-	-
<i>Tristriata elegans</i>	29.8	4.8	1-16	135	11.1	4.5	1-12	18	-	-	-	-
<i>Eucootyle castanea</i>	2.1	1.0	1	2	11.1	2.0	1-3	8	4.4	10.0	1-19	20
<i>Gymnophallus bursicola</i>	10.6	5.6	1-20	56	5.6	3.5	1-6	7	8.9	9.0	1-36	46

TABLE 4. CONTINUED.

DIGENEA	SURF SCOTER				WHITE-WINGED SCOTER				BLACK SCOTER			
	A	B	C	D	A	B	C	D	A	B	C	D
<i>Gymnophallus choledeus</i>	7.4	5.3	1-20	37	8.3	5.4	1-12	16	2.2	1.0	1	1
<i>Gymnophallus deliciosus</i>	14.9	4.6	1-18	64	30.6	8.8	1-26	35	8.9	6.3	1-24	69 68ad, 1imm.
<i>Gymnophallus</i> sp. Type 1	3.2	6.0	1-15	18	5.6	6	12	12	2.2	6.0	6	6
<i>Gymnophallid</i> sp.	1.1	2.0	2	2	-	-	-	-	2.2	2.0	2	2
<i>Gymnophallus</i> spp.	2.1	2.5	1-4	5	8.3	9.0	4-14	27imm.	11.1	7.0	1-26	35imm.
<i>Echinostoma revolutum</i>	-	-	-	-	-	-	-	-	2.2	42.0	42	42imm.
<i>Dietsiella egregia</i>	-	-	-	-	22.2	69.8	4-240	⁵⁵⁹ 328ad, 231imm.	-	-	-	-
<i>Himasthla incisa</i>	-	-	-	-	2.8	1.0	1	1	2.2	3.0	3	3

TABLE 4. CONTINUED.

DIGENEA	SURF SCOTER				WHITE-WINGED SCOTER				BLACK SCOTER			
	A	B	C	D	A	B	C	D	A	B	C	D
<i>Acanthoparyphium melarittae</i>	1.1	3.0	3	3	-	-	-	-	-	-	-	-
<i>Peiloetomum</i> sp.	7.4	5.0	1-20	35	-	-	-	-	-	-	-	-
<i>Ascorhytis charadriiformis</i>	3.2	10.3	1-23	31	2.8	1.0	1	1	4.4	82.5	63-102	165
<i>Microphallus pygmaeus</i>	-	-	-	-	-	-	-	-	2.2	8.0	8	8
<i>Pseudospelotrema japonicum</i>	4.3	4.8	1-13	19	16.7	5.7	1-27	34	2.2	191.0	191	191
<i>Reticola</i> sp.	2.1	3.0	1-5	6	2.8	1.0	1	1	-	-	-	-

TABLE 4. CONTINUED.

DIGenea	SURF SCOTER				WHITE-WINGED SCOTER				BLACK SCOTER				
	A	B	C	D	A	B	C	D	A	B	C	D	
<i>Cotylurus strigeoides</i>	-	-	-	-	-	-	-	-	-	2.2	11.0	11	11
<i>Apatemon burti</i>	-	-	-	-	-	-	-	-	-	2.2	61.0	61	61
<i>Apatemon gracilis</i>	1.1	19.0	19	19	-	-	-	-	-	-	-	-	-

A = prevalence (% of ducks infected)
 B = intensity (mean no. of parasites per infected bird)
 C = range of parasite numbers
 D = total no. of parasites collected
 no. of males, no. of females, no. of immatures
 or
 no. of adults, no. of immatures

McDonald (1969) listed 4 species of *Austrobilharzia* Johnston, 1917 from anatids (*A. canadensis* McLeod, 1937; *A. chapini* Price, 1929; *A. manitobensis* McLeod, 1939 and *A. variglandis* (Miller & Northup, 1926) while Farley (1971) synonymized all four species with *A. terrigenensis*.

Zygocotyle lunata (Diesing, 1836) occurred in 2 (2.1%) of the surf scoters examined. Mettrick (1959) reports this parasite as a cause of mortality in domestic ducks. The finding of this trematode in the surf scoter constitutes a new host record. Table 3 gives further details on this infection of surf scoters. The specimens recovered in this study all come from the caeca of their hosts.

Notocotylus attenuatus (Rudolphi, 1809) infected all three hosts examined in this study (Table 4). It was recovered from 1 (1.1%, 2.8%, 2.2%) of each of the surf, white-winged and black scoters examined respectively. Serafin (1957, *vide* McDonald, 1969) reports this parasite as a cause of mortality in domestic geese. The occurrence of this parasite in the surf scoter constitutes a new host record.

Catantropis verrucosa (Froelich, 1789) was recovered from the caeca of 3 (8.3%) of the white-winged scoters examined (Table 4).

Tristriata elegans Filimonova, 1971 was found in 28 (29.8%) of the surf and 4 (11.1%) of the white-winged scoters examined (Table 4). The presence of *T. elegans* in the surf scoter constitutes a new host record. The parasites were recovered from the large intestine, caeca and cloaca of their hosts.

There are presently two valid species in the genus *Tristriata* Belopol'skaia, 1953 (*T. Anatis* Belopol'skaia, 1953 and *T. elegans*) which are distinguished from other Notocotyliidea by having two to four rod shaped ridges on their ventral surfaces (Yamaguti, 1971). *Tristriata elegans* is morphologically quite similar to *N. attenuatus* (Figure 1), except for the configuration of its ventral ridges. Methods of fixation alter the configuration of the aforementioned ridges in *Tristriata* which could lead to confusion with *Notocotylus*. It is consequently almost impossible to distinguish between these 2 genera after making permanent mounts of specimens. Frame (1969) states "techniques used by previous workers appear to have importantly influenced conclusions" when discussing fixation and mounting of specimens of *f. anatis*. Further, he notes that the ventral body ridges are difficult to see when methods of preparation that exert pressure on the worm, or involve excessive clearing are utilized. He also noted that the ridges were more readily visible on unstained specimens or on those stained

X
FIGURE 1.

Showing the similarity in the morphology
of (A.) *Trietriata elegans* and (B.)
Notocotylus attenuatus.




Figure 1

with Ehrlich's acid haemotoxylin. The ventral ridges of specimens obtained during this study were quite apparent upon removal from the frozen hosts. Light staining with Acid Carmine, or*Trichrome stain, and mounting in glycerine aided in microscopic examination of the specimens.

Eucotyle castenea Jensen, 1971 was recovered from 2 (2.1%) of the surf, 4 (11.1%) of the white-winged and 2 (4.4%) of the black scoters examined (Table 4). The finding of this parasite in the surf and black scoter both constitute new host records. A single specimen was recovered from the kidney of a white-winged scoter, while all other specimens were recovered from the caeca and body cavity. Jensen (1971) reports finding specimens in the intrarenal branches of ureters in *M. deglandi*. The location that they were recovered from in this study may have been due to post-mortem migration.

The family Gymnophallidae presently comprises 5 or 6 genera depending on the classification scheme one chooses to follow. Ching (1973) recognizes 6 genera: namely, *Gymnophallus* Odhner (1900), *Gymnophalloides* Fujita (1925), *Lacunovermis* Ching (1975), *Meiogymnophallus* Ching (1965), *Paragymnophallus* Ching (1973) and *Parvatrema* Cable (1953), while Yamaguti (1971) recognizes four genera in the family: namely, *Gymnophallus*, *Gymnophalloides* (synonym *Lacunovermis*), *Meiogymnophallus* and *Parvatrema* with *Paragymnophallus* being described in 1973. He

*Gomori's Trichrome

lists 10, 2, 7 and 10 species in each genus respectively, and in addition lists 12 species which cannot be placed in any genus due to inadequate descriptions. It is of interest to note that there are 8 species of *Gymnophallus* alone recorded from the black scoter. The author questions this multitude of species from a single genus occurring in a single host species. It is much more likely that the "species" described represent a "species-complex" with a number of morphological variations (possibly host induced - see Blankespoor, 1974) as suggested by Stunkard & Uzmann (1958). Due to a lack of life-cycle work the author feels that no new species should be described, unless it is an unique form, until the family is revised and augmented with life-cycle data, particularly as even the generic designations are open to interpretation. Deposition of more representative specimens in officially recognized depositories would have greatly aided this work. It is thus the opinion of the author that the family Gymnophallidae is in dire need of revision.

Members of the genus *Gymnophallus* were found in 27 (28.5%) surf, 13 (36.1%) white-winged and 10 (22.2%) of the black scoters examined.

Gymnophallus bursticola Odhner, 1900 was found to inhabit all three species of scoters examined, being recovered from 10 (10.6%) of the surf, 2 (5.6%) of the white-winged and

4 (8.9%) of the black scoters examined (Table 4). These parasites were recovered from the duodenum, caeca, Bursa of Fabricius and cloaca of their hosts.

Gymnophallus choledocus Odhner, 1900 was recovered from 7 (7.4%) surf, 3 (8.3%) white-winged and 1 (2.2%) of the black scoters examined (Table 4). The occurrence of this parasite in the surf and black scoters both constitute new host records. The sites of infection by this parasite included the gall bladder, Bursa of Fabricius and cloaca.

Gymnophallus delictosus (Olsson, 1893) was recovered from 14 (14.9%) of the surf, 11 (30.6%) of the white-winged and 4 (8.9%) of the black scoters examined (Table 4). Prior to this study *G. delictosus* had only been recovered from the black scoter, thus the present findings constitute new host records for the surf and white-winged scoters. These parasites were recovered from the gall bladder, small intestine (sections 1 and 3) and the Bursa of Fabricius. In a number of cases, it proved impossible to assign a specific name to some gymnophallids. The reasons for this include: (1) some specimens did not take stain very well; (2) others were immature; (3) others were too gravid; (4) some were damaged and (5) some did not fit the description of valid species well enough to warrant assigning them to a species.

Gymnophallus sp. (Type 1) was found in, 3 (3.2%) surf, 2 (5.6%) white-winged and 1 (2.2%) black scoter examined (Table 4). The specimens were recovered from the caeca, bursa and cloaca of their hosts. The range of measurements for the 33 specimens recovered are shown in Table 5. These specimens most closely resemble *G. choledochus*; however, the criteria used to key other specimens of this species did not fit the aforementioned specimens well enough to assign them definitely to this species. While the specimens possessed ova, often the uterus contained less than 20 ova whereas gravid specimens may contain several hundred ova.

Gymnophallus spp. refers to damaged or immature specimens that due to their condition could not be assigned a specific name. These specimens were recovered from 2 (2.1%) surf, 3 (8.3%) white-winged and 5 (11.1%) of the black scoters examined (Table 4). Sites of infection were the ureters, gall bladder, small intestine (sections 2 and 3), large intestine, bursa and cloaca.

A gymnophallid sp. was recovered from the caeca of 1 (1.1%) surf and from the small intestine (section 3) of 1 (2.2%) black scoter (Table 4). Table 5 gives the range of measurements for the 3 specimens measured. These specimens appear to be members of the genus *Gymnophalloides*.

Table 5. Range of measurements of *Gymnophallus* sp. (Type 1) and the gymnophallid sp. found in this study.

MEASUREMENT	<i>Gymnophallus</i> , sp Type 1	Gymnophallid sp.
Total Length	.650-.980	.440-.530
Total Width	.210-.480	.150
Oral Sucker Length	.100-.135	.90
Oral Sucker Width	.90-.110	.100-.105
Ventral Sucker Length	.95-.105	57
Ventral Sucker Width	.90-.105	-
Eggs	.020-.025	.020
Position of the Uterus	forebody*	hindbody

*In 2 of the 11 specimens the uterus extended to the hindbody.

Echinostoma revolutum (Froelich, 1802) a parasite that can cause disease and mortality in anatids (McDonald, 1969b), was recovered from 1 (2.2%) of the black scoters examined (Table 4). The 42 specimens recovered were located in the anterior small intestine (sections 1 and 2).

Dietziella egregia (Dietz, 1909) was recovered from 8 (22.2%) of the white-winged scoters, thus constituting a new host record. The sites of infection were the duodenum and small intestine (sections 1 and 2). The occurrence of this parasite in waterfowl is puzzling as it has only previously been recorded from a stork (*Harpiprion caeruleescens* Gibson, 1880) in Brazil. Its presence in the white-winged scoter may be considered accidental; however, it occurred in all the white-winged scoters collected in British Columbia in February, 1978. Conversely, its occurrence in *H. caeruleescens* may be accidental. The life-cycle of this parasite is unknown; however, the number of immature parasites collected (Table 4) suggests it has a marine intermediate host.

Himasthya incisa Linton, 1928 was located in the small intestine (section 3) of 1 (2.8%) white-winged and 1 (2.2%) black scoter (Table 4). Yamaguti (1958, 1971) lists this parasite as occurring in the black scoter, with McDonald (1969) calling the 1958 record erroneous.

Acanthoparyphium melanittae Yamaguti, 1939 was recovered from the duodenum of a single surf scoter (1.1%), for the first time, in this study (Table 4).

Specimens of a *Peilostomum* sp. were taken from the duodenum and small intestine (sections 1-3) of 7 (7.4%) of the surf scoters examined in this study (Table 4). The specimens recovered resemble most closely *P. magniovum* presently being described by H. L. Ching (pers. comm. H. L. Ching).

Ascorhytis charadriiformis Ching, 1965 was recovered from 3 (3.2%) of the surf, 1 (2.8%) of the white-winged and 2 (4.4%) of the black scoters examined (Table 4). The occurrence of this parasite in all three hosts constitutes new host records. The sites of infection were the small intestine (sections 1-3), large intestine, cloaca and caeca of the hosts.

Microphallus pygmaeus (Levinsen, 1881), a potential pathogen of eiders (Belopol'skaia 1953 *vide* McDonald 1969), was recovered from the small intestine (section 3) of 1 (2.2%) black scoter. Yamaguti (1971) uses the generic name *Spelotrema* for this parasite (*Spelotrema pygmaeus*), reserving the generic name *Microphallus* for digenea from fishes. The author prefers to follow the more logical scheme of Deblock (1971), which bases scientific names on morphology rather than host group.

Pseudospelotrema japonicum Yamaguti, 1939 was recovered from 4 (4.3%) surf, 6 (16.7%) white-winged and 1 (2.2%) of the black scoters examined (Table 4). The presence of this parasite in the surf and black scoters constitutes new host records. The parasites were recovered from the duodenum, small intestine (section 3), large intestine and caeca of the infected birds.

Specimens of a *Renicola* sp. were recovered from the kidneys of 2 (2.1%) surf and 1 (2.8%) of the white-winged scoters examined (Table 4). The occurrence of this genus in the white-winged scoter constitutes a new host record. Yamaguti (1971) lists *R. dollfusi* Odning, 1962 as being found in the genus *Melanitta* but does not note the species of scoter, while Oshmarin (1963) lists a *Renicola* sp. from white-winged scoters. Due to the gravid uteri of the specimens recovered few measurements were obtainable. The specimens resemble *R. mollisimus* Kulachkova, 1957 most closely, when taking egg size (average 36 μ m for surf scoter specimens and 42.5 μ m for specimens recovered from white-winged scoter) and host group into consideration.

Cotylurus strigeoides Dubois, 1958 was recovered from the small intestine (sections 2 and 3) of 1 (2.2%) black scoter. This constitutes a new host record. The measurements of the

TABLE 6. Comparison of measurements of *Cotylurus strigeoides* and *Cotylurus raabet* with specimens recovered in this study.

	<i>C. strigeoides</i> vide Dubois (1968)	<i>C. strigeoides</i> present study	<i>C. raabet</i> (Bezubik, 1958) vide Dubois (1968)
Total length	1.9mm	2.28-2.60mm	5mm
Length of Anterior Segment	0.44-0.71mm	0.780-0.900mm	1.1-1.8mm
Length of Posterior Segment	1.03-1.19mm	1.50-1.75mm	1.7-3.2mm
Length of Anterior Segment Length of Posterior Segment	1.5-2.3mm	1.8-2.2mm	1.46-1.85mm
Eggs	0.085-0.099mm 24 eggs	0.092-0.100mm 12 eggs	0.120-0.139mm numerous

specimens recovered in this study (Table 6) differ slightly from those of the only two known species of *Cotylurus* Szidat, 1928 with vitellaria in their forebody. However, accepting that egg size is the least variable character of a trematode, considering host-induced variation (Blankespoor, 1974), then the present specimens are undoubtedly *C. strigeoides*.

Apatemon burti (Miller, 1933) was located in the duodenum and small intestine (section 1) of 1 (2.2%) black scoter, for the first time.

Apatemon gracilis (Rudolphi, 1819), a parasite that can cause mortality in anatids (Zajicek 1963, vide McDonald 1969.), was recovered from the small intestine (sections 2 and 3) of 1 (1.1%) surf scoter (Table 4). The present finding constitutes a new host record.

Cestoda

Three genera of cestodes comprising an unknown number of species were recovered from the scoters examined in this study. A total of 56 (59.6%) of the surf, 23 (63.8%) of the white-winged and 29 (64.4%) of the black scoters were infected with Cestoda. Due to the lack of rostellar hooks and the poor condition of some of the specimens it was hard in many cases to even assign a generic name, let alone a specific one.

Hymenolepis spp. were recovered from 55 (58.5%) of the surf, 23 (63.9%) of the white-winged and 26 (57.8%) of the black scoters (Table 7). The cestodes were recovered from the duodenum, small intestine (sections 1-3), caeca and large intestine of their hosts. While the specific identification of these helminths fell outside the bounds of the present study, the author believes that approximately 10 species were present in the samples examined. The genus *Hymenolepis* s. l. is extremely large and taxonomically confused, with some 3, 26 and 24 species of *Hymenolepis* s. l. being recorded from surf, white-winged and black scoters respectively (Appendix 1).

Cloacotaenia megalops (Nitzsch, 1829) was recovered from the bursa of 1 (2.8%) of the white-winged scoters examined (Table 7). While this genus is considered by McDonald (1969) to be part of *Hymenolepis* s. l. the author diverges from him and accepts *Cloacotaenia* due to its lack of rostellar hooks.

A member of the sub-family *Fimbriariinae* was recovered from the duodenum and small intestine (sections 1 and 2) of 13 (13.8%) of the surf and 6 (13.3%) of the common scoters examined (Table 7). Due to the presence of a poorly developed pseudoscolex and internal segmentation, the author believes these specimens to belong to the genus *Fimbriarioides*.

TABLE 7. Details of infection of Surf, White-Winged and Black Scoters with Cestoda.

CESTODA	SURF SCOTER			WHITE-WINGEDSCOTER			BLACK SCOTER		
	A	B	C	A	B	C	A	B	C
<i>Hymenolepis</i> spp.	58.5	5-10	1-5 to 50-55	63.9	25-30	1-5 to 305-310	57.8	40-45	1-5 to 670- 675
<i>Cloacotaenia megalops</i>	-	-	-	2.8	3	3	-	-	-
Sub. family <i>Fimbriariinae</i>	13.8	5-10	1-5 to 25-30	-	-	-	13.3	50-55	1-5 to 140- 145

A = prevalence (% of ducks infected)

B = intensity (mean no. of parasites per infected bird)

C = range of parasite numbers

Nematoda

During this study 8 genera and 14 species of Nematoda were recovered from the three species of scoters; surf scoters contained 8 genera (13 species), white-winged scoters 4 genera (8 species), black scoters 6 genera (9 species) (Table 8). A total of 77 (81.9%) surf, 22 (61.1%) white-winged and 26 (57.8%) black scoters were infected. The absence of the proventriculus and/or gizzard, and removal of food-stuffs, in some birds may have influenced the indicated incidence of all Nematoda, except the capillarids and *Sarcofilaria* Wehr, 1939.

A single *Capillaria anatis* (Schrank, 1790) was located in the body cavity of 1 (1.1%) of the surf scoters examined (Table 8). The presence of the parasite in this host constitutes a new host record. The specimen probably migrated to the body cavity from the intestine after death of the host.

Capillaria nyrocinarium Madsen, 1945 was recovered from the small intestine (sections 1-3), Bursa of Fabricius and caeca of 20 (21.3%) surf, 12 (33.3%) white-winged and 2 (4.4%) black scoters (Table 8). The occurrence of this parasite in the first named species constitutes a new host record.

Amidostomum acutum (Lundahl, 1848) the Nematode with the highest prevalence, was present in 48 (51.1%) surf, 11 (30.5%) of the white-winged and 22 (48.9%) of the black scoters examined (Table 8). The occurrence of this parasite in the surf

TABLE 8. Details of infection of surf, white-winged and black scoters with Nematoda.

PARASITE	SURF SCOTER				WHITE-WINGED SCOTER				BLACK SCOTER			
	A	B	C	D	A	B	C	D	A	B	C	D
<i>Capillaria anatis</i>	1.1	1	1	1,0,0	-	-	-	-	-	-	-	-
				46				30				10
<i>Capillaria nyrocinarum</i>	21.3	2.3	1-8	20,18,8	33.3	2-5	1-5	13,14,3	4.4	5	4-6	3,6,1
				307				52				94
<i>Amidostomum acutum</i>	51.1	6.2	1-53	126,174,7	30.5	4.7	1-27	21,31,0	48.9	4.3	1-15	45,46,3
				8				-				-
<i>Amidostomum sp.</i>	5.3	1.6	1-2	0,0,8	-	-	-	-	-	-	-	-
				-				-				10
<i>Epididostomum uncinatum</i>	-	-	-	-	-	-	-	-	4.4	5	5	5,5,0
				38				1				156
<i>Streptocara californica</i>	21.3	1.9	1-6	14,23,1	2.8	1	1	1,0,0	17.8	19.5	1+90	36,119,1
				10				13				4
<i>Streptocara a. crassicauda</i>	7.4	1.4	1-2	4,5,1	2.8	3.3	1-6	2,11,0	8.9	1	1	4,0,0
				18				6				-
<i>Streptocara formosensis</i>	16.0	1.2	1-4	6,12,0	2.2	6	6	1,5,0	-	-	-	-
				2				-				-
<i>Streptocara sp.</i>	2.1	1	1	0,0,2	-	-	-	-	-	-	-	-
				2				-				-
<i>Echinuria borealis</i>	1.1	2	2	2,0,0	-	-	-	-	-	-	-	-
				186				-				16
<i>Echinuria hypognatha</i>	22.3	8.9	1-54	93,61,32	-	-	-	-	15.6	2.3	1-5	5,4,7
				37				4				32
<i>Tetrameres fistispina</i>	7.4	6.2	1-22	37,0,0	2.8	4	4	4,0,0	6.7	10.7	1-22	32,0,0

TABLE 8. Details of infection of surf, white-winged and black scoters with Nematoda.
CONTINUED

PARASITE	SURF SCOTER				WHITE-WINGED SCOTER				BLACK SCOTER			
	A	B	C	D	A	B	C	D	A	B	C	D
<i>Tetrameres somaterias</i>	46.8	4.6	1-27	203,0,0	25.0	5.4	1-26	49,0,0	2.2	3	3	3,0,0
<i>Tetrameres spinosa</i>	3.1	2	1-4	6,0,0	2.8	1	1	1,0,0	2.2	1	1	1,0,0
<i>Tetrameres</i> sp.	5.3	1	1	0,4,1	5.6	1.5	1-2	3,0,0	-	-	-	-
<i>Sarcofaga</i> sp.	1.1	1	1	0,1,0	-	-	-	-	-	-	-	-
<i>Nematode</i> sp. Type 1	1.1	4	4	4	-	-	-	-	-	-	-	-

A = prevalence (percent of ducks infected)
 B = intensity (mean no. of parasites per infected duck)
 C = range of parasite
 D = total no. of parasites collected
 no. of males, no. of females, no. of immatures

scoter constitutes a new host record. If the prevalence was calculated for those missing their gizzard it would be 55.2%, 34.4% and 53.3% respectively. This parasite was characteristically found under the gizzard lining of the host, where it can cause pathological changes (Zajiček 1960, *vide* McDonald 1969). Zajiček (*op. cit.*) worked on *A. boschadis* which is now considered a synonym of *A. acutum* (*vide* McDonald 1969).

An *Amidostomum* sp. (*acutum?*) was located under the gizzard lining of 5 (5.3%) of the surf scoters examined (Table 8). The specimens were either immature or damaged, and hence could not be identified to the specific level. All these infections occurred in surf scoters that were infected with *A. acutum*.

Epomidiostomum uncinatum (Lundahl, 1848), a parasite which causes pathological changes in the host (Beaudette 1939, *vide* McDonald 1969) was noted under the gizzard lining of 2 (4.4%) of the black scoters examined (Table 8).

The genus *Streptocara* (Raillet, Henry et Sisoff, 1912) had a prevalence of 39.4% (37 birds), 13.9% (5 birds) and 24.4% (11 birds) in the surf, white-winged and black scoters respectively. If the prevalence is recalculated omitting those birds lacking a gizzard, the gizzard is the preferred site of infection for members of this genus of nematode, then it is 42.5%, 16.1%, 24.4% respectively.

Streptocara californica (Geddes, 1919) infected 20 (21.3%), 1 (2.8%) and 8 (17.8%) of the surf, white-winged and black scoters respectively (Table 8). The preferred habitat of these worms was under the gizzard lining. If the prevalence was calculated omitting those birds that had their gizzard removed the figures were a little higher than noted above (23.0%) for the surf scoter and (3.1%) for the white-winged scoter.

Streptocara crassicauda crassicauda (Creplin, 1829) infected 7 (7.4%) surf, 1 (2.8%) white-winged and 4 (8.9%) of the black scoters (Table 8). Garkavi (1950) reports this parasite to cause pathological conditions and death in anatids. Once again the preferred habitat of these worms was under the gizzard lining, although some parasites were removed from the proventriculus. If the prevalence is recalculated omitting the birds that had their gizzard removed, then it is 8.1% for the surf and 3.1% for the white-winged scoter.

Streptocara formosensis Sugimoto, 1930 was located under the gizzard lining and in the proventriculus of 15 (16.0%) surf scoters, and 1 (2.8%) white-winged scoter (Table 8). If the prevalence is recalculated omitting those birds that had their gizzards removed, then the result is 15.0% for the surf scoter.

Several specimens of a *Streptocara* sp. were taken from the gizzard and proventriculus of 2 (2.2%) of the surf scoters from Labrador. It seems likely that they are either *S. c. crassicauda*, *S. formosensis*, or *S. californica*.

The genus *Echinuria* (Soloviev, 1912) infected 21 (22.3%) surf and 7 (15.6%) of the black scoters examined in this study. If the prevalence is calculated omitting the birds missing their proventriculus (McDonald (1969) notes this organ to be the preferred habitat for members of this genus), then the figures are 22.8% and 15.6% for the surf and black scoters respectively.

Echinuria borealis (Mawson, 1956) was taken from the proventriculus of 1 (1.1%) surf scoter for the first time (Table 8).

Echinuria hypognatha (Wehr, 1937) infected 21 (22.3%) surf and 7 (15.6%) of the black scoters examined (Table 8), with the former constituting a new host record. The preferred site of infection for this parasite was the proventriculus with infections in the esophagus and gizzard being noted. The true prevalence (omitting birds missing their proventriculus) is 22.8% for surf and 15.6% for black scoters.

The genus *Tetrameres* was located in 44 (46.8%) surf, 11 (30.6%) white-winged and 4 (8.9%) black scoters. If the prevalence is recalculated omitting those birds missing their proventriculus, then the prevalence is 47.8%, 31.4% and 8.9% respectively.

Tetrameres fissiapina (Diesing, 1861) was recovered from the esophagus or proventriculus of 7 (7.4%), 1 (2.8%) and 3 (6.7%) of the surf, white-winged and black scoters respectively (Table 8). This worm has been reported to be the cause of pathological conditions (Swales 1963a,b) and death (Bittner 1923, *vide* McDonald 1969) in anatids. This is the first record for this parasite in the surf scoter. If the prevalence is recalculated omitting birds missing their proventriculus, then 7.6%, 2.9% and 6.7% were infected respectively.

Tetrameres somateriae Ryzhikov, 1963 was taken from 44 (46.8%) surf, 9 (25.0%) white-winged and 1 (2.2%) of the black scoters, for the first time (Table 8). The major site of infection was the proventriculus, and discounting those hosts with this organ missing the prevalence was 47.8%, 25.8% and 22.2% respectively.

Tetrameres spinosa (Maplestone, 1931) was found in the esophagus and/or proventriculus of 3 (3.1%) surf, 1 (2.8%)

white-winged and 1 (2.2%) of the black scoters examined (Table 8). If birds missing their proventriculus are omitted in calculating the prevalence, then it is as follows 3.3%, 2.9% and 2.2% for the three host species respectively. The finding of *T. spinosa* in these hosts constitutes new host records.

Specimens of a *Tetrameres* sp. were recovered from 5 (5.3%) surf and 2 (5.6%) white-winged scoters. In the surf scoters all infections involved a single worm (4 female and 1 immature male), while in the white-winged scoters 3 damaged male parasites were removed from the 2 infected hosts. McDonald (1974) notes that it is not possible to key female *Tetrameres* to the species level. The 4 females recovered in this study were found in birds harbouring only *T. somateriae* males; thus it is not unreasonable to suspect that the females recovered were also *T. somateriae*. The single immature male worm was recovered from a surf scoter harbouring adult males of *T. fassispina* and *T. somateriae*. The 3 damaged male worms recovered from the 2 infected white-winged scoters were the only individuals of their genus to be recovered from these birds.

A single gravid specimen of the genus *Sarcocoma* was recovered from 1 (1.1%) of surf scoters examined (Table 8). This parasite was recovered from the air sacs of its host and constitutes a new host record.

An unidentifiable nematode sp designated Type 1 was recovered from the proventriculus of 1 (1.1%) of surf scoters examined (Table 8). The specimens recovered were immature, but resembled members of the genus *Paracuarua* Krishna Rao, 1951.

Acanthocephala

Acanthocephala belonging to three genera and four species were recovered in this study; Surf scoters harboured 3 genera (3 species), white-winged scoters 1 genus (1 species) and black scoters 2 genera (2 species). A total of 19 (20.2%) surf, 7 (19.4%) white-winged and 7 (15.6%) black scoters were infected with Acanthocephala.

Polymorphus botulus (Van Cleave, 1916) was found in 6 (6.25%) surf, 7 (19.4%) white-winged and 7 (15.6%) black scoters examined (Table 9). This parasite has been reported as causing pathological changes in infected animals and has been the cause of heavy mortality and epizootics in *Somateria mollissima* (Linnaeus, 1758) (Clark et al. 1958, Garden et al. 1964). Single infections of up to 654 worms, in the latter host, were reported by Bishop (1971). In this study a single immature black scoter harboured 171 worms. Scars (referred to as nodular taeniasis (Harrison, 1955)) are host reactions to the embedded proboscis of this worm (Figure 2). The presence of *P. botulus* in the small and large intestines and caeca, of each species of host constitutes a new host record.

Falsifilicollis altmani (Perry, 1942) was detected in a single (1.1%) surf scoter (Table 9). Perry (1942) in the

FIGURE 2.

Showing nodular taeniasis on the intestine
of a black scoter.



Figure 2

TABLE 9. Details of infection of surf, white-winged and black scoters with *Acanthocephala*.

PARASITE	SURF SCOTER				WHITE-WINGED SCOTER				BLACK SCOTER			
	A	B	C	D	A	B	C	D	A	B	C	D
<i>Falsificollis altmani</i>	1.1	2	2	2,0,0	-	-	-	-	-	-	-	-
<i>Polymorphus latulus</i>	6.3	5.7	1-23	34 16,18,0	19.4	10.6	1-37	74 34,21,19	15.6	45.3	1-171	317 104,156,57
<i>Corynosoma constrictum</i>	-	-	-	-	-	-	-	-	2.2	7	7	7 6,1,0
<i>Corynosoma sudsuche</i>	13.8	4.2	1-14	55 25,18,12	-	-	-	-	-	-	-	-

A = prevalence (percent of birds infected with parasite)
 B = intensity (mean no. of parasites per infected bird)
 C = range of parasite
 D = total no. of parasites collected
 no. of males, no. of females, no. of immatures.

only other record of this parasite reported it as a cause of mortality in *M. f. deglandi* and *M. perspicillata*. The parasites in this study were located in the small intestine (section 2).

Seven *Corynosoma constrictum* (Van Cleave, 1918) were found in the small intestine (section 3) of 1 (2.2%) of the black scoters (Table 9).

Corynosoma sudsuche Belopol'skaia, 1959 was removed from the small and large intestines and caeca of 13 (28.9%) surf scoters (Table 9), this constituting a new host record.

Mallophaga

Four genera and 6 species of Mallophaga were recovered during this study. Surf scoters bore 4 genera (6 species), white-winged scoters 3 genera (4 species) and black scoters 3 genera (5 species) (Tables 10, 11, 12). A total of 34 (70.8%) surf, 16 (94.1%) white-winged and 16 (94.1%) black scoters were infested.

The lice collected in this study belong to the Superfamily Ischnocera (Family Philopteridae) and the Superfamily Amblycera (Family Menoponidae). The philopterid population on a bird can be divided into two basic morphological types: "On the head and neck is found a short, round-bodied type not greatly flattened dorso-ventrally and with a large head" (Clay, 1957) that is adapted to movement through and among the shorter feathers

TABLE 10. Details of Infestation of surf scoters with Mallophaga.

PARASITE	No. (%) of birds infested	NO. OF PARASITES				MEAN	RANGE
		MALE	FEMALE	NYMPH	TOTAL		
<i>Anatocous dentatus</i> (males)	4 (8.3%)	5	-	-	5	1.3	1-2
<i>Anatocous teterodes</i> (males)	11 (22.9%)	17	-	-	17	1.5	1-3
<i>Anatocous</i> spp. (females and nymphs)	20 (41.7%)	-	39	3	42	2.1	1-7 (females) 1 (nymphs)
<i>Anaticola crassicornis</i>	28 (58.3%)	24	44	64	132	4.7	1-22
<i>Holomenopon leucomanthum</i>	5 (10.4%)	1	7	-	8	1.6	1-3
<i>Holomenopon loomisi</i>	3 (6.3%)	-	5	-	5	1.7	1-2
<i>Holomenopon nymphs</i>	4 (8.3%)	-	-	6	6	1.5	1-2
<i>Trinoton querquedulae</i>	1 (2.1%)	-	-	1	1	1	1

TABLE 11. Details of infestation of white-winged scoters with Mallophaga.

PARASITE	No. (%) of birds infested	NO. OF PARASITES				MEAN	RANGE
		Male	Female	Nymph	Total		
<i>Anatoecus dentatus</i> (males)	4 (23.5%)	15	-	-	15	3.8	1-8
<i>Anatoecus iterodes</i> (males)	4 (23.5%)	9	-	-	9	2.3	1-4
<i>Anatoecus</i> spp. (females and nymphs)	5 (29.4%)	-	16	5	21	4.2	1-5 (female) 1 (nymph)
<i>Anaticola crassirostris</i>	7 (41.2%)	52	43	129	224	32	1-91
<i>Holomenopon loomisi</i>	4 (23.5%)	-	1	5	6	1.5	1-2

TABLE 12. Details of infestation of black scoters with Mallophaga.

PARASITE	NO. (%) of birds infested	NO. OF PARASITES				MEAN	RANGE
		Male	Female	Nymph	Total		
<i>Anatocercus dentatus</i> (males)	4 (23.5%)	9	-	-	9	2.3	1-5
<i>Anatocercus icterodes</i> (males)	7 (41.2%)	16	-	-	16	1.8	1-4
<i>Anatocercus</i> spp. (females and nymphs)	9 (53.0%)	-	32	32	64	7.1	1-9 (female) 4-16 (nymph)
<i>Anaticola crassicornis</i>	17 (100%)	33	39	93	165	9.7	1-22
<i>Holomenopon leucoxanthum</i>	4 (23.5%)	7	7	-	14	3.5	2-7
<i>Holomenopon loomisi</i>	5 (29.4%)	4	11	-	15	3	1-6
<i>Holomenopon</i> nymphs	7 (41.2%)	-	-	22	22	3.1	1-7

of the head and neck e.g. *Anatoecus* (Cummings, 1916). On the larger, broader feathers of the wings and back are found more elongate forms which have undergone severe dorso-ventral flattening to enable them to better evade preening e.g. *Anaticola* (Clay, 1936). The Amblycera, being a less specialized group seem to rely on speed to avoid the preening of their host as they have diverged little from their psocid ancestors (Clay, 1957) e.g. *Holomenopon* Eichler, 1941 and *Trinoton* Nitzsch, 1818. Amblycerans are found mainly on the breast and wings of their hosts (Bourgeois and Threlfall, 1979; Fitzpatrick and Threlfall, 1977).

The occurrence of *Anatoecus dentatus* (Scopoli, 1763), *A. iterodes* (Nitzsch, 1818), *Holomenopon leuocanthum* (Burmeister, 1838) and *H. loomisti* (Kellogg, 1896) on the surf scoter and *A. iterodes* and *H. loomisti* on the black scoter constitute new host records.

The males of *A. dentatus* and *A. iterodes* are separated from the *Anatoecus* nymphs and females in Tables 9, 10 and 11 due to the fact that at present the females of the two species are indistinguishable, if indeed two species exist. Emerson (1972) states "whether or not *Anatoecus iterodes* is a dimorphic form of *Anatoecus dentatus* has not been settled satisfactorily." Thus to assign females or nymphs to either species could be a misinterpretation of data. Given the number of mixed infections of *A. dentatus* and *A. iterodes* males (surfs-2, white-winged-3

and black scoters-4) one cannot assume that because one type of male is absent that the females are of the other species. *Anitococcus* species were found to infest 25 (52.0%) surf, 10 (58.8%) white-winged and 13 (81.25%) of the black scoters examined.

Anaticola crassicornis (Kellogg, 1896) infested 28 (58.3%) surf, 7 (41.2%) white-winged and 17 (100%) of the black scoters examined.

Holomenopon species infested 8 (16.7%) surf, 4 (23.5%) white-winged and 9 (56.3%) black scoters examined in this study. The presence of *H. loomisi* on the black scoter (Table 12) indicates that two species from this genus are found on black scoters. With neither species of *Holomenopon* being previously recorded from the surf scoter makes the assigning of either species as a straggler difficult if indeed one is. However, while collecting specimens for this study mixed moulting flocks of surf and black scoters were observed regularly. Body contact of two hosts could facilitate transfer of lice; given the close phylogenetic relationships of the hosts, it would not seem too unlikely that transferred lice could establish a population on a host. Also the possibility of shed feathers, with lice on them, coming into contact with other birds may be a more likely method of transfer in mixed moulting flocks.

Trinoton querquedulae (Linnaeus, 1758) had been recovered prior to this study, from all three scoters. However, the finding of only one female nymph on a single surf scoter is suggestive that this louse may be a straggler on scoters. A possible explanation for its low incidence may lie in the number of moulting birds examined i.e. the majority of the *Trinoton* population may have been lost with the moulted feathers.

Acarina

A single acarine, *Freyana anatinae* (Koch, 1844) was located on 6 (12.5%) of the surf and 6 (35.3%) of the black scoters examined. No data is available on intensity of infestation as only presence or absence was noted. A possible explanation for its low prevalence concerns its primary habitat i.e. the primary and secondary feathers of the wing. The fact that many birds were in heavy moult may have reduced its prevalence.

GENERAL DISCUSSION

A total of 16 (9.1%) of the 175 scoters examined were helminth free (8(8.5%) of the surf, 3(8.3%) of the white-winged and 5(11.1%) of the black scoters).

Surf scoters had the highest prevalence of nematodes and acanthocephalans, while white-winged scoters had the highest prevalence of trematodes and black scoters had the highest prevalence of cestodes. The Labrador sample of birds showed the highest prevalence for most helminths in all three host species. However, this latter fact may be misleading due to the small sample size. Table 13 details the infection of scoters by the major parasite groups and by sample area. In all cases, except two, Western Atlantic birds contained more species than the other sample areas (Table 14).

PARASITE DISTRIBUTION

"Parasitologists have known for a long time that individual species of parasites are found only in restricted, specific sites (or microhabitats) within their hosts, that the microhabitats occupied by some species are more restricted than those of others" (Holmes, 1973). It is well known that each organ and sometimes even different tissues may be regarded as unique environments for helminths due to morphological, biochemical and physiological differences. Further "site selection is predominately an active process on the part of the parasite; for some parasites at least, site selection is a continuing process, which may result in para-

TABLE 13. DETAILS OF INFECTION OF SCOTERS AND INDIVIDUAL SAMPLE AREAS BY THE MAJOR HELMINTH GROUPS.

PARASITE GROUP	ALL SCOTERS	SURF SCOTERS				WHITE-WINGED SCOTERS					BLACK SCOTERS			
	A	A	B	C	D	A	B	C	D	E	A	B	C	D
Trematoda	100 (57.1)	55 (58.5)	3 (12.0)	29 (64.4)	23 (95.8)	28 (97.8)	1 (33.3)	17 (77.3)	5 (100.0)	5 (83.3)	17 (37.8)	7 (25.9)	5 (38.5)	5 (100.0)
Cestoda	108 (61.7)	56 (59.6)	11 (44.0)	25 (55.6)	21 (87.5)	23 (63.8)	-	15 (68.2)	5 (100.0)	3 (50.0)	29 (64.4)	18 (66.7)	6 (46.2)	5 (100.0)
Nematoda	125 (71.4)	77 (81.9)	10 (40.0)	44 (97.8)	23 (95.8)	22 (61.1)	1 (33.3)	12 (54.5)	5 (100.0)	4 (66.7)	26 (57.8)	10 (37.0)	13 (100.0)	3 (60.0)
Acanthocephala	33 (18.9)	19 (20.2)	-	18 (40.0)	1 (4.2)	7 (19.4)	-	6 (27.3)	-	1 (16.7)	7 (15.6)	-	6 (46.2)	1 (20.0)

A = No. (%) infected

B = No. (%) of New Brunswick sample infected

C = No. (%) of British Columbia sample infected

D = No. (%) of Labrador sample infected

E = No. (%) of Norway sample infected

Table 14. Number of species of each major helminth group in birds from different sample areas.

PARASITE GROUP	SURF SCOTER		WHITE-WINGED SCOTER			BLACK SCOTER	
	A	B	A	B	C	A	B
Trematoda	11	10	8	9	3	5	12
Cestoda	1	2	2	1	1	2	2
Nematoda	10	11	5	6	2	7	8
Acanthocephala	2	2	1	0	1	1	2

A = Pacific sample

B = Western Atlantic sample (New Brunswick & Labrador samples combined)

C = Eastern Atlantic

NOTE: *Gymnophallus* spp. are excluded from this Table.

Also the no. of cestode species is a minimum.

sites alternating between different microhabitats; and the presence of other parasites may modify the microhabitat selected by a parasite." (Holmes, 1973). Thus, it is not too surprising to find a parasite in more than one area within a host or hosts.

Many authors list the area(s) parasites are recovered from but few list together all the parasites found in a given area. This is deceiving in that it masks the fact that parasites are found in communities. It is not a goal of this thesis to discuss the various communities that can occur in scoters but merely to give an idea that parasites are not solitary organisms, as they are often depicted. Therefore, the parasites will be listed with their respective environments to give an idea of the possible parasite communities as well as to indicate the areas in which they occurred and the preferred area/s if any (Table 15).

A further interesting aspect of this work relates to the geographical distribution of the parasites found. Not all parasites are evenly distributed throughout the geographical range of a host. The reasons for this include (1) absence of one or more intermediate hosts and (2) biotic factors e.g. water pH, temperature, water salinity, etc..

In this study, (see Table 16) 2 trematodes (*G. choledocus* and *G. deliciosus*), 1 genus of cestodes (*Hymenolepis*), 2 nematodes (*A. acutum* and *T. somateriae*) and 1 acanthocephalan (*P. botulus*) occurred across the geographical range of the hosts sampled. Four trematodes, 1 cestode and 2 nematodes were recovered only from the Pacific sample. Eight trematodes, 4 nematodes and 2 acanthocephalans were harboured only by Western Atlantic hosts. A single trematode, *C. verrucosa* was the only parasite restricted to the Eastern

108

TABLE 15. DETAILS OF PARASITE LOCATIONS AND PARASITE COMMUNITIES POSSIBLE IN SCOTERS EXAMINED IN THIS ST

	Esophagus	Proventriculus	Gizzard	Duodenum	SI ₁	SI ₂	SI ₃	Caecae	
<i>A. terrigalensis</i>	-	-	-	1(7.7)	2(15.4)	1(7.7)	2(15.4)	-	3
<i>Z. lunata</i>	-	-	-	-	-	-	-	2(100)	
<i>N. attenuatus</i>	-	-	-	-	-	-	-	4(100)	
<i>C. verrocuea</i>	-	-	-	-	-	-	-	3(100)	
<i>T. elegans</i>	-	-	-	-	-	-	-	27(81.8)	4
<i>E. castanea</i>	-	-	-	-	-	-	-	2(28.6)	
<i>G. burticola</i>	-	-	-	1(4.8)	-	-	-	4(19.0)	
<i>G. choledocus</i>	-	-	-	-	-	-	-	-	
<i>G. deliciosus</i>	-	-	-	1(3.6)	-	-	1(3.6)	-	
<i>Gymnophallus</i> sp. Type 1	-	-	-	-	-	-	-	6(66.7)	
<i>Gymnophallus</i> spp.	-	-	-	-	-	1(10.0)	1(10.0)	-	1
<i>Gymnophallid</i> sp.	-	-	-	-	-	-	1(50.0)	1(50.0)	
<i>E. revolutum</i>	-	-	-	-	1(50.0)	1(50.0)	-	-	
<i>D. egregia</i>	-	-	-	-	-	-	-	-	
<i>H. inoisa</i>	-	-	-	-	-	-	2(100)	-	
<i>A. melanittae</i>	-	-	-	1(100.0)	-	-	-	-	
<i>Psilostomum</i> sp.	-	-	-	1(11.1)	1(11.1)	3(33.3)	4(44.4)	-	
<i>A. okavadiiformis</i>	-	-	-	-	1(7.1)	4(28.6)	1(7.1)	5(35.7)	2
<i>M. pygmaeum</i>	-	-	-	-	-	-	1(100.0)	-	
<i>P. japonicum</i>	-	-	-	1(7.1)	-	-	1(7.1)	8(57.1)	2
<i>Rantocla</i> -sp.	-	-	-	-	-	-	-	-	
<i>C. strigatodes</i>	-	-	-	-	-	1(50.0)	1(50.0)	-	
<i>A. burti</i>	-	-	-	1(50.0)	1(50.0)	-	-	-	
<i>A. gracilis</i>	-	-	-	-	-	1(50.0)	1(50.0)	-	
<i>Hymenolepis</i> spp.	-	-	-	16(8.8)	64(35.4)	53(29.3)	27(14.9)	14(7.7)	2
<i>G. megalops</i>	-	-	-	-	-	-	-	-	
Primbriariinae	-	-	-	12(46.2)	7(26.9)	5(19.2)	-	-	
<i>D. anatis</i>	-	-	-	-	-	-	-	26(88.4)	
<i>C. nyroodinarium</i>	-	-	-	-	2(5.3)	4(10.5)	4(10.5)	-	
<i>A. acutum</i>	1(1.3)	5(7.9)	6(86.8)	-	-	-	1(1.3)	-	
<i>Amidostium</i> sp.	-	-	(85.7)	-	-	-	-	-	
<i>E. uncinatum</i>	-	-	(100.0)	-	-	-	-	-	
<i>S. californica</i>	-	1(3.4)	8(96.6)	-	-	-	-	-	
<i>S. c. crassicauda</i>	-	1(7.7)	1(84.6)	-	-	-	1(7.7)	1(1.6)	
<i>S. formosensis</i>	-	1(6.3)	5(93.7)	-	-	-	-	-	
St +	-	1(50.0)	(50.0)	-	-	-	-	-	

<i>G. ohledocus</i>	-	-	-	-	-	-	-	-	-	-
<i>G. deliciosus</i>	-	-	-	1(3.6)	-	-	-	-	1(3.6)	-
<i>Gymnophallus</i> sp. Type 1	-	-	-	-	-	-	-	-	-	6(66.7)
<i>Gymnophallus</i> spp.	-	-	-	-	-	-	1(10.0)	-	1(10.0)	-
<i>Gymnophallid</i> sp.	-	-	-	-	-	-	-	-	1(50.0)	1(50.0)
<i>E. revolutum</i>	-	-	-	-	-	1(50.0)	1(50.0)	-	-	-
<i>D. egregia</i>	-	-	-	-	-	-	-	-	-	-
<i>H. incisa</i>	-	-	-	-	-	-	-	-	2(100)	-
<i>A. melanittas</i>	-	-	-	1(100.0)	-	-	-	-	-	-
<i>Filostomum</i> sp.	-	-	-	1(11.1)	1(11.1)	3(33.3)	4(44.4)	-	-	-
<i>A. ohradiformis</i>	-	-	-	-	1(7.1)	4(28.6)	1(7.1)	5(35.7)	2(14.3)	2(14.3)
<i>M. pygmaeum</i>	-	-	-	-	-	-	-	1(100.0)	-	-
<i>P. japonicum</i>	-	-	-	1(7.1)	-	-	-	1(7.1)	8(57.1)	2(14.3)
<i>Remicola</i> sp.	-	-	-	-	-	-	-	-	-	-
<i>C. strigoides</i>	-	-	-	-	-	-	1(50.0)	1(50.0)	-	-
<i>A. burti</i>	-	-	-	1(50.0)	1(50.0)	-	-	-	-	-
<i>A. gracilis</i>	-	-	-	-	-	-	1(50.0)	1(50.0)	-	-
<i>Hymenolepis</i> spp.	-	-	-	16(8.8)	64(35.4)	53(29.3)	27(14.9)	14(7.7)	2(1.1)	2(1.1)
<i>C. megalops</i>	-	-	-	-	-	-	-	-	-	-
Fimbriariinae	-	-	-	12(46.2)	7(26.9)	5(19.2)	-	-	-	-
<i>C. aratis</i>	-	-	-	-	-	-	-	-	-	26(88.4)
<i>C. nyroonarium</i>	-	-	-	-	-	2(5.3)	4(10.5)	4(10.5)	-	-
<i>A. acutum</i>	1(1.3)	6(7.9)	66(86.8)	-	-	-	-	1(1.3)	-	-
<i>Amidoctium</i> sp.	-	-	6(85.7)	-	-	-	-	-	-	-
<i>E. uncinatum</i>	-	-	2(100.0)	-	-	-	-	-	-	-
<i>S. californica</i>	-	1(3.4)	28(96.6)	-	-	-	-	-	-	-
<i>S. o. crassicauda</i>	-	1(7.7)	11(84.6)	-	-	-	-	1(7.7)	1(1.6)	-
<i>S. formosensis</i>	-	1(6.3)	15(93.7)	-	-	-	-	-	-	-
<i>Streptocoara</i> sp.	-	1(50.0)	1(50.0)	-	-	-	-	-	-	-
<i>E. borealis</i>	-	1(100.0)	-	-	-	-	-	-	-	-
<i>E. hypognatha</i>	1(3.3)	25(83.3)	4(13.3)	-	-	-	-	-	-	-
<i>T. fissispina</i>	1(10.0)	9(90.0)	-	-	-	-	-	-	-	-
<i>T. somateriae</i>	6(9.8)	45(73.8)	3(13.1)	1(1.6)	-	-	-	-	-	-
<i>T. spinosa</i>	1(20.0)	4(80.0)	-	-	-	-	-	-	-	-
<i>Tetrameres</i> sp.	1(14.3)	5(71.4)	1(14.3)	-	-	-	-	-	-	-
<i>Sarcoema</i> sp.	-	-	-	-	-	-	-	-	-	-
Nematode sp. (Type 1)	-	1(100.0)	-	-	-	-	-	-	-	-
<i>P. altmani</i>	-	-	-	-	-	-	1(100.0)	-	-	-
<i>P. botulus</i>	-	-	-	-	-	6(15.4)	15(38.5)	13(33.3)	-	-
<i>C. constrictum</i>	-	-	-	-	-	-	-	1(100.0)	-	-
<i>C. sudsucho</i>	-	-	-	-	-	1(5.9)	3(17.6)	11(64.7)	1(5.9)	-

TABLE 16. GEOGRAPHICAL DISTRIBUTION OF HELMINTH PARASITES RECOVERED FROM SCOTERS IN THIS STUDY.

PARASITE	A	B	C
<i>A. terrigalensis</i>	-	+	-
<i>Z. lunata</i>	+	-	-
<i>N. attenuatus</i>	+	+	-
<i>C. verrucosa</i>	-	-	+
<i>T. elegans</i>	+	+	-
<i>E. oastenea</i>	+	+	-
<i>G. bursericola</i>	+	+	-
<i>G. choledocus</i>	+	+	+
<i>G. deliciosus</i>	+	+	+
<i>Gymnophallus</i> sp. Type 1	-	+	-
<i>Gymnophallid</i> sp.	-	+	-
<i>Gymnophallus</i> spp.	+	-	-
<i>E. revolutum</i>	-	+	-
<i>D. ergregia</i>	+	-	-
<i>H. incisa</i>	-	+	-
<i>A. melanittae</i>	+	-	-
<i>Psilostomum</i> sp.	+	-	-
<i>A. charadriiformis</i>	+	-	-
<i>M. pygmaeum</i>	-	+	-
<i>P. japonicum</i>	+	-	-
<i>Renicola</i> sp.	+	+	-
<i>C. strigeoides</i>	-	+	-
<i>A. burtti</i>	-	+	-
<i>A. gracilis</i>	+	+	-
<i>Hymenolepis</i> spp.	+	+	+
<i>C. megalops</i>	+	-	-
Fimbriariinae	+	+	-
<i>C. anatis</i>	-	+	-
<i>C. nyrocinorum</i>	+	+	-

TABLE 16. CONTINUED

PARASITE	A	B	C
<i>A. acutum</i>	+	+	+
<i>Amidostomum</i> sp.	+	+	-
<i>E. uncinatum</i>	+	+	-
<i>S. californica</i>	+	-	-
<i>S. c. crassicauda</i>	+	+	-
<i>S. formosensis</i>	+	+	-
<i>Streptocara</i> sp.	+	-	-
<i>E. borealis</i>	+	-	-
<i>E. hypognatha</i>	+	+	-
<i>T. fistipina</i>	+	+	-
<i>T. somateriae</i>	+	+	+
<i>T. spinosa</i>	+	+	-
<i>Tetrameres</i> sp.	+	-	+
<i>Sarconema</i> sp.	-	+	-
Nematode sp. Type 1	-	+	-
<i>P. altmani</i>	-	+	-
<i>P. botulus</i>	+	+	+
<i>C. constrictum</i>	-	+	-
<i>C. sudsuche</i>	+	+	-

A = Pacific sample
 B = Eastern Atlantic sample
 C = Western Atlantic sample

Atlantic hosts. It is apparent from Table 16 that the hosts in the Western Atlantic carried the greatest number of species.

A number of new host records were established during the course of this work (Table 17). *Tristriata elegans*, *D. egregia*, *P. japonicum* and *C. sudsuche* are reported from North America for the first time. In addition, 4 parasites were recorded from Canada for the first time; namely, *C. strigeoides*, *A. burti*, *C. constrictum* and *F. altmani*. *Tetrameres somateriae* were recorded from Europe for the first time.

To further address parasite distribution cluster analysis was performed on the data to determine if birds from a certain locality were more similar to birds from the same locality with respect to their parasite fauna than to birds from other localities. This would tell the author whether or not a bird or a sample of birds could be separated from birds in other sample areas examined on the basis of their helminth fauna. Due to the fact that Hymerolepids and some specimens of *Gymnophallus* were not identified to the species level these groups of parasites were omitted from the analysis. It is unfortunate that the hymenolepids were not identified to the species level as it is likely that the different geographical areas sampled would have yielded different species within the genus. If this hypothesis is true then the hymenolepids would have caused birds from different geographical areas to be more similar to other birds from their area than to birds from any of the other areas sampled.

TABLE 17. NEW HOST RECORDS NOTED IN THIS STUDY.

SURF SCOTER	WHITE-WINGED SCOTER	BLACK SCOTER
<i>A. terrigalenste</i>	<i>A. terrigalenste</i>	<i>A. terrigalenste</i>
<i>Z. lunata</i>	<i>G. deliciosus</i>	<i>E. castenea</i>
<i>N. attenuatus</i>	<i>D. egregia</i>	<i>G. choledocus</i>
<i>T. elegans</i>	<i>A. charadriiformis</i>	<i>A. charadriiformis</i>
<i>E. castenea</i>	<i>Reticola</i> sp.	<i>P. japonicum</i>
<i>G. choledocus</i>	<i>T. somateriae</i>	<i>C. strigoides</i>
<i>G. deliciosus</i>	<i>T. spinosa</i>	<i>A. byrti</i>
<i>A. melanittas</i>	<i>P. botulus</i>	<i>T. somateriae</i>
<i>A. charadriiformis</i>		<i>T. spinosa</i>
<i>P. japonicum</i>		<i>P. botulus</i>
<i>A. gracilis</i>		<i>A. icterodes</i>
<i>C. nyrocinarum</i>		<i>H. loomisi</i>
<i>A. acutum</i>		
<i>E. borealis</i>		
<i>E. hypognatha</i>		
<i>T. fiesispina</i>		
<i>T. somateriae</i>		
<i>T. spinosa</i>		
<i>Sarcönema</i> sp.		
<i>P. botulus</i>		
<i>C. sudsuche</i>		
<i>A. dehtatus</i>		
<i>A. icterodes</i>		
<i>H. leurocanthum</i>		
<i>H. loomisi</i>		

"Cluster analysis is a multivariate approach to sorting data into groups so that members of any particular group are more similar to each other than to non members" (Bush, 1980). In this study, the average linkage method was employed with Jaccard's coefficient (see Wishart, 1978). Cluster analysis, in this study, is used as a technique to reduce data with the resulting dendrogram depicting the similarity of individual birds based on presence of helminth species.


Cluster analysis was performed individually on each of the three bird species and on the three bird species as a group. The reason for the latter was to see if the three species of birds could be separated on the basis of presence of specified helminths in their parasite faunas.

Figures 3, 4 and 5 depict the result of cluster analysis on surf, white-winged and black scoters respectively. Due to the very low levels of similarity displayed in each dendrogram, it is, therefore, not possible to separate birds of either species on the basis of presence of helminths in their parasite faunas by geographical areas. This, however, is not to say that certain individual birds would not be able to be separated by a certain parasite e.g. the presence of *Perilostomum* sp. would strongly suggest that the bird came from British Columbia. Overall, one can state that due to the high levels of similarity that the parasite fauna of scoters, for the most part, is independent of geography.

The cluster analysis performed on all three species of birds together did not allow the author to separate either of the three



FIGURE 3.
Cluster analysis comparing individual surf scoters with respect to presence of helminth species.



104

- .05

- .1

- .2

- .3

- .4

- .5

- .6

5 6 7 8 9

- LA7
- LA17
- BC19
- BC39
- BC15
- LA5
- NB9
- NB3
- NB2
- BC27
- BC17
- BC11
- BC12
- BC4
- BC3
- BC8
- BC2
- BC1
- LA9
- BC33
- BC30
- BC44
- BC34
- BC26
- BC31
- LA16
- BC40
- BC35
- BC24
- LA14
- LA1
- BC42
- BC36
- BC32
- BC38
- BC18
- BC23
- BC16
- NB11
- LA11
- LA21

LA11
LA24
LA19
LA22
LA13
LA18
LA20
LA12
LA10
LA15
LA4
LA3
LA8
LA2
LA23
BC9
BC37
BC25
BC20
BC6
BC41
BC22
BC21
BC29
BC14
BC7
BC43
BC28
NB7
NB4
NB12
NB6
LA6
BC10
NB10
NB5
BC13
BC5
NB8
NB1

FIGURE 4.


Cluster analysis comparing individual white-winged scoters with respect to presence of helminth species.

10f

0
1
2
3
4
5
6
7
8

FIGURE 5.

Cluster analysis comparing individual black scoters
with respect to presence of helminth species.



10f

- .05

- .1

- .2

- .3

- .4

- .5

- .6

- .7

- .8

- .9

242

3

4

5

6

7

8

9

10

- LA5
- LA4
- LA2
- LA1
- BC4
- BC2
- LA3
- BC13
- BC9
- BC10
- BC8
- BC7
- BC3
- BC1
- BC11
- BC6
- NB9
- BC12
- BC5
- NB13
- NB11
- NB10
- NB8
- NB7
- NB5
- NB4
- NB3
- NB12
- NB6
- NB2
- NB1



host species by presence of helminths in their parasite fauna. This is what one might expect when one considers the close phylogenetic relationship of the three hosts. While they do occupy different niches and their diets differ (see Cottam, 1939), they have very similar parasite faunas, which can be used as further evidence for the suppression of the genus *Oidemia* as discussed in the Introduction.

HOST SPECIFICITY

Researchers have long recognized two basic types of parasite-host specificity i.e. phylogenetic and ecological specificity. Phylogenetic specificity deals with those parasites that have evolved with their hosts and form a very close relationship. Often this relationship is so intimate that the parasite can only develop in its "natural host". Ecological specificity refers to parasites that develop in many hosts (their host requirements are not stringent) with the limiting factor most often being the chance of a host consuming the right intermediate host. Rysavý (1966) states "The composition of the fauna of cestodes, parasitizing birds of various orders or families, does not only depend on their phylogenetic relations but also on the way of living and the composition of their food." Therefore, the biology of the host is as important as or more important in ecological specificity than the diet of the host.

In addition, an animal that is a member of a large group of closely related animals (eg. an anatid) has the possibility of more potential parasites than does an animal that belongs to a small group. In ecological specificity the possibility of infection rests

largely on a host organism eating an infected intermediate host which in turn depends largely on the number of parasites seeding an area. Most animal species have a primary food item with their diet being supplemented by secondary and tertiary food items. These secondary and tertiary food items in the diet of one anatid may be the primary food item in the diet of another anatid, thus, making it possible for a parasite to have many natural hosts, with certain hosts having a higher prevalence of certain parasites.

This leads to the problem of trying to classify the parasites of a host. Given the present knowledge of the life-cycles and ecology of many parasites, it is probably easier to classify a parasite as an inhabitant of a higher taxon rather than a species. A modification of the system devised by McDonald (1969b) that categorizes parasites as accidental, rare, infrequent, frequent, common or very common will be employed in this work. A scheme such as this is much more accurate as it can take many studies into account. As in this study, the classifications of a parasite that was only recovered from a single host is questionable.

Table 18 lists the category that each parasite falls into and compares it to McDonald's (1969b) scheme. The status of some parasites (as given in this study) may change as further work is done. In addition to categorizing the parasites for each host species, they are also classified according to geographical origin, as their status changes with the environment.

TABLE 18. CATEGORIZATION OF PARASITES RECOVERED IN THIS STUDY.

PARASITE	SURF SCOTER			WHITE-WINGED SCOTER				BLACK SCOTER			MCDONALD'S CATEGORIZATION FOR WATERFOWL
	A	B	D	A	B	C	D	A	B	D	
<i>A. terrigenensis</i>	-	freq	inf.	-	freq	-	inf.	-	inf	inf	rare
<i>Z. lunata</i>	rare	-	rare	-	-	-	-	-	-	-	comm. (char.)
<i>N. attenuatus</i>	-	rare	rare	rare	-	-	rare	-	rare	rare	V. comm. (char.)
<i>C. verrucosa</i>	-	-	-	-	-	comm	inf.	-	-	-	-
<i>T. elegans</i>	freq	comm	freq	-	comm	-	inf.	-	-	-	-
<i>E. oastenea</i>	rare	rare	rare	inf.	inf.	-	inf.	inf.	rare	rare	-
<i>G. bureicoela</i>	freq	inf.	inf.	rare	inf.	-	inf.	freq	rare	inf.	inf.
<i>G. oholodocus</i>	rare	inf.	inf.	rare	inf.	freq	inf.	rare	-	rare	freq.
<i>G. deliciosus</i>	rare	freq	inf.	freq	comm	comm	comm.	-	freq	inf.	inf.
<i>Gymnophallus</i> sp. Type 1	-	inf.	rare	-	comm	-	inf.	-	freq	rare	-
<i>gymnophallid</i>	-	rare	rare	-	-	-	-	-	freq	rare	-
<i>E. revolutum</i>	-	-	-	-	-	-	-	-	rare	rare	V. comm. (char.)
<i>D. egregia</i>	-	-	-	comm	-	-	freq	-	-	-	-
<i>H. incisa</i>	-	-	-	-	inf.	-	rare	-	rare	rare	rare
<i>A. melanittae</i>	rare	-	rare	-	-	-	-	-	-	-	rare
<i>Psilostomum</i> sp.	freq	-	inf.	-	-	-	-	-	-	-	-
<i>A. charadriiformis</i>	inf.	-	rare	rare	-	-	rare	freq	-	rare	acc.
<i>M. pygmaeum</i>	-	-	-	-	-	-	-	-	rare	rare	comm.

TABLE 18. CONTINUED.

PARASITES	SURF SCOTER			WHITE-WINGED SCOTER				BLACK SCOTER			McDONALD'S CATEGORIZATION FOR WATERFOWL
	A	B	D	A	B	C	D	A	B	D	
<i>P. japonicum</i>	inf.	-	rare	freq.	-	-	inf.	inf.	-	rare	-
<i>Renicola</i> sp.	inf.	-	rare	freq.	-	-	rare	-	-	-	-
<i>C. strigeoides</i>	-	-	-	-	-	-	-	-	rare	rare	rare
<i>A. burtti</i>	-	-	-	-	-	-	-	-	rare	rare	rare
<i>A. gracilis</i>	-	rare	rare	-	-	-	-	-	-	-	V. comm. (char.)
<i>C. megalops</i>	-	-	-	rare	-	-	rare	-	-	-	V. comm. (char.)
Fimbriariinae	-	freq.	inf.	-	-	-	-	rare	freq.	inf.	freq.
<i>C. anatis</i>	-	rare	rare	-	-	-	-	-	-	-	V. comm. (char.)
<i>C. nyrocinarum</i>	inf.	rare	inf.	comm.	V. comm.	-	comm.	inf.	rare	rare	probably freq.
<i>A. aoutum</i>	comm.	comm.	comm.	freq.	comm.	comm.	comm.	V. comm.	comm.	comm.	V. comm. (char.)
<i>E. uncinatum</i>	-	-	-	-	-	-	-	inf.	rare	rare	V. comm. (char.)
<i>S. californica</i>	comm.	-	freq.	rare	-	-	rare	V. comm.	-	freq.	inf.
<i>S. c. crassicauda</i>	inf.	rare	inf.	-	inf.	-	rare	freq.	rare	inf.	V. comm. (char.)
<i>S. formosensis</i>	inf.	inf.	inf.	-	inf.	-	rare	-	-	-	inf.
<i>E. borealis</i>	rare	-	rare	-	-	-	-	-	-	-	rare
<i>E. hypognatha</i>	comm.	inf.	freq.	-	-	-	-	comm.	inf.	freq.	inf.

TABLE 18. CONTINUED.

PARASITE	SURF SCOTER			WHITE-WINGED SCOTER				BLACK SCOTER			MCDONALD'S CATEGORIZATION FOR WATERFOWL
	A	B	D	A	B	C	D	A	B	D	
<i>T. flastospina</i>	inf.	inf.	inf.	rare	-	-	rare	inf.	inf.	inf.	V. comm. (char.)
<i>T. somaterias</i>	comm.	comm.	comm.	freq.	comm.	comm.	freq.	-	rare	rare	inf.
<i>T. spinosa</i>	rare	rare	rare	-	inf.	-	rare	-	rare	rare	comm. in Athyini in N.A.
<i>Sarcónema</i> sp.	-	rare	rare	-	-	-	-	-	-	-	-
Nematode sp. Type 1	-	rare	rare	-	-	-	-	-	-	-	-
<i>F. altmani</i>	-	rare	rare	-	-	-	-	-	-	-	rare
<i>F. botulus</i>	inf.	-	inf.	freq.	-	freq.	freq.	comm.	rare	freq.	freq.
<i>G. constrictum</i>	-	-	-	-	-	-	-	-	rare	rare	inf.
<i>C. sudsuche</i>	freq.	rare	inf.	-	-	-	-	-	-	-	-

V. comm. - Very Common 60% A - Pacific
 Comm. - Common 30% 60% B - Western Atlantic
 freq. - frequent 15% 30% C - Eastern Atlantic
 inf. - infrequent 5% 15% D - Overall
 rare - rare 5%
 acc. - accidental (Parasite does not mature in host)
 char. - characteristic for waterfowl

NOTE: *Gymnophallus* spp. and *Hymenolepis* spp. are omitted.

LIFE-CYCLES

The genus *Melanitta*, Tribe Mergini, are known to breed in a freshwater environment and inhabit the marine environment for their non-breeding season. The males of the genus spend very little time in the freshwater environment as they depart from the breeding grounds after the nest is built whence they undergo a migration to the sea to moult.

Our lack of knowledge of the life-cycle of many anatid parasites makes it difficult to assess the percentage of scoter parasites that carry out their larval life-cycle in the marine or freshwater environment; although it is possible, in many cases to predict with some degree of accuracy the most likely environment in which the parasite completes its life-cycle. Table 19 details the environment in which certain known life-cycles are completed and gives speculations on others. The life-cycle is presently known for 22 of the parasites recovered in this study; of these 22, 13 (59.0%) can complete their life-cycle in the marine environment and the same number in the freshwater environments (Note: 4 (18.0%) can complete their life-cycle in either environment). Thus, the scoters examined harboured more parasites which cycle through the marine environment than the freshwater environment. Of the 15 parasites believed to have freshwater life-cycles, 5 (*E. revolutum*, *C. strigatodes*, *A. burti*, *E. borealis* and *C. constrictum*) were only located in immature females and 2 (*A. gracilis* and *C. megalops*) only in immature males. This means that adult males harboured 8 (53.0%)

TABLE 19. ENVIRONMENT IN WHICH PARASITE LIFE-CYCLES ARE COMPLETED.

MARINE	FRESHWATER	BOTH	UNKNOWN but probably	
			MARINE	FRESHWATER
<p>Trematoda:</p> <p><i>A. terrigalensis</i></p> <p><i>G. bursicola</i></p> <p><i>G. choledocus</i></p> <p><i>G. delictosus</i></p> <p><i>A. melanittae</i></p> <p><i>A. charadriiformis</i></p> <p><i>M. pygmaeum</i></p> <p><i>Gymnophallus</i> sp. Type 1</p> <p><i>Gymnophallid</i> sp.</p>	<p>Trematoda:</p> <p><i>Z. lunata</i></p> <p><i>N. attenuatus</i></p> <p><i>E. revolutum</i></p> <p><i>A. gracilis</i></p> <p><i>C. verrucosa</i></p> <p>Cestoda:</p> <p><i>C. megalops</i></p> <p>Fimbriariinae</p> <p>Nematoda:</p> <p><i>T. spinosa</i></p> <p>Acanthocophala:</p> <p><i>C. constrictum</i></p>	<p>Nematoda:</p> <p><i>C. anatis</i></p> <p><i>S. c. grassicauda</i></p> <p><i>S. formosensis</i></p> <p><i>T. fissisepina</i></p>	<p>Trematoda:</p> <p><i>T. elegans</i></p> <p><i>D. ergregia</i></p> <p><i>Palostomum</i> sp.</p> <p><i>H. incisa</i></p> <p><i>Reticola</i> sp.</p> <p>Nematoda:</p> <p><i>C. nyrocinarum</i></p> <p><i>S. californica</i></p> <p><i>T. somateriae</i></p> <p>Acanthocephala:</p> <p><i>F. altmani</i></p> <p><i>C. sudsuche</i></p>	<p>Trematoda:</p> <p><i>C. strigeoides</i></p> <p><i>A. burti</i></p> <p>Cestoda:</p> <p><i>Hymenolepis</i> s.l.</p> <p>Nematoda:</p> <p><i>E. hypognatha</i></p> <p><i>E. borealis</i></p> <p><i>E. uncinatum</i></p>

Life-cycle unknown:

Trematoda:

E. castanea
P. japonicum

Nematoda:

Sarconema sp. utilizes a dipteran vector
- Nematode sp. (Type 1)

of the freshwater species of parasites while males in general harboured 10 (66.7%) and females 13 (87.0%). Of the marine parasites males contained 21 (91.0%) and females 19 (83.0%). This is possibly the result of the male moult migration which causes them to spend more time in the sea.

THE INFLUENCE OF AGE & SEX OF THE HOST ON PARASITE BURDEN

Buscher (1965) stated that "the extensity and intensity of invasion of parasites is greater in juvenile ducks than in adults." The reason for this was believed to be that young ducks probably did not acquire an age immunity to parasite infection for some time after hatching (within their first year of life). The immature birds necropsied in this study were prebreeding birds (as stated in the Introduction), however, they all contained a Bursa of Fabricius. In the strictest sense the results of Buscher (1965) and other workers in this area do not apply to the immature birds in this study. However, the author feels that it is important that this theory be checked utilizing two recognizable age classes; namely, prebreeding (bursate) and breeding adult birds.

Bychovskaya-Pavlovskaya (1962, *vide* Graham, 1966) stated that "the trematode fauna is more diverse in adults than in immatures", with Graham (1966) noting that this fact can be extended to include the total helminth fauna.

Table 20 lists the parasites found only in immature and adult birds respectively. A total of 43 species of helminths were re-

TABLE 20. Parasites found to occur in only immature or adult birds.

IMMATURE BIRDS	ADULT BIRDS
<i>E. revolutum</i>	<i>A. terrigenensis</i>
<i>G. choledocus</i>	<i>D. egregia</i>
<i>C. strigeoides</i>	<i>H. incisa</i>
<i>A. burti</i>	<i>A. melanittae</i>
<i>A. gracialis</i>	<i>Psilostomum</i> sp.
<i>C. anatis</i>	<i>C. verrucosa</i>
<i>E. borealis</i>	
Nematode sp. I	
<i>Sarconema</i> sp.	
<i>F. altmani</i>	
<i>C. constrictum</i>	
<i>C. megalops</i>	

covered in this study with immatures harbouring 37 species and adults 30 species. This contradicts both Bychovskaja-Pavlovskaja (1962) and Graham (1966) as the trematode fauna of immature birds was as diverse as that of adults and immature birds had a more diverse nematode, cestode and acanthocephalan fauna than adult birds.

To further address the question of age and sex influences on parasite G-tests (see Sokal and Rohlf, 1969) were performed on the data in Appendix 3 (i.e. the number of infected and uninfected adult males was run against the number of infected and uninfected females for each age class by individual parasites). The same procedure was carried out for total adults versus total immatures, total males versus total females and total immature males versus total immature females.

The result of this would tell if a parasite occurred in significantly more adult male than adult female birds, more immature male than immature female birds, more adult than immature birds or more male than female birds. All the aforementioned tests were carried out for the surf scoters with only the total immature male versus total immature not being performed on the black scoter and total adult male versus total adult female and total immature female versus total immature male not being performed on the white-winged scoters. The reason for these tests not being performed on the white-winged and black scoters was that the sample size of birds was less than 5.

The following parasites infected significantly ($G = 3.814$ with 1 degree of freedom at the .05 level) more immature surf scoters than adult scoters; *G. choledocus*, *G. bureicola*, *Hymenolepis* spp., *Frimbriariinae*, *S. c. crassicauda*, *S. formosensis*, *E. hypognatha*, *C. nyrocinarum*, *T. fistispina* and *T. somateriae*. *Hymenolepis* spp. also preferred immature males to immature females. *Gymnophallus choledocus* preferred immature white-winged scoters to adult white-winged scoters and *Frimbriariinae* and *C. nyrocinarum* preferred immature black scoters to adult black scoters, while *S. californica* preferred female black scoters to male black scoters. In no instance did any parasite infect significantly more adult than immature birds. These data indicate that the extensity of invasion for immature birds was greater than that of adults for the forementioned parasites and species.

The reasons for this include the following possibilities:

- (1) immature (prebreeding) birds have not yet acquired the level of immunity to parasites that adult birds have (as yet there has been no physiological study to prove or disprove this statement).
- (2) that their diets differ from breeding birds - different ecological habitats i.e. breeding birds tend to spend more time on the breeding grounds which restricts their foraging range or (3) differences in their biology.

Since there was a significant difference in the extensity of invasion of certain parasites in certain host species of given age and sex, the author used a Mann-Whitney U-test (see Sokal and Rohlf, 1969) to determine if there was a significant difference in the in-

tensity of helminth infection between immatures and adults. The results of these tests all proved to be insignificant. This suggests that the observed significant differences in extensity of invasion of certain species and sexes may more likely be due to dietary differences.

SUMMARY

1. One hundred and seventy-five scoters from 4 locations (British Columbia, New Brunswick, Labrador and Norway) were necropsied for metazoan parasites. A total of 94 surf, 36 white-winged and 45 black scoters were examined.
2. Ninety-one percent (86 surf, 33 white-winged and 40 black scoters) of the specimens examined contained parasites.
3. A total of 37 genera and approximately 50 species of metazoan parasites were recovered. More specifically, Necropsies yielded 29 genera (39 species), 21 genera (28 species) and 25 genera (33 species) of parasites from the surf, white-winged and black scoters respectively.
4. Parasite species are considered individually and their prevalence and intensity of infection is noted for each host species.
5. The locations in which parasites were found are discussed and are presented in tabular form with the number and percent of times a parasite occurred in a given area.
6. The geographical distribution of helminth parasites found in this study is presented in tabular form.
7. Cluster analysis revealed that species of hosts nor individuals of a species of a certain host could not be separated geographically based on the degree of similarity in their parasite faunas.
8. A total of 45 new host records (25 for the surf, 8 for the white-winged and 12 for the black scoter) were noted as well as 4 North American, 4 Canadian and 1 European Record.
9. The helminth parasites were categorized for each host and this categorization was compared to that of McDonald's (1969) categorization for waterfowl.
10. It was shown that there was no difference in the intensity of helminth infection in immature versus adult scoters.
11. Finally, a checklist of parasites which have been recovered from these hosts was compiled and included as an appendix to this manuscript.

LITERATURE CITED

- Andrews, S. E. 1974. Helminth parasites of the Common (North American) Crow (*Corvus brachyrhynchos* Brehm, 1822) in insular Newfoundland. Unpublished M.Sc. thesis, Memorial University of Newfoundland. 82pp.
- Beaudette, F. R. 1939. Flukes in the respiratory tract of ducks. J. Am. Vet. Med. Ass., 47: 44.
- Belopol'skaia, M. M. 1952. The trematode family Microphallidae Travassos, 1920. IN: Skrjabin, K. I., Trematody Zhivotnykh i cheloveka, Osnovy trematodologii, 6: 619-756. (Russ. text).
- Belopol'skaia, M. M. 1953. *Balanus balanoides* L. as an intermediate host for some parasitic worms. Doklady AN SSSR, W.S. 91: 437-440. (Russ. text).
- Bezubik, B. 1956. The helminth fauna of wild ducks (subfamily Anatinae) of the Lublin and Bialystok districts. Acta Parasit. Polon., 4: 408-510.
- Bishop, C. A. 1971. Helminth parasites of the Common Eider Duck (*Somateria mollissima* L.) in Newfoundland and Labrador. Unpublished M.Sc. thesis, Memorial University of Newfoundland. 82pp.
- Bittner, H. 1923. *Schistogonimus rarus* (Braun), ein seltener Trematode in der Bursa Fabricii einer an *Tetrameres*-Invasion gestorbenen Hausente. Arch. Wissensch. u. Prakt. Tierh., 50: 253-261.

- Blankespoor, H. D. 1974. Host-induced variation in *Plagiorchis noblei* Park, 1936 (Plagiorchidae: Trematoda). Amer. Midl. Nat., 92: 415-433.
- Bourgeois, C. E. & W. Threlfall, 1979. Parasites of the greater shearwater (*Puffinus-gravis*) from Newfoundland, Canada. Can. J. Zool., 57: 1355-1357.
- Bowers, E. A. 1966. A description of *Metogymnophallus jamesoni* sp. nov. from the intestine of the common scoter, *Melanitta nigra* L. Ann. Mag. Nat. Hist. s. XIII, 8: 277-283.
- Bowers, E. A. & B. L. James. 1967. Studies on the morphology, ecology, and life cycle of *Metogymnophallus minutus* (Cobbold, 1859) comb. nov. (Gymnophallidae). Parasitol. 57: 281-300.
- Boscher, H. N. 1965. Dynamics of the intestinal helminth fauna in three species of ducks. J. Wildl. Mgmt., 29: 772-781.
- Bush, A. D. 1980. Faunal similarity and infracommunity structure in the helminths of Lesser Scaup. (Unpublished) Phd. thesis University of Alberta. 254 pp.
- Bykhovskaia-Pavlovskaiâ, I.E. 1962. Trematodes of the bird fauna of USSR. Izdat. AN SSSR, Moskva, 407pp. (Russ. text, Eng. Summary).
- Carney, A. & A. Geis. 1964 (MS). Preliminary key, age and sex of duck wings. Can. Wild. Ser., (Unpublished) M.Sc. thesis. 56 pp.
- Cheatum, E. L. 1938. *Tanaisia pelidnae* n. sp. and *Ochtipedum tracheicola* (Trematoda). J. Par., 24: 135-141.

- Ching, H. L. 1965a. Life cycles of *Lacunovermis conspicuus* n. gen., n. sp. and *Metogymnophallus multigemmulus* n. gen., n. sp. (Gymnophallidae: Trematoda) from *Macoma inconspicua* and diving ducks from Vancouver, Canada. Proc. Helm. Soc. Wash., 32: 53-63.
- Ching, H. L. 1965b. Systematic notes on some North American microphallid trematodes. Proc. Helm. Soc. Wash., 32: 140-148.
- Ching, H. L. 1973. Description of *Gymnophallus somateriae* (Levinsen, 1881) from *Macoma inconspicua* and diving ducks from Vancouver, Canada. Can. J. Zool., 51: 801-806.
- Clark, G. M., D. O'Meara, & J. W. Van Weelden. 1958. An epizootic among eider ducks involving an acanthocephalid worm. J. Wildl. Mgmt., 22: 204-205.
- Clay, T. 1957. The Mallophaga of birds. 1st symposium on Host Specificity among Parasites of Vertebrates. Paul Atlinger S. A., Neuchatel, pp. 120-158.
- Cornwell, G. B. and A. B. Cowan. 1963. Helminth populations of the canvasback (*Aythya valisineria*) and host-parasite-environmental interrelationships. Trans. 28th N. Am. Wildl. & Nat. Res. Conf., pp. 173-198.
- Cottam, C. 1939. Food habits of North American diving ducks. U.S. Dep. Agric. Tech. Bull. No. 643. 140 pp.
- Czapliński, B. 1956. Hymenolepidae: Fuhrmann, 1907 (Cestoda) in some domestic and wild Anseriformes in Poland. [Abstr.] Wiadom. Parazytol., 2, Suppl.: 269-270. [Pol. text, Eng. summary].

- Czaplinski, B. 1962. Nematodes and acanthocephalans of domestic and wild Anseriformes in Poland. I. Revision of the genus *Amidostomum* Railliet et Henry, 1909. Acta Parasit. Polon., 10: 125-164.
- Czaplinski, B. 1973. Redescription of *Sobolevtoanthus kenaiensis* (Schiller, 1952) comb. n. (syn. *S. gladium* Spassky et Bobova, 1962) (Cestoda, Hymenolepidae). Acta Parasit. Polon., 24: 251-261.
- Deblock, S. 1971. Contribution à l'étude des Microphallidae Travassos, 1920 XXIV. Tentative de phylogénie et de taxonomie. Bull. Mus. Natl. Hist. Nat. Zool., 7: 1-468.
- Deblock, S. & P. Tran Van Ky. 1966. Contribution à l'étude des Microphallidae Travassos, 1920 (Trematoda). XII. Espèces d'Europe occidentale. Création de *Sphairiotrema* nov. gen.; considerations diverses de systematique. Ann. Parasitol., 41: 23-60.
- Delacour, J. 1954-64. The waterfowl of the world. Country Life, Ltd. London, Vol. 3, 270 pp.
- Denny, M. 1969. Life-cycles of helminth parasites: *Hymenolepis albertensis* using *Gammarus lacustris* as an intermediate host in a Canadian Lake. Parasitol., 59: 795-827.
- Dubois, G. 1967. Notes helminthologiques. I: Strigeidae Railliet (Trematoda). Rev. Suisse Zool., 74: 693-700.
- Dubois, G. 1968. Du statut de quelques Strigeata La Rue, 1926. III. Bull. Soc. Neuch. Sc. Nat., 91: 5-19.
- Emerson, K. C. 1972. Checklist of the Mallophaga of North America (North of Mexico) Parts I and II. Dugway Proving Ground, Dugway, Utah, Part I 171 pp., Part II 118 pp.

- Eveleigh, E. S. 1974. A study of the ectoparasites of alcids in Newfoundland. (Unpublished) M.Sc. thesis, Memorial University of Newfoundland. 176 pp.
- Farley, J. 1971. A Review of the Family Schistosomatidae: excluding the Genus *Schistosoma* from Mammals. J. Helminthol., XLV: 289-320.
- Fitzpatrick, C. & W. Threlfall, 1977. The ectoparasites of three species of seabirds from Newfoundland, Canada. Can. J. Zool., 55: 1205-1209.
- Filimonova, L. V. 1971. New species of Notocotylidae from Anseriformes in the Yakutsk ASSR. Trudy Gel'mint. Lab., 22: 211-214. (Russ. text).
- Frame, G. W. 1969. New definitive host, range extension, and notes on the morphology of *Tristriata anatis* Belopolskaia, 1953 (Trematoda, Notocotylidae) from southeast Alaska. Can. J. Zool., 47: 265-266.
- Fuhrmann, O. 1913. Nordische Vogelcestoden aus dem Museum von Göteborg. Göteborgs K. Vetensk. - O. Vitterhets - Samh. Handl., 4 f. (1911-12), V. 14-15, Medd. Göteborgs Mus. Zool. Avd. (1), 419 pp.
- Gallimore, J. R. 1964. Taxonomy and ecology of helminths of grebes in Alberta. (Unpublished) M.Sc. thesis, University of Alberta. 167 pp.
- Garden, E. A., C. Rayski & V. M. Thom. 1964. A parasitic disease in eider ducks. Bird Study, 11: 280-287.
- Garkavi, B. L. 1950. Streptocariasis of ducks. Veterinariia, 27: 30 (Russian text).

- Gibson, G. G. 1964. Taxonomic and biological observation on *Streptocara californica* (Gedoeelst, 1919) Gedoeelst + Liegeois, 1922 and the genus *Streptocara* (Nematoda: Acuariidae). Can. J. Zool., 42: 773-783.
- Gibson, G. B. 1968. Species composition of the genus *Streptocara* Railliet et al., 1912 and the occurrence of these avian Nematodes (Acuariidae) on the Canadian Pacific Coast. Can. J. Zool. 46: 629-645.
- Graham, L. C. 1966. The ecology of helminths in breeding populations of Lesser Scaup (*Aythya affinis*, Eyton) and Ruddy ducks (*Oxyura jamaicensis*, Gmelin). (Unpublished) M.Sc. thesis, University of Alberta. 90 pp.
- Harrison, J. M. 1955. A case of nodular taeniasis due to *Fillicollis anatis* in an eider duck *Somateria mollissima* (Linnaeus). Bull. Br. Orn. Club., 75: 121-123.
- Holmes, J. C. 1973. Site selection by parasitic helminths: inter-specific interactions, site segregation, and their importance to the development of helminth communities. Can. J. Zool., 51: 333-347.
- Jameson, H. L. & W. Nicoll. 1913. On some parasites of the scoter duck (*Oedemia nigra*) and their relation to the pearl-inducing trematode in the edible mussel (*Mytilus edulis*). Proc. Zool. Soc. London, 12: 53-63.
- Jerisen, D. N. 1971. *Eucotyle castenea* n. sp. and other trematodes of the family Eucotylidae from birds in British Columbia. Can. J. Zool., 49: 1053-1058.
- Johnsgard, P. A. 1978. Ducks, geese and swans of the world. University of Nebraska, Press: Lincoln. 404 pp.

- Kossack, W. F. K. 1911. Über Monostomiden. Zool. Jahrb. Syst.,
31: 491-590.
- Leonov, V. A., K. M. Tyzhikov, A. K. Tsimballuk and O. I. Belogunv.
1963. Trematodes of anseriformes from Kamchatka. Trudy Gel'm.
Lab. Akad. Nauk SSSR, 15: 196-207. (Russian text).
- Linton, E. 1928. Notes on trematode parasites of birds. Proc.
U.S. Nat. Mus., 73: 1-36.
- Loos-Frank, B. 1968. *Psilostomum* (Crepl., 1829). Z. Parasitenk.,
30: 185-191.
- Loos-Frank, B. 1969. Zur Kenntnis des gymnophalliden Trematoden
des Nordseesaumes. I. Die Alternation Zyklus von
Gymnophallus choleocheus Odhner, 1900. Ztschr. Parasitenk.,
32: 135-156.
- Loos-Frank, B. 1971. Zur Kenntnis des gymnophalliden Trematoden
des Nordseeraumes. III. *Gymnophallus gibberosus* n. sp.
and seine Metacercarie. Ztschr. Parasitenk., 35: 270-281.
(Eng. Summ.).
- Madsen, H. 1945. The species of *Capillaria* (Nematodes, Trichinell-
oides) parasitic in the digestive tract of Danish gallinaceous
and anatine game birds, with a revised list of species of
Capillaria in birds. Danish Rev. Game Biol., 1, 112 pp.
- McDonald, M. E. 1969. Catalogue of helminths of waterfowl (Anatidae).
U.S. Dept. Int., Spec. Sci. Rep. - Wildl. No. 126. 692 pp.
- McDonald, M. E. 1974. Key to the Nematodes reported in Waterfowl.
Department of Int., Bur. Sport. Fish Wildl. Res. Pub. 122.
44 pp.

- Mettrick, D. F. 1959. *Zygoootyde lunata*. A redescription of *Zygoootyde lunata* (Diesing, 1836) Stunkard, from *Anas platyrhynchos domesticus* in southern Rhodesia. Rhodesia Agric. J., 56: 197-198.
- Mehlis, E. 1831. Novae observationes de entozois. Auctore Dr. F. C. H. Creplin. Isis (Oken), 1: 68-99, 2: 166-199.
- Odening, K. 1963. Echinostomatoidea, Notocotylata and Cyclocoelida (Trematoda, Digenea, Redicoinnei) aus Vogeln des Berliner Tierparks. Bijdr. Dierkunde, 33: 37-60.
- Oshmarin, P. G. 1963. Parasitic worms of mammals and birds in the Maritime Territory. Izd-vo AN SSSR, Moskva, 322 pp. (Russian text).
- Palma, R. L. 1978. Slide-mounting of Lice: A detailed description of the Canada Balsam technique. N.Z. Entomol., 6: 432-436.
- Palmer, R. S. (ed.) 1976. Handbook of North American birds. Vol. 3. New Haven, Yale University Press. 560 pp.
- Perry, M. L. 1942. A new species of the Acanthocephalan genus *Filicollis*. J. Parasitol., 28: 385-388.
- Petrochenko, V. I. 1949. New species of acanthocephala from birds of central Asia. Trudy Gel'mint Lab. AN SSSR, 2: 114-127.
- Podesta, R. B. & I. C. Holmes, 1970. Hymenolepid cysticercoids in *Hyalella asteca* of Cooling Lake, Alberta: Life cycles and descriptions of 4 new spp.. J. Parasitol., 56: 1124-34.
- Price, R. D. 1976. A review of the genus *Holomenopon* (Mallophaga: Menoponidae) from the Anseriformes. Ent. Soc. Amer. Annals, 64: 633-646.

- Rayski, C and M. A. Fahmy, 1962. Investigation on some trematodes of birds from East Scotland. Z. Par., 22: 186-195.
- Reimer, L. 1963. Zur Verbreitung der Adulti and Larvenstadien der Familie Microphallidae Viana, 1924, (Trematoda, Digenea) in der Mittleren Ostsee. Z. Par., 23: 253-273.
- Rhode, K. 1964. Some current problems in helminth-taxonomy. Mimeographed Material from Proc. First. Internat. Congr. Parasitol., Rome.
- Ryšavý, B. 1966. The occurrence of cestodes in the individual orders of birds and the influence of food on the composition of the fauna of bird cestodes. Folia Parasitol. (Prague), 13: 158-169.
- Ryzhikov, K. M. 1962. *Microsomoganthus melanittae*, a new cestode species from the white-winged scoter (*Melanitta deglandi*). Trudy Gel'mint. Lab. AN SSSR, 12: 102-105. (Russ. text).
- Ryzhikov, K. M. 1963a. Helminth fauna of wild and domestic anserine birds of the Far East. Trudy Gel'mint. Lab. AN SSSR, 13: 78-132. (Russ. text).
- Ryzhikov, K. M. 1963b. *Poiloostoma borealis* sp. nov. and *Gymnophallus minor* sp. nov. - New trematodes from birds of the order Anseriformes. Helminthologia, 4: 424-429. (Russ. text; Eng., Fr., Ger. summaries).
- Ryzhikov, K. M., V. A. Isomov & A. K. Tsimbaliuk. 1964. New helminth of Anseriform birds - *Australapatemon skrjabini* sp. nov. (Trematoda: Strigeidae). Trudy Gel'mint. Lab. AN SSSR, 14: 182-186. (Russian text).

- Schmidt, G. D. 1965a. *Polymorphus swartzii* sp. n. and other Acanthocephala of Alaskan ducks. J. Parasitol., 51: 809-813.
- Schmidt, G. D. 1965b. *Corynosoma bipapillum* sp. n. from Bonaparte's gull *Larus philadelphia* in Alaska, with a note on *C. constrictum* Van Cleave, 1918. J. Parasitol., 51: 814-816.
- Serafin, C. 1957. On the occurrence of the fluke *Notocotylus attenuatus* Rudolphi, 1809 in geese. Medycyna Wet., 13: 398-399.
- Sokal, R. R. and F. J. Rohlf 1969. Biometry. W. H. Freeman and Company: San Francisco. 776 pp.
- Spasskaia, L. P. 1966. Cestodes of birds of the USSR. Hymenolepidae "Nauka", AN SSSR, AN Moldav. SSR, Inst. Zool., Moskva, 689 pp. (Russ. text).
- Stunkard, H. W. and J. R. Uzzmann, 1958. Studies on digenetic trematodes of the genera *Gymnophallus* and *Parvatrema*. Biol. Bull., 115: 276-302.
- Swales, W. E. 1963a. Morphological and biological studies on *Tetrameres crami* Swales, 1933, an important nematode parasite of ducks. J. Parasitol., 22: 528.
- Swales, W. E. 1963b. *Tetrameres crami* Swales, 1933 an important parasite of ducks in North America. Proc. North Am. Wildl. Wildl. Conf. P. 491-493.
- Tolkacheva, L. M. 1966. The cestode fauna of anserine birds of the Lower Yenisei and Noril Lake. Trudy Gel'mint. Lab. AN SSSR, 17: 211-239. (Russ. text).

- Tsimbaljuk, A. K. & V. A. Leonov. 1963. Two new species of trematodes from the diving ducks of Kamchatka. *Trudy Gel'mint. Lab. AN SSSR*, 13: 216-219. (Russ. text).
- Turner, B. C. 1974. The parasite fauna of the green-winged teal (*Anas crecca* L.) and blue-winged teal (*Anas discors* L.). (Unpublished) M. Sc. Thesis, Memorial University of Newfoundland. 63 pp.
- Webster, J. D. 1943. A revision of the Fimbriariinae (Cestoda, Hymenolepidae). *Tr. Am. Micr. Soc.*, 62: 390-397.
- Wishart, D. 1978. Cluster user manual. 3rd Edition. Univ. College, London, London. 175 pp.
- Xuan, S. 1962. Gel'minthofauna exotnitsy - promislivix ptic Niseego Amura. *Gelan*, XII: 284-300. (Russ. text).
- Yamaguti, S. 1934. Studies on the helminth fauna of Japan, Part 3, Avian trematodes II. *Jap. J. Zool.*, 5: 543-583.
- Yamaguti, S. 1939. Studies on the helminth fauna of Japan. 25. Trematodes of birds, IV. *Jap. J. Zool.*, 8: 129-210.
- Yamaguti, S. 1958. *Systema helminthum*. Vol. 1. The digenetic trematodes of vertebrates. Parts I and II. Interscience Publ. Co., N.Y. Part I p.1-979, Part II. pp. 980-1575.
- Yamaguti, S. 1961. *Systema helminthum*. Vol. III. The nematodes of vertebrates. Parts I and II. Interscience Publ. Co., N.Y. Part I pp.1-679, Part II. pp. 680-1261.
- Yamaguti, S. 1963. *Systema helminthum*. Vol. V. Acanthocephala. Interscience Publ. Co. N.Y., 423 pp.

- Yamaguti, S. 1971. Synopsis of Digenetic Trematodes of Vertebrates. Vol. I and II. Keigaku Publishing Co., Tokyo, Japan. Vol. I: 1-1074; Vol. II pp. 1-349.
- Zajiček, D. 1960. Příspěvek k druhovému výskytu a patogenitě vlasovek rodu *Amidostomum* Railliet et Henry, 1909 (Nematoda). Sborn. Českoslov. Akad. Zeměděl. Věd., Rada Vet. Med., 5: 775-788.
- Zajiček, D. 1963. The pathogenic effect of some species of flukes in the digestive tract of birds. Sborn. Českoslov. Akad. Zeměděl. Věd., Rada Vet. Med., 36: 263-266. (Czechoslovakia text, Russ. Eng. summaries).
- Zlotoryzcka, J. 1970. Studien an den mitteleuropaischen Arten der Gattung *Anatoecus* Cum. (Esthiopteridae, Mallophaga).

ADDENDUM

- Sinclair, N. R. 1971. A review of *Odhneria odhneri* Travassos, 1921 (Trematoda: Microphallidae). J. Parasitol., V. 57 (5) pp. 980-982.

Appendix 1. Checklist, compiled from original papers and other checklists, of Metazoan parasites recovered from the Surf Scoter, White-Winged Scoter and Black Scoter (excluding Acarina).

SURF SCOTER	
PARASITE	REFERENCE
<i>Anaticola crassicornis constrictus</i> (Kellogg, 1896)	Emerson, 1972
<i>Trinoton querquedulae</i> (Linnaeus, 1758)	Emerson, 1972
<i>Tristriata anatis</i> Belopol'skaia, 1953	Frame, 1969
<i>Gymnophallus bursicola</i> Odhner, 1900	Yamaguti, 1939
<i>Gymnophallus somaterias</i> Ching, 1973	Ching, 1973
<i>Meiogymnophallus multigemulus</i> Ching, 1975	Ching, 1975
<i>Parvatrema lintoni</i> James, 1964	Linton, 1928
<i>Myzolepis collaris</i> (Batsch, 1786)	McDonald, 1969
<i>Dicranotaenia (H.) coronula</i> (Dujardin, 1845)	Spasskaia, 1966
<i>Fimbriaria fasciolaris</i> (Pallas, 1781)	McDonald, 1969
<i>Fimbriarioides lintoni</i> (Webster, 1943)	Linton, 1927
<i>Hymenolepis tenuirostris</i> (Rudolphi, 1819)	Linton, 1927

Appendix 1. Continued

SURF SCOTER	
PARASITE	REFERENCE
<i>Streptocara californica</i> (Geddes, 1919)	Gibson, 1968
<i>Streptocara crassicauda crassicauda</i> (Creplin, 1829)	Gibson, 1968
<i>Streptocara formosensis</i> Sugimoto, 1930	Gibson, 1968
<i>Streptocara recta</i> (Linstow, 1879)	Gibson, 1968
<i>Paracuaria tridentata</i> (Linstow, 1877)	Gibson, 1968
<i>Falsificollis altmani</i> (Perry, 1942)	Perry, 1942

Appendix 1. Continued.

WHITE-WINGED SCOTER	
PARASITE	REFERENCE
<i>Anaticola crassicornis constrictus</i> (Kellogg, 1896)	Emerson, 1972
<i>Anaticola crassicornis punctalatus</i> (Rudow, 1869)	Emerson, 1972
<i>Anatoecus dentatus</i> (Eichler, 1946)	Zlotoryzcka, 1970
<i>Anatoecus laterodes</i> (Nitzsch, 1818)	Zlotoryzcka, 1970
<i>Holomenopon loomisi</i> (Kellogg, 1896)	Price, 1971
<i>Trinoton querquedulae</i> (Linnaeus, 1758)	Emerson, 1972
<i>Notocotylus attenuatus</i> (Rudolphi, 1809)	Ryzhikov, 1963a
<i>Notocotylus</i> sp. Diesing, 1839	Xuan, 1962
<i>Catantopis verrucosa</i> (Froelich, 1789)	Ryzhikov, 1963a
<i>Paramonostomum alveatum</i> (Mehlis in Creplin, 1846)	McDonald, 1969
<i>Tristriata elegans</i> Filimonova, 1971	Filimonova, 1971
<i>Cycloaxanthum mutabile</i> (Zeder, 1800)	McDonald, 1969
<i>Hyptiasmus arcuatus</i> (Brandes, 1892)	McDonald, 1969

Appendix 1. Continued

WHITE-WINGED SCOTER	
PARASITE	REFERENCE
<i>Hyptiasmus laevigatus</i> Kossack, 1911	Kossack, 1911
<i>Hyptiasmus theodori</i> Witenberg, 1928	Ryzhikov, 1963a
<i>Typhlocoelum cucumerinum</i> (Rud. 1809)	Mehlis, 1831
<i>Tracheophilus sisovi</i> Skrjabin, 1913	McDonald, 1969
<i>Eucotyle castenea</i> Jensen, 1971	Jensen, 1971
<i>Eucotyle cohni</i> Skrjabin, 1924	Ryzhikov, 1963a
<i>Eucotyle popowi</i> Skrjabin & Euranova, 1942	Ryzhikov, 1963a
<i>Eucotyle zakharovi</i> Skrjabin, 1920	McDonald, 1969b
<i>Gymnophallus bursteola</i> Odhner, 1900	Yamaguti, 1939
<i>Gymnophallus gibberosus</i> Loos-Frank, 1971	Loos-Frank, 1971a
<i>Meiogymnophallus macroporus</i> (Jameson & Nicoll, 1913)	Bykhovskaia-Pavlovskaja, 1962
<i>Echinostoma revolutum</i> (Froelich, 1802)	McDonald, 1969b
<i>Echinostoma</i> sp. (Rudolphi, 1809)	Ryzhikov, 1963a
<i>Echinoparyphium baculus</i> (Diesing, 1850)	McDonald, 1969c

Appendix 1. Continued

WHITE-WINGED SCOTER	
PARASITE	REFERENCE
<i>Echinoparyphium recurvatum</i> (Linstow, 1873)	Odening, 1963
<i>Himasthala incisa</i> (Linton, 1928)	McDonald, 1969
<i>Acanthoparyphium marilae</i> Yamaguti, 1934	McDonald, 1969b
<i>Acanthoparyphium melanittae</i> Yamaguti, 1939	McDonald, 1969b
<i>Acanthoparyphium tyrosense</i> Yamaguti, 1939	McDonald, 1969
<i>Pollostomum brevicolle</i> (Creplin, 1829)	Loos-Frank, 1968
<i>Prosthogonimus cuneatus</i> (Rudolphi, 1803)	McDonald, 1969
<i>Prosthogonimus pellucidus</i> (Linstow, 1873)	Ryžhikov 1963a
<i>Schistogonimus rarus</i> (Braun, 1901)	McDonald, 1969
<i>Orchipedum tracheicola</i> Braun, 1901	Cheatum, 1938
<i>Bulbovitellus fusiformis</i> (Reimer, 1963)	Reimer, 1963
<i>Levinseniella brachysoma</i> (Creplin, 1837)	Reimer, 1963
<i>Microphallus fusiformis</i> Reimer, 1963	Reimer, 1963
<i>Microphallus papillorobustum</i> (Rankin, 1940)	Reimer, 1963
<i>Microphallus pizum</i> (Afanas'ev, 1941)	Ching, 1965B

Appendix 1. Continued

WHITE-WINGED SCOTER	
PARASITE	REFERENCE
<i>Microphallius pygmaeus</i> (Levinsen, 1881)	Belopol'skaia, 1952
<i>Maritrema japonicum</i> (Yamaguti, 1939)	McDonald, 1969
<i>Maritrema subdolum</i> Jagerskiold, 1908	Reimer, 1963
<i>Odhneria odhneri</i> Travassos, 1921	Sinclair, 1971
<i>Pseudoapeloitrema japonicum</i> Yamaguti, 1939	Yamaguti, 1939
<i>Reticola</i> sp. Cohn, 1904	Oshmarin, 1963
<i>Cyathocotyle melanittas</i> Yamaguti, 1934	Yamaguti, 1934
<i>Cyathocotyle prussica</i> Mühling, 1896	Yamaguti, 1971
<i>Cotylurus erraticus</i> (Rudolphi, 1809)	McDonald, 1969
<i>Cotylurus japonicus</i> Ischii, 1932	Ryzhikov, 1963
<i>Apatemon canadensis</i> Dubois et Rausch, 1950	Dubois, 1967
<i>Apatemon fulgulus</i> Yamaguti, 1933	Ryzhikov, 1963
<i>Apatemon gracilis</i> (Rudolphi, 1819)	Ryzhikov, 1963
<i>Apatemon somateriae fiqcheri</i> Dubois, 1968	Dubois, 1968

Appendix 1. Continued

WHITE-WINGED SCOTER	
PARASITE	REFERENCE
<i>Australapatemon skrjabini</i> Ryzhikov et al, 1964	Ryzhikov et al., 1964
<i>Plagiurochis nanus</i> (Rudolphi, 1802)	McDonald, 1969
<i>Plagiurochis potanini</i> (Skrjabin, 1928)	McDonald, 1969
<i>Anomotaenia ciliata</i> Fuhrmann, 1913	McDonald, 1969
<i>Anomotaenia citrus</i> (Krabbe, 1869)	McDonald, 1969
<i>Lateriporus biuterinus</i> Fuhrmann, 1908	McDonald, 1969
<i>Lateriporus mathevosiands</i> Ryzhikov & Gubanov, 1962	Denny, 1969
<i>Lateriporus skrjabini</i> Matevosian, 1946	Tolkacheva, 1966
<i>Lateriporus teres</i> (Krabbe, 1869)	McDonald, 1969
<i>Gastrotasnia dogieli</i> (Gieneksinskaja, 1944)	Spasskaaa, 1966
<i>Fimbriaria fasciolaris</i> (Pallas, 1781)	Czapliński, 1956
<i>Fimbriariella falsiformis</i> (Linton, 1927)	Webster, 1943
<i>Fimbriarioides intermedia</i> (Fuhrmann, 1913)	Spasskaaa, 1966
<i>Fimbriarioides Lintoni</i> Webster, 1943	McDonald, 1969

Appendix 1. Continued

WHITE-WINGED SCOTER	
PARASITE	REFERENCE
<i>Aploparaksis furetigera</i> (Nitzsch, 1819)	Spasskaia, 1966
<i>Diorchis bulbodes</i> Mayhew, 1929.	Spasskaia, 1966
<i>Diorchis ransomi</i> Johri, 1939	Spasskaia, 1966
<i>Microsomacanthus</i> (H.) <i>abortiva</i> (Linstow, 1904)	Tolkacheva, 1966
<i>Drepanidolepis</i> (H.) <i>anatina</i> (Krabbe, 1869)	Spasskaia, 1966
<i>Monosacanthus</i> (H.) <i>andrejewoi</i> (Matevosian, 1945)	Spasskaia, 1966
<i>Wardium</i> (H.) <i>arctica</i> (Schiller, 1955)	Spasskaia, 1966
<i>Microsomacanthus</i> (H.) <i>arcuata</i> (Kowalewski, 1904)	Spasskaia, 1966
<i>Myzolepis</i> (H.) <i>collaris</i> (Batsch, 1786)	McDonald, 1969
<i>Microsomacanthus</i> (H.) <i>compressa</i> (Linton, 1892)	Tolkacheva, 1966
<i>Dicranataenia</i> (H.) <i>coronula</i> (Dujardin, 1845)	Spasskaia, 1966
<i>Dicranataenia</i> (H.) <i>deglandi</i> Skrjabin & Matevosian, 1942	McDonald, 1969
<i>Microsomacanthus</i> (H.) <i>fausti</i> (Shentseff, 1932)	Ryzhikov, 1963
<i>Microsomacanthus</i> (H.) <i>formosoides</i> Spasskaia & Spasskii, 1961	Tolkacheva, 1966

Appendix 1. Continued

WHITE-WINGED SCOTER	
PARASITE	REFERENCE
<i>Sobolevicanthus kenatensis</i> (Schiller, 1952)	Czapliński, 1973
<i>Microsomacanthus (H.) hygie</i> Spasskaia & Spasskii, 1961	Tolkacheva, 1966
<i>Microsomacanthus (H.) jaegerskioeldi</i> (Fuhrmann, 1913)	Fuhrmann, 1913
<i>Sobolevicanthus (H.) krabbeella</i> (Hughes, 1940)	Tolkacheva, 1966
<i>Tschertkovilepis (H.) krabbei</i> (Kowalewski, 1895)	McDonald, 1969
<i>Cloacotaenia megalops</i> (Nitzsch, 1829)	McDonald, 1969
<i>Microsomacanthus (H.) melanittae</i> Ryzhikov, 1962	Ryzhikov, 1962
<i>Microsomacanthus (H.) microskrjabini</i> Spasskii & Turpalova, 1964	Tolkacheva, 1966
<i>Microsomacanthus (H.) microsoma</i> (Creplin, 1829)	Fuhrmann, 1913
<i>Microsomacanthus (H.) mirabilis</i> Spasskii & Turpalova, 1964	Spasskii, 1966
<i>Wardium nyrocae</i> Ryzhikov & Gubanov, 1959	McDonald, 1969
<i>Sobolevicanthus (H.) octacanthoides</i> Fuhrmann, 1906	McDonald, 1969
<i>Microsomacanthus (H.) pachycephala</i> (Linstow, 1872)	Ryzhikov, 1963
<i>Echinatrium skrjabini</i> Spasskaia & Turpalova, 1965	Spasskaia, 1966

Appendix 1. Continued

WHITE-WINGED SCOTER	
PARASITE	REFERENCE
<i>Hamatolepis (H.) teresoides</i> (Fuhrmann, 1906)	Czaplinski, 1962
<i>Hymenolepis turicensis</i> Podestra and Holmes, 1970	Podesta & Holmes, 1970
<i>Capillaria nyrocinarum</i> Madsen, 1945	Madsen, 1945
<i>Capillaria mergi</i> Madsen, 1945	Czaplinski, 1962
<i>Capillaria s. l.</i> sp. Zeder, 1800	Ryzhikov, 1963b
<i>Amidostomum acutum</i> (Lundahl, 1848)	Czaplinski, 1962
<i>Amidostomum ansersis</i> (Zeder, 1800)	Czaplinski, 1962
<i>Epimodostium oriepinum</i> (Molin, 1861)	Czaplinski, 1962
<i>Epimodostium uncinatum</i> (Lundahl, 1848)	McDonald, 1969
<i>Streptocara kalifornica</i> (Gedoelst, 1919)	Gibson, 1964
<i>Streptocara crassicauda crassicauda</i> (Creplin, 1829)	Gibson, 1968
<i>Streptocara formosensis</i> Sugimoto, 1930	Gibson, 1968
<i>Paracuaría somaterias</i> (Ryzhikov, 1960)	McDonald, 1969

Appendix 1. Continued

WHITE-WINGED SCOTER	
PARASITE	REFERENCE
<i>Echinuria hypognatha</i> Wehr, 1937	McDonald, 1969
<i>Tetrameres fissispina</i> (Diesing, 1861)	Ryzhikov, 1963b
<i>Trichostrongylus tenuis</i> (Mehlis, 1846)	Czapliński, 1962
<i>Fillicolis anatis</i> (Schrank, 1788)	Czapliński, 1962
<i>Polymorphus altmani</i> (Perry, 1942)	Perry, 1942
<i>Polymorphus diploinflatus</i> Lundstrom, 1942	McDonald, 1969
<i>Polymorphus kostylewi</i> Petrochenko, 1949	Petrochenko, 1949
<i>Polymorphus magnus</i> Skrjabin, 1913	Schmidt, 1965a
<i>Polymorphus marilis</i> Van Cleave, 1939	Schmidt, 1965a
<i>Polymorphus matheoassianae</i> Petrochenko, 1949	Petrochenko, 1949
<i>Polymorphus minutus</i> (Goetze, 1782)	Petrochenko, 1949
<i>Polymorphus strumosoides</i> Lundstrom, 1942	Petrochenko, 1949
<i>Polymorphus trochus</i> Van Cleave, 1945	Petrochenko, 1949
<i>Polymorphus paradoxus</i> Conell & Corner, 1957	Denny, 1969
<i>Corynopoma constrictum</i> Van Cleave, 1945	Schmidt, 1965b

Appendix 1. Continued

BLACK SCOTER	
PARASITE	REFERENCE
<i>Anaticola crassicornis angustalimbatus</i> (Giebel, 1866)	Emerson, 1972
<i>Anatococcus dentatus</i> (Eichler, 1946)	Zlotorzyska, 1970
<i>Holomenopon leucoxanthum</i> (Burmeister, 1838)	Price, 1976
<i>Trinoton querquedulae</i> (Linnaeus, 1758)	Emerson, 1972
<i>Notocotylus attenuatus</i> (Rudolphi, 1809)	Ryzhikov, 1963
<i>Notocotylus dafillae</i> Harwood, 1939	Leonov et al., 1963
<i>Notocotylus gibbus</i> (Mehlis in Creplin, 1846)	Leonov et al., 1963
<i>Catantropis verrucosa</i> (Froelich, 1789)	McDonald, 1969
<i>Paramonostomum alveatum</i> (Mehlis in Creplin, 1846)	McDonald, 1969
<i>Cyclocoelum mutabile</i> (Zeder, 1800)	McDonald, 1969
<i>Hyptiasmus arcuatus</i> (Brandes, 1892)	McDonald, 1969
<i>Hyptiasmus laevigatus</i> Kossack, 1911	Kossack, 1911
<i>Eucotyle cohni</i> Skrjabin, 1924	Ryzhikov, 1963
<i>Gymnophallus affinis</i> James & Nicoll, 1913	Jameson & Nicoll, 1913

Appendix 1. Continued

BLACK SCOTER	
PARASITE	REFERENCE
<i>Gymnophallus bureicola</i> Odhner, 1900	Jameson & Nicoll, 1913
<i>Gymnophallus choledocus</i> Odhner, 1900	Loos-Frank, 1968
<i>Gymnophallus dapsilis</i> Nicoll, 1907	Jameson & Nicoll, 1913
<i>Gymnophallus deliciosus</i> (Olsson, 1893)	Loos-Frank, 1969
<i>Gymnophallus geratostomus</i> Tsimbaliuk & Leonov, 1963	Tsimbaliuk & Leonov, 1963
<i>Gymnophallus gibberosus</i> Loos-Frank, 1971	Loos-Frank, 1971
<i>Gymnophallus somateriae</i> Ching, 1973	Ching, 1973
<i>Gymnophalloides conspicuus</i> (Ching, 1965)	Ching, 1965a
<i>Meiogymnophallus jamesoni</i> Bowers, 1966	Bowers, 1966
<i>Meiogymnophallus macroporus</i> (Jameson & Nicoll, 1913)	Jameson & Nicoll, 1913
<i>Meiogymnophallus macrostoma</i> (Yamaguti, 1939)	Yamaguti, 1971
<i>Meiogymnophallus minutus</i> (Cobbold, 1859)	Bowers & James, 1967
<i>Meiogymnophallus multigemmulus</i> Ching, 1965	Ching, 1965
<i>Parvatrema ovoplenum</i> (Jameson & Nicoll, 1913)	Yamaguti, 1971

Appendix 1. Continued

BLACK SCOTER	
PARASITE	REFERENCE
<i>Echinostoma revolutum</i> (Froelich, 1802)	McDonald, 1969
<i>Echinoparyphium aceniatum</i> Dietz, 1909	Ryzhikov, 1963
<i>Echinoparyphium flexum</i> (Linton, 1892)	McDonald, 1969
<i>Echinoparyphium recurvatum</i> (Linstow, 1873)	McDonald, 1969b
<i>Stephanoprora denticulata</i> (Baer, 1959)	McDonald, 1969b
<i>Himastha inoisa</i> Linton, 1928	Linton, 1928
<i>Acanthoparyphium kurógamo</i> Yamaguti, 1939	Yamaguti, 1939
<i>Acanthoparyphium marilae</i> Yamaguti, 1934	Yamaguti, 1934
<i>Acanthoparyphium tyrosenense</i> Yamaguti, 1939	Yamaguti, 1939
<i>Peilostomum brevicolle</i> (Creplin, 1829)	Loos-Frank, 1968
<i>Peilochasmus oxyurus</i> (Creplin, 1825)	McDonald, 1969b
<i>Peilotrema oligoon</i> (V. Linstow, 1887)	McDonald, 1969b
<i>Peilotrema spiculigerum</i> (Muhl, 1898)	Ryzhikov, 1963
<i>Prosthogonimus pellucidus</i> (v. Linstow, 1887)	Ryzhikov, 1963

Appendix 1. Continued

BLACK SCOTER	
PARASITE	REFERENCE
<i>Wetzelitrema melanittae</i> Rayski & Fahmy, 1962	Rayski & Fahmy, 1962
<i>Levinseniella brachysomum</i> (Creplin, 1837)	Reimer, 1963a
<i>Microphallus excellens</i> (Nicoll, 1907)	McDonald, 1969a
<i>Microphallus primas</i> (Jagerskiold, 1908)	Reimer, 1963a
<i>Microphallus pygmaeum</i> (Levinsen, 1881)	Deblock & Tran Van Ky, 1966a
<i>Maritrema graciosum</i> Nicoll, 1907	Deblock & Tran Van Ky, 1966a
<i>Maritrema subdolum</i> (Jagerskiold, 1908)	Reimer, 1963a
<i>Cryptocotyle concava</i> (Creplin, 1825)	McDonald, 1969b
<i>Cotylurus ornatus</i> (Rudolphi, 1808)	Ryzhikov, 1963
<i>Apatemon gracilis</i> (Rudolphi, 1819)	Ryzhikov, 1963
<i>Australapatemon skrjabini</i> Ryzhikov et al., 1964	Ryzhikov et al., 1964
<i>Plagiorhynchus laticola</i> Skrjabin, 1924	Leonov et al., 1963
<i>Plagiorhynchus obtusus</i> Shtrom, 1940	Leonov et al., 1963
<i>Metorhynchus xanthosomus</i> (Creplin, 1846)	McDonald, 1969b

Appendix 1. Continued

BLACK SCOTER	
PARASITE	REFERENCE
<i>Anomotaenia ciliata</i> Fuhrmann, 1913	McDonald, 1969
<i>Lateriporus teres</i> (Krabbe, 1869)	McDonald, 1969
<i>Lateriporus skrjabini</i> Matevosian, 1946	Tolkacheva, 1966
<i>Pactiterotaenia borealis</i> (Linstow, 1905)	McDonald, 1969
<i>Gastrotaenia dogieli</i> (Ginetsinskaiá, 1944)	Spasskáia, 1966
<i>Diploposthe laevis</i> (Bloch, 1872)	Spasskáia, 1966
<i>Fimbriaria fasciolaris</i> (Pallas, 1781)	Czapliński, 1962
<i>Fimbriarioidea intermedia</i> (Fuhrmann, 1913)	Spasskáia, 1966
<i>Diorchis stefanski</i> Czapliński, 1956	McDonald, 1969
<i>Microsomacanthus (H.) abortiva</i> (Linstow, 1904)	Tolkacheva, 1966
<i>Microsomacanthus (H.) acus</i> Spasskii & Jurpalova, 1965	Spasskáia, 1966
<i>Microsomacanthus (H.) compressa</i> (Linton, 1892)	Spasskáia, 1966
<i>Dicranotaenia (H.) coronula</i> (Dujardin, 1845)	McDonald, 1969
<i>Microsomacanthus (H.) formosa</i> (Dubinia, 1954)	Tolkacheva, 1966

Appendix 1. Continued

BLACK SCOTER	
PARASITE	REFERENCE
<i>Retinomera</i> (H.) <i>macracanthus</i> (Linstow, 1877)	Tolkacheva, 1966
<i>Echinatrium mellitas</i> Tolkacheva, 1966	Tolkacheva, 1966
<i>Hymenolepis micraneristrotta</i> (Wedl, 1855)	McDonald, 1969
<i>Microsomacanthus microskrjabina</i> Spasskii & Iurpalova	Tolkacheva, 1966
<i>Microsomacanthus</i> (H.) <i>miensoma</i> (Creplin, 1829)	Bowers, 1966
<i>Microsomacanthus mirabilis</i> Spasskii & Iurpalova, 1964	Tolkacheva, 1966
<i>Wardoidea</i> (H.) <i>nyrocae</i> (Yamaguti, 1935)	McDonald, 1969
<i>Microsomacanthus oidemiae</i> Spasskii & Iurpalova, 1964	Spasskaia, 1966
<i>Wardoidea oidemiae</i> Spasskii, 1966	Spasskaia, 1966
<i>Microsomacanthus</i> (H.) <i>pachycephala</i> (Linstow, 1872)	Tolkacheva, 1966
<i>Microsomacanthus</i> (H.) <i>paracompressa</i> (Czaplinski, 1956)	Tolkacheva, 1966
<i>Microsomacanthus</i> (H.) <i>paramicrosoma</i> (Gasowska, 1931)	Tolkacheva, 1966
<i>Amphipetrovia</i> (H.) <i>retracta</i> (Linstow, 1905)	Tolkacheva, 1966
<i>Echinatrium skrjabini</i> Spasskii, 1965	Spasskaia, 1966

Appendix 1. Continued

BLACK SCOTER	
PARASITE	REFERENCE
<i>Microsomacanthus</i> (H.) <i>jaegerskioeldi</i> (Fuhrmann, 1913)	Spasskaia, 1966
<i>Microsomacanthus</i> <i>spasskii</i> Tolkacheva, 1965	Tolkacheva, 1966
<i>Sobolevicanthus</i> <i>kenafensis</i> (Schiller, 1952)	McDonald, 1969
<i>Anatinella</i> <i>spinulosa</i> (Dubinina, 1954)	Tolkacheva, 1966
<i>Microsomacanthus</i> <i>tuvensis</i> Spasskaia & Spasskii, 1961	McDonald, 1969
<i>Capillaria</i> <i>nyroccinarum</i> Madsen, 1945	Madsen, 1945
<i>Amidostomum</i> <i>acutum</i> (Lundahl, 1845)	Ryzhikov, 1963b
<i>Amidostomum</i> <i>anseris</i> (Zeder, 1800)	Czaplinski, 1962
<i>Epomidostium</i> <i>orispinum</i> (Molin, 1861)	Czaplinski, 1962
<i>Epomidostium</i> <i>uncinatum</i> (Lundahl, 1848)	Czaplinski, 1962
<i>Streptocara</i> <i>californica</i> (Geddelst, 1919)	Gibson, 1964
<i>Streptocara</i> <i>crassicauda</i> <i>crassicauda</i> (Replin, 1829)	Gibson, 1968
<i>Streptocara</i> <i>formosensis</i> Sugimoto, 1930	Gibson, 1968

Appendix 1. Continued

BLACK SCOTER	
PARASITE	REFERENCE
<i>Streptocara dogieli</i> Belopol'skaya, 1952	Ryzhikov, 1963b
<i>Echinuria borealis asiatica</i> Ryzhikov, 1962	Ryzhikov, 1963b
<i>Echinuria hypognatha</i> Wehr, 1937	Ryzhikov, 1963b
<i>Tetrameres fissipina</i> (Diesing, 1861)	Ryzhikov, 1963b
<i>Fillicolis anatis</i> (Schrank, 1788)	McDonald, 1969
<i>Polymorphus diploinflatus</i> Lundstrom, 1942	McDonald, 1969
<i>Polymorphus magnus</i> Skrjabin, 1913	McDonald, 1969
<i>Polymorphus minutus</i> (Goeze, 1782)	McDonald, 1969
<i>Polymorphus strumosoides</i> Lundstrom, 1942	McDonald, 1969
<i>Polymorphus trochus</i> Van Cleave, 1945	McDonald, 1969
<i>Corynosoma constrictum</i> Van Cleave, 1945	McDonald, 1969

APPENDIX-2. Comparison of the helminths recovered from 6
M. f. fusca to those recovered from the 30
M. f. deglandi.

PARASITE	<i>M. f. deglandi</i>	<i>M. f. fusca</i>
<i>Austrobilharzia terrigenensis</i>	✓	-
<i>Notocotylus attenuatus</i>	✓	-
<i>Tristriata elegans</i>	✓	-
<i>Catantropis verrucosa</i>	-	✓
<i>Eucotyle castanea</i>	✓	-
<i>Gymnophallus basicola</i>	✓	-
<i>Gymnophallus choledocus</i>	✓	✓
<i>Gymnophallus delictosus</i>	✓	✓
<i>Gymnophallus</i> sp. Type I	✓	-
<i>Dietsiella egregia</i>	✓	-
<i>Himasthala incisa</i>	✓	-
<i>Ascorhyttis charadriiformis</i>	✓	-
<i>Pseudospelotrema japonicum</i>	✓	-
<i>Renticola</i> sp.	✓	-
<i>Capillaria nyrocinarum</i>	✓	✓
<i>Amidostomum acutum</i>	✓	✓

APPENDIX 2. CONTINUED

PARASITE	<i>M. f. deglandi.</i>	<i>M. f. fusca</i>
<i>Streptocara californica</i>	✓	-
<i>Streptocara</i> s. <i>crassicauda</i>	✓	-
<i>Streptocara formosensis</i>	✓	-
<i>Tetrameres fissispina</i>	✓	-
<i>Tetrameres somaterias</i>	✓	✓
<i>Tetrameres spinosa</i>	✓	-
<i>Tetrameres</i> sp.	✓	✓
<i>Hymenolepis</i> sp.	✓	✓
<i>Cloacotaenia megalopa</i>	✓	-
<i>Polymorphus botulus</i>	✓	✓

1
 Details of Infection of Scoters with helminths by age and sex.
 APPENDIX 3. Details of infection of surf scoters with Trematoda by age and sex.

PARASITE	ADULTS			IMMATURES			TOTAL	TOTAL
	MALE	FEMALE	TOTAL	MALE	FEMALE	TOTAL	MALES	FEMALES
<i>A. terrigalensis</i>	8(14.3)	-	8(11.1)	-	-	-	8(11.1)	-
<i>Z. lunata</i>	1(1.8)	-	1(1.4)	1(6.3)	-	1(4.5)	2(2.8)	-
<i>N. attenuatus</i>	-	-	-	1(6.3)	-	1(4.5)	1(1.4)	-
<i>T. elegans</i>	15(26.8)	5(31.3)	20(27.8)	3(18.8)	3(50.0)	6(27.3)	18(25.0)	8(36.4)
<i>E. castanea</i>	-	1(6.3)	1(1.4)	1(6.3)	-	1(4.5)	1(1.4)	1(4.5)
<i>G. bureicola</i>	2(3.6)	1(6.3)	3(4.2)	6(37.5)	1(16.7)	7(31.8)	8(11.1)	2(9.1)
<i>G. choledocus</i>	-	-	-	6(37.5)	1(16.7)	7(31.8)	6(8.3)	1(4.5)
<i>G. delticopus</i>	5(8.9)	1(6.3)	6(8.3)	4(25.0)	4(66.7)	8(36.4)	9(12.5)	5(22.7)
<i>Gymnophallus</i> sp. Type 1	1(1.8)	-	1(1.4)	1(6.3)	1(16.7)	3(13.6)	2(2.8)	1(4.5)
<i>Gymnophallus</i> spp.	1(1.8)	-	1(1.4)	1(6.3)	-	1(4.5)	2(2.8)	-
<i>Gymnophallid</i> sp.	-	-	-	1(6.3)	-	1(4.5)	1(1.4)	-
<i>A. melanittae</i>	1(1.8)	-	1(1.4)	-	-	-	1(1.4)	-
<i>Patlostomum</i> sp.	5(8.9)	2(12.5)	7(9.7)	-	-	-	5(6.9)	2(9.1)
<i>A. charadriiformis</i>	1(1.8)	1(6.3)	2(2.8)	1(6.3)	-	1(4.5)	2(2.8)	2(9.1)
<i>P. japonicum</i>	1(1.8)	1(6.3)	2(2.8)	2(12.5)	-	2(9.1)	3(4.2)	1(4.5)
<i>Rentocla</i> sp.	1(1.8)	-	1(1.4)	1(6.3)	-	1(4.5)	2(2.8)	-
<i>A. gracilis</i>	-	-	-	1(6.3)	-	1(4.5)	1(1.2)	-

NOTE: The first figure represents the no. of birds of that class infected and the no. in brackets represents the percentage of that class infected.

Details of infection of white-winged scoters with Trematoda by age and sex.

PARASITE	ADULTS			IMMATURES			TOTAL	TOTAL
	MALE	FEMALE	TOTAL	MALE	FEMALE	TOTAL	MALE	FEMALE
<i>A. terrigalensis</i>	2(7.7)	-	2(6.9)	-	-	-	2(6.7)	-
<i>N. attenuatus</i>	1(3.8)	-	1(3.4)	-	-	-	1(3.3)	-
<i>C. verrucosa</i>	2(7.7)	1(33.3)	3(10.3)	-	-	-	2(6.7)	1(20.0)
<i>T. elegans</i>	3(11.5)	-	3(10.3)	1(25.0)	-	1(16.6)	4(13.3)	-
<i>E. castanea</i>	4(15.4)	-	4(13.8)	-	-	-	4(13.3)	-
<i>G. bursicola</i>	1(3.8)	-	1(3.4)	-	1(50.0)	1(16.6)	1(3.3)	1(20.0)
<i>G. choledocus</i>	-	-	-	2(50.0)	1(50.0)	3(50.0)	2(6.7)	1(20.0)
<i>G. deliciosus</i>	7(26.9)	1(33.3)	8(27.6)	2(50.0)	-	2(6.7)	9(30.0)	1(20.0)
<i>Gymnophallus</i> spp.	2(7.7)	-	2(6.9)	1(25.0)	-	1(16.6)	3(10.0)	-
<i>Gymnophallus</i> sp. 1	1(3.8)	-	1(3.4)	1(25.0)	-	1(16.6)	2(6.7)	-
<i>D. egregia</i> *	6(23.1)	1(33.3)	7(24.1)	-	-	-	6(20.0)	1(20.0)
<i>H. incisa</i>	1(3.8)	-	1(3.4)	-	-	-	1(3.3)	-
<i>A. charadriiformis</i>	1(3.8)	-	1(3.4)	-	-	-	1(3.3)	-
<i>P. japonicum</i>	4(15.4)	1(33.3)	5(17.2)	1(25.0)	-	1(16.6)	5(16.7)	1(20.0)
<i>Renicola</i> sp.	1(3.8)	-	1(3.4)	-	-	-	1(3.3)	-

**D. egregia* infected 1 white-winged scoter that could not be aged or sexed.

Details of infection of black scoters with Trematoda by age and sex.

PARASITE	ADULTS			IMMATURES			TOTAL	TOTAL
	MALE	FEMALE	TOTAL	MALE	FEMALE	TOTAL	MALE	FEMALE
<i>A. terrigalensis</i>	2 (6.3)	2 (25.0)	4 (10.0)	-	-	-	2 (6.1)	2 (16.7)
<i>N. attenuatus</i>	-	-	-	-	1 (25.0)	1 (20.0)	-	1 (8.3)
<i>E. caetenea</i>	2 (6.3)	-	2 (5.0)	-	-	-	2 (6.1)	-
<i>G. bursicola</i>	2 (6.3)	-	2 (5.0)	-	2 (50.0)	2 (40.0)	2 (6.1)	2 (16.7)
<i>G. choledocus</i>	-	-	-	1 (100.0)	-	1 (20.0)	1 (3.0)	-
<i>G. delticosus</i>	2 (6.3)	-	2 (5.0)	-	2 (50.0)	2 (40.0)	2 (6.1)	2 (16.7)
<i>Gymnophallus</i> sp. Type 1	-	-	-	1 (100.0)	-	1 (20.0)	1 (3.0)	-
<i>Gymnophallus</i> spp.	4 (13.0)	-	4 (10.0)	-	1 (25.0)	1 (20.0)	4 (12.2)	1 (8.3)
<i>Gymnophallid</i> sp.	1 (3.1)	-	1 (2.5)	-	-	-	1 (3.0)	-
<i>E. revolutum</i>	-	-	-	-	1 (25.0)	1 (20.0)	-	1 (8.3)
<i>H. inoisa</i>	1 (3.1)	-	1 (2.5)	-	-	-	1 (3.0)	-
<i>A. charadiformis</i>	1 (3.1)	-	1 (2.5)	-	1 (25.0)	1 (20.0)	1 (3.0)	1 (8.3)
<i>M. pygmaeum</i>	-	-	-	-	1 (25.0)	1 (20.0)	-	1 (8.3)
<i>P. japonicum</i>	-	-	-	-	1 (25.0)	1 (20.0)	-	1 (8.3)
<i>C. strigeoides</i>	-	-	-	-	1 (25.0)	1 (20.0)	-	1 (8.3)
<i>A. burti</i>	-	-	-	-	1 (25.0)	1 (20.0)	-	1 (8.3)

Details of infection of scoters with Cestoda by age and sex.

HOST	PARASITE	ADULTS			IMMATURES			TOTAL	TOTAL
		MALE	FEMALE	TOTAL	MALE	FEMALE	TOTAL	MALE	FEMALE
<i>M. perspicillata</i>	<i>Hymenolepis</i> spp.	26(46.4)	10(62.5)	36(38.3)	16(100.0)	3(50.0)	19(86.4)	42(58.3)	13(59.1)
	Frimbriariinae	6(10.7)	-	6(8.3)	5(31.3)	2(33.3)	7(31.8)	11(15.3)	2(9.1)
<i>M. fusca</i>	* <i>Hymenolepis</i> spp.	16(61.5)	2(66.6)	18(62.1)	2(50.0)	2(100.0)	4(66.7)	18(60.0)	4(80.0)
	<i>C. megalops</i>	-	-	-	1(25.0)	-	1(16.7)	1(3.3)	-
<i>M. nigra</i>	<i>Hymenolepis</i> spp.	18(56.3)	5(62.5)	23(57.5)	1(100.0)	2(50.0)	3(60.0)	19(57.6)	7(58.3)
	Frimbriariinae	3(9.4)	-	3(7.5)	1(100.0)	2(50.0)	3(60.0)	4(12.1)	2(16.7)

**Hymenolepis* spp. infected 1 *M. fusca* that could not be aged or sexed.

Details of infection of surf scoters with Nematoda by age and sex.

PARASITE	ADULTS			IMMATURES			TOTAL	TOTAL
	MALE	FEMALE	TOTAL	MALE	FEMALE	TOTAL	MALE	FEMALE
<i>C. anatis</i>	-	-	-	-	1(16.7)	1(4.5)	-	1(4.5)
<i>C. nyrocinarum</i>	5(8.9)	1(6.3)	6(8.3)	10(62.5)	4(66.7)	14(63.6)	15(20.8)	5(22.7)
<i>A. acutum</i>	28(50.0)	8(50.0)	36(50.0)	9(56.3)	3(50.0)	12(54.5)	37(51.4)	11(50.0)
<i>S. californica</i>	10(17.9)	6(37.5)	16(22.2)	3(18.8)	1(16.7)	4(18.2)	13(18.1)	7(31.8)
<i>S. o. crassicauda</i>	-	2(12.5)	2(2.8)	5(31.3)	-	5(22.7)	5(6.9)	2(9.1)
<i>S. formosensis</i>	4(7.1)	1(6.3)	5(6.9)	5(31.3)	3(50.0)	8(36.4)	9(12.5)	4(18.2)
<i>E. borealis</i>	-	-	-	-	1(16.7)	1(4.5)	-	1(4.5)
<i>E. hypognatha</i>	7(12.5)	5(31.3)	12(16.7)	8(50.0)	1(16.7)	9(40.9)	15(20.8)	6(27.3)
<i>T. fissiepinia</i>	1(1.8)	-	1(1.4)	5(31.3)	1(16.7)	6(27.3)	6(8.3)	1(4.5)
<i>T. somateriae</i>	20(35.7)	8(50.0)	28(38.9)	13(81.3)	3(50.0)	16(72.7)	33(45.8)	11(50.0)
<i>T. spinosa</i>	-	1(6.3)	1(1.4)	2(12.5)	-	2(9.1)	2(2.8)	1(4.5)
<i>Amidostomum</i> sp.	4(7.1)	1(6.3)	5(6.9)	-	-	-	4(5.6)	1(4.5)
<i>Streptocara</i> sp.	2(3.6)	-	2(2.8)	-	-	-	2(2.8)	-
<i>Tetrameres</i> sp.	1(1.8)	-	1(1.4)	1(6.3)	3(50.0)	4(18.2)	2(2.8)	3
<i>Sarcoconema</i> sp.	1(1.8)	-	1(1.4)	-	-	-	1(1.4)	-
Nematode sp. II	-	-	-	-	1(16.7)	1(4.5)	-	1(4.5)

Details of infection of white-winged scoters with Nematoda by age and sex.

PARASITE	ADULTS			IMMATURES			TOTAL	TOTAL
	MALE	FEMALE	TOTAL	MALE	FEMALE	TOTAL	MALE	FEMALE
<i>C. nyrocinarum</i>	10(38.5)	1(33.3)	11(37.9)	-	1(50.0)	1(16.7)	10(33.3)	2(40.0)
<i>A. acutum</i>	8(30.7)	-	8(27.6)	1(25.0)	2(100.0)	3(50.0)	9(30.0)	2(40.0)
<i>S. californica</i>	1(2.7)	-	1(3.4)	-	-	-	1(3.3)	-
<i>S. c. crassicauda</i>	1(2.7)	-	1(3.4)	-	-	-	1(3.3)	-
<i>S. formosensis</i>	1(2.7)	-	1(3.4)	-	-	-	1(3.3)	-
<i>T. fissispina</i>	-	-	-	-	1(50.0)	1(16.7)	-	1(20.0)
<i>T. somateriae</i>	7(26.9)	-	7(24.1)	1(25.0)	1(50.0)	2(33.3)	8(26.7)	1(20.0)
<i>T. spinosa</i>	1(2.7)	-	1(3.4)	-	-	-	1(3.3)	-
<i>Tetrameres</i> sp.	2(7.7)	-	2(6.9)	-	-	-	2(6.7)	-

Details of infection of black scoters with Nematoda by age and sex.

PARASITE	ADULTS			IMMATURES			TOTAL	TOTAL
	MALE	FEMALE	TOTAL	MALE	FEMALE	TOTAL	MALE	FEMALE
<i>C. nyrocinorum</i>	-	-	-	-	2(50.0)	2(40.0)	-	2(16.7)
<i>A. acutum</i>	17(53.1)	4(50.0)	21(52.5)	1(100.0)	2(50.0)	3(60.0)	18(54.5)	6(50.0)
<i>E. uncinatum</i>	1(3.1)	-	1(2.5)	-	1(25.0)	1(20.0)	1(3.0)	1(8.3)
<i>S. californica</i>	5(15.6)	1(12.5)	6(15.0)	-	2(50.0)	2(40.0)	5(15.2)	8(66.7)
<i>S. c. crassicauda</i>	3(9.4)	-	3(7.5)	-	1(25.0)	1(20.0)	3(9.1)	1(8.3)
<i>E. hypognatha</i>	4(12.5)	1(12.5)	5(12.5)	-	2(50.0)	2(40.0)	4(12.1)	3(25.0)
<i>T. fissispina</i>	1(3.1)	-	1(2.5)	-	2(50.0)	2(40.0)	1(3.1)	2(16.7)
<i>T. somaterias</i>	-	-	-	-	1(25.0)	1(20.0)	-	1(8.3)
<i>T. spinosa</i>	-	-	-	1(100.0)	-	1(20.0)	1(3.0)	-

Details of infection of scoters with *Acanthocephala* by age and sex.

HOST	PARASITE	ADULTS			IMMATURES			TOTAL	TOTAL
		MALE	FEMALE	TOTAL	MALE	FEMALE	TOTAL	MALE	FEMALE
<i>M. perspicillata</i>	<i>F. almani</i>	-	-	-	1(6.3)	-	1(4.5)	1(1.4)	-
	<i>P. botulus</i>	4(7.1)	1(6.3)	5(6.9)	1(6.3)	-	1(4.5)	5(6.9)	1(4.5)
	<i>C. sudsuche</i>	6(10.7)	2(12.5)	8(11.1)	5(31.3)	-	5(22.7)	11(15.3)	2(9.1)
<i>M. fusca</i>	<i>P. botulus</i>	3(11.5)	1(33.3)	4(13.8)	2(50.0)	1(50.0)	3(50.0)	5(16.7)	2(40.0)
<i>M. nigra</i>	<i>P. botulus</i>	4(12.5)	1(12.5)	5(12.5)	-	2(50.0)	2(40.0)	4(12.1)	3(25.0)
	<i>C. constrictum</i>	-	-	-	-	1(25.0)	1(20.0)	-	1(8.3)



