STRATIGRAPHY AND TERRESTRIAL PALYNOLOGY OF LATE CRETACEOUS ECLIPSE GROUP STRATA, BYLOT ISLAND, NORTHWEST TERRITORIES, CANADA



KERRY ELWYN SPARKES, B. Sc. (Honours)









National Library of Canada Bibliothèque nationale du Canada

Canadian Theses Service Service des thèses canadiennes

Ottawa Canada K1A 0N4

The author has granted an irrevocable nonexclusive licence allowing the National Library of Canada to reproduce, loan, distribute or sell copies of his/her thesis by any means and in any form or formal, making this thesis available to interested persons.

The author retains ownership of the copyright in his/her thesis. Neither the thesis nor substantial extracts from it may be printed or oth.rwise reproduced without his/her permission. L'auteur a accordé une licence irrévocable et non exclusive permettant à la Bibliothèque nationale du Canada de reproduire, prêter, distribuer ou vendre des copies de sa thèse de quelque manière et sous quelque forme que ce soit pour mettre des exemplaires de cette thèse à la disposition des personnes inféressées.

L'auteur conserve la propriété du droit d'auteur qui protège sa thèse. Ni la thèse ni des extraits substantiels de celle-ci ne doivent être imprimés ou autrement reproduits sans son autorisation.

ISBN 0-315-54999-8



STRATIGRAPHY AND TERRESTRIAL PALYNOLOGY

OF

LATE CRETACEOUS ECLIPSE GROUP STRATA,

BYLOT ISLAND,

NORTHWEST TERRITORIES, CANADA

by

Kerry Elwyn Sparkes, B.Sc. (Honours)

A THESIS SUBMITTED TO THE SCHOOL OF GRADUATE STUDIES IN PARTIAL FULFILLMENT OF THE REQUIREMENTS FOR THE DEGREE OF MASTER OF SCIENCE

> DEPARTMENT OF EARTH SCIENCES MEMORIAL UNIVERSITY OF NEWFOUNDLAND APRIL 1989

ST.JOHN'S NEWFOUNDLAND

Ser case increased and

ABSTRACT

Upper Cretaceous " Kanguk " strata on Bylot Island have been informally divided into three new, lithologically distinct formations: the Byam Martin Formation (mid to late Campanian), the Sermilik Formation (late Campanian to early late Maastrichtian) and the Bylot Island Formation (late Campanian to late Maastrichtian). These formations and Tertiary formations (not studied herein), comprise the Eclipse Group.

The Byam Martin Formation consists of mudstone, sandy mudstone and minor sandstone, and originated as basin plain and turbidite deposits followed by regressive marginal marine deposits of deltaic? origin. The Byam Martin Formation is conformably overlain by the Sermilik Formation in the south and the Bylot Island Formation in the north. The Sermilik Formation represents a major late Cretaceous regressive event, and is comprised of coarse-grained sandstones that were shed from the adjacent Byam Martin Mountains. These sandstones are preserved as braid delta and submarine fan deposits. The Bylot Island Formation conformably overlies the Sermilik Formation, and is in part a lateral equivalent to the Sermilik Formation. The Bylot Island Formation is comprised of mudstone and minor sandstone, representing slope, basin plain and turbidite deposits. Uppermost Bylot Island Formation strata conformably overlie the Sermilik Formation and represent a

ii

late Maastrichtian transgression within Eclipse Trough.

Three informal palynomorph assemblage zones are recognized in these rocks: Gleicheniidites sp. cf. G. circinidites - Antulsporites distaverrucosus (GA) zone, the Porosipollis porosus ~ Aquilapollenites scabridus (PA) zone and the Singularia aculeata - Pesavis parva (SP) zone. These zones support lithostratigraphic correlations between south coast and Twosnout Creek sections. The GA zone of mid to late Campanian age is characterized by the occurrence of late Cretaceous palynomorphs Carpinipites ancipites. Hazaria sheopiarii, and Polyatriopollenites stellatus. Its age is in part based on the overlying PA zone. The PA zone, of late Campanian to mid Maastrichtian age is characterized by the first occurrence of several biostratigraphically important late Cretaceous palynomorphs. Diagnostic species include Ceratiopsis diebelii, Palaeoperidinium kozlowskii, Momipites wyomingensis, Wodehouseia gracile and Porosipollis porosus. The SP zone of late Maastrichtian age is characterized by the first occurrence of Pesavis parva and Paraalnipollenites alterninorus.

These pollen floras are similar to other late Campanian to Maastrichtian assemblages from Banks Island, Horton River and Police Island, N.W.T..

iii

man in a survey a start and a survey of the second

ACKNOWLEDGEMENTS

I wish to thank Dr. Elliott Burden for the opportunity to study under his supervision. Joshua Enookolook provided able field assistance while on Bylot Island and is duly credited for the most northerly discovery of dinosaur remains in the eastern Canadian Arctic. My Tertiary counterpart, James Waterfield, is thanked for numerous helpful discussions and his companionship while on Bylot Island.

Financial and logistical support was provided by the Polar Continental Shelf Project at its base in Resolute, the Arctic Research Establishment at Pond Inlet, a Texaco Research Grant, Department of Indian Affairs and Northern Heritage Society - Northern Scientific Training Program Grant, and a Career Development Award.

Acknowledgements are also addressed to D. Russell, A. King, B. Stevens, R. Ellis Hayes, H. Williams, A. Aksu, D. Wilton and P. O'Neill. Brian Sears deserves a special thanks for photographic services and editorial suggestions. Final thanks are addressed to my fiancee, Anne Hogan, for her understanding and encouragement; and to my parents for their unending support.

iv

TABLE OF CONTENTS

Abstract	ii
Acknowledgements	iv
Table of Contents	v
List of Figures	vii
List of Tables	ix
CHAPTER 1 INTRODUCTION	1
1.1 Purpose and Objectives	1
1.2 Location and Access	1
1.3 Weather Conditions	3
1.4 Field Methods	3
1.5 Previous Studies	6
1.7 Geologic Setting	12
CHAPTER 2 STRATIGRAPHY	18
2.1 Introduction	18
2.2 Byam Martin Formation	18
2.3 Sermilik Formation	23
2.4 Bylot Island Formation	34
CHAPTER 3 PALYNOLOGY	42
3.1 Introduction	42
3.2 Methods	42
3.2.1 Processing	42
3.2.2 Identification	44
3.3 Systematics	45

v Page

	vi
CHAPTER 4 DISCUSSION	129
4.1 Biostratigraphy	129
4.1.1 GA Zone	131
4.1.2 SP Zone	133
4.1.3 PA Zone	137
4.2 Environmental Interpretation	141
4.3 Summary and Conclusions	148
REFERENCES CITED	151
PLATES	168
APPENDIX $A_1 - A_4$	186
APPENDIX B1 - B3	191
APPENDIX C1	194
APPENDIX C2	(in pocket)

LIST OF FIGURES (Short Titles)

Figure	Page
1.1 Regional Setting of Eclipse Trough	2
1.2 Local transportation methods	4
1.3 Location map of stratigraphic sections	5
1.4 Cretaceous / Tertiary geology of Bylot Island	11
1.5 Structural trends in the North Baffin Rift Zone	13
1.6 Location of Arctic Platform	14
2.1 Byam Martin For.ation - Type Section	19
2.2 Byam Martin Formation - Type Section	19
2.3 Detrital plots for sandstones	22
2.4 Sermilik Formation - Type Section	25
2.5 Sermilik Formation - Lithotype 3	25
2.6 Sermilik Formation - Channel cut-and-fill	28
2.7 Sermilik Formation - Unimodal planar crossbedding	28
2.8 Sermilik Formation - Carbonate cemented concretions	31
2.9 Bylot Island Formation - Type Section	35
2.10 Bylot Island Formation interfingering with	
Sermilik Formation	36
2.11 Bylot Island Formation / Pond Inlet Formation	
contact	38
2.12 Gradational contact between Bylot Island Formation	
and Sermilik Formation strata	38
2.13 Bylot Island Formation - Bioturbation	40
2.14 Bylot Island Formation - Tabular sandstones	40
4.1 Palynology Range Chart	130

vii

		viii
4.2	Paleogeographic reconstruction of Eclipse Trough	146
4.3	Stratigraphic cross-section	150

LIST OF TABLES (Short Titles)

Tab	le								I	age
1.1	Мар	Units	of	Bylot	Island	(after	Jackson	and		
	Day	vidson	1	975)						8
1.2	Map	Units	of	Bylot	Island	(after	Miall <u>e</u>	<u>t</u> al.,	1980)	10

4.1 Characteristics of Fan-deltas and Braid deltas 145

CHAPTER 1 INTRODUCTION

1.1 Purpose and Objectives

Exposures of Cretaceous and Tertiary strata are rare in the eastern Canadian Arctic (Miall <u>et al.</u>, 1980). Thus, a thick sequence of Cretaceous and Tertiary strata preserved in Eclipse Trough, Bylot Island (Figure 1.1) contains important evidence for interpreting the geologic history of the eastern Canadian Arctic. This study focuses on late Cretaceous "Kanguk Formation" strata of Bylot Island. The research involves the identification of stratigraphic units, an interpretation of their depositional environments, and the use of palynological data to correlate and date these units.

1.2 Location and Access

Bylot Island is situated at the northeastern tip of Baffin Island in the Northwest Territories (Figure 1.1), and encompasses an area of approximately 14,400 sq. km. The study area lies on southwest Bylot Island, where Cretaceous and Tertiary strata are exposed within the Eclipse Trough (Figure 1.1).

Canadian Airlines provided transportation north from Montreal to Pond Inlet, on nearby Baffin Island. Bylot Island is uninhabited; thus plans had to be finalized and sufficient supplies acquired prior to departure from Pond Inlet. The 4 man field party, including equipment, was



Figure 1.1 Regional Setting of Eclipse Trough (after Miall <u>et</u> <u>al.</u>, 1980). Inset shows location of Bylot Island in the southeastern Canadian Arctic. transported across the sea ice on Eclipse Sound with a komatik and skidoo (Figure 1.2). Ice conditions were favourable for the trip, although breakup occurred rapidly in the days following. Helicopter support throughout the field season (July 6th - August 6th, 1987) provided food, supplies, camp moves, support during traverses and final transport back to Pond Inlet.

1.3 Weather Conditions

Weather conditions were monitored on a daily basis with a comprehensive report being submitted to Polar Continental Shelf Project (PCSP) at the culmination of the field season.

Temperatures varied between -1° C and 6° C, with high winds gusting continuously for days. Few sunny days were recorded; overcast conditions prevailed, with precipitation being predominantly in the form of fog and drizzle. Early August saw colder conditions and light snowfall.

1.4 Field Methods

Four stratigraphic sections along the southern shore of Bylot Island and on western Bylot Island at Twosnout Creek were measured for palynology and sedimentology (Figure 1.3). Sections were generally well exposed, with lateral facies changes being unobscured by post depositional features such as faulting or slumping. Weathering of the exposures and cryoturbation was responsible for the distortion of some



Figure 1.2 Traditional transportation methods used by local Inuit provided initial access to the field area.



Figure 1.3 Location map of stratigraphic sections.

structures.

Sedimentology was carried out noting lithology, bed thickness, cyclicity, color, grain size, bed contacts, sedimentary structures and fossil content. Seventeen sandstones were collected and thin sectioned for petrography.

One hundred fifty palynomorph samples were collected and forty were processed. Samples were normally collected at 15 metre intervals; where tighter control was warranted, samples were collected at 5 metre spacings.

No systematic search was conducted for vertebrate or invertebrate remains. Discoveries of vertebrates during routine measuring and collecting were recorded for proper collection and preservation at some later date. Only small isolated bones, bone fragments, and teeth, in danger of destruction were collected for preservation. A small suite of invertebrates was also collected for identification and preservation.

1.5 Previous Studies

The earliest documented record of the geological history of Bylot Island was submitted in a report to the Department of Marine and Fisheries, Ottawa, by A.P. Low (1906), a geologist with D.G.S. Neptune. He reported that the northern and eastern coasts of Bylot Island appeared to be wholly formed of crystalline rocks.

The first extensive geological study of the coast of Bylot Island was administered and recorded by Captain J.E. Bernier (1910) of the D.G.S. Arctic. During routine prospecting at Canada Point on Bylot Island (Figure 1.3), Bernier's second officer, R.S. Janes, noted a similarity between these rock formations and those observed at the Salmon River coal deposit, near Pond Inlet (North Baffin Island). Further investigations at this locality uncovered fossilized trees and buds. An extensive coal seam was also discovered, thus confirming the relationship with the Salmon River coal beds.

The first systematic geological investigation of the entire island was carried out by the Geological Survey of Canada in 1968 (Jackson, 1969). The project resulted in the publication of two reports, each with a geological map at a 1:250,000 scale. The first was by Jackson and Davidson (1975) for the area north of 73° latitude, the other, by Jackson <u>et al.</u>, (1975), described the southern half of Bylot Island and northern Baffin Island.

Jackson and Davidson (1975) introduced the epithet Eclipse Group for at least 1067 metres of Cretaceous -Tertiary sediments on Bylot Island. Sediments within the group were divided into four map units (Table 1.1). Palynology by W.S. Hopkins and D.C. McGregor (Jackson and Davidson, 1975) indicated an early Cretaceous to Eocene age.

TABLE 1.1

Map units of Jackson and Davidson (1975)

	Fissile shale,
	muuscone, siitstone
2	Arkosic sandstone, siltstone, rare coal
	Subgreywacke, quartzwacke, mudstone, siltstone
	Orthoquartzite, arkosic sandstone, coal

These strata were examined by B. Clarke of Shell Oil Company in 1973. Clarke (in Jackson <u>et al.</u>, 1975) indicated the Eclipse Group may be much thicker than originally thought, possibly 1830 metres thick, and that the section may thin westward. A suite of samples from the Eclipse Group submitted to Aquitaine Company of Canada Ltd. for hydrocarbon studies (Jackson <u>et al.</u>, 1975), showed that the organic matter was incapable of sourcing viable quantities of hydrocarbons.

Cretaceous - Tertiary strata were investigated by Miall st al., (1980). From sedimentology and palynology, Miall <u>et</u> al., (1980) revised the map units established by Jackson and Davidson (1975) and Jackson <u>et</u> al., (1975). Miall <u>et</u> al., (1980) recognized three formations: Hassel - Early Cretaceous; Kanguk - Late Cretaceous; Eureka Sound -Tertiary (Table 1.2), (Figure 1.4). These formations are widespread throughout the Arctic Islands (Miall <u>et al.</u>, 1980). In reclassifying the Cretaceous - Tertiary stratigraphy, Miall <u>et al.</u>, (1980) dropped the term Eclipse Group. Miall (1986) redefined the Eureka Sound Formation as a group, combining the Te¹ and Te⁴ members into the Mount Lawson Formation, and the Te³ and Te⁴ members into the Mokka Fiord Formation.

Ioannides (1986) used dinoflagellate cysts to provide a more concise age assignment, (late Campanian to Maastrichtian) for the Kanguk Formation and (late

Table 1.2

Map units of Bylot Island

(units mapped by Miall et al., 1980)

Map Unit	Formation	Thickness (m)	Lithology
Te ⁴		200+	mudstone, minor sandstone
Те ³		1370+	immature sandstone, minor siltstone, mudstone
Te ²	Eureka Sound	80-500+	mudstone, jarosite, minor sandstone
Te'		0-200+	glauconitic sandstone
Kk²	Kanguk	0-540	immature sandstone, minor siltstone, mudstone
Kk ¹		560-590	mudstone
Kh	Hassel	10-120	sandstone, minor mudstone



Figure 1.4 Cretaceous - Tertiary geology of southwest Bylot Island (after Miall <u>et al.</u>, 1980).

Maastrichtian to Paleocene) for the Eureka Sound Group. The study suggested that revisions to the current stratigraphy are necessary.

1.6 Geologic Setting

Eclipse Trough is one of a series of fault-controlled basins in the North Baffin Rift Zone (Niall <u>et al.</u>, 1980) (Figure 1.5). Extensional deformation propagated northwestward along pre-existing structural trends inherited from the Precambrian crystalline basement (Kerr, 1980). Approximately 1200 metres of Upper Cretaceous strata were deposited in response to faulting and extensional deformation during the Eurekan Rifting Episode (Kerr, 1980).

The Eurekan Rifting Episode occurred with the compressional deformation event (Eurekan Orogeny), which affected the Sverdrup Basin several hundred kilometres to the northwest (Thorsteinsson and Tozer, 1970). Both events, collectively known as the Eurekan Deformation, affected the Canadian Arctic in the late Cretaceous (Kerr, 1980). Structural and stratigraphic evidence also indicates that a physical barrier (Arctic Platform) separated the Sverdrup Basin and Eclipse Trough (Meneley <u>et al.</u>, 1975; Balkwill, 1978; Kerr, 1980 and Miall, 1986), (Figure 1.6).

Prior to and during the onset of the Eurekan Deformation, marine sediments were thought to be accumulating in a widespread shallow sea that covered much



Figure 1.5 Dominant structural trends of Northern Baffin Island and Bylot Island, inherited from northwest trending structures in Precambrian crystalline basement (after Jackson and Iannelli, 1981).





of the Arctic during the late Cretaceous (Kerr, 1980; Miall <u>et al.</u>, 1980). The resultant sedimentary package has come to be known as the Kanguk Formation (Souther, 1963; Kerr, 1980; Miall <u>et al</u>., 1980).

The Kanguk Formation (Souther, 1963) was initially named for a 364 m thick marine succession of grey shale, in part silty, with minor amounts of sandstone, bentonitic shale and tuffaceous beds. Souther (1963) designated the type locality within the Sverdrup Bosin as the Kanguk Peninsula, Axel Heiberg Island. Kanguk equivalent strata have been recognized elsewhere within the Sverdrup Basin on Ellef Ringnes Island (Grenier, 1963; Stott, 1969), Lougheed Island (Balkwill <u>et al.</u>, 1982), Amund Ringnes Island (Balkwill, 1983), within the Banks Basin on Banks Island (Jutard and Plauchut, 1973; Miall, 1975; Miall, 1979) and within the Eclipse Trough on Bylot Island (Miall <u>et al.</u>, 1980).

On Lougheed Island, Balkwill <u>st al.</u>, (1982) assigned 200 metres of intercalated black, coaly, pyritic shale and fine-grained buff carbonaceous sandstone to the Kanguk Formation. Balkwill <u>et al.</u>, (1982) noted that strata of Lougheed Island differed lithologically when compared to Kanguk strata in the type locality and its immediate vicinity (i.e. Amund Ringnes Island and Ellef Ringnes Island). Nevertheless, due to underlying and overlying relationships with assumed Hassel and Eureka Sound Formation

strata, the rocks were assigned to the Kanguk Formation.

On Banks Island, 1000 km southwest of the type section, beds assigned to the Kanguk Formation have been divided into two members (Jutard and Plauchut, 1973): (1) a lower bituminous member, consisting of black bituminous shale, coal and carbonaceous shale; (2) and an upper shale member, consisting mainly of light grey shale, minor sandstone and rare jarosite layers. Miall (1979) further subdivided the Kanguk Formation on Banks Island into five members: a lower bituminous member; a lower salty shale; a lower sand; a silty shale; and an upper sand member.

The structural setting of the Banks Basin is similar to that for Eclipse Trough. The area underwent extensional deformation, contrasting in structural style with that of the Sverdrup Basin (Eurekan Oroge.y). The Banks Basin was similarly separated from the Sverdrup Basin by part of the Sverdrup Rim (Storkerson Uplift) (Meneley <u>et al.</u>, 1975) and part of the Prince Patrick Uplift (Cape Crozier Anticline) (Miall, 1979). Petrographic evidence indicates that sandstones within the Banks Basin were derived locally from the nearby Storkerson Uplift and the Cape Crozier Anticline, with the shale being derived from adjacent low-lying land areas (Miall, 1979).

Upper Cretaceous rocks on Bylot Island are very distinctive. They consist of up to 100 metres of grey lithified mudstone and muddy sandstone, 450 metres of black

unconsolidated silty mudstone with randomly distributed sandstone beds and a 540 metre package of very coarsegrained marine sandstone. Petrographic evidence (Chapter 3) strongly suggests a local sediment source. The rocks are clearly distinct from strata identified as Kanguk Formation elsewhere in the arctic.

From comparisons in the literature, it is evident that several lithofacies variations exist between "Kanguk formation" strata of the Sverdrup Basin, Banks Baein and Eclipse Trough. As required by the North American Stratigraphic Code, a formation should be based strictly on description. Therefore, from descriptions, Upper Cretaceous "Kanguk formation" strata within the Eclipse Trough, the Sverdrup Basin and the Banks Basin comprise several distinct lithological and mappable formation rank units.

In addition to the lithostratigraphic evidence, structural and stratigraphic evidence suggests that "Kanguk" strata in the Eclipse Trough were affected by different tectonic regimes and are from different provenance than that for rocks of the Sverdrup Basin (Meneley <u>et al.</u>, 1975; Balkwill, 1978; Kerr, 1980; and Miall <u>et al.</u>, 1980).

To assign late Cretaceous strata of Bylot Island to the "Kanguk formation" of the Sverdrup basin would clearly be in violation of the North American Stratigraphic Code (Article 22a). In keeping with this code, late Cretaceous "Kanguk formation" strata on Bylot Island will be revised.

CHAPTER 2 STRATIGRAPHY

2.1 Introduction

From arguments presented in the preceding section, and in keeping with the North American Code on Stratigraphic Nomenclature, late Cretaceous strata on Bylot Island have been informally divided into three new and lithologically distinct formations. The epithet "Eclipse Group" of Jackson <u>et al.</u> (1975) is herein reinstated. It is comprised of three Cretaceous formations, herein named: the Byam Martin Formation, the Sermilik Formation, the Bylot Island Formation and Tertiary formations named and discussed by Waterfield (personal communication, 1989).

2.2 Byam Martin Formation

Definition, Distribution and Thickness

The Byam Martin Formation (named after the adjacent Byam Martin Mountains) is the lowest stratigraphic unit examined in this study. It consists of grey, well lithified mudstone; thin sandstone beds comprise less than 1% of the formation. The thickness at its type section at Twosnout Creek is 105 metres (Section A, Appendix A; Figures 2.1, 2.2), no other complete sections were measured. Talus and frost heave indicate that the formation extends south along the front of the Byam Martin Mountains to the Sermilik Glacier, one poorly exposed section in an outcrop near the Sermilik Glacier is 12 metres thick (Section B,



Figure 2.1 Type Section of the Byam Martin Formation.



Figure 2.2 Byam Martin Formation at Twosnout Creek. Close up of Type Section (largest interval of staff is 0.5 metres).

Appendix A,).

Synonyms

The Byam Martin Formation includes the lower 100 metres of the Kanguk Formation (KK¹ mudstone member) of Miall <u>et</u> <u>al.</u>, (1980) at Twosnout Creek. It is also equivalent to the entire Kanguk Formation (KK¹ mudstone member) exposed on the south coast of Bylot Island.

Contacts

At the type locality, the Byam Martin Formation unconformably overlies 14.5 metres of a basal sandstone, identified (<u>in Miall et al.</u>, 1980) as Hassel Formation strata. Contact with the overlying Bylot Island Formation is gradational over several metres; the contact between these formations is placed on the upper contact of the last occurrence of lithified grey mudstone.

Poorly exposed strata of interbedded sandstone and muddy sandstone along the south coast suggest a transitional and probable conformable relationship with the overlying Sermilik Formation. The basal contact is not exposed in this area.

Lithology

The formation consists of well lithified grey (occasionally grey-black to red) mudstone with rare, centimetre to decimetre thick beds of medium-grained, white,

weakly lithified subarkosic sandstone. Sandstone beds are laterally continuous for several tens of metres and frequently contain mudstone rip-up-clasts. Other sedimentary features include sandstone dykes and flame structures. Bioturbation is sporadic (1-5% - based on classification by Reineck and Singh, 1973). Burrows are horizontal and very small (1-2mm wide); they resemble Chondrites (Plate 1, Figure 1).

Immediately west of the Sermilik glacier, a 12 metre section outcrops adjacent to the Byam Martin Mountains. The sediments, where exposed in section, consist of poorly lithified, interbedded, very fine- to fine-grained white sandstone and brown muddy sandstone, and lesser coarser sands. Abundant, grey, lithified mudstone fragments are scattered about this hillside as frost heave, but nowhere are they exposed in section. Sedimentary structures are exclusively burrows; these vertical shafts (Plate 1, Figure 2) are confined to the muddier brown substrate. The section is only weakly bioturbated (5-30%), with burrows being infilled by coarser sand.

Petrology

Sandstones from the Byam Martin Formation straddle the boundary between the subarkose and quartzarenite fields (Figure 2.3; Appendix B₁, B₂, B₃). Orthoclase and andesine are the dominant feldspar types. Most feldspars are clear



Figure 2.3 Detrital plots for sandstones.
and unaltered. However, two specific alteration types are recognized in the altered feldspars. An alteration assemblage of calcite and epidote is recognized as replacing some plagioclase grains, while sericite is noted as replacing the alkali grains. Heavy minerals identified in thin section include epidote and rutile. The adjacent Byam Martin Mountains provide an obvious source of detritus.

Clay matrix composes less than 5% of the total rock volume. Grains are rounded to well rounded, and moderately to well sorted.

Age

Palynomorph analysis (see Chapter 3) suggests a middle to late Campanian age for the Byam Martin Formation. This age assignment is based primarily on the absence of characteristic early Maastrichtian assemblages of <u>Caratiopsis</u>, and <u>Palaeoperidinium</u>, and the conformable nature of the formation with the overlying early Maastrichtian strata.

Angiosperms recovered from the Byam Martin Formation included such genera as <u>Carpinipites</u>, <u>Pseudoplicapollis</u>, and <u>Polyatriopollenites</u>.

2.3 Sermilik Formation

Definition, Distribution and Thickness

The Sermilik Formation (named after the adjacent

Sermilik Glacier) conformably overlies the Byam Martin Formation. It is named after the Sermilik glacier, where the type section occurs (Section B, Appendix λ_2), and is exceptionally well exposed (Figure 2.4). It is comprised of three lithotypes: a unit of weakly lithified, calcareous, buff colored, very coarse-grained sandstone; a green to buff unconsolidated, coarse-grained sandstone unit; and a grey, coarse-grained to conglomeratic unit. The thickness of the Sermilik Formation at is type section is 472 metres.

It also outcrops at Twosnout Creek; lenticular sandstone tongues of the Sermilik Formation (Lithotype 3) measuring between 17 and 40 metres in thickness, occur within the Bylot Island Formation (Figure 2.5). A reference section is herein included to demonstrate Lithotype 3 characteristics (Section C, Appendix A).

Synonyms

At its type section, the Sermilik Formation is equivalent to the KR^2 sandstone member of the Kanguk Formation of Miall <u>et al.</u>, (1980). At Twosnout Creek, the Sermilik Formation is a series of tongues in the Bylot Island Formation. These sandstone tongues were not mentioned in Miall <u>et al.</u>, (1980). They were recognized by Jackson and Davidson (1975), and incorporated into their Unit K of the Eclipse Group.



Figure 2.4 Type Section of the Sermilik Formation. Note the stacked channels at the center of the photo. Each channel is approximately 10 metres thick.



Figure 2.5 Sermilik Formation: Lithotype 3. Note the interfingering with Bylot Island Formation strata.

Contacts

The contact of the Sermilik Formation with the underlying Byam Martin Formation is not exposed. A river valley separates Byam Martin strata from the Sermilik Formation strata: nevertheless, a conformable contact is inferred within the stratigraphic succession. The basal contact is placed at the point where the first coarsegrained sandstone bed is observed. The upper contact with the overlying Bylot Island Formation is conformable and gradational over 10 metres, where coarse-grained and interbedded with mudstones of the Bylot Island Formation. The contact is arbitrarily placed at the point where the average thickness of mudstone beds exceeds the thickness of sandstone beds.

At Twosnout Creek, lenticular sandstone pods are enveloped by Bylot Island Formation mudstone. These tongues suggest the Sermilik Formation is in part laterally equivalent with the Bylot Island Formation.

Lithology

The basal unit (Lithotype 1) is composed of 316 metres of well-sorted, buff to less commonly grey, medium- to very coarse-grained pebbly, subarkosic sandstone.

Beds are lenticular, with a maximum width of 200 metres. Thickness varies from decimetre to metre-thick

units, with some individual beds reaching a maximum of ten metres (Figure 2.4). Contacts are always scoured. Some are incised up to 3 metres into underlying strata. The channeled bases contain rip-up-clasts, woody debris and well rounded pebbles or cobbles (Figure 2.6). Sediment grain size almost always decreases upwards to a medium-grained, occasionally bioturbated sandstone. Channels commonly erode previously deposited channel-fill to produce a series of stacked or multi-story sands which preserve only the coarse portion of the channel sequence.

Poor to well-defined planar-tabular and planartangential cross-stratification is preserved in channel sandstone bodies. Sets average 10 centimetres in thickness, the maximum set thickness is 30 centimetres. Cosets average one metre. However, one coset measured in the south coast (Section B, Figure 2.7) is 16 metres thick. Cross-bedding is unimodal; Miall <u>et al.</u>, (1980) found a paleocurrent azimuth of 292°.

Other sedimentary structures in channel deposits include poorly defined trough cross-stratification averaging 15 centimetres in thickness and 50 centimetres in length. Packages of channelized beds exhibit fining and coarsening cycles over several metres. Diagenetic features evident include discontinuous centimetre to decimetre beds of round to platy calcite cemented concretions. Inter-channel strata consist of fining-upwards, decimetre thick beds of medium-



Figure 2.6 Channel cut-and-fill in Lithotype 1.



Figure 2.7 Sermilik Formation: Lithotype 1. Planar crossbedding with unimodal orientation.

grained argillaceous sandstones. Lateral tracing indicates beds thin and fine outwards from the channel margin. Some groups of stacked sandstone beds exhibit reactivation surfaces. Overall coarsening upward sequences are a common feature. Locally, ripple cross-stratification is preserved, but is rare.

Marine macrofossil detris is uncommon, only scattered pelecypod fragments were recovered, although, palynological evidence indicates an abundance of well preserved, late Cretaceous dinoflagellates.

 λ 156 metre thick unit (Lithotype 2) of green to buff, weakly consolidated, subarkosic sandstone directly overlies the lower buff unit. Beds from Lithotype 1 are gradational with beds of Lithotype 2.

Beds of Lithotype 2 are dominantly lenticular, with widths averaging between 10 and 20 metres. Decimetre thick units within beds are usual, though metre size beds are also present. Channelized bases frequently contain rip-up clasts with well rounded pebbles, cobbles, or boulders, and fineupwards to a coarse-grained and muddy top. Channels are commonly stacked, and exhibit coarsening- and fining-cycles as in Lithotype 1.

Beds transitional with the Bylot Island Formation are finer-grained, and are lithified. Here, channelized sandstones exhibit a variety of sedimentary structures such as angular rip-up-clasts, poorly defined trough cross-

bedding, climbing ripple cross-laminations, parallellaminations and large, round, metre size, carbonate cemented concretions (Figure 2.8).

Centimetre to decimetre beds of fine- to mediumgrained, parallel bedded, muddy sandstome represent interchannel areas. A variety of trace fossils, which include ' Diplocraterion, Skolithos, and Thalassinoides, have been identified (Plate 1, Figure 3). Marine macrofossil debris is abundant and diverse; fossils include bivalves, gastropods, marine reptile vertebrae (Plate 1, Figure 4), and shark teeth (Plate 1, Figure 5).

Lithotype 3 sediments are comprised of lentils of grey, coarse-grained pebbly and congloweratic, lithic subarkose; and occur within the Bylot Island Formation at Twosnout Creek. These lentils are offshore extensions of the Semulik Formation. Beds are lenticular; averaging between 5 and 20 metres in width and are preserved as decimetre to metre thick units.

Lithotype 3 is characterized by a polymodal, matrix supported, pebble to boulder conglomerate which fines upwards. Clasts are sub-angular to rounded. The average size is 40 centimetres and the maximum size is 80 centimetres. The clasts are composed of granitic, metamorphic and intraformational rock fragments. Coarsely crystalline bivalve debris is preserved in some intraformational clasts. Sedimentary structures include



Figure 2.6 Channel cut-and-fill in Lithotype 1.



Figure 2.7 Sermilik Formation: Lithotype 1. Planar crossbedding with unimodal orientation.

normal to reverse grading, trough cross-stratification, parallel laminations, flame structures and contorted beds. fossil invertebrates also occur as scattered debris and coquina of abraded bivalves, gastropods, echinoids, and scaphopods in sandstone beds.

Petrology

Sandstones from Lithotypes 1 and 2 of the Sermilik Formation fall clearly within the subarkose field (Figure 2.3), while sandstones from Lithotype 3 at Twosnout Creek fall within the lithic subarkose field. Feldspars are unaltered; the dominant types are orthoclase and andesine. Quartz grains are dominantly unstrained and monocrystalline. The dominant rock fragment recognized is a quartz rich plutonic rock. Other recognizable rock fragments include non-foliated metamorphic clasts and sedimentary clasts. Heavy minerals include biotite, garnet, tourmaline and pyroxene.

During routine measuring and sampling, pebbles and cobbles were visually examined for gross composition. Plutonic pebbles are the most common. These are followed by metamorphic pebbles or cobbles of mylonitic and gneissic origin. Sedimentary and volcanic pebbles were thirdly observed.

Porosity in lithified samples (Appendix B,) is consistently less than 5%. Some sandstone beds are cemented

with calcite, which may comprise up to 10% of the total rock volume. Clay matrix accounts for less than 1% of the total rock volume. Both sorting and roundness are highly variable; grains range from poor to well-sorted and subangular to well-rounded.

The angularity and freshness of the plutonic rock fragments clearly suggests a first-cycle origin. Unnamed and undivided Archean-Aphebian plutonic rocks of the Byam Martin High are an obvious localized detrital source. The presence of sedimentary pebbles, volcanic pebbles, and accessory minerals (pyroxene) suggests that the sandstones were in part derived from rocks of the Mary River Group, adjacent the Semilik Glacier.

Age

The age of the Sermilik Formation is designated by both macrofossil and microfossil evidence. The recovery of a dinosaur toe bone (Hadrosaur) (Plate 1, Figure 6) near the base of the formation (Lithotype 1) constrains the age, as Hadrosaurs lived between 65 and 75 million years ago (Russell, personal communication, 1987). One bivalve recovered from Lithotype 3 at Twosnout Creek vas identified as <u>Nemdon</u> sp. by J. Haggart of the Geological Survey of Canada. This particular bivalve is restricted to the late Cretaceous (Cenomanian to Masstrichtian) of North America, Japan, and Madagascar (Cox <u>et al.</u>, 1969). From palynology,

the age of the Sermilik Formation is late Campanian to early late Maastrichtian. Characteristic palynomorphs include <u>Modehouseia, Momipites, Ceratiopsis</u> and <u>Palaeoperidinium</u>.

2.4 Bylot Island Formation

Definition, Distribution and Thickness

The Bylot Island Formation is a sequence of black, friable silty mudstone, with subordinate medium-grained, subarkose and quartzarenite. At its type section at Twosnout Creek (Section A, Appendix A,; Figure 2.9) the Bylot Island Formation is 420 metres thick. An additional section (Section C, Appendix A,), incorporating Lithotype 3 strata of the Sernilik Formation, measured 80 metres in thickness. Along the south coast, a tongue of Bylot Island Formation extends into Sermilik Formation (Section D, Appendix A, Figure 2.10).

Synonyms

The Bylot Island Formation has been called the KK¹ mudstone member of the Kanguk Formation of Miall <u>et al.</u>, (1980). In south coast sections, a tongue of Bylot Island strata was previously recognized as part of the KK² sandstone member of the Kanguk Formation. Waterfield (personal communication, 1989) has also shown that the Bylot Island Formation along the south coast is equivalent to the Te² mudstone member of Miall <u>et al.</u>, (1980).



Figure 2.9 Type Section of the Bylot Island Formation.



Figure 2.10 Tongue of Bylot Island Formation strata interfingering with Sermilik Formation (Lithotype 1 and Lithotype 2).

Contacts

The contact with the underlying Byam Martin Formation is gradational. Lithified grey mudstones of the Byam Martin Formation become interbedded with black, friable silty mudstones of the Bylot Island Formation. The contact between these formations is placed directly on the upper contact of the last occurrence of lithified grey mudstone.

Overlying the Bylot Island Formation is the Pond Inlet Formation (Waterfield, personal communication, 1989). Although the lateral transition from the Pond Inlet Formation to the Bylot Island Formation is considered gradational (Waterfield, personal communication, 1989), the contact is locally sharp and erosional. Locally, the Pond Inlet Formation is thought to be a submarine fan. Lensoid deposits contain erosional disconformities in a regionally conformable sequence (Waterfield, personal communication, 1989). The contact is placed at the base of the first white sandstone bed more than 1 metre thick (Figure 2.11).

Along the south coast, tongues of Bylot Island Formation strata interfinger laterally with Sermilik Formation sandstones. Here, contacts consist of interbedded sandstones and siltstones (Figures 2.12).

Lithology

The Bylot Island Formation is composed of black, friable silty mudstone. Locally, the mudstone contains



Figure 2.11 Bylot Island Formation / Pond Inlet Formation contact.



Figure 2.12 Interbedded sandstones (Sermilik Formation) and mudstones (Bylot Island Formation) indicating the gradational contact between these formations. siderite concretions which are up to a metre in diameter. Bioturbation consisting of <u>Chondrites</u>? is sporadic at Twosnout Creek, in contrast, the tongue of Bylot Island Formation mudstone along the south coast is completely bioturbated by <u>Thalassinoides</u> (Section D, Appendix λ_{i} , Figure 2.13). These burrows are silt filled and/contrast little with material adjacent to the burrow.

Sandstone beds of the Bylot Island Formation are similar to sandstone beds of the Byam Martin Formation. Beds are tabular in appearance and average 50 centimetres in thickness (Figure 2.14); they display normal grading and frequently contain small (<1 cm thick) rip-up-clasts. Sedimentary structures consist of contorted beds and flame structures.

Petrology

Sandstones of the Bylot Island Formation are similar in composition to those of the Byam Martin Formation. All samples straddle the boundary between the subarkose and quartzarenite fields (Figure 2.3). Feldspars are unaltered, with the major types being orthoclase and andesine. Rock fragments consist of plutonic and metamorphic fragments. Heavy minerals include biotite, garnet, sphene, tourmaline and muscovite. Unnamed and undivided Archean-Aphebian Plutonic rocks of the Byam Martin High represent a local detrital source.



Figure 2.13 Close-up of Bylot Island Formation strata in Figure 2.10. Note the extensive bioturbation in the mudstone (staff is marked in 0.1 metre intervals).



Figure 2.14 Subordinate, tabular sandstones occurring within the Bylot Island Formation.

Clay matrix was consistently less than 6% of the total rock volume. Where present, calcite cement was less than 10%. All samples contain moderately to well-sorted, and subrounded to well-rounded grains.

Age

From palynology, this marine unit is very late Campanian to early late Maastrichtian in age (Chapter 3). Characteristic genera include <u>Aquilapollenites</u>, <u>Porosipollis</u>, <u>Pesavis</u> and <u>Ceratiopsis</u>.

CHAPTER 3 PALYNOLOGY

3.1 Introduction

Terrestrial palynology has been utilized to develop the first detailed palynomorph biostratigraphy for late Cretaceous strata on Bylot Island. Reconnaissance biostratigraphic studies (Jackson and Davidson, 1975, Miall <u>et al.</u>, 1980, Ioannides, 1986) were inconclusive; stated problems include reworking, corrosion of palynomorphs, an abundance of long-ranging species and a lack of distinctive taxa such as those recorded by Felix and Burbridge (1973); McIntyre (1974); and Doerenkamp <u>et al.</u>, (1976) for other Arctic localities.

3.2 Methods

3.2.1 Processing

One hundred twenty samples were collected from strata on Bylot Island. To limit contamination, unconsolidated sediment was freshly exposed before sampling. Forty of these samples have been processed and examined for palynology (Appendix C.).

Labelled samples were placed in a clear plastic container, and covered with a lint-free tissue (for 24hrs) to air dry. The samples were then weighed (15-20 grams for siltstone and shale; 30-35 grams for sandstones) and placed in labelled 250 ml beakers. To each sample, 12,100 ± 400 <u>kroepodium</u> grains were added to provide an estimate of

palynomorph concentrations, and to identify possible processing errors (Stockmarr, 1971).

To remove carbonate, 150 ml of 20% Hydrochloric acid (RC1) was added to each sample. Methanol was used to control any excessive effervescence. The sample was periodically stirred and left for 12 hours. The samples were then centrifuged and the acid decanted. Distilled water was added, centrifuged, and decanted to remove all remaining acid.

Hydrofluoric (HF) acid was used to remove silicates. Samples were immersed in (HP) for 12 hours. Periodic stirring was necessary. The acid was then decanted and the sample washed with distilled water, and centrifuged until neutral. At this stage, the first of 5 slides was prepared from the unoxidized, unsieved portion.

The unoxidized, unsieved portion was then sieved through a 10µm screen using a technique described by Cwynar <u>et al.</u>, (1979). Prior to oxidation, slides 2 and 3 were prepared from the sieved sample. In addition, approximately 15 mls of residue was placed in a labelled vial for storage. The preparation of slides at various increments and the storage of residue allows for mistakes to be found and corrected without the entire preparation having to be restarted.

For oxidation, 20 ml of schultz solution was added to the sample. Samples were stirred thoroughly, and left for

4-5 minutes. The sample was then washed and centrifuged three times. Prior to staining, the sample was sieved once again.

Three drops of Safranin O, were added to the sample, mixed, and left for five minutes. The sample was then centrifuged and decanted until the liquid was clear. Slides 4 and 5 of the oxidized, sieved, stained material were prepared.

To mount the slides, one or two drops of the sample were placed on a cover slip with two drops of polyvinal alcohol, and spread evenly with a toothpick. Samples were dried with a heat lamp to avoid prolonged exposure to possible contaminants. Once dry, two drops of Elvacite (Dupont) were placed on the residue on the cover slip. The coverslip was then mounted on a glass slide. The slide was left to dry overnight (8-12 hrs). The remaining residue was placed in a labelled vial and stored.

3.2.2 Identification

Two microscopes were used for palynology. Taxonomy and counts were completed with a Reichert Zetopan microscope, serial number 341717. Photographs were taken with a Zeiss Photomicroscope III, using Kodak, T-Max 100 E.I., with the optics set for interference contrast.

Slides were scanned at 0.5 mm intervals at 400X. Excluding the <u>Lycopodium</u> spike, 200 grains were counted per

slide. In all samples a minimum of two slides were needed to achieve this number.

3.3 Systematics

For presentation purposes, proper classification of fossil palynomorph data is of utmost importance. Two of the more widely used systems are: the "Turma" system (Potonié and Kremp, 1954; Potonié, 1956; 1958; 1960; and Dettmann, 1963); and the "Natural" classification system used by Couper (1958), Filatoff (1975) and Singh (1964; 1971).

The "Turma" system is a morphological classification system; phyletic relationships are not implied. In "Natural" classification, genera are assigned to higher rank according to their morphologic similarities (Burden, 1982).

The classification system used in this study is a morphological one, based in part on that of Burden (1982). Palynomorphs are divided into the following groups and subgroups:

TERRESTRIAL PALYNOMORPHS

Trilete Spores Monolete Spores Bisaccate Pollen Monosulcate Pollen Tricolpate Pollen Tricolporate Pollen Triporate Pollen Stephanoporate Pollen Binigeminate Pollen Triprojectate Pollen Syncolpate Pollen Unknown Specimens FUNGAL REMAINS MARINE MICROPLANKTON

Proximate Dinoflagellates

The entry for each taxon includes a photograph, selected synonymy, a short description (where appropriate) and data on the relative abundance and occurrence in this study. Age and distribution data from the literature are included for those species which are biostratigraphically significant.

TERRESTRIAL PALYNOMORPHS

Trilete Spores

Genus Annulispora de Jersey, 1959

Type Species: <u>Annulispora folliculosa</u> (Rogalska) de Jersey, 1959.

Annulispora sp. 1 Pl. 2, Fig. 1

Description: Trilete; amb convexly subtriangular with a 4µm

wide cingulum and a 4µm wide circum polar thickening. Size Range: Equatorial diameter of two measured specimens is 34µm and 36µm.

Remarks: The two identified specimens were restricted to the Bylot Island Formation and the Byam Martin Formation.

Genus <u>Cinqutriletes</u> Pierce emend. Dettmann, 1963 Type Species: <u>Cinqutriletes congruens</u> Pierce, 1961.

Cingutriletes pocockii (Burger) Burden, 1982

Pl. 2, Fig. 2

Synonymy:

- 1980 <u>Stereisporites pocockii</u> Burger, p. 58, pl. 12, figs. 14-17.
- 1982 <u>Cingutriletes pocockii</u> (Burger) Burden, p. 191, pl. 10, figs. 1, 2.

Remarks: Burden (unpublished Ph.D thesis, 1982) recognized that grains described by Burger (1980) as <u>Stereisporites</u> <u>pocockii</u> were clearly cingulate, whereas the genus <u>Stereisporites</u> Pflug, 1953 is acingulate. Species observed in this study have a well developed cingulum. Sporadic occurrences were recorded in the Bylot Island and Sermilik formations, however, this species occurs consistently within the Byam Martin Formation along the south coast. Genus Densoisporites Weyland and Kreiger

emend. Dettmann, 1963

Type Species: <u>Densoisporites</u> velatus Weyland and Kreiger, emend. Krasnova, 1961

Densoisporites velatus Weyland and Kreiger,

emend. Krasnova, 1961

Pl. 2, Fig. 3

Synonymy:

- 1963 <u>Densoisporites velatus</u> (Weyland and Kreiger) Krasnova; Dettmann, p. 84, pl. 19, figs. 4-8.
- 1971 <u>Densoisporites velatus</u> (Weyland and Kreiger) Krasnova; Singh, p. 46, pl. 3, figs. 9, 10.
- 1980 <u>Densoisporites velatus</u> (Weyland and Kreiger) Krasnova; Wingate, p. 12, pl. 2, fig. 10.
- 1981 <u>Densoisporites velatus</u> Weyland and Kreiger; Ferngreen and Chlonova, p. 447, pl. 1, fig. 6.
- 1982 <u>Densoisporites velatus</u> (Weyland and Kreiger) Krasnova; Burden, p. 305, pl. 24, figs. 5-10.
- 1986 <u>Densoisporites velatus</u> (Weyland and Kreiger) Krasnova; Ricketts and Sweet, p. 19, pl. 1, fig. 14.

Remarks: This species is restricted to the Sermilik Formation (Lithotype 1) along the south coast and to the Bylot Island Formation at Twosnout Creek. Genus <u>Antulsporites</u> Archangelsky and Gamerro, 1966

Type Species: <u>Antulsporites baculatus</u> (Archangelsky and Gamerro) Archangelsky and Gamerro, 1966.

Antulsporites distaverrucosus (Brenner) Archangelsky and Gamerro, 1966

Pl. 2, Fig. 4

Synonymy:

- 1963 <u>Cingulatisporites distaverrucosus</u> Brenner, p. 58, pl. 13, figs. 6, 7; pl. 14, fig. 1.
- 1971 <u>Antulsporites distaverrucosus</u> (Brenner) Archangelsky and Gamerro; Singh, p. 109, pl. 15, figs. 6, 7.
- 1980 <u>Antulsporites distaverrucosus</u> (Brenner) Archangelsky and Gamerro; Wingate, p. 23, pl. 9, figs. 7, 8.

Remarks: This species is restricted to rocks of the Byam Martin Formation on the south coast.

Genus <u>Cirratriradites</u> Wilson and Coe, 1940 Type Species: <u>Cirratriradites saturni</u> (Ibrahim) Schopf, Wilson and Bentall, 1944.

Cirratriradites teter Norris, 1967

Pl. 2, Fig. 5

Synonymy:

1967 Cirratriradites teter Norris, p. 98, pl. 14, figs. 6-10.

- 1971 <u>Cirratriradites teter</u> Norris; Singh, p. 118, pl. 15, figs. 14, 15.
- 1975 <u>Cirratriradites</u> <u>teter</u> Norris; Brideaux and McIntyre, p. 56, pl. 2, fig. 36.
- 1975 <u>Cirratriradites teter</u> Norris; Norris <u>et al.</u>, p. 338, pl. 1, fig. 15.

Age and Distribution: The range of this species is Aptian-Albian (Brideaux and McIntyre, 1975), middle to late Albian (Singh, 1971), and Albian ? to Cenomanian (Norris, 1967).

Remarks: This species is restricted to the Sermilik Formation (Lithotype 1) along the south coast. The presence of <u>Cirratriradites teter</u> in late Cretaceous strata of Bylot Island may be a result of reworking. Specimens of this species were, however, very well preserved; alternatively, this may indicate a range extension of the species into the Maastrichtian of Bylot Island.

Genus <u>Appendicisporites</u> Weyland and Krieger, 1953 Type Species: <u>Appendicisporites tricuspidatus</u> Weyland and Greifeld, 1953.

> Appendicisporites bifurcatus Singh, 1964 Pl. 2, Fig. 6

Synonymy:

1964 Appendicisporites bifurcatus Singh, p. 54, pl. 5,

figs. 1-5.

- 1971 <u>Appendicisporites bifurcatus</u> Singh; Singh, p. 56, pl. 4, figs. 3-5.
- 1971 <u>Appendicisporites bifurcatus</u> Singh; Playford, p. 544, pl. 104, fig. 10.
- 1975 Appendicisporites bifurcatus Singh; Brideaux and McIntyre, p. 15, pl. 2, fig. 10.
- 1975 <u>Appendicisporites bifurcatus</u> Singh; Srivastava, p. 12, pl. 3, figs. 8-10; pl. 4, figs. 1-8; pl. 5, figs. 1-3.
- 1987 <u>Appendicisporites bifurcatus</u> Singh; Langille, p. 60-61, pl. 2, fig. 5.

Remarks: This species is very rare, a total of three specimens were observed; one specimen from the Sermilik Formation (Lithotype 2) and one each from the Byam Martin and Bylot Island formations.

Appendicisporites erdtmanii Pocock, 1964

Pl. 2, Fig. 7

Synonymy:

- 1964 <u>Appendicisporites erdtmanii</u> Pocock; Singh, p. 48, pl. 2, figs. 5-7.
- 1971 <u>Appendicisporites</u> erdtmanii Pocock; Singh, p. 58, pl. 4, fig. 11.
- 1975 <u>Appendicisporites erdtmanii</u> Pocock; Brideaux and McIntyre, p. 2, pl. 2, fig. 17.

- 1980 <u>Appendicisporites</u> <u>erdtmanii</u> Pocock; Wingate, p. 15, pl. 3, figs. 13, 14.
- 1982 <u>Appendicisporites erdtmanii</u> Pocock; Burden, p. 204, pl. 11, fiqs. 5, 6.
- 1988 <u>Appendicisporites</u> <u>erdtmanii</u> Pocock; Sweet and McIntyre, p. 503, fig. 5, No. 5, 8.

Remarks: This species is characterized by 4 to 6 distal ribs in each interradial region. Ribs are parallel to one another and to the side of the spore where they form a triangular area centered on the distal pole (Singh, 1971). This species is restricted to the Bylot Island Formation.

Genus Camarozonosporites Pant ex Potonié emend.

Klaus 1960

Type Species: <u>Camarozonosporites</u> <u>cretaceous</u> (Weyland and Krieger) Potonié 1956.

Camarozonosporites ambigens (Fradkina) Playford, 1971 Pl. 2, Fig. 8

Synonymy:

- 1967 <u>Camarozonosporites insignis</u> Norris, p. 96-97, pl. 13, figs. 12-16.
- 1971 <u>Camarozonosporites insignis</u> Norris; Singh, p. 111, pl. 15, figs. 8-13.
- 1971 Camarozonosporites ambigens (Fradkina) Playford, p. 546, pl. 104, figs. 22, 23

- 1975 <u>Camarozonosporites ambigens</u> (Fradkina) Playford; Brideaux and McIntyre, p. 15, pl. 2, fig. 25.
- 1980 Camarozonosporites ambigens (Fradkina) Playford; Wingate, p. 10, pl. 1, fig. 10.

Remarks: This genus has interradial crassitudes and is readily distinguished from the genus <u>Hamulatisporis</u> which lacks interradial crassitudes (Farabee and Canright, 1986).

Genus <u>Hamulatisporis</u> Krutzsch emend. Srivastava, 1972 Type Species: <u>Hamulatisporis</u> <u>hamulatis</u> Krutzsch, 1959.

Hamulatisporis amplus Stanley, 1965

Pl. 2, Fig. 9

Synonymy:

- 1965 <u>Hamulatisporis amplus</u> Stanley, p. 242, pl. 29, figs. 1-6.
- 1969 <u>Hamulatisporis</u> <u>amplus</u> Stanley, Oltz, p. 119, pl. 39, fig. 18.
- 1974 <u>Hamulatisporis amplus</u> Stanley; McIntyre, pl. 14, figs. 24, 25.
- 1986 <u>Hamulatisporis amplus</u> Stanley; Farabee and Canright, p. 19, pl. 4, figs, 1, 2.

Age and Distribution: This species has been reported from the Maastrichtian of the Hell Creek Formation, South Dakota, U.S.A. (Stanley, 1965); early to mid Maastrichtian of the Horton River Section (McIntyre, 1974); Maastrichtian of the Lance Formation, Wyoming, U.S.A. (Farabee and Canright, 1986); and Maastrichtian of the Bearpaw Shale, east-central Montana, (Oltz, 1969).

Remarks: This species is distinguished from <u>H. hamulatis</u> by its distinctly larger size; (52µm to 83µm) as opposed to (24µm to 36µm) for <u>Hamulatisporis hamulatis</u> (Farabee and Canright, 1986). <u>Hamulatisporis amplus</u> is restricted to the Sermilik Formation and the Bylot Island Formation.

Hamulatisporis sp. cf. <u>H. loeblichii</u> Srivastava, 1972 Pl. 2, Fig. 10

Remarks: A single specimen found in the Bylot Island Formation at Twosnout Creek, is cluttered with organic debris; thus a positive identification is not possible. The finely punctate proximal face is not visible; the distal surface which is visible lacks the characteristic punctae. The specimen has an equatorial diameter of 101µm, thus, fitting the specified size range of <u>Hamulatisporis</u> <u>loeblichii</u>. <u>Hamulatisporis</u> <u>loeblichii</u> is the largest species of the genus <u>Hamulatisporis</u>. The largest known equatorial diameter of <u>Hamulatisporis</u> (83µm) was recorded by Farabee and Canright (1986). <u>Hamulatisporis</u> <u>loeblichii</u> has been reported from the Edmonton Formation

(Maastrichtian) of Alberta (Srivastava, 1972).

Genus <u>Gleicheniidites</u> Ross 1949 ex Delcourt and Sprumont

emend. Dettmann, 1963

Type Species: <u>Gleicheniidites senonicus</u> Ross, 1949 <u>Gleicheniidites senonicus</u> Ross, 1949

Pl. 2, Fig. 11

Synonymy:

- 1958 <u>Gleicheniidites senonicus</u> Ross; Couper, p. 138, pl. 19, figs. 13-15.
- 1964 <u>Gleicheniidites senonicus</u> Ross: Singh, p. 69, pl. 8, figs. 10, 11.
- 1965 <u>Gleicheniidites senonicus</u> Ross: McGregor, pl. 10, fig. 6.
- 1971 <u>Gleicheniidites senonicus</u> Ross; Rouse <u>et al.</u>, p. 218, pl. 1, figs. 1, 2.
- 1971 <u>Gleicheniidites</u> <u>senonicus</u> Ross; Singh, p. 97, pl. 14, fig. 1.
- 1973 <u>Gleicheniidites senonicus</u> Ross; Hopkins and Balkwill, p. 14, pl. 1, fig. 23.
- 1974 <u>Gleicheniidites senonicus</u> Ross; McIntyre, pl. 14, figs. 4, 5.
- 1986 <u>Gleicheniidites senonicus</u> Ross; Farabee and Canright, p. 19, pl. 3, figs. 5-7.
- 1986 <u>Gleicheniidites senonicus</u> Ross; Ricketts and Sweet, p. 20, pl. 1, fig. 21.

Remarks: This species occurs (2-30 specimens/sample) in every sample.

<u>Gleicheniidites</u> sp. cf. <u>G. circinidites</u> (Cookson) Dettmann, 1963 Pl. 2, Fig. 12

Synonymy:

- 1964 <u>Gleicheniidites</u> sp. cf. <u>G. circinidites</u> (Cookson) Dettmann; Singh, p. 39, pl.8, figs. 10, 11.
- 1971 <u>Gleicheniidites</u> sp. cf. <u>G. circinidites</u> (Cookson) Dettmann; Singh, p. 97, pl. 14, figs. 2, 3.
- 1982 <u>Gleicheniidites</u> sp. cf. <u>G. circinidites</u> (Cookson) Dettmann; Burden, p. 224, pl. 13, fig. 21.

Remarks: <u>Gleicheniidites</u> sp. cf. <u>G. circinidites</u> can be distinguished from <u>Gleicheniidites</u> <u>senonicus</u> by its more pointed apices and thicker interradial crassitudes (Singh, 1964; 1971). The species is restricted to the Byam Martin Formation, with the exception of one occurrence in the Bylot Island Formation at Twosnout Creek.

Genus <u>Asbeckiasporites</u> von der Brelie, 1964 Type Species: <u>Asbeckiasporites</u> <u>wirthi</u> von der Brelie, 1964.

Asbeckiasporites wirthi von der Brelie, 1964

Pl. 2, Fig. 13

Synonymy:

- 1963 Cinqulatisporites sp. B; Davis, p. 78, pl. 5, fig. 4.
- 1964 <u>Asbeckiasporites</u> <u>wirthi</u> von der Brelie, p. 142, pl. 8, figs. 7, 8; p. 142, pl. 9, figs. 1-6.
- 1971 Murospora truncata Singh, p. 128, pl. 18, figs. 1, 2.
- 1971 Asbeckiasporites wirthi von der Brelie; Playford,

p. 545, pl. 105, figs. 11, 12.

Remarks: This species is rare, a total of three specimens were observed from the Sermilik Formation (Lithotype 2) and the Bylot Island Formation at Twosnout Creek.

Genus <u>Ornamentifera</u> Bolkhovitina, 1966

Type Species: <u>Ornamentifera echinata</u> (Bolkhovitina) Bolkhovitina, 1966.

Ornamentifera baculata Singh, 1971

Pl. 2, Fig. 14

Synonymy:

- 1971 <u>Ornamentifera baculata</u> Singh, p. 100, pl. 14, figs. 4, 5.
- 1975 <u>Ornamentifera baculata</u> Singh; Brideaux and McIntyre, p. 15, pl. 2, fig. 29.

Remarks: This species is restricted to the Byam Martin Formation. Only two specimens were observed. Genus Tappanispora Srivastava, 1972

Type Species: Tappanispora loeblichii Srivastava, 1972

Tappanispora reticulata (Singh) Srivastava, 1975

Pl. 2, Fig. 15

Synonymy:

- 1971 <u>Tigrisporites reticulatus</u> Singh, p. 139, pl. 18, figs. 17, 18.
- 1975 <u>Tappanispora reticulata</u> (Singh) Srivastava, p. 64, pl. 29, figs. 7-9.
- 1981 <u>Tappanispora reticulata</u> (Singh) Srivastava; Srivastava, pl. 8, figs. 9, 10.
- 1982 <u>Tigrisporites reticulatus</u> Singh; Burden, p. 230, pl. 14, figs. 1-3.
- 1987 <u>Tappanispora</u> <u>reticulata</u> (Singh) Srivastava; Langille, p. 72, pl. 3, fig. 8.

Remarks: <u>Tappanispora</u> <u>reticulata</u> is restricted to the Bylot Island Formation, except for one isolated occurrence within the Byam Martin Formation.

Genus <u>Lycopodiacidites</u> Couper emend. Potonié, 1956 Type Species: <u>Lycopodiacidites bullerensis</u> Couper, 1953 Lycopodiacidites canaliculatus Singh, 1971

Pl. 2, Fig. 16

Synonymy:

1971 Lycopodiacidites canaliculatus Singh, p. 38, pl. 1,
fig. 15.

1987 Lycopodiacidites canaliculatus Singh; Langille, p. 66, pl. 2, fig. 13.

Remarks: This species is characteristic of the Byam Martin Formation. It is absent from the Sermilik Formation (Lithotype 1 and 2), and occurs very rarely (3 isolated occurrences) in the Bylot Island Formation.

Lycopodiacidites sp. 1

Pl. 2, Fig. 17

Description: Trilete; amb rounded to convexly triangular; proximal and distal surface ornamented by long narrow rugulae or ridges, separated by narrow grooves. The sculpture is composed of 0.5 to 1µm wide rugulae of uneven width and 0.5 to 1µm wide grooves. The exine has a smoothly orenulated surface and is 2µm thick.

Size Range: Equatorial diameter (12 specimens) 24 (28) 32µm.

Remarks: This species can be distinguished from <u>Hamulatisporis hamulatis</u> Krutzsch, by the presence of well developed sculpture on the proximal surface and the absence of tortuous muri. Lycopodiacidites <u>canaliculatus</u> and <u>Lycopodiacidites caperatus</u> Singh are distinct from this species, by their larger size; 36µm to 55µm for <u>Lycopodiacidites canaliculatus</u> and 48µm to 63µm for <u>Lycopodiacidites caperatus</u>.

Genus <u>Cicatricosisporites</u> Pflug and Thomson, 1953 Type Species: <u>Cicatricosisporites</u> dorogensis Potonié and Gelletich, 1933.

<u>Cicatricosisporites australiensis</u> (Cookson) Potonié, 1956 Pl. 2, Fig. 18

- 1963 <u>Cicatricosisporites australiensis</u> (Cookson) Potonié; Dettmann, p. 53, pl. 9, figs. 10-16.
- 1967 <u>Cicatricosisporites australiensis</u> (Cookson) Potonié; Norris, p. 92, pl. 11, fig. 14.
- 1971 <u>Cicatricosisporites</u> australiensis (Cookson) Potonié; Singh, p. 69, pl. 7, figs. 12-15.
- 1974 <u>Cicatricosisporites</u> <u>australiensis</u> (Cookson) Potonié; Hopkins, p. 15, pl. 3, fig. 32.
- 1975 <u>Cicatricosisporites</u> australiensis (Cookson) Potonié; Brideaux and McIntyre, p. 15, pl. 1, fig. 37.
- 1981 <u>Cicatricosisporites australiensis</u> (Cookson) Potonié; Herngreen and Khlonova, p. 483, pl. 9, fig. 2.
- 1982 <u>Cicatricosisporites australiensis</u> (Cookson) Potonié; Burden. p. 239, pl. 14, figs. 21, 22.
- 1986 Cicatricosisporites australiensis (Cookson) Potonié;

Ricketts and Sweet, p. 18, pl. 1, figs. 5, 6. 1987 <u>Cicatricosisporites australiensis</u> (Cookson) Potonié; Langille, pl. 3, fig. 1.

Remarks: <u>Cicatricosisporites australiensis</u> occurs sporadically in all sections. Nevertheless, there appears to be a conspicuous absence of this species from (Lithotype 1) of the Sermilik Formation and lowermost strata of the Bylot Island Formation; a total of 5 specimens were recorded. In contrast, the frequency of this species in the Byam Martin Formation, (Lithotype 2) of the Sermilik Formation and uppermost strata of the Bylot Island Formation is higher (1 to 9 specimens/sample) for most samples.

Cicatricosisporites sp. 1

Pl. 3, Fig. 1

Description: This species is characterized by its exceptionally large size (approximately 65-70 μ m in equatorial diameter) and large ribs (3.5 to 4 μ m wide) separated by (2 to 3 μ m wide) canals. Each interradial region on the proximal surface has between 6 and 8 ribs, which are parallel to the side and oblique to the laesurae. Ribs appear to terminate at the laesurae, however, the specimen is poorly preserved, and it is not possible to indicate whether the ribs continue around the apices. The distal surface has 12 to 13 subparallel ribs.

Remarks: The 3 specimens described here, resemble in part those described by Singh (1971) as <u>Cicatricosisportes</u> sp. cf. <u>C. potomacensis</u>. The species described by Singh (1971) differs, in possessing a greater number (17) of narrow (2m wide) distal ribs, with narrow canals (.5m wide).

This species was restricted to the Sermilik Formation (Lithotype 2) and the Bylot Island Formation at Twosnout Creek.

Genus Radialisporis Krutzsch, 1967

Type Species: <u>Radialisporis</u> radiatus (Krutzsch) Krutzsch, 1967

Radialisporis radiatus (Krutzsch) Krutzsch, 1967

Pl. 3, Fig. 2

- 1965 <u>Anemia radiatus</u> (Krutzsch) Stanley, p. 258, pl. 33, figs. 6, 7.
- 1972 <u>Radialisporis radiatus</u> (Krutzsch) Krutzsch; Rouse and Srivastava, p. 1177, pl.5, figs. 54, 55.
- 1972 <u>Radialisporis radiatus</u> (Krutzsch) Krutzsch; Srivastava, p. 29, pl. 24, figs. 8-13, pl. 25, figs. 1, 2.
- 1974 <u>Radialisporis radiatus</u> (Krutzsch) Krutzsch; McIntyre, p. 36, pl. 14, figs. 15, 16.

Age and Distribution: This species has been reported from the Edmonton Formation (Maastrichtian) of Alberta (Srivastava, 1972); the Bonnet Plume Formation (Paleocene) of the Yukon Territory (Rouse and Srivastava, 1972); the Horton River Section (late Campanian to mid Maastrichtian) of the North West Territories, Canada, (McIntyre, 1974).

Remarks: This species is found in the Sermilik Formation and the Bylot Island Formation.

Genus <u>Retitriletes</u> (van der Hammen, 1956 ex Pierce, 1961) emend. Doring, Krutzsch, Mai and Schultz, 1963 Type Species: <u>Retitriletes globosus</u> Pierce, 1961.

Retitriletes austroclavatides (Cookson) Doring, Krutzsch, Mai, and Schultz, 1963

Pl. 3, Fig. 3

- 1965 <u>Lycopodiumsporites austroclavatides</u> (Cookson) Potonië; McGregor, p. 24, pl. 7, fig. 11.
- 1966 <u>Lycopodiumsporites austroclavatides</u> Cookson; Burger, p. 247, pl. 15, fig. 2.
- 1971 <u>Lycopodiumsporites</u> <u>austroclavatides</u> (Cookson) Potonié; Singh, p. 40, pl. 2, fig. 1.
- 1971 <u>Lycopodiumsporites</u> austroclavatides (Cookson) Potonié; Playford, p. 537, pl. 103, fig. 2.
- 1972 Retitriletes austroclavatides (Cookson) Krutzsch;

Srivastava, p. 30, pl. 25, figs. 5-9, pl. 26, figs. 1-3.

- 1974 <u>Lycopodiumsporites austroclavatides</u> (Cookson) Potonié; Hopkins, p. 30, pl. 2, fig. 14.
- 1975 <u>Lycopodiumsporites austroclavatides</u> (Cookson) Potonié; Brideaux and McIntyre, p. 56, pl. 1, fig. 21.
- 1981 <u>Retitriletes austroclavatides</u> (Cookson) Krutzsch; Srivastava, p. 19, pl. 7, figs. 10, 11.

Remarks: Only one specimen of <u>Retitriletes</u> <u>austroclavatides</u> was observed in the study area. This specimen appears to be reworked due to its consistantly poor preservation.

Genus Foveotriletes Potonié, 1956

Type Species: <u>Foveotriletes</u> <u>scrobiculatus</u> (Ross) Potonié, 1956.

Foveotrilites subtriangularis Brenner, 1963

Pl. 3, Fig. 4

Sy ionymy:

- 1966 Foveotriletes subtriangularis Brenner; Burger, p. 246-247, pl. 14, fig. 1.
- 1971 <u>Foveotriletes</u> <u>subtriangularis</u> Brenner; Singh, p. 122, pl. 17, figs. 4, 5.
- 1980 Foveotriletes subtriangularis Brenner; Wingate, p. 25, pl. 10, fig. 1.
- 1982 Foveotriletes subtriangularis Brenner; Burden, p. 258,

pl. 17, figs. 13, 14.

Description: Specimens are convexly triangular, having an equatorial diameter of 42µm-45µm (5 specimens observed). The grains have a foveolate exine with evenly distributed round lumina.

Remarks: All grains are pitted and corroded, indicating that they are possibly reworked. This species was restricted to the Bylot Island Formation and the Sermilik Formation.

Genus Stereisporites Pflug, 1953

Type Species: <u>Stereisporites</u> <u>stereoides</u> (Potonié and Venitz) Pflug, 1953.

Stereisporites antiquasporites (Wilson and Webster)

Dettmann, 1963

Pl. 3, Fig. 5

- 1963 <u>Stereisporites</u> <u>antiquasporites</u> (Wilson and Webster) Dettmann, p. 25, pl. 1, figs. 20, 21.
- 1965 Sphagnum antiquasporites (Wilson and Webster) Stanley, p. 236, pl. 27, figs. 1-5.
- 1971 <u>Stereisporites</u> <u>antiquasporites</u> (Wilson and Webster) Dettmann; Singh, p. 33-34, pl. 1, figs. 4-5.
- 1974 Stereisporites antiquasporites (Wilson and Webster)

Dettmann; McIntyre, pl. 14, fig. 1.

- 1975 <u>Stereisporites antiquasporites</u> (Wilson and Webster) Dettmann; Brideaux and McIntyre, p. 14, pl. 1, fig. 6.
- 1982 <u>Stereisporites antiquasporites</u> (Wilson and Webster) Dettmann; Jarzen, pl. 1, fig. 2.
- 1983 <u>Stereisporites antiquasporites</u> (Wilson and Webster) Dettmann; Truswell, p. 142, pl. 1, figs. 1, 2.
- 1987 <u>Stereisporites antiquasporites</u> (Wilson and Webster) Dettmann; Langille, p. 75, pl. 3, fig. 15.

Remarks: This species was observed in all samples, in quantities greater than (5-9 specimens/sample) for most samples. <u>Stereisporites antiquasporites</u> was the most abundant trilete spore.

Stereisporites regium (Drozhastchich) Drugg, 1967

Pl. 3, Fig. 6

- 1965 Sphagnum regium Drozhastchich; Stanley, p. 238, pl.27, figs. 12-17.
- 1967 <u>Stereisporites regium</u> (Drozhastchich) Drugg, p. 37, pl. 6, fig. 20.
- 1974 <u>Stereisporites regium</u> (Drozhastchich) Drugg; McIntyre, p. 36, pl. 14, figs. 2, 3.
- 1978 <u>Stereisporites regium</u> (Drozhastchich) Drugg; Wilson, p. 109, pl. 1, fig. 4.

Genus <u>Cingulatisporites</u> Pflug emend. Potonié, 1956 Type Species: <u>Cingulatisporites</u> <u>levispeciosus</u> Pflug, 1953.

Cingulatisporites dakotaensis Stanley, 1965

Pl. 3, Fig. 7

Synonymy:

- 1965 <u>Cingulatisporites dakotaensis</u> Stanley, p. 243, pl. 30, figs. 1-8.
- 1969 <u>Cingulatisporites</u> <u>dakotaensis</u> Stanley; Norton and Hall, p. 15, pl. 2, fig. 6.
- 1969 <u>Cingulatisporites dakotaensis</u> Stanley; Oltz, p. 120, pl. 39, fig. 22.
- 1973 <u>Cingulatisporites dakotaensis</u> Stanley; Stone, p. 66, pl. 10, fig. 56.
- 1975 <u>Cingulatisporites dakotaensis</u> Stanley; Norris <u>et al.</u>, p. 349, pl. 3, fig. 2.
- 1986 <u>Cingulatisporites</u> dakotaensis Stanley; Farabee and Canright, p. 15, pl. 2, fig. 1.

Age and Distribution: This species has been reported from the Fort Union Formation (Maastrichtian) of South Dakota, (Stanley, 1965); the Tullock Member of the Fort Union Group (Paleocene), Montana, (Oltz, 1969); the Almond Formation (late Campanian) of Wyoming (Stone, 1973); and the Lance Formation (Maastrichtian) of Wyoming (Farabee and Canright, 1986).

Remarks: The characteristic "Y " - shaped polar thickening is preserved on the distal surface. This species is differentiated from <u>C. radiatus</u> which has a homogeneous and transparent cingulum that lacks radiating striations (Stanley, 1965). Three poorly preserved specimens were recovered; one from the Byam Martin Formation, one from the Bylot Island Formation at Twosnout Creek, and the other from Sormalik Formation (Lithotype 1).

Genus Deltoidospora Miner, 1935

Type Species: Deltoidospora hallii Miner, 1935.

Deltoidospora hallii Miner, 1935

Pl. 3, Fig. 8

- 1964 <u>Deltoidospora hallii</u> Miner; Singh, p. 80, pl. 9, figs. 13, 14.
- 1971 <u>Deltoidospora hallii</u> Miner; Singh, p. 118, pl. 16, fig. 8.
- 1980 <u>Deltoidospora hallii</u> Miner; Wingate, p. 24, pl. 9, fig. 14.
- 1982 <u>Deltoidospora hallii</u> Miner; Burden, p. 261, pl. 17, fig. 23.
- 1986 Deltoidospora hallii Miner; Boland, p. 106, pl. 1,

fig. 11.

1987 <u>Deltoidospora hallii</u> Miner; Langille, p. 76, pl. 3, fig. 16.

Remarks: This species is sporadic in distribution and rare in abundance (1-3 specimens/sample).

Genus Cibotiumspora Chang, 1965

Type species: <u>Cibotiumspora parodoxa</u> (Malyavkina) Chang, 1965.

Cibotiumspora juncta (Kara-Murza) Singh, 1983

Pl. 3, Fig. 9

- 1964 <u>Concavisporites jurienensis</u> Balme; Balme, p. 77, pl. 6, fig. 2.
- 1965 Deltoidospora juncta (Kara-Murza) Singh; McGregor, p. 22, pl. (fig. 4.
- 1966 Concavisporites jurienensis Balme; Burger, p. 237, pl. 4, fig. 6.
- 1971 <u>Deltoidospora juncta</u> (Kara-Murza) Singh; Singh, p. 119-120, pl. 16, figs. 10, 11.
- 1975 <u>Cibotiumspora jurienensis</u> (Balme) Filatoff, p. 61, pl. 10, figs. 8-13.
- 1980 <u>Concavisporites</u> jurienensis (Balme) Wingate, p. 25, pl. 9, fig. 12.
- 1982 Concavisporites jurienensis Balme; Burden, p. 267,

pl. 18, figs. 7-10.

1983 <u>Cibotiumspora sinuata</u> (Couper) Fensome, p. 403-409, pl. 15, figs. 7-14, 16, 18.

Remarks: Fensome (1983) spent considerable time sorting out systematic problems and emending the genus <u>Cibotiumspora</u>. His unpublished study (Fensome, 1983), suggested that this taxon should be transferred to the species designation <u>sinuata</u>. The basis for this transfer was his recognition of a highly variable morphology that had lead to well over 30 distinct names or combinations to the genus and species level.

Genus Cyathidites Couper, 1953

Type Species: Cyathidites australis Couper, 1953.

Cyathidites minor Couper, 1953

Pl. 3, Fig. 10

Synonymy:

- 1958 <u>Cyathidites minor</u> Couper; Couper, p. 139, pl. 20, figs. 9, 10.
- 1963 <u>Cyathidites minor</u> Couper; Dettmann, p. 22, pl. 1, figs. 4, 5.

1963 <u>Cyathidites minor</u> Couper; Davis, p. 39, pl. 1, fig. 2.
1964 <u>Cyathidites minor</u> Couper; Singh, p. 71, pl. 8, fig. 13.
1965 <u>Cyathidites minor</u> Couper; McGregor, pl. 7, figs. 3, 4.
1967 <u>Cyathidites minor</u> Couper; Srivastava, pl. 1, fig. M.

- 1971 <u>Cyathidites minor</u> Couper; Singh, p. 101, pl. 14, fig. 9.
- 1973 <u>Cyathidites minor</u> Couper; B.D. Tschudy, p. 6, pl. 1, fig. 1.
- 1974 Cyathidites minor Couper; McIntyre, pl. 14, fig. 8.
- 1975 <u>Gyathidites minor</u> Couper; Brideaux and McIntyre, p. 14, pl. 1, fig. 3.
- 1987 <u>Cyathidites minor</u> Couper; Langille, p. 77, pl. 4, fig. 1.

Remarks: This species occurs in all sections.

Genus Gemmatriletes Pierce, 1961

Type Species: Gemmatriletes morulus Pierce, 1961.

Gemmatriletes clavatus Brenner, 1968

Pl. 3, Fig. 11

Synonymy:

- 1968 Gemmatriletes clavatus Brenner, p. 353, pl. 1, fig. 7.
- 1983 <u>Gemmatriletes</u> <u>clavatus</u> Brenner; Singh, p. 30, pl. 1, figs. 2, 3.
- 1988 <u>Gemmatriletes</u> clavatus Brenner; Sweet and McIntyre, fig. 6, No. 11.

Age and Distribution: This species is reported from Albian to Cenomanian strata of northeastern Peru (Brenner, 1968); Cenomanian strata of Alberta (Singh, 1983); and late Turonian strata of the Alberta Foothills (Sweet and McIntyre, 1988).

Remarks: <u>Gemmatriletes clavatus</u> is distinguished from <u>Gemmatriletes densegemmatus</u> Brenner, by its smaller size (40 (46) 54 µm), and more videly spaced gemmae .Singh, 1983). A single specimen was recovered from the Bylot Island Formation.

Genus Osmundacidites Couper, 1953

Type Species: <u>Osmundacidites</u> wellmanii</u> Couper, 1953 <u>Osmundacidites wellmanii</u> Couper, 1953

Pl. 3, Fig. 12

- 1958 Osmundacidites wellmanii Couper; Couper, p. 1¹, pl. 16, figs. 4, 5.
- 1963 <u>Osmundacidites wellmanii</u> Couper; Dettmann, p. 32, pl. 3, figs. 19-21.
- 1963 <u>Osmundacidites wellmanii</u> Couper; Davis, p. 51, pl. 2, fig. 7.
- 1964 <u>Osmundacidites wellmanii</u> Couper; Singh, p. 44, pl. 1, fig. 20.
- 1965 <u>Osmundacidites wellmanii</u> Couper; McGregor, pl. 2, fig. 12.
- 1966 <u>Osmundacidites wellmanii</u> Couper; Burger, p. 251, pl. 20, fig. 3.

- 1967 <u>Osmundacidites wellmanii</u> Couper; Norris, p. 88, pl. 10, fig. 14.
- 1971 <u>Osmundacidites wellmanii</u> Couper; Singh, p. 251, pl. 20, f⁴g. 3.
- 1972 <u>Osmundacidites wellmanii</u> Couper; Srivastava, p. 27, pl. 23, figs. 1-3.
- 1974 <u>Osmundacidites wellmanii</u> Couper; McIntyre, p. 36, pl. 14, figs. 10, 11.
- 1974 <u>Osmundacidites wellmanii</u> Couper; Hopkins, p. 13, pl. 2, fig. 24.
- 1982 <u>Osmundacidites wellmanii</u> Couper; Burden, p. 268, pl. 18, figs. 14, 15.
- 198" <u>Osmundacidites wellm/nii</u> Couper; Langille, p. 80, pl. 4, fig. 6.

Remarks: This species occurs in the Byam Martin Pormation, the Bylot Islard Formation and Lithotype 3 of the Sermilik Formation.

Genus <u>Baculatisporites</u> Thomson and Pflug, 1953 Type Species: <u>Baculatisporites</u> primarius (Wolff) Thomson and Pflug, 1953.

Baculatisporites comaumensis (Cookson) Potonié, 1956

Pl. 3, Fig. 13

Synonymy:

1963 Baculatisporites comaumensis (Cookson) Potonie;

Dettmann, p. 35, pl. 3, figs. 22, 23.

- 1965 <u>Baculatisporites</u> <u>comaumensis</u> (Cookson) Potonié; McGregor, pl. 7, fig. 10.
- 1971 <u>Baculatisporites comaumensis</u> (Cookson) Potonié; Hopkins, pl. 20, figs. 10, 11.
- 1971 <u>Baculatisporites</u> <u>comaumensis</u> (Cookson) Potonié; Singh, p. 48, pl. 3, fig. 14.
- 1974 <u>Baculatisporites comaumensis</u> (Cookson) Potonié; McIntyre, p. 36, pl. 14, fig. 9.
- 1975 <u>Baculatisporites comaumensis</u> (Cookson) Potonié; Brideaux and McIntyre, p. 56, pl. 1, fig. 12.
- 1983 <u>Baculatisporites comaumensis</u> (Cookson) Potonié; Truswell, p. 141, pl. 1, figs. 3, 4.
- 1987 <u>Baculatisporites comaumensis</u> (Cookson) Potonié; Langille, p. 80-81, pl. 4, fig. 7.

Remarks: This species is restricted to the Bylot Island and Byam Martin formations.

Monoleta Spores

Genus Laevigatosporites Ibrahim emend. Schopf,

Wilson and Bentall, 1944

Type Species: Laevigatosporites vulgarig (Ibrahim) Ibrahim, 1933.

Laevigatosporites haardti (Potonié and Venitz)

Thomson and Pflug, 1953

Pl. 3, Fig. 14

Synonymy:

- 1965 <u>Laevigatosporites haardti</u> (Potonié and Venitz) Thomson and Pflug; Stanley, p. 252, pl. 32, figs. 1-3.
- 1972 <u>Laevigatosporites haardti</u> (Potonié and Venitz) Thomson Pflug; Srivastava, p. 232, pl. 5, figs. 9, 10.
- 1978 <u>Laevigatosporites haardti</u> (Potonie and Venitz) Thomson and Pflug; Wilson, p. 117-118, pl. 4, figs. 7, 8.
- 1981 <u>Laevigatosporites haardti</u> (Potonié and Venitz) Thomson and Pflug; Srivastava, p. 8, pl. 7, fig. 4.
- 1986 <u>Laevigatosporites haardti</u> (Potonié and Venitz) Thomson and Pflug; Boland, p. 135, pl. 3, fig. 4.

Description: Grains assigned to this species are distinct)y bean shaped, thick walled (1-2µm), psilate, and range from 30-70µm in diameter.

Remarks: Wilson (1978) indicated that any distinction between Laevigatosporites haardtii and Laevigatosporites ovatus Wilson and Webster was purely artificial. Srivastava (1972) also grouped these two species, stating that similar morphologies, shapes, and size ranges exist. Waterfield (personal communication, 1989) noted a morphological gradation, and both species were categorized as Lasvigatosporites haardti. This study has two distinct end members; no gradation was noted.

```
Laevigatosporites sp. cf. <u>L. ovatus</u> Wilson and Webster,
1946
Pl. 3, Fig. 15
```

Description: Specimens have a faint monolete mark preserved on the proximal surface. The exine is translucent, thin (<1µm), psilata; and the length of the polar axis ranges between 24-28µm.

Remarks: Differential diagnosis of <u>Laevigatosporites</u> sp. cf. <u>ovatus</u> from <u>Laevigatosporites</u> <u>ovatus</u> is based on a faint monolete mark and a slightly thinner exine. All other characteristics compare favourably.

Genus <u>Hazaria</u> Srivastava, 1971 Type Species: <u>Hazaria sheopiarii</u> Srivastava, 1971 <u>Hazaria sheopiarii</u> Srivastava, 1971

Pl. 3, Fig. 16

Synonymy:

1971 <u>Hazaria sheopiarii</u> Srivastava, p. 258, pl. 2, figs. 1-4.

1972 <u>Hazaria sheopiarii</u> Srivastava; Rouse and Srivastava, p. 1172, pl. 2, fiqs. 15, 16.

- 1974 <u>Hazaria sheopiarii</u> Srivastava; McIntyre, p. 37, pl. 14, fiqs. 26, 27.
- 1986 <u>Hazaria sheopiarii</u> Srivastava; Jerzykiewicz and Sweet, p. 1364, pl. 1, fig. 4.

Age and Distribution: This species is reported from the Edmonton Formation (Maastrichtian) of Alberta (Srivastava, 1971); the Bonnet Plume Formation (Maastrichtian) of the Yukon Territory (Rouse and Srivastava, 1972); the Horton River section (late Campanian to late Maastrichtian) of the Northwest Territories (McIntyre, 1974); the Coalspur Formation (Maastrichtian to Paleocene) of the Alberta Foothills (Jerzykiewicz and Sweet, 1986).

Remarks: The minimum size of this taxon, according to Srivastava (1971) is 40µm. Specimens in this study were seen to be as small as 37µm. This species is present in all sections. It is most frequently observed (1-3 specimens/sample) in Lithotype 3 of the Sermilik Formation.

Inaperturate Pollen

Genus <u>Taxodiaceaepollenites</u> Kremp, 1949 ex Potonié, 1958 Type Species: <u>Taxodiaceaepollenites</u> <u>hiatus</u> Potonié ex Potonié, 1958. Taxodiaceaepollenites hiatus (Potonié) Kremp, 1949

Pl. 3, Fig. 17

Synonymy:

- 1965 <u>Thuja</u> ? <u>hiatus</u> (Potonié) Stanley, p. 273, pl. 38, figs. 1-3.
- 1967 <u>Taxodiaceaepollenites hiatus</u> (Potonié) Kremp; Srivastava, pl. 2, fig. C.
- 1971 <u>Taxodiaceaepollenites hiatus</u> (Potonié) Kremp; Singh, p. 158, pl. 22, fig. 7.
- 1975 <u>Taxodiaceaepollenites hiatus</u> (Potonié) Kremp; Brideaux and McIntyre, p. 17, pl. 4, fig. 19.
- 1980 <u>Taxodiaceaepollenites hiatus</u> (Potonié) Kremp; Wingate, p. 37, pl. 13, fig. 15.
- 1986 <u>Taxodiaceaepollenites</u> <u>hiatus</u> (Potonié) Kremp; Farabee and Canright, p. 35, pl. 10, fig. 2.
- 1987 Taxodiaceaepollenites hiatus (Potonié) Kremp; Langille, p. 85, pl. 4, fig. 14.

Remarks: <u>Taxodiaceaepollenites hiatus</u> is present (2-15 specimens/sample) in every sample.

Taxodiaceaepollenites vacuipites (Wodehouse) Wingate, 1980 Pl. 3, Fig. 18

Synonymy:

1980 Taxodiaceaepollenites vacuipites (Wodehouse) Wingate, p. 37, pl. 13, fig. 16. Remarks: <u>Taxodiaceaepollenites vacuipites</u> is distinguished from <u>Taxodiaceaepollenites hiatus</u> by its larger size, fusiform shape, and exinal folds adjacent to the area of dehiscence (Wingate, 1980).

Bisaccate Pollen

Genus <u>Podocarpidites</u> Cookson, 1947, ex Couper 1953 Type Species: <u>Podocarpidites</u> ellipticus Cookson, 1947.

Podocarpidites multesimus (Bolkovitina) Pocock, 1962

Pl. 4, Fig. 1

- 1964 <u>Podocarpidites multesimus</u> (Bolkhovitina) Pocock; Singh, p. 116, pl. 15, figs. 12, 13.
- 1971 <u>Podocarpidites multesimus</u> (Bolkhovitina) Pocock; Singh, p. 166, pl. 24, fig. 2.
- 1973 <u>Podocarpidites multesimus</u> (Bolkhovitina) Pocock; B.D. Tschudy, p. 16, pl. 7, figs. 1, 2.
- 1975 <u>Podocarpidites multesimus</u> (Bolkhovitina) Pocock; Brideaux and McIntyre, p. 16, pl. 4, figs. 3, 4.
- 1980 <u>Podocarpidites multesimus</u> (Bolkhovitina) Pocock; Wingate, p. 38, pl. 14, figs. 7, 8.
- 1982 <u>Podocarpidites multesimus</u> (Bolkhovitina) Pocock; Burden, p. 314, pl. 25, fig. 16.

Remarks: This species was observed only rarely (2 to 4 specimens / sample) at Twosnout Creek and along the south coast.

Podocarpidites sp. 1

Pl. 4, Fig. 2

Description: Bisaccate; The central body is granulate; the bladders are coarsely reticulate (1µm to 2µm wide). The bladders have a length / breadth ratio of 1.85 / 1. The bladders appear to be distally pendant, however, the poor preservation of the grain makes this remark tenuous.

Size Range: Total breadth of the grain 140 μm Breadth of the central body 60 μm Breadth of the bladders 40 μm Length of the bladders 74 μm

Remarks: This grain is noted for its exceptionally large size. The single grain observed is torn and distorted and was recovered from the top of the Sermilik Formation (Lithotype 2).

Genus <u>Pityosporites</u> Seward emend. Manum, 1960 Type Species: <u>Pityosporites</u> <u>antarcticus</u> Seward, 1914. <u>Pityosporites</u> <u>constrictus</u> Singh, 1964 Synonymy:

- 1964 <u>Pityosporites constrictus</u> Singh, p. 122, pl. 16, figs. 8, 9.
- 1966 <u>Pityosporites constrictus</u> Singh; Srivastava, p. 524, pl. 6, fig. 2.
- 1967 <u>Pityosporites constrictus</u> Singh; Srivastava, p. 140, pl. 2, fig. M.
- 1971 <u>Pityosporites constrictus</u> Singh; Singh, p. 167, pl. 25, fig. 10.
- 1973 <u>Pityosporites</u> sp. cf <u>P. constrictus</u> Singh; B.D. Tschudy, p. 15, pl. 5, fig. 9.
- 1980 <u>Pinuspollenites constrictus</u> Singh; Wingate, p. 40, pl. 15, figs. 6, 7.
- 1982 Pityosporites constrictus Singh; Burden, p. 317, pl. 15, figs. 2, 3.

Remarks: Specimens of this species were exceptionally well preserved. The species occurred in Lithotype 1 of the Sermilik Formation and the Bylot Island Formation

<u>Pityosporites alatipollenites</u> (Rouse) Singh, 1964 Pl. 4, Fig. 4

Synonymy:

1964 Pityosporites alatipollenites (Rouse) Singh, p. 123, pl. 16, fig. 10.

- 1971 Pityosporites alatipollenites (Rouse) Singh; Singh, p. 173, pl. 25, fig. 9.
- 1973 <u>Pitvosporites alatipollenites</u> (Rouse) Singh; B.D. Tschudy, p. 15, pl. 6, fig. 1.
- 1982 <u>Pitvosporites</u> <u>alatipollenites</u> (Rouse) Singh; Burden, p. 315, pl. 26, fig. 1.
- 1987 <u>Pityosporites</u> <u>alatipollenites</u> (Rouse) Singh; Langille, p. 89, pl. 5. fig. 5.

Remarks: Specimens of <u>Pityosporites alatipollenites</u> were poorly preserved. They are restricted to the Bylot Island and Sermilik formations.

Pitvosporites sp. 1

Pl. 4, Fig. 5

Description: Bisaccate; bladders distally pendant and moderately constricted towards the base. The bladders exhibit a fine to coarse reticulum (.5µm to 1µm wide). The central body is finely granulate with a 1-2µm thick granulose proximal cap.

Size Range: Proximal cap 1-2µm thick Total breadth of the grain 50-90µm Height of the central body 16-20µm Breadth of the central body 32-50µm Breadth of the bladders 24-40µm Width of the distal furrow 10-14µm (51 grains examined)

Remarks: A concise description was hampered by poor preservation and a lack of specimens in polar view. This species was frequently observed in south coast sections (1-5 specimens/sample) for most samples; it is less common (1-2 specimens/sample) at Twosnout Creek.

Pityosporites sp. 2

Pl. 4, Fig. 6

Description: Bisaccate; bladders are circular in outline, coarsely reticulate and widely separated from one another. The central body is sub-circular with a width greater than its length. The grain exhibits a wide distal furrow.

Size Range: Total breadth of the grain 64-67µm Breadth of the central body 35-36µm Length of the central body 28-30µm Breadth of the bladders 23-25µm Length of the bladders 26-31µm Width of the distal furrow 16-18µm (37 specimens examined) Remarks: This species occurs in most samples along the south coast. The distribution at Twosnout Creek is sporad c.

Pityosporites sp. 3

Pl. 4, Fig. 7

Description: Bisaccate; bladders are distally pendant, much shorter than the central body and coarsely reticulate (2 μ m wide). In lateral view the bladders and the central body are in the same sweeping curve. The proximal cap is $\leq 1 \mu$ m thick, and the central body is finely rugulate (<.5 μ m wide).

Size Range: Total breadth of the central body 66-72µm Breadth of the central body 40-48µm Height of the central body 30-40µm Breadth of the bladders 14-20µm Width of the distal furrow 5-6µm (5 specimens examined)

Remarks: This species appears most like the Eocene bisaccate <u>Pinug coloradensis</u> Ting (pl. 2, figure 11). This species was restricted to the Sermilik Formation and the Bylot Island Formation. Genus <u>Alisporites</u> Daugherty emend. Jansonius, 1971 Type Species: <u>Alisporites</u> <u>opii</u> Daugherty, 1941.

Alisporites bilateralis Rouse, 1959

Pl. 4, Fig. 8

Synonymy:

Parana a san

i

- 1962 <u>Alisporites thomasii</u> (Couper) Pocock, p. 62, pl. 14, fig. 143.
- 1964 <u>Alisporites thomsaii</u> (Couper) Pocock; Singh, p. 109, pl. 14, figs. 11, 12.
- 1965 <u>Alisporites thomasii</u> (Couper) Nilsson; McGregor, p. 24, pl. 7, fig. 39.
- 1966 <u>Alisporites thomasii</u> (Couper) Pocock; Burger, p. 259, pl. 35, fig. 2.
- 1971 <u>Alisporites bilateralis</u> Rouse; Singh, p. 169, pl. 24, fig. 9.
- 1975 <u>Alisporites bilateralis</u> Rouse; Brideaux and McIntyre, p. 16, pl. 3, fig. 32.
- 1976 <u>Alisporites bilateralis</u> Rouse; Scott, p. 592, pl. 9, fig. 5.
- 1987 <u>Alisporites bilateralis</u> Rouse; Langille, p. 91, pl. 5, fig. 8.

Remarks: This species was observed in all sections.

Alisporites grandis (Cookson) Dettmann, 1963

Pl. 4, Fig. 9

Synonymy:

- 1957 <u>Alisporites rotundus</u> Rouse, p. 371, pl. 1, figs. 15, 16.
- 1962 <u>Alisporites rotundus</u> Rouse; Pocock, p. 61-62, pl. 9, figs. 140, 141.
- 1964 <u>Alisporites rotundus</u> Rouse; Singh, pl. 14, figs. 13, 14; p. 110, pl. 15, figs. 1, 2.
- 1967 <u>Alisporites grandis</u> (Cookson) Dettmann; Srivastava, p. 140, fig. pl. 2, fig. K.
- 1971 <u>Alisporites grandis</u> (Cookson) Dettmann; Singh, p. 170, pl. 25, figs. 1,2.
- 1973 <u>Alisporites grandis</u> (Cookson) Dettmann; B.D. Tschudy, p. 14, pl. 5, figs. 2, 3.
- 1980 <u>Alisporites grandis</u> (Cookson) Dettmann; Wingate, p. 39, pl. 15, fig. 1.
- 1982 <u>Alisporites grandis</u> (Cookson) Dettmann; Burden, p. 318, pl. 26, fig. 7.

Remarks: <u>Alisporites</u> grandis was observed in all formations.

Genus Cedripites Wodehouse, 1933

Type Species: <u>Cedripites eocenicus</u> Wodehouse, 1933 <u>Cedripites canadensis</u> Pocock, 1962

Pl. 4, Fig. 10

- 1964 Cedripites canadensis Pocock; Singh, p. 112, pl. 15, fig. 6.
- 1967 <u>Cedripites canadensis</u> Pocock; Norris, p. 102, pl. 15, figs. 149, 150.
- 1971 <u>Cedripites canadensis</u> Pocock; Singh, p. 171, pl. 25, figs. 4, 5.
- 1975 <u>Cedripites canadensis</u> Pocock; Brideaux and McIntyre, p. 16, pl. 3, fig. 37.
- 1982 <u>Cedripites canadensis</u> Pocock; Burden, p. 325, pl. 27, figs. 1, 2.
- 1987 <u>Cedripites canadensis</u> Pocock; Langille, p. 93, pl. 5, fig. 12.

Remarks: According to Singh (1964) the size for the " total breadth of the grain " ranges between 60 µm and 90 µm. Specimens in this study reach a maximum size of 100 µm. <u>Cadripites canadensis</u> can be distinguished from <u>Cadripites</u> <u>creataceous</u> Pocock by a notable difference in the thickness of the proximal cap (Singh, 1964).

<u>Cedripites</u> <u>canadensis</u> is the most abundant bisaccate taxon at Twosnout Creek, with as many as 10-14 specimens/sample. In south coast sections <u>Cedripites</u> <u>canadensis</u> is as abundant, although other bisaccate grains frequently occur. Genus <u>Pristinuspollenites</u> B.D. Tschudy, 1973 Type Species: <u>Pristinuspollenites microsaccus</u> (Couper) B.D. Tschudy, 1973.

Pristinuspollenites sp. cf. <u>P. inchoatus</u> (Pierce) B.D. Tschudy, 1973

Pl. 4, Fig. 11

Remarks: Except for the shape of the bladders, this taxon resembles <u>Fristinuspollenites</u> inchoatus. Singh (1964) indicates that the bladders are elongate with a length to width ratio of 2:1. The one specimen recorded in this study has bladders that are crescent shaped, with a length to width ratio of 1:1.

Size Range: Total breadth of the grain 50µm Length of the central body 43µm Exine thickness 2µm Breadth of the bladders 20µm Length of the bladders 20µm

Monosulcate Pollen

Genus <u>Liliipollis</u> Krutzsch, 1970 Type Species: <u>Liliipollis lilioides</u> Krutzsch, 1970. <u>Liliipollis</u> sp. 1 ? Pl. 4. Fig. 12

88

Description: Monosulcate pollen with two-layered exine. Nexine is $\leq 1\mu$ m thick; sexine with baculae less than 1μ m high. Baculae cover the entire surface and are arranged in a crotonid reticulate pattern. The sulcus extends the entire length of the grain.

Size Range: Length x Width: 38 (42) 44 x 22 (24) 25 μ m, (five measured specimens)

Remarks: The morphological description of specimens in this study resemble in detail those of Farabee and Canright (1986). However, the described specimeens are on average 10µm larger than those described by Farabee and Canright (1986).

In south coast sections, this species is restricted to Lithotype 1 of the Sermilik Formation, and a tongue of Bylot Island Formation strata (Section D, Appendix λ_i). At Twosnout Creek, this species is confined to samples occuring around the Byam Martin - Bylot Island Formation contact.

Genus <u>Liliacidites</u> Couper, 1953 Type Species: <u>Liliacidites kaitangataensis</u> Couper, 1953. <u>Liliacidites variegatus</u> Couper, 1953

Pl. 4, Fig. 13

Synonymy:

1957 Liliacidites variegatus Couper; Rouse, p. 368, pl. II,

figs. 29, 30.

- 1966 Liliacidites variegatus Couper; Srivastava, p. 25, pl. 4, figs. 15, 16.
- 1967 Liliacidites variegatus Couper; Drugg, p. 46-47, pl. 7, fig. 23.
- 1968 <u>Liliacidites variegatus</u> Couper; Elsik, p. 312, pl. 15, fig. 1.
- 1973 <u>Liliacidites variegatus</u> Couper; B.D. Tschudy, p. 20, pl. 8, fig. 12.
- 1980 <u>Liliacidites variegatus</u> Couper; Wingate, p. 43, pl. 16, fig. 11.
- 1986 <u>Liliacidites variegatus</u> Couper; Farabee and Canright, p. 53, pl. 19, fig. 5.

Description: Reticulate, monosulcate pollen. The sulcus extends almost the entire length of the grain, and is narrow. The reticulum is coarse (about 2µm wide), and the muri are simplibaculate.

Size Range: Length x breadth (39.6 x 24.2) μm Length / breadth ratio 1.6 (3 specimens examined)

Age and Distribution: Early to middle Oligocene of New Zealand (Couper, 1953); late Campanian of North Central Montana (B.D. Tschudy, 1973); Albian of Oklahoma (Wingate, and supplicity and supplicity and

1980); Maastrichtian of Wyoming (Farabee and Canright, 1986).

Remarks: This species is rare, a total of three specimens were observed from the Sermilik Formation (Lithotype 1), the Bylot Island Formation and the Byam Martin Formation.

Genus Cycadopites Wodehouse, 1933, ex Wilson and

Webster, 1946

Type Species: <u>Cycadopites</u> <u>follicularis</u> Wilson and Webster, 1946.

Cycadopites fragilis Singh, 1964

Pl. 4, Fig. 14

Synonymy:

- 1962 <u>Monosulcites minimus</u> Cookson; Pocock, p. 77, pl. 13, figs. 206-208.
- 1964 Cycadopites fragilis Singh, p. 103, pl. 14, fig. 2.
- 1965 Cycadopites fragilis Singh; McGregor, p. 24, pl. 7, fig. 25.
- 1980 <u>Cycadopites fragilis</u> Singh; Wingate, p. 36, pl. 13, fig. 9.

Remarks: A single specimen was found in the Sermilik Formation (Lithotype 1). Genus <u>Monosulcites</u> Cookson ex Couper, 1953 Type Species: <u>Monosulcites</u> <u>minimus</u> Couper, 1953.

> Monosulcites sp. 1 Pl. 4, Fig. 15

Description: Monosulcate; sulcus running entire length of grain, and narrowing at the longitudinal ends. The exine is psilate and less than lµm thick.

Size Range: Length 60µm Width 18µm Length to Breadth ratio 3 / 1

Remarks: Two specimens were recovered; one from the Sermilik Formation (Lithotype 1), and the other from Bylot Island Formation strata located along the south coast (Section D, Appendix λ_i).

Genus <u>Entylissa</u> Naumova, 1939, ex Ishchenko, 1952 Type Species: <u>Entylissa</u> <u>caperatus</u> (Luber) Potonié and Kremp, 1954.

Entylissa nitidus Balme, 1957

Pl. 5, Fig. 1

Synonymy:

1962 Ginkgocycadophytus nitidus (Balme) de Jersey, p. 12, pl. 5, figs. 1-3.

- 1963 <u>Ginkgocycadophytus nitidus</u> (Balme) de Jersey; Dettmann, p. 104, pl. 26, figs. 8, 9.
- 1971 <u>Ginkgocycadophytus nitidus</u> (Balme) de Jersey; Singh, p. 155, pl. 22, fig. 3.
- 1980 <u>Ginkgocycadophytus nitidus</u> (Balme) de Jersey; Wingate, p. 36, pl. 13, fig. 12.

1982 <u>Entylissa nitidus</u> Balme; Burden, p. 340, pl. 30, figs. 4, 5.

Remarks: This species is sporadic in distribution, but occurs in all formations.

Tricolpate Pollen

Genus <u>Tricolpites</u> Cookson ex Couper 1953 emend. Potonié 1960 Type Species: <u>Tricolpites reticulatus</u> Cookson, 1947. <u>Tricolpites</u> sp. 1 Pl. 5, Fig. 2

Description: Prolate tricolpate pollen; characterized by 4µm high club shaped pila. The body is granular, with a length / width ratio of 1.4 to 1.

Remarks: A single specimen was found in the Sermilik Formation (Lithotype 2). Genus Rousea Srivastava, 1969

Type Species: Rousea subtillis Srivastava, 1969.

Rousea georgensis (Brenner) Dettmann, 1973

PL. 5, Fig. 3

Synonymy:

- 1968 <u>Retitricolpites georgensis</u> Brenner; Hedlund and Norris, p. 145, pl. 6, figs. 1, 3.
- 1971 <u>Retitricolpites georgensis</u> Brenner; Singh, p. 200, pl. 30, figs. 1-6.
- 1971 <u>Retitricolpites georgensis</u> Brenner; Playford, p. 561, pl. 107, figs. 11, 12.
- 1973 <u>Retitricolpites georgensis</u> (Brenner) Burger, p. 7, pl. 3, figs. 2-4.
- 1975 <u>Retitricolpites georgensis</u> Brenner; Brideaux and McIntyre; p. 17, pl. 4, figs. 31, 32.
- 1975 <u>Rousea georgensis</u> (Brenner) Dettmann; Srivastava, p. 92, pl. 43, figs. 10, 11.
- 1980 <u>Tricolpites</u> sp. cf. <u>Retitricolpites</u> <u>georgensis</u> Brenner; Wingate, p. 45, pl. 17, figs. 3, 4.
- 1981 <u>Rousea georgensis</u> (Brenner) Dettmann; Srivastava, p. 27, pl. 11, fig. 3.
- 1988 Rousea georgensis (Brenner) Dettmann; Sweet and McIntyre, p. 506, fig. 7, No. 15, 16.

Remarks: This species is characterized by the reduction in size of the lumina towards the poles and the pilate
structure of the muri (Singh, 1971). This species occurs in the Sermilik and Bylot Island Formations.

Genus <u>Cupuliferoidaepollenites</u> Potonié, Thomson & Thiergart 1950 ex Potonié, 1960

Type Species: <u>Cupuliferoidaepollenites liblarensis</u> (Thomson) Potonié, 1960.

Cupuliferoidaepollenites parvulus (Groot and Penny)

Dettmann, 1973

Pl. 5, Fig. 4

Synonymy:

- 1960 <u>Tricolpopollenites parvulus</u> Groot and Penny, p. 232, pl. 2, figs. 8, 9.
- 1961 <u>Tricolpopollenites</u> parvulus Groot and Penny; Groot <u>et</u> <u>al.</u>, p. 132, pl. 26, figs. 3, 4.
- 1967 <u>Psilatricolpites parvulus</u> (Groot and Penny) Norris, p. 107, pl. 17, figs. 5-7.
- 1971 <u>Psilatricolpites parvulus</u> (Groot and Penny) Norris; Singh, p. 198, pl. 29, figs. 15, 16.
- 1973 <u>Cupuliferoidaepollenites</u> sp. cf. <u>C. parvulus</u> (Groot and Penny) Dettmann, p. 12, pl. 2, figs. 11-15.
- 1986 <u>Cupuliferoidaepollenites</u> parvulus (Groot and Penny) Dettmann; Farabee and Canright, p. 46, pl. 17, figs. 5, 6.
- 1988 <u>Cupuliferoidaepollenites</u> parvulus (Groot and Penny) Dettmann; Sweet and McIntyre, p. 508, fig. 7, No. 8, 9.

Remarks: This species is rare, a total of 7 specimens were observed throughout the Bylot Island, Sermilik and Byam Martin formations.

Genus <u>Fraxinoipollenites</u> Potonié ex Potonié, 1960 Type Species: <u>Fraxinoipollenites</u> pudicus (Potonié) Potonié, 1960.

Fraxinoipollenites variabilis Stanley, 1965

Pl. 5, Fig. 5

Synonymy:

- 1965 <u>Fraxinoipollenites</u> variabilis Stanley, p. 306, pl. 45, figs. 29-35.
- 1971 <u>Fraxinoipollenites variabilis</u> Stanley; Leffingwell, p. 45, pl. 8, figs. 8-10.
- 1972 <u>Fraxinoipollenites variabilis</u> Stanley; Rouse and Srivastava, p. 1178, figs. 71-73.
- 1973 <u>Fraxinoipollenites</u> variabilis Stanley; Stone, p. 88, pl. 18, figs. 116, 117.
- 1975 <u>Fraxinoipollenites variabilis</u> Stanley; Srivastava, p. 88, pl. 42, figs. 3-5.
- 1981 <u>Fraxinoipollenites variabilis</u> Stanley; Srivastava, pl. 10, figs. 9, 10.
- 1986 Fraxinoipollenites variabilis Stanley; Farabee and Canright, p. 49, pl. 18, fig. 1.

Age and Distribution: Fraxinoipollenites variabilis is

found in the Almond Formation (late Campanian) of Wyoming (Stone, 1973); the Lance Formation (Maastrichtian) of Wyoming (Farabee and Canright, 1986; the Fort Union Formation (Paleocene) of South Dakota (Stanley, 1965); and the Fort Union Formation (Paleocene) of Wyoming (Leffingwell, 1971).

Remarks: This species is restricted to the Byam Martin Formation and lowermost strata of the Bylot Island Formation.

Genus <u>Tubifloridites</u> Cookson ex Potonié, 1960 Type species: <u>Tubifloridites antipodica</u> (Cookson) Potonié, 1960.

Tubifloridites lilliei (Couper) Farabee and Canright, 1986 Pl. 5, Fig. 6

Synonymy:

- 1953 <u>Tricolpites lilliei</u> Couper, p. 43, pl. 8, figs. 116, 117.
- 1973 <u>Tricolporites lilliei</u> (Couper) Stover and Evans, p. 65, pl. 1, fig. 8.
- 1973 <u>Tricolpites lilliei</u> Couper; Stone, p. 86, pl. 18, fig. 111.
- 1969 <u>Tricolpites lilliei</u> Couper; Norton and Hall, p. 46, pl. 7, fig. 6.
- 1986 Tubifloridites lilliei (Couper) Farabee and Canright,

p. 70, pl. 24, fig. 15.

Age and Distribution: This species was reported from the Almond Formation (Campanian) of Wyoming (Stone, 1973); the Lebo Formation (Paleocene) of Montana (Norton and Hall, 1969); the Lance Formation (Maastrichtian) of Wyoming (Farabee and Canright, 1986)

Remarks: The species <u>lilliei</u> was placed in the genus <u>Tubiflorides</u> because it possessed an echinate to granulate sexine with conspicuous colpi (Farabee and Canright, 1986). A single specimen was recovered from the Sermilik Formation (Lithotype 1).

Syncolpate Pollen

Genus Porosipollis Krutzsch, 1969

Type Species: <u>Porosipollis porosus</u> Krutzsch (Mchedlishvili 1969).

Porosipollis porosus Krutzsch (Mchedlishvili) 1969

Pl. 5, Fig. 7

Synonymy:

- 1969 Porosipollis porosus Krutzsch, p. 407, pl. 1, figs. 10-14.
- 1974 <u>Syncolpites porosus</u> Mchedlishvili; McIntyre, p. 51, pl. 21, figs. 4, 5.

98

1976 <u>Syncolpites porosus</u> Mchedlishvili; Doerenkamp <u>et al.</u>, p. 407, pl. 2, figs. 1, 2.

1989 Porosipollis porosus Krutzsch; Sweet et al., p. 94, pl. 1, fig. 29.

Age and Distribution: This species was reported from Paleocene strata of Siberia (Krutzsch, 1969); late Campanian to mid Maastrichtian strata of Horton River (McIntyre, 1974); Kanguk Formation strata (early to mid Maastrichtian) of Banks Island and Adjacent Areas (Doerenkamp et al., 1976); and mid or late Maastrichtian strata of Police Island, N.W.T. (Sweet et al., 1989).

Remarks: In their Police Island Section, Sweet <u>et al.</u>, (1989), informally defined a <u>Porosicollis porosus</u> zone of mid or late Maastrichtian age. Specimens of this species were restricted to south coast sections. Only two specimens were recovered; one from the Bylot Island Formation (Section D, Appendix A_k), and the other from the Sermilik Formation (Lithotype 1).

Tricolporate Pollen

Genus <u>Caprifoliipites</u> Wodehouse, 1933 Type Species: <u>Caprifoliipites viridifluminis</u> Wodehouse, 1933. Caprifoliipites longus Stanley, 1965

Pl. 5, Fig. 8

Synonymy:

1965 Caprifoliipites longus Stanley, p. 295, pl. 44, figs. 10-14.

Age and Distribution: This species is reported from the Fort Union Formation (Paleocene) of South Dakota (Stanley, 1965).

Remarks: This species is characterized by meridionally elongated pores, having a length of 8 to 10µm (Stanley, 1965). A single specimen was recovered from the Sermilik Formation (Lithotype 1).

Genus <u>Cranwellia</u> Srivastava emend. Srivastava, 1969 Type Species: <u>Cranwellia striata</u> (Couper) Srivastava, 1966. <u>Cranwellia</u> sp. cf. <u>C. striata</u> Srivastava, 1966 Pl. 5, Fig. 9

Remarks: The specimen in this study is slightly different from <u>Cranvellia striata</u>, in having a less pronounced finely granulate sculpture. The equatorial view shown in the photo is a poor orientation for an exact species designation. A single specimen was observed from the Bylot Island Formation along the south coast. Triporate Pollen

Genus <u>Pseudoplicapollis</u> Krutzsch ex Goczan <u>et al.</u>, 1967 emend. Christopher 1979

Type Species: <u>Pseudoplicapollis palaeocaenicus</u> Krutzsch, 1967.

Pseudoplicapollis longiannulata Christopher, 1979 Pl. 5, Fig. 10

Synonymy:

1979 Pseudoplicapollis longiannulata Christopher,

p. 114-115, pl. 5, figs. 3,4; pl. 9, figs. 1-9.

Age and Distribution: Late Turonian to Santonian of the Magothy Formation, New Jersey (Christopher, 1979). ' Remarks: Christopher emended the genus <u>Bseudoplicanollis</u> to include species that have protruding annuli and/or endannuli. Three specimens were recovered from the Byam Marti: Formation on the south coast; a single specimen was recovered from the Sermilik Formation (Lithotype 2).

> Pseudoplicapollis sp. G of Christopher, 1979 Pl. 5. Figs. 11, 12

Synonymy:

1979 <u>Pseudoplicapollis</u> sp. G Christopher, pl. 7, figs. 4, 5. Description: Grains in polar view were convex-triangular. Endoplicae are Y-shaped structures centered over the poles, and terminate as points within the germinals. The exogerminals are very weakly annulate. The surface sculpture is scabrate.

Size Range: Equatorial diameter 19µm (2 specimens examined)

Age and Distribution: Santonian of the Magothy Formation, New Jersey (Christopher, 1979).

Remarks: Two specimens were recovered; one from the Byam Martin Formation and one from the Sermilik Formation (Lithotype 2).

Genus <u>Carpinipites</u> Srivastava, 1966 Type Species: <u>Carpinipites</u> ancipites (Wodehouse) Srivastava, 1966.

Carpinipites ancipites (Wodehouse) Srivastava, 1966

Pl. 5, Fig. 13

Synonymy:

1933 Carpinus ancipites Wodehouse, p. 510, fig. 42.

- 1966 <u>Carpinus ancipites</u> Wodehouse; Martin and Rouse, p. 197, pl. 8, figs. 74-76.
- 1967 <u>Carpinites ancipites</u> (Wodehouse) Srivastava; Srivastava, p. 141, pl. 2, fig. R.

1971 Carpinus ancipites Wodehouse; Rouse et al., p. 236, pl. 8, figs. 12, 15.

1978 <u>Carpinites ancipites</u> (Wodehouse) Srivastava; Wilson, p. 142, pl. 10, fig. 2.

Age and Distribution: This species has been reported from the Lebo Formation (late Campanian to Maastrichtian) of the Yukon Territory (Wilson, 1978); and the Burrard Formation (Eccene) of British Columbia (Rouse <u>et al.</u>, 1971.

Remarks: The genus <u>carpinipites</u> is differentiated from other similar genera by the lack of a thickened region in the pore area, and having pores that are slightly or not at all protruding (Wodehouse, 1959). This species is very rare; one or two specimens were recovered from the Sermilik, Byam Martin and Bylot Island formations.

Genus <u>Paraalnipollenites</u> Hills and Wallace, 1969 Type Species: <u>Paraalnipollenites</u> <u>alterniporus</u> (Simpsom) Srivastava, 1975.

> <u>Paraalnipollenites alterniporus</u> (Simpson) Srivastava, 1975 Pl. 4, Fig. 14

Synonymy:

1967 Triatriopollenites confusus Zaklinskaia; Bratzeva,

p. 123, pl. 1, fig. F.

- 1969 <u>Paraalnipollenites confusus</u> (Zaklinskaia) Hills and Wallace, p. 141, pl. 17, fiqs. 1-8.
- 1972 <u>Paraalnipollenites confusus</u> (Zaklinskaia) Hills and Wallace; Rouse and Srivastava, p. 1177, fig. 59.
- 1974 <u>Paraalnipollenites confusus</u> (Zaklinskaia) Hills and Wallace; McIntyre, p. 53, pl. 22, fig. 14.
- 1977 Paraalnipollenites confusus (Zalinskaia) Hills and Wallace; Rouse, p. 51, pl. 1, fig. 2.
- 1978 <u>Paraalnipollenites confusus</u> (Zaklinskaia) Hills and Wallace; Wilson, p. 144, pl. 10, fig. 25.
- 1983 <u>Paraalnipollenites confusus</u> (Zaklinskaia) Hills and Wallace; Wingate, p. 118, pl. 5, fig. 18.
- 1986 <u>Paraalnipollenites alterniporus</u> (Simpson) Srivastava; Jerzykiewicz and Sweet, p. 1371, pl. 4, fig. 1.
- 1988 <u>Paraalnipollenites</u> <u>confusus</u> (Zaklinskaia) Hills and Wallace; Frederiksen <u>et al.</u>, p. 521, pl. 1, figs. 8, 9.
- 1989 <u>Paraalnipollenites alterniporus</u> (Simpson) Srivastava; Sweet <u>et al.</u>, p. 98, pl. 2, fig. 10.
- 1989 <u>Paraalnipollenites alterniporus</u> (Simpson) Srivastava; McIntyre, p. 195, pl. 2, fig. 5.

Age and Distribution: This species has been reported from the Eureka Sound Formation (Maastrichtian to Paleocene) of Bathurst Island, N.W.T., Canada (Hills and Wallace, 1969); the Bonnet Plume Formation (early Paleocene) of northeastern Yukon (Rouse and Srivastava, 1972); late Maastrichtian strata of Horton River (McIntyre, 1974); the Eureka Sound Formation (Paleocene) of Ellesmere Island (Rouse, 1977); Maastrichtian strata of the Yukon Territory (Wilson, 1978); the Elko Formation (Eocene) of Nevada (Wingate, 1983); the Coalspur Formation (late Maastrichtian to Paleocene) of central Alberta (Jerzykiewicz and Sweet, 1986); mid Maastrichtian strata of north Alaska (Frederiksen <u>et al.</u>, 1988); late Maastrichtian to Paleocene strata of Police Island, N.W.T. (Sweet <u>et al.</u>, 1989) and the Eureka Sound Group (Paleocene) of Somerset Island, N.W.T. (McIntyre).

Remarks: <u>Paraalnipollenites alterniporus</u> is distinguished from other triporates by the presence of false pores, the arci extending between the true pores arc out as if to form a pore (Wilson, 1978). This species is restricted to the Bylot Island Formation and the Sermilik Formation (Lithotypes 1 and 2).

Genus <u>Momipites</u> (Wodehouse) Frederiksen and Christopher, 1978

Type Species: <u>Momipites coryloides</u> Wodehouse, 1933. <u>Momipites wyomingensis</u> Nichols and Ott, 1978 Pl. 5, Fig. 15

Synonymy:

1978 Momipites wyomingensis Nicholls and Ott, p. 100, pl. 1, figs. 1-4. 1986 <u>Momipites wyomingensis</u> Nicholls and Ott; Farabee and Canright, p. 55, pl. 19, fig. 15.

1989 <u>Momipites wyomingensis</u> Nicholls and Ott; McIntyre, p. 195, pl. 1, fig. 3.

Age and Distribution: This species has been reported from the Fort Union Formation (Paleocene) in the Wind River Basin, Wyoming (Nicholls and Ott, 1978); the Lance Formation (Maastrichtian) of Wyoming (Farabee and Canright, 1986); and the Eureka Sound Group (late Paleocene) of Somerset Island, N.W.T..

Remarks: This species is restricted to the Sermilik Formation and the Bylot Island Formation. <u>Momipites</u> <u>wyomingensis</u> is most abundant (5 to 9 specimens/sample) in Lithotype 3 of the Sermilik Formation.

Genus Triatriopollenites Pflug, 1953

Type Species: Triatriopollenites rurensis Pflug and Thomson, 1953.

Triatriopollenites rurensis Pflug and Thomson, 1953 Pl. 5, Fig. 16

Synonymy:

1978 <u>Triatriopollenites rurensis</u> Pflug and Thomson; Schumaker-Lambry, pl. 13, figs. 4, 8. 107

Age and Distribution: Faleocene of Belgium (Schumacker-Lambry, 1978).

Remarks: This species first occurred in the Sermilik Formation (Lithotype 2) in the south coast sections. At Twosnout Creek, the species was by and large restricted to upper strata of the Bylot Island Formation. Isolated occurrences were recorded in lower strata of the Bylot Island Formation and the Byam Martin Formation.

Genus <u>Trudopollis</u> Pflug emend. Krutzsch, 1967 Type Species: <u>Trudopollis pertrudens</u> (Pflug) Pflug, 1953.

Trudopollis variabilis R.H. Tschudy, 1975

Pl. 5, Fig. 17

Synonymy:

1975 <u>Trudopollis variabilis</u> R.H. Tschudy, p. 25, pl. 16, figs. 13-22.

1976 <u>Trudopollis variabilis</u> R.H. Tschudy; Wolfe, p. 13, pl. 1, fig. 28.

Age and Distribution: This species has been reported from the Ripley Formation (mid Campanian to early Maastrichtian) of the Mississippi Embayment (R.H. Tschudy, 1975); and the Englishtown Formation (mid Campanian) of the middle Atlantic States (Wolfe, 1976). "A tay has manufactured bare and an and barrant over a set of the set

1. 18 -1

Remarks: This taxon is characterized by a verrucate ektexine, a greatly thickened and angular annulus and the absence of an interloculum (R.H. Tschudy, 1975). <u>Trudopollis conrector</u> Pflug differs from <u>Trudopollis</u> <u>variabilis</u> by the presence of a prominent vestibulum and a scabrate exine.

Trudopollis conrector Pflug, 1953

Pl. 5, Fig. 18

Synonymy:

- 1961 <u>Trudopollis conrector</u> Pflug; Groot <u>et. al.</u>, p. 137, pl. 26, fig. 52.
- 1981 <u>Trudopollis conrector</u> Pflug; Pacltová, p. 205, pl. 17, figs. 2, 2b, 2c.
- 1981 <u>Trudopollis conrector</u> Pflug; Mikhelis, p. 229, pl. 13, figs. 5-10.

Age and Distribution: This species has been reported from the Magothy Formation (Santonian) of the Eastern United States (Groot <u>et al.</u>, 1961); Campanian to Paleocene strata of the Sea of Azov (Mikhelis, 1981); Santonian strata of the Bohemian Massif, Eastern Europe (Pacltová, 1981).

Remarks: Only 5 specimens of this species were recovered. Four of the specimens were from the Bylot Island Formation at Twosnout Creek and a single specimen from the Byam Martin Formation at Twosnout Creek.

Cenus <u>Megatriopollis</u> Goczan and Krutzsch, 1967 Type Species: <u>Megatriopollis santonius</u> Goczan and Krutzsch, 1967.

Megatriopollis sp. 1

Pl. 6, Fig. 1

Synonymy:

- 1974 <u>Extratriporopollenites</u> sp. 2 McIntyre (pars) p. 52, pl. 22, fig. 2 only.
- 1976 <u>Extratriporopollenites</u> sp. 2 McIntyre; Doerenkamp <u>et</u> <u>al.</u>, p. 405, pl. 1, fig. 16.

Description: Pollen grains have a triangular to convex triangular shape. Nall is two layered (lµm thick), with a slightly thickened exine (1.5µm thick) bordering the germinal. Surface sculpture is finely reticulate (lumina ≤ 0.5µm wide). The commonly observed internal structured layer is absent.

Size Range: Polar diameter 28-30µm (2 specimens measured)

Remarks: The generic diagnosis by Goczan and Krutzsch, (1967; <u>in</u> Jansonius and Hills, 1976) indicates that the internal structured layer may or may not be present. A specimen labelled <u>Extratriporopollenites</u> sp. 2 by McIntyre (1974) from Horton River appears identical to specimens found in the Sermilik Formation (Lithotype 1). As well, the specimen labelled <u>Extratriporopollenites</u> sp. 2 McIntyre by Doerenkamp <u>et al.</u>, (1976) also appears identical to those observed in this study.

Age and Distribution: Specimens labelled <u>Extratriporopollenites</u> sp. 2 McIntyre have been reported from early to mid Maastrichtian strata of Horton River (McIntyre, 1974) and from late Campanian to Paleocene strata of Banks Island (Doerenkamp <u>et al.</u>, (1976).

Genus <u>Neotriangulipollis</u> Goczan and Krutzsch, 1967 Type Species: <u>Neotriangulipollis piolencensis</u> Krutzsch, 1967.

Neotriangulipollis sp. 1 of Azema, 1981

Pl. 6, Fig. 2

Synonymy:

1980 Vacuopollis microconcavus Pacltova; Medus et al.,

p. 144, pl. 11, figs. 11-13, 26.

1981 Neotriangulipollis sp. 1 Azema; p. 263, pl. 5, fig. 13.

Description: This species is similar to <u>Neotriangulipollis</u> sp. 1 of Azema (1981), however, there is a slight variation in the pore area. With the latter species, the ectexine is thickened around the pore to form a 2.5µm annulus. The single specimen recovered from this study exhibited a thickened region (approximately 1.5μm wide) around the pore. The specimen measured 24 μm in polar diameter.

Age and Distribution: Santonian strata of France (Azema <u>et al.</u>, 1981); and late Turonian strata of Portugal (Medus <u>et al.</u> 1980).

Remarks: Azema (1981) indicates that <u>Vacuopollis</u> <u>microconcavus</u> by Medus <u>et al.</u>, (1980) from the Turonian of Portugal is identical. A single specimen was recovered from the Sermilik Formation (Lithotype 1).

Genus <u>Extratriporopollenites</u> Pflug emend. Skarby, 1968 Type Species: <u>Extratriporopollenites</u> fractus Pflug, 1953. <u>Extratriporopollenites</u> sp. 2 of McIntyre, 1974 Pl. 6, Fig. 3

Synonymy:

1974 Extratriporopollenites sp. 2 McIntyre (pars) p. 52, pl. 22, figs. 3 and 4 only.

Description: Triporate; triangular to convex triangular; surface finely reticulate (lumina $\leq 0.5\mu\pi$ wide); ektexine uniformly thin (0.5 $\mu\pi$ thick) and unthickened in the aperatural region. Size Range: Polar diameter 21-24µm (5 specimens measured)

Remarks: This species is identical in all respects to that of McIntyre (1974). Specimens were restricted to the Sermilik Formation (Lithotype 2) and the Bylot Island Formation (Section C).

Age and Distribution: Early to mid Maastrichtian strata of Horton River (McIntyre, 1974).

Stephanoporate Pollen

Genus Polyvestibulopollenites Pflug, 1953

Type Species: <u>Polyvestibulopollenites verus</u> (Potonie) Thomson and Pflug, 1953.

Polyvestibulopollenites trinus (Stanley) Norris, 1986 Pl. 6, Fig. 4

Synonymy:

1965 Alnus trina Stanley, p.289, pl. 43, figs. 4-6.

- 1969 <u>Alnipollenites trina</u> Stanley; Norton and Hall, p. 42, pl. 5, fig. 20.
- 1978 <u>Alnipollenites versus</u> Potonie; Wilson, p. 145, pl. 11, figs. 9, 10.
- 1980 Jarzenipollenites trinus Stanley; Kedves, p. 172, pl. II, figs. 4-6.

1986 Alnus trina Stanley; Jerzykiewicz and Sweet, pl. 1,

fig. 19.

- 1986 <u>Alnipollenites trina</u> (Stanley) Norton; Farabee and Canright, p. 36, pl. 10, figs. 11-13.
- 1986 Polyvestibulopollenites trinus (Stanley) Norris, p. 41, pl. 10, figs. 46, 53.

Age and Distribution: This species has been reported from the Fort Union Formation (early Paleocene) of South Dakota (Stanley, 1965); the Tullock Formation (Paleoce', 3) of Montana (Norton and Hall, 1969); the Lance Formation (Maastrichtian) of Wyoming (Farabee and Canright, 1986); the Paskapoo Formation (Paleocene) of the Alberta Foothills (Jerzykiewicz and Sweet, 1986); and the Mackenzie Bay and Kugmallit Formations (Paleocene to Miocene. (Norris, 1986).

Remarks: <u>Polyvestibulopollenites trinus</u> is readily distinguished from other polyporate grains by the presence of curved arci joining the pores (Wilson, 1978). This species is also characterized by a thickened exine in the pore region, forming an annulus and a slight labrum (Stanley, 1965).

Specimens of this species were recovered from the Bylot Island Formation at Twosnout Creek.

113

Genus Polyatriopollenites Pflug, 1953

Type Species: <u>Polyatriopollenites</u> <u>stellatus</u> (Potonié) Pflug, 1953.

Polyatriopollenites stellatus (Potonié) Pflug, 1953

Pl. 6, Fig. 5

Synonymy:

- 1966 Pterocarva stellatus (Potonié); Martin and Rouse, p. 196, pl. 8, figs. 79, 80.
- 1969 Pterocarya stellarus (Potonié) Martin and Rouse; Hopkins, p. 1121, pl. 9, figs. 122-124.
- 1969 Pterocaryapollenites stellatus (Potonié) Thiergart; Norton and Hall, p. 42, pl. 5, fig. 23.
- 1971 <u>Pterocarya stellatus</u> (Potonié) Martin and Rouse; Piel, p. 1910, pl. 13, figs. 113, 114.
- 1978 Pterocaryapollenites stellatus (Potonié) Raatz; Wilson, p. 146, pl. 9, fig. 7; pl. 10, figs. 13, 17.
- 1986 <u>Polyatriopollenites stellatus</u> (Potonié) Pflug; Norris, p. 42, pl. 11, figs. 5-7.
- 1987 <u>Polvatriopollenites stellatus</u> (Potonié) Pflug; Langille, p. 101, pl. 6, fig. 12.

Age and Distribution: Mid Maastrichtian of the Hell Creek Formation, Montana (Norton and Hall, 1969); Maastrichtian to Paleocene of the Bonnet Plume Formation, Northeastern Yukon (Rouse and Srivastava, 1972); Maastrichtian strata of the Mackenzie Delta (Wilson, 1978); Eocene to Pliocene of the Kugmallit and Richards Formations, Mackenzie Delta (Norris, 1986); Paleocene strata of Baffin Island (Langille, 1987).

Remarks: Specimens in this study have slightly raised labra. Four, five and six pored pollen were observed. This species was very rare; one or two specimens were recovered from all three formations (Byam Martin, Setmilik, and the Bylot Island Formation).

Genus <u>Tetraporites</u> Samoliovich, 1965 Type Species: <u>Tetraporites</u> pravus Samoliovich, 1965. <u>Tetraporites</u> sp. 1 of McIntyre, 1974 Pl. 6, Fig. 6

Synonymy:

1974 Tetraporites sp. 1 McIntyre, pl. 22, fig. 9, 10.

Description: Only two specimens were observed. The grains are granulate and measured 26 and 30 μ m in diameter. The exine is thin (< 1 μ m thick) and slightly thickened (1.5 μ m thick) in the pore region.

Age and Description: Very late Campanian to mid Maastrichtian strata of Horton River (McIntyre, 1974).

Remarks: <u>Tetraporites</u> sp. 1, is identical in nearly all respects to <u>Tetraporites</u> sp. 1 of McIntyre (1974).

<u>Tetraporites</u> sp. 2 differs in being 8µm to 9µm smaller in diameter, and possessing a psilate exime. This species was very rare; a single specimen was recovered from the Bylot Island Formation along the south coast (Section D, Appendix A) and the other from the Sermilik Formation (Lithotype 1).

Tetraporites sp. 2

Pl. 6, Fig. 7

Description: The exine is psilate, thin (< lµm thick) and slightly thickened (1.5µm thick) in the pore region.

Size Range: Equatorial diarater 18-21µm (6 specimens measured)

Remarks: This species is restricted to the Sermilik Formation (Lithotype 2) and uppermost strata of the Bylot Island Formation at Twosnout Creek.

Binigeminate Pollen

Genus <u>Wodehouseia</u> Stanley, 1961 Type Species: <u>Wodehouseia gpinata</u> Stanley, 1961. <u>Wodehouseia gracile</u> (Samoilovitch) Pokrovaskaya, 1966 Pl. 6, Fig. 8

Synonymy:

- 1967 Wodehouseia gracile (Samoilovitch) Samoilovitch, p. 132, pl. 2, fig. 6.
- 1974 <u>Wodehousiea</u> gracile (Samoilovitch) Pokrovaskāya; McIntyre, pl. 15, figs. 15, 16.
- 1976 <u>Wodehouseia</u> <u>gracile</u> (Samoilovitch) Pokrovaskaya; Doerenkamp <u>et al.</u>, pl. 3, figs. 18, 19.
- 1976 <u>Wodehouseia gracile</u> (Samoilovitch) Pokrovaskaya; Wiggins, p. 65, pl. 3, figs. 3, 4.
- 1989 Wodehouseia gracile (Samoilovitch) Pokrovaskaya; Sweet et. al., p. 95, pl. 1, fig. 20.

Age and Distribution: Very late Campanian to mid Maastrichtian strata of Horton River (McIntyre, 1974); Maastrichtian of the Kanguk Formation, Banks Island and adjacent areas (Doerenkamp <u>et al.</u> 1976); late Maastrichtian strata of Alaska (Wiggins, 1976); and mid(?) Maastrichtian strata of Police Island, N.W.T. (Sweet <u>et al.</u>, 1989).

Remarks: Wiggins (1976) indicated that the size range for specimens in his Alaskan study was (28-31 µm in length) and (39-41 µm in transverse equatorial projection). Specimens in this study were as small as 20µm in length and ranged between 26-34 µm in transverse equatorial projection.

In south coast sections, <u>Wodehouseia gracile</u> is restricted to the Sermilik Formation (Lithotype 1) and a tongue of Bylot Island Formation strata (Section D, Appendix A_i). At Twosnout Creek, this species is restricted to (Section C, Appendix A₃), which incorporates both Bylot Island Formation strata and Lithotype 3 of the Sermilik Formation.

> Wodehouseia sp. 1 Pl. 6, Fig. 9

Description: Isopolar fossil pollen with bilateral symmetry. Equatorial pollen profile, circumscribing the longest pollen axis, suboblate to oblate. Apertures binigeminate, large, elliptical colpi, approximately 10µm long and 4µm wide. The colpi are at right angles to the longest pollen axis. The wall is double layered, composed of a smooth endexine which forms a central pollen body, and a slightly punctate flange with external spines. The spines are restricted to the flange meridian. The flange is 1µm wide at the termini of the longest equatorial axis.

Size Range: Polar Length (2 specimens) 34-38 µm Transverse equatorial projection 50-52 µm Central body length 13-15 µm Flange length 9-10 µm Length of spines 1.5-2 µm

Remarks: This species is differentiated from Wodehousia

118

<u>spinata</u> Stanley and <u>Wodehousia octospina</u> Wiggins by the absence of spines on the central body. It differs from <u>Wodehousia edmontonicola</u> Wiggins by its distinctly larger size, and smooth central body. This species is restricted to the Sermilik Formation (Lithotype 2) and the Bylot Island Formation.

Genus <u>Singularia</u> Samoilovitch, 1961 Type Species: <u>Singularia aculeata</u> Samoilovitch, 1961. <u>Singularia aculeata</u>, Samoilovitch, 1961 Pl. 6, Fig. 10

Synonymy:

1974 <u>Singularia aculeata</u> Samoilovitch; McIntyre, p. 39, pl. 15, fig. 17.

1976 Singularia aculeata Samoilovitch; Wiggins, p. 68.

Description: The equatorial profile is oblate. The flange is very poorly developed with the ektexine and endexine being closely appressed. The central body is characterized by numerous short spines. The spines are developed along the meridian as well. The flange or meridian is $\leq 1 \mu m$.

Size Range: (1 Specimen measured) Transverse equatorial projection 30μm Polar length 16μm Age and Distribution: Maastrichtian-Paleocene? strata of the Western Siberian Lowland (Chlonova, 1961); late Maastrichtian strata of Alaska (Wiggins, 1976); and very late Campanian to mid Maastrichtian strata of Horton River, (McIntyre, 1974).

Remarks: This species is very rare; two specimens were recovered from the Sermilik Formation (Lithotype 2). Wiggins (1976) encountered too few specimens to make a precise morphologic evaluation of the genus, but indicated that its stratigraphic value could be important.

Triprojectate Pollen

Genus Mancicorpus Mtchedlishvili emend.

Srivastava, 1968

Type Species: <u>Mancicorpus anchoriforme</u> Mtchedlishvili, 1961. <u>Mancicorpus trapeziforme</u> Mtchedlishvili, 1961

Pl. 6, Fig. 11

Synonymy:

1974 <u>Mancicorpus trapeziforme</u> Mtchedlishvili; McIntyre, p. 49, pl. 20, figs. 5, 6.

1976 <u>Mancicorpus trapeziforme</u> Mtchedlishvili; Doerenkamp <u>et</u> <u>al.</u>, p. 409, pl. 3, fig. 2.

Age and Distribution: Very late Campanian to mid

Maastrichtian of Horton River (McIntyre, 1974); Maastrichtian of Banks Island and adjacent areas (Doerenkamp <u>et al.</u>, 1976).

Remarks: B.D. Tschudy (1969) indicates that <u>Mancicorpus</u> <u>trapeziforme</u> resembles <u>Aquilapollenites senonicus</u> Mtchedlishvili; the two species are only differentiated by a variation in the size of the angle formed at the contact of the equatorial projections. For <u>Mancicorpus trapeziforme</u> the equatorial projection juns the body nearly at right angles, for <u>Aquilapollenites senonicus</u> the equatorial projection joins the hody at an acute angle. A single specimen was recovered from the Sermilik Formation (Lithotype 2).

Genus <u>Aquilapollenites</u> (Rouse) Funkhouser, 1961 Type Species: <u>Aquilapollenites guadrilobus</u> Rouse, 1957. <u>Aquilapollenites scabridus</u> B.D. Tschudy, 1969 Pl. 6, Fig. 12

Synonymy:

1969 Aquilapollenites scabridus B.D. Tschudy, p. All, pl. 13, figs. 1-11, pl. 14, figs. 1, 2.

Age and Distribution: <u>Aquilapollenites</u> scabridus B.D. Tschudy, was recovered with other mid Campanian to Maastrichtian species of <u>Aquilapollenites</u> at Nation River,

121

Alaska. Only one sample was collected at that locality, hence a more precise age was not possible.

Remarks: This species is diagnostic for its rough surface sculpture (B.D. Tschudy, 1969). A single specimen was recovered from the Bylot Island Formation along the south coast.

Aquilapollenites sp. 1

Pl. 6, Fig. 13

Remarks and Description: λ single poorly preserved specimen was recovered from the Sermilik Formation (Lithotype 2). The surface is ornamented with numerous short spines (< 1 μ m high); the grain has an equatorial diameter of 30 μ m.

Unknown Specimens

Specimen 1 Pl. 6, Fig. 14

Description: Spore or pollen of unknown affinity. Grains circular: surface ornament psilate to finely granulate. Diameter of 25 measured specimens 26 (30) 33 µm.

Remarks: Specimens were observed in the Sermilik Formation

(Lithotype 1) and lowermost strata of the Bylot Island Formation at Twosnout Creek.

> Specimen 2 Pl. 6, Fig. 15

Description: Monosulcate pollen ? of unknown affinity. Grain circular; surface ornament rugulate (<1µm wide). Ektexine consists of 2µm high simple, slender spines. The dimensions of the only recorded specimen are; Length - 49.5 µm, Breadth - 44 µm.

Remarks: A single specimen was recovered from uppermost strata of the Bylot Island Formation at Twosnout Creek.

FUNGAL REMAINS

Genus <u>Pesavis</u> Elsik and Jansonius, 1974 Type Species: <u>Pesavis tagluensis</u> Elsik and Jansonius, 1974. <u>Pesavis parva</u> Kalqutkar and Sweet, 1988

Pl. 6, Fig. 16

Synonymy:

- 1974 <u>Pesavis tagluensis</u> Elsik and Jansonius (pars), p. 955, pl. 1, fig. 10 only.
- 1976 "<u>Pesavis parva</u>" Jansonius, p. 133, pl. 1, fig. 2. 1978 <u>Pesavis</u> sp. Sweet, p. 36, pl. 6.2, fig. 15.

1988 <u>Pesavis parva</u> Kalgutkar and Sweet, p. 123, pl. 6.1, figs. 6-12.

Description: The overall diameter is 26 µm; there are 6 lateral arm cells; and a central cavity is absent (one specimen measured).

Age and Distribution: Late Maastrichtian to Lower Paleocene strata of the Yukon Coastal Plain (Kalgutkar and Sweet, 1988); late Maastrichtian strata of the Bonnet Plume Basin (Kalgutkar and Sweet, 1988).

Remarks: <u>Pesavis parva</u> is a biostratigraphically important fungal spore described by Kalgutkar and Sweet (1988). Two specimens of <u>Pesavis parva</u> were recorded in this study, both from uppermost strata of the Bylot Island Formation at Twosnout Creek.

MARINE MICROPLANKTON

Introduction

This project is a stratigraphic and terrestrial palynomorph study; however, selected biostratigraphically significant dinoflagellates were incorporated to help establish age relationships. The dinoflagellates were not counted, thus, the range chart (Appendix C_2) represents only the presence or absence of a taxon.

Proximate Dinoflagellates

Genus <u>Ceratiopsis</u> Vozzhenikova emend. Bujak, Downie, Eaton and Williams, 1980

Type Species: <u>Ceratiopsis leptoderma</u> Vozzhennikova, 1963

Ceratiopsis diebelii (Alberti) Vozzhencikova, 1967.

Pl. 6, Fig. 17

Synonymy:

- 1967 <u>Deflandrea diebelii</u> sp. cf. <u>D. Diebelii</u> Alberti; Drugg, p. 16, pl. 2, fig. 6.
- 1974 <u>Deflandrea diebelii</u> Alberti; McIntyre, p. 17, pl. 4, figs. 4, 5.
- 1975 <u>Deflandrea diebelii</u> Alberti; McIntyre, p. 67, pl. 4, figs. 1, 2.
- 1976 <u>Deflandrea diebelii</u> Alberti; Doerenkamp <u>et al.</u>, pl. 4, fig. 7.
- 1977 <u>Deflandrea diebelii</u> Alberti; Williams and Bujak, p. 46, pl. 5, fig. 1.
- 1978 <u>Deflandrea diebelii</u> Alberti; Wilson, p. 151, pl. 11, fig. 15.
- 1980 <u>Deflandrea diebelii</u> Alberti; May, p. 75, pl. 8, fig. 16.
- 1980 Deflandrea diebelii Alberti; Croxton, p. 25, pl. 4,

fig. 6; p. 27, pl. 5, fig. 6.

- 1985 <u>Ceratiopsis diebelii</u> (Alberti) Vozzhennikova; Williams and Bujak, p. 870, fig. 23, No. 17.
- 1986 <u>Deflandrea</u> <u>diebelii</u> Alberti; Ioannides, p. 19, pl. 11, figs. 6, 7, 10, 11.
- 1988 <u>Ceratiopsis diebelii</u> (Alberti) Vozzhennikova; Shaozhi and Norris, p. 78, pl. 9, fiq. 6.

Age and Distribution: This species has been reported from Maastrichtian strata of Horton River (McIntyre, 1974; 1975); Maastrichtian of Banks Island and Adjacent Areas (Doerenkamp <u>et al.</u>, 1976); Maastrichtian of offshore southeastern Canada, (Bujak and Williams, 1978); Maastrichtian of the Atlantic Highlands, New Jersey (May, 1980); Maastrichtian and Danian of Escarpado Canyon, California (Drugg, 1967); Maastrichtian - Paleocene ?, of West Central Greenland (Croxton, 1980).

Remarks: Lentin and Williams (1975, p. 2153); considered this genus to be a junior synonym of <u>Deflandrea</u>. This view was later rejected by Lentin and Williams (1977, p. 20); because of the nature of the hexa 2a archeopyle.

<u>Ceratiopsis diebelii</u> is the most significant marine palynomorph marking the Maastrichtian worldwide. Extensive compilations by Wilson (1971), Bujak and Williams (1978), McIntyre (1974; 1975), May (1980) and Williams and Bujak (1985) indicate that the presence of <u>Caratiopsis diebelii</u> marks the first occurrence of <u>Maastrichtian strata</u>. The first occurrence of <u>Caratiopsis diebelii</u> is subsequently used to indicate the presence of <u>Maastrichtian rocks</u> on Bylot Island. This species was recovered from the Sermilik Formation and the Bylot Island Formation.

Genus <u>Palaeoperidinium</u> Deflandre emend. Sarjeant, 1967 Type Species: <u>Palaeoperidinium pyrophorum</u> (Ehrenberg) Sarjeant, 1967.

Palaeoperidinium kozlowskii (Gorka) Davey, 1970 Pl. 6, Fig. 18

Synonymy:

- 1974 <u>Lejeunia kozlowskii</u> Gorka; McIntyre, p. 17, pl. 4, figs. 2, 3.
- 1975 <u>Leieunia kozlowskii</u> Gorka; McIntyre, p. 68, pl. 4, figs. 3, 4.
- 1976 Lejeunia kozlowskii Gorka; Doerenkamp et al., p. 411, pl. 4, fig. 8.

Age and Distribution: This species has been reported from the late Campanian to early Maastrichtian strata of Horton River (McIntyre, 1974; 1975); and late Campanian to early Maastrichtian strata of Banks Island and adjacent areas (Doerenkamp <u>et al.</u>, (1976). Remarks: McIntyre (1974; 1975), Doerenkamp <u>at al.</u>, (1976), and Bujak and Williams (1978) suggest that the coexistance of <u>Ceratopsis</u> disbelii and <u>Palaeoperidinium kozlowskii</u> are excellent indicators for Maastrichtian strata. A single specimen of <u>Palaeoperidinium kozlowskii</u> was recovered from the Sermilik Formation (Lithotype 1).

CHAPTER 4 DISCUSSION

4.1 Biostratigraphy

Palynological analyses of Cretaceous strata from Eclipse Trough are divided into three palynological assemblage zones, herein called the <u>Gleicheniidites</u> sp. cf. <u>G. circinidites</u> - <u>Antulsporites distaverrucosus</u> (GA) zone, the <u>Porosipollis porosus</u> - <u>Aquilapollenites scabridus</u> (PA) zone and the <u>Singularia aculeata</u> - <u>Posavis parva</u> (SP) zone. These units are informally established in accordance with guidelines in the North American Stratigraphic Code (NACSN, 1983) and the International Stratigraphic Guide (Hedberg, 1976).

Zones are characterized by the occurrence of selected index species. It is not necessary for every index species to be present in a sample, nor is it necessary for a species to be restricted to a particular zone (Hedberg, 1976). The boundary between two zones is placed at the median between the uppermost sample of the lower zone and the lowermost sample of the overlying zone. Those long ranging taxa that are in abundance and characterize no particular zone are recorded on the range chart only (Appendix C₂). Figure 4.1 indicates the known ranges of the significant taxa within these zones.

For each assemblage zone, those species which define or characterize the zone are identified under the following subheadings: species which are restricted to the biozone

129





Carlos a la la la
(both biostratigraphic and long ranging taxa), species which characterize the zono, but overlap with other zones (both biostratigraphic and long ranging taxa), those species which are considered to be new species or unknown taxa, and those species which are common in all zones, but are biostratigraphically significant for late Cretaceous and younger strata.

4.1.1 <u>Gleicheniidites</u> sp. cf. <u>G. circinidites</u>, <u>Antulsporites</u> <u>distaverrucosus</u> (GA) zone

Species which are restricted to the biozone include: <u>Gleicheniidites</u> sp. cf. <u>G. circinidites</u> (Cookson) Dettmann, <u>Ornamentifera</u> baculata Sinsh, <u>Antuleporites</u> distaverrucosus (Brenner) Archangelsky and Gamerro and <u>Gemmatriletes</u> <u>clavatus</u> Brenner. Those species which are most common in the <u>GA</u> zone but overlap with other zones include: <u>Lycopodiacidites</u> <u>canaliculatus</u> Sinsh, <u>Pseudoplicapollis</u> <u>lonsiannulata</u> Christopher and <u>Fraxinoipollenites</u> <u>variabilis</u> Stanley. New species which first occur in this assemblage zone and are locally restricted to strata of Eclipse Trough are: <u>Lycopodiacidites</u> sp. 1, <u>Pityosporites</u> sp. 1, <u>Pityosporites</u> sp. 2, <u>Annulispora</u> sp. 1 and <u>Liliipollis</u> sp. 1.

Species that are biostratigraphically important, but not definitive of the GA zone include: <u>Cingulatisporites</u> <u>dakotaensis</u> Stanley, <u>Hazaria sheopiarii</u> Srivastava, <u>Carpinipites</u> an<u>cipites</u> (Wodehouse) Srivastava, <u>Pseudoplicapollis</u> Sp. G of Christopher, <u>Polyatriopollenites</u> <u>stellatus</u> (Potonié) Pflug, <u>Trudopollis</u> <u>conrector</u> Pflug and <u>Tristriopollenites</u> <u>rurensis</u> Pflug.

Distribution

The stratotype for the GA biozone is the interval which spans 01-SC-22 to 02-SC-23, 0 to 12 metres, of the Byam Martin Formation on the south coast (Section B, Appendix λ_2). At Twosnout Creek, the GA zone is equivalent with the interval which spans 01-TC-85 to 04-TC-98, 14.5 to 220 metres, (Section A, Appendix λ_1). This corresponds with the Byam Martin Formation and the lower 100 metres of the Bylot Island Formation, and is herein designated as a parastratotype for the biozone.

Age

The age of the GA zone is not certain. Terrestrial palynomorph diversity is relatively low and age is in part inferred from associations with conformably overlying zones. Species occurring in this zone which indicate a general late Cretaceous age include: <u>Pseudoplicapollis longiannulata</u> Christopher (late Turonian to Santonian ?), <u>Pseudoplicapollis</u> pp. G of Christopher (Santonian ?), and <u>Trudopollis contector</u> Pflug (Santonian to Paleocene). Those species which are biostratigraphically restricted include: <u>Cingulatisporites</u> <u>dakotaensis</u> Stanley (late Campanian to Paleocone), <u>Hazaria</u> <u>sheopiarii</u> Srivastava (late Campanian to Paleocene), <u>Carpinipites</u> <u>ancipites</u> (Wodehouse) Srivastava (late Campanian to Eocene), <u>Polyatriopollenites</u> <u>stellatus</u> (Potonié) Pflug (Maastrichtian to Paleocene) and Triatriopollenites <u>rurensis</u> Pflug and Thomson (Paleocene).

The presence of characteristic late Campanian to Maastrichtian taxa in the overlying PA biozone, and in particular, <u>Ceratiopsis diebelii</u> (Alberti) Vozhennikova and <u>Palaeoperidinium kozlowskii</u> (Ehrenberg) Davey, coupled with the conformable relationship of the two zones, suggests a mid to late Campanian age for the GA biozone.

The paucity of diverse late Cretaceous terrestrial palynomorphs within the GA biozone makes correlations with other Arctic (Campanian age) palynofloral zones difficult.

4.1.2 <u>Porosipollis porosus</u> - <u>Aquilapollenites scabridus</u> (PA) zone

Species restricted to the biozone include: <u>Densoisporites velatus</u> (Weyland and Kreiger) Krasnova, <u>Cirratriradites teter Norris, Pityosporites constrictus</u> Singh, <u>Occadopites fragilis</u> Singh, <u>Pityosporites</u> <u>alatipollenites</u> (Rouse) Singh, <u>Caprifolibites longus</u> Stanley, <u>Rousea georgensis</u> (Brenner) Dettmann, <u>Neotriangulipollis</u> sp. 1 of Azema, <u>Tubifloridites lilliei</u> (Couper) Farabee and Canright, <u>Porosipollis porosus</u> Mchedlishvili, <u>Aquilapollenites scabridus</u> Tschudy and <u>Palaeoperidinium kozlowskii</u> (Ehrenberg) Davey.

Those species that characterize the PA biozone, but which overlap with other zones include: Wodehouseia gracile (Samoilovitch) Pokrovaskava, Tetraporites sp. 1 of McIntvre, Cibotiumspora juncta (Kara-Murza) Singh and Fovestriletes subtriangularis Brenner. New species and unknown taxa occurring within the PA zone are: Lycopodiacidites sp. 1. Pityosporites sp. 1, 2, and 3, Cicatricosisporites sp. 1, Cycadopites sp. 1, Lilipollis sp. 1, Megatropollis sp. 1. and Unknown Specimen 1. Of the aforementioned species. Lilipollis sp. 1, Megatropollis sp. 1, and Unknown Specimen 1 are locally important for biostratigraphy. Species that are biostratigraphically important, but not definitive of the PA zone include: Hamulatisporis amplus Stanley. Radialisporis radiatus (Krutzsch) Krutzsch, Cinquiatisporites dakotaensis Stanley, Carpinipites ancipites (Wodehouse) Srivastava, Polvatriopollenites stellatus (Potonié) Pflug, Trudopollis conrector Pflug, Momipites wyomingensis Nichols and Ott. Polyvestibulopollenites trinus (Stanley) Norris. Triatriopollenites rurensis Pflug and Thomson, and Ceratiopsis diebelii (Alberti) Vozzhennikova.

Distribution

The stratotype for the PA biczone incorporates the interval 03-SC-24 to 03-SC-36, 30 to 370 metres, of the Sermilik Formation (Section B, Appendix h_2). The PA zone is also found in a 70 metre thick section of Bylot Island Formation strata and basal Lithotype 2 strata (Section D, Appendix λ_2), which occurs directly adjacent to and overlying the stratotype.

At Twosnout Creek, the FA zone is the interval which spans 05-TC-720 to 09-TC-109, 220 to 420 metres (Section A, Appendix A_1) of the Bylot Island Formation. This portion of strats is designated the parastratotype for the PA biozone.

Age

Several biostratigraphically important Late Cretaceous and younger palynomorphs first occur in the PA zone. These include: <u>Radialisporis radiatus</u> (Krutzsch) Krutzsch (late Campanian to Paleocene), <u>Hamulatisporis amplus</u> Stanley (Maastrichtian), <u>Kodehouseia gracile</u> (late Campanian to late Maastrichtian), <u>Komipites wyomingensis</u> (Maastrichtian to Paleocene), <u>Polywstibulopolenites trinus</u> (Stanley) Norris (Maastrichtian to Miocene), <u>Forosipollis porosus</u> (Mchedlishvili) Krutzsch (late Campanian to late Maastrichtian), <u>Falaeoperidinium kozlowskii</u> (Ehrenberg) Davey (late Campanian to mid Maastrichtian) and <u>Cerstiopsis</u> diebelii (Alberti) Vozzhennikova (Maastrichtian to Danian).

of these species, <u>Porosipollis porosus</u> (Mchedlishvili) Krutzsch is not known to range above the mid Maastrichtian of the Horton River Section (McIntyre, 1974), and mid Maastrichtian strata of Banks Island (Doerenkamp <u>et al.</u>, 1976). In the western Canadian Arctic, this species is reported from mid or late Maastrichtian strata of Police Island (Sweet et al., 1989).

Palaeoperidinium kozlowskii (Ehrenberg) Davey first appears in the late Campanian of Horton River (McIntyre, 1974) and Banks Island (Doerenkamp <u>et al.</u>, 1976), and ranges no higher than early to mid Maastrichtian in either of these areas. Extensive compilations by Wilson (1971), Bujak and Williams (1978), McIntyre (1974; 1975), May (1980) and Williams and Bujak (1985) indicate that <u>Ceratiopsis diebelii</u> marks the base of the Maastrichtian. The first occurrence of <u>Ceratiopsis diebelii</u> is subsequently used to indicate Maastrichtian rocks on Bylot Island.

The ranges of taxa in the PA zone suggest a very late Campanian to mid Maastrichtian age. The PA biozone correlates with microfloral division H3 of the CR 16B N-68 Section (late Campanian to Maastrichtian) at Horton River (MCIntyre, 1974), the (CVIc) subzone of the <u>Expressipollis</u>, <u>Orbiculapollis</u>, <u>Deflandrea</u> <u>biapertura</u> (CVI) zone (late Campanian to mid Maastrichtian) at Banks Island (Doerenkamp <u>at al.</u>, 1976), the <u>Kodehouseia gracile / Atuilapollenites</u> <u>parallelus</u> (mid Maastrichtian) and the <u>Porcesipollis</u> porceus

zones (mid or late Maastrichtian) at Police Island (Sweet <u>et</u> <u>al.</u>, 1989).

4.1.3 Singularia aculeata - Pesavis parva (SP) zone

Species restricted to this biozone include: Appendicisporites erdtmanii Pocock, Asbeckiasporites wirthi von der Brelie, Extratriporopollenites sp. 2 of McIntyre, Singularia aculeata Samoilovitch, Mancicorpus trapeziforme Mchedlishvili, Paraalnipollenites alterniporus (Simpson) Srivastava, <u>Resavis parva</u> Kalgutkar and Sweet and Trudopollis variabilis Tschudy. Species characterizing the SP zone but overlapping with other zones, include: Hamulatisporis mappus Stanley, <u>Radialisporis radiatus</u> (Krutzsch) Krutzsch, <u>Hazaria sheopiarii</u> Srivastava, Memipies wyomingensis Nicholls and Ott, <u>Triatriopollenites</u> rurensis Pflug and Thomson, <u>Carpinipites ancipites</u> (Modehouse) Srivastava and <u>Ceratiopsis diebelii</u> (Alberti) Vozzhennikova.

New species and unknown taxa which occur in this assemblage zone and are locally restricted to strata of Eclipse Trough are: <u>Lycopodiacidites</u> sp. 1, <u>Pityosporites</u> sp. 1, 2 and 3, <u>Annulispora</u> sp. 1, <u>Cicatricosisporites</u> sp. 1, <u>Cycadopites</u> sp. 1, <u>Podocarpidites</u> sp. 1, <u>Aquilapollenites</u> sp. 1, <u>Kodehouseia</u> sp. 1, <u>Tricolpites</u> sp. 1, and Unknown Specimen 2. Of these palynomorphs, <u>Podocarpidites</u> sp. 1, <u>Aquilapollenites</u> sp. 1, <u>Modehouseia</u> sp. 1, <u>Tricolpites</u> sp. Pristinuspollenites sp. cf. P. inchoatus, Hamulatisporis
pristinuspollenites sp. cf. C. striata and
Unknown Specimen 2 are restricted to the zone. Species that
are biostratigraphically important, but not definitive of
the SP zone include: <u>Cingulatisportes dakotaensis</u> Stanley,
Pseudoplicapollis longiannulata Christopher,
Polyatricpollenites stellatus (Potonie) Pflug,
Polyvestibulopollenites trinus (Stanley) Norris,
Tetraporites sp. 1 of McIntyre and Trudopollis conrector
Pflug.

Distribution

The stratotype for the SP biozone incorporates the interval which spans 11-SC-201 to 14-SC-209, 370 to 500 metres of the Sermilik Formation (Section B, Appendix λ_2). At Twosnout Creek, the SP zone is equivalent to the interval which spans 10-TC-116 to 13-TC-114, 420 to 540 metres (Section A, Appendix λ_1) of the Bylot Island Formation, and 01-TC-142 to 03-TC-156, 0 to 82 metres, (Section C, Appendix λ_3) of the Bylot Island Formation and Lithotype 3 of the Sermilik Formation. The former (Section A, Appendix λ_1) is designated as a parastratotype for the SP biozone.

Age

Biostratigraphicelly important palynomorphs making their first appearance in the SP zone include: Extratriporopollenites sp. 2 of McIntyre (early to mid Maastrichtian), <u>Singularia aculeata</u> Samoilovitch (late Campanian to Paleocene), <u>Mancicorpus trapeziforme</u> Mchedlishvili (late Campanian to Maastrichtian), <u>Paraalnipollenites alterniporus</u> (Simpson) Srivastava (late Maastrichtian to Eocene), <u>Pesavis parva</u> Kalgutkar and Sweet (late Maastrichtian to early Paleocene) and <u>Trudopollis</u> variabilis Tschudy (late Campanian to mid Maastrichtian).

Of these species, Pesavis parva Kalgutkar and Sweet is not known to range below the late Maastrichtian in Arctic Canada (Kalgutkar and Sweet, 1988). Similarly, Paraalnipollenites alterniporus (Simpson) Srivastava is not known to range below the late Maastrichtian of northwestern Canada (McIntyre, 1974). This species is also not known to range below late Maastrichtian strata of Police Tsland (Sweet et al., 1989). Late Cretaceous taxa that are not known to range into the Paleocene, include: Wodehouseia gracile (late Campanian to late Maastrichtian). Hamulatisporis amplus (Maastrichtian), Mancicorpus trapeziforme Mchedlishvili (late Campanian to Maastrichtian) and Trudopollis variabilis Tschudy (late Campanian to mid Maastrichtian). The continued presence of Ceratiopsis diebelii (Alberti) Vozzhennikova also supports a Maastrichtian age for the SP biozone.

The ranges of the taxa within the SP zone strongly suggest an early late Maastrichtian age for the strata. The SP zone approximately correlates with microfloral division H3 of Section CR 17A N-68 (mid to late Maastrichtian), of Horton River (McIntyre, 1974), and the "<u>Deflendrea</u>" <u>diebelii, Palaeoperidinium pyrophorum</u> (CVII) zone (late Maastrichtian) of Banks Island (Doerenkamp <u>et al.</u>, 1976).

4.2 Environmental Interpretation

Byam Martin Formation

Based on sedimentological characteristics and abundant marine palynomorphs, the depositional environment is a basin plain setting, with medium-grained sandstones representing distal turbidites.

Criteria which characterize a basin plain setting have only been formulated since the dual model of turbidite fan and basin plain was introduced (Mutti and Ricci-Lucchi, 1972 in Reading, 1978). A number of features observed in this study are consistent with criteria outlined by Walker (1967), Mutti and Ricci-Lucchi (1972), Walker and Mutti (1973), and Reading (1978) for a basin plain environment. These features include: (a) well developed mudstone layers with a low sand/mud ratio; (b) a lack of abrupt lateral and vertical facies changes; (c) regular bedding with alternating sandstone and mudstone beds; (d) no distinct vertical thickening or thinning sequences; (e) good lateral continuity; (f) a dominantly thin bed size for sandstones, as noted by the preponderance of centimetre to decimetre scale bedding; and (g) sharp based sandstones with few scours and no channels. Sandstones which display these characteristics in association with thick successions of mudstone have been termed basin plain turbidites (Nelson and Nilson, 1974). Flame structures and sandstone dykes are all evidence of rapid deposition associated with distal

turbidites (Ayres, 1986; Allen, 1984).

Near the Sermilik glacier, the depositional environment of the Byam Martin Formation is more problematic. The relative isolation from the overlying Sermilik Formation, the absence of other intrastratal facies and its generally poor exposure, hinders a confident assessment. The mixed bioturbated and coarser non-bioturbated beds of one small section indicate intermittent and rapid sedimentation, as might occur seaward of a delta distributary mouth (Young <u>et</u> <u>al.</u>, 1963). Its stratigraphic position (approximately 16 metres below the base of the Sermilik Formation), suggests a transitional environment between typical Byam Martin strata and the marginal marine conditions of the Sermilik Formation. This coarsening of the strata may represent the initial stage of a regressive cycle, as suggested by Miall <u>at</u> <u>al.</u> (1980).

Sermilik Formation

Lithotypes 1 and 2

Lenticular beds with scoured bases, rip-up-clasts, coarse pebbly lag, planar tabular and trough cross-beds and normal to reverse grading indicate a subaqueous origin in channels. Large stacked sets of planar tabular cross-beds are commonly formed in high energy channel-bar complexes (Williams and Rust, 1969; Miall, 1977). Coarsening upward strata proximal to channel margins represent crevases splay

and associated overbank deposits (Harms and Fahnestock, 1965).

Channelized deposits and their inter-channel strata have been examined in many sedimentary environments (eg: Mutti and Ricci Lucchi, 1972; Ricci Lucchi, 1975; Miall, 1978; Putnam and Oliver, 1980; Walker and Cant, 1984; Swift <u>et al.</u>, 1987; McPherson <u>et al.</u>, 1987; and Shanmugan and Moiola, 1988). The combination of sedimentologic characteristics displayed by Lithotypes 1 and 2 suggest deposition in a high energy braided distributary system, such as that described by McPherson <u>et al.</u>, (1987) for a braid delta deposit.

The braid delta model is a relatively new model. It was introduced by McPherson <u>et al.</u>, (1987) to encompass all coarse-grained deltas formed by the progradation of a braided fluvial system into a standing body of water. Coarse-grained deltas are distinguished from normal finegrained deltas by the presence of coarser sand sizes and few facies changes (McPherson <u>et al.</u>, 1987). Another type of coarse-grained delta is a fan-delta. Fan-deltas are formed by the progradation of an alluvial fan into a standing body of water (Holmes, 1965). Hence, the most diagnostic feature distinguishing braid deltas from fan-deltas is the presence of their respective subaerial components. In the absence of these components other diagnostic characteristics are used to distinguish between both types of coarse-grained deltas

(Table 4.1). Based on the lithological characteristics outlined in Chapter 2, and the data presented in Table 4.1, characteristics of the braid delta model are more appropriate in defining the environment for Sermilik strata (Figure 4.2).

Within the study area, it is not clear if a subaerial component is present. Lithotype 1 displays all of the characteristics necessary to define a high energy braided river system. Scattered and abraded dinosaur bones are suggestive of a nearby terrestrial source. However, pelecypod fragments and abundant dinoflagellates suggest a marine depositional setting. Typical shoreline strata (i.e. beaches) are not present in these deposits. McPherson <u>et</u> <u>al.</u>, (1987) indicates that beaches are highly modified by the high rates of coarse sediment input into braid deltas. By channel switching and migration, beach deposits are subject to constant erosion, and may therefore be erased from the record.

Lithologic evidence suggests that Lithotype 1 represents a marginal marine environment within the braid delta system, whereas Lithotype 2 is the offshore equivalent to Lithotype 1. Evidence supporting a subaerial component is not exposed.

Lithotype 3

Channel deposits in the lentils at Twosnout Creek

	Table	4.1		
Generalized	characteristics of	Fan-deltas	and Braid	deltas
(afte	r McPherson et al.,	1987), com	pared with	
	Bylot Islar	nd strata		

Characteristics	Fan-delta	Braid delta	Sermilik Formation	
Subaerial component	Alluvial fan	Braided river	Braided River ?	
Paleocurrents	Semi-radial and complex	Unimodal and simple	Unimodal (Miall et al., 1980)	
Maximum grain size	Boulders and cobbles very common	Boulders and cobbles uncommon	Boulders and cobbles uncommon	
Sorting	Poor, grading uncommon	Moderate-good, grading common	Moderate-good, grading common	
Clast shape	Angular-subrounded	Subrounded-rounded	Sub-angular to well rounded	
Lateral continuity	Low	Moderate-high	Moderate	
Facies changes (vertical and lateral)	Complex, numerous, sharp	Simple, few, gradational	Simple, few, both gradational and sharp	
Soils and oxidation	Common	Uncommon	Not present	



Figure 4.2 Schematic paleogeographic reconstruction of late Cretaceous Eclipse Trough strata, illustrating the depositional setting.

(Lithotype 3) suggest deposition in a submarine fan complex, within the upper- to mid-fan region (Shanmugam and Moiola, 1988). The lack of large-scale slumps or deep canyons argues against the interpretation that these are innermost fan channels (Watts, 1987). Inverse to normally graded conglomeratic units, like beds observed in this study, suggest deposition within the main feeder channels of the upper fan (Walker, 1975; 1977). Comparative marine macrofossils, shark teeth, and petrological data suggest that Litnotype 3, is in part, laterally equivalent with Lithotype 2.

Bylot Island Formation

The characteristics of this unit at Twosnout Creek (Section A, Appendix A₁) indicate deposition within a basin plain environment similar to that inferred for the Byam Martin Formation. The fine grain size shows deposition was predominantly by suspension (Arndorfer, 1973; and Reineck and Singh, 1973). Subordinate, medium-grained sandstones preserverd within the Bylot island Formation, are typical of distal turbidites.

Bylot Island Formation strata on the south coast are strongly bioturbated where they interfinger with the Sermilik Formation. Bioturbated mudstone suggests that sediments were rich in nutrients (Young <u>et al.</u>, 1983) and oxygenated (Ekdale <u>et al.</u>, (1984). Bioturbation is known to

occur where intrastratal sediment feeders can thoroughly rework the sediment before it is too deeply buried. In the Sacramento Basin (California), bioturbated shales interfingering with and occurring seaward of deltaic distributaries are recognized as lower slope shales (Cherven, 1993). The extent of bioturbation and the proximity to Lithotype 2, suggests that Bylot Island Formation mudstones along the south coast are lower slope sediments that have formed under similar conditions to those of the Sacramento Basin.

4.3 Summary and Conclusions

Upper Cretaceous strata on Bylot Island are lithologically distinct from formations of the Sverdrup Basin. Additional structural and petrographic evidence suggests that strata in Eclipse Trough were affected by different tectonic regimes and are of different provenance than rocks of the Sverdrup basin. To assign Upper Cretaceous strata of Eclipse Trough to units of the Sverdrup Basin would clearly be a violation of the North American Stratigraphic Code (Article 22a). As a result, strata within Eclipse Trough have been divided into three new, lithologically distinct formations: the Byam Martin Formation, the Sermilik Formation and the Bylot Island Formation. Three informal palynomorph assemblage zones of mid Campanian to late Maastrichtian age provide important intrabasinal correlations and constrains the timing for the initiation of basin fill within Eclipse Trough.

As a result, a revised stratigraphic reconstruction for Eclipse Trough is proposed (Figure 4.3). During the late Campanian to early Maastrichtian, a local sediment source (undivided Archean - Aphebian rocks and the Mary River Group) gave rise to the Sermilik Formation through a major regressive event. Several phases of coarsening-upward cycles within the Sermilik Formation imply a complex history of basin fill, with both sea level fall and source terrane uplift intricately related. Falynomorph assemblages suggest that this regressive event was terminated by a late Maastrichtian transgression, represented by the Bylot Island Formation.





REFERENCES CITED

- Allen, J.R.L. (1984) Developments in Sedimentology 30: Sedimentary Structures, their Characteristics and Physical Basis. Elsevier, New York, 663 p..
- Arndorfer, D.J. (1973) Discharge Patterns in Two Crevasses of the Mississippi River Delta. Marine Geology, Volume 15, pp. 269-287.
- Ayers, W.B., Jr. (1986) Lacustrine and Fluvial-Deltaic Depositional Systems, Fort Union Formation (Paleocene), Powder River Basin, Wyoming and Montana. The American Association of Petroleum Geologists Bulletin, Volume 70, Number 11, pp. 1651-1673.
- Azema, C., Fauconnier, D. and Viaud, J.M. (1981) Microfossils from the Upper Cretaceous of Vendee (France). Review of Palaeobotany and Palynology, Volume 35, pp. 237-281.
- Balkwill, H.R. (1978) Evolution of the Sverdrup Basin, Arctic Canada. American Association of Petroleum Geologists Bulletin, Volume 62, Number 6, pp. 1004-1028.
- Balkwill, H.R. (1983) Geology of Amund Ringnes, Cornwall and Haig-Thomas Islands, District of Franklin (Parts of 59C, 59F, 69D, 69E). Geological Survey of Canada, Memoir 190, 76 p..
- Balkwill, H.R., Hopkins, W.S. and Wall, J.W. (1982) Geology of Lougheed Island and Nearby Small Islands, District of Franklin (Parts of 69C, 79D). Geological Survey of Canada, Memoir 395, 22 p.
- Balme, B.E. (1964) The Palynological record of Australian Pre-Tertiary Floras. <u>In</u>: Ancient Pacific Ploras; University of Hawaii Press, pp. 49-80.
- Bernier, J.E. (1910) Report on the Dominion Government Expedition to the Northern Waters and Arctic Archipelago of the D.G.S. Arctic. Department of Marine and Fisheries, Ottawa, Canada, 161 p...
- Boland, D.C. (1986) Upper Jurassic (Portlandian) Sedimentology and Palynofacies of Cabo Espichel, Portugal. Unpublished M.Sc. Thesis, Memorial University of Newfoundland, 261 p..
- Bratzeva, G.M. (1967) The Problem of the Tasagainsk Flora with regard to Spore-Pollen Analytical Data. Review of Palaeobotany and Palynology. Volume 2, pp. 119-126.

- Brenner, G.J. (1963) The Spores and Pollen of the Potomac Group of Maryland. Maryland Department of Geology. Mines and Water Resources, Bulletin 27, 215 p.
- Brenner, G.J. (1968) Middle Cretaceous Spores and Pollen from Northeastern Peru. Pollen et Spores, Volume 10, pp. 341-383.
- Brideaux, W.W. and McIntyre, D.J. (1975) Miospores and Microplankton from Aptian-Albian Rocks along Horton River, District of MacKenzie. Geological Survey of Canada, Bulletin 252, 85 p..
- Bujak, J.P. and Williams, G.L. (1978) Cretaceous Palynostratigraphy of Offshore Eastern Canada. Geological Survey of Canada, Bulletin 297, 19 p..
- Burden, E.T. (1982) Lower Cretaceous Terrestrial Palynomorph Biostratigraphy of the McMurray Formation, Northeastern Alberta. Ph.D Thesis, University of Calgary, 422 p..
- Burden, E.T. (1984) Terrestrial Palynomorph Biostratigraphy of the Lower Part of the Mannville Group (Lower Cretaceous), Alberta and Montana. In The Messozio of Middle North America. <u>Edited by</u> D.F. Stott and D.J. Glass. Canadian Society of Petroleum Geologists, Memoir 9, pp. 249-269.
- Burger, D. (1966) Palynology of Uppermost Jurassic and Lowermost Cretaceous Strata in the Eastern Netherlands. Leidse Geologische Mededelingen, Deel 35, pp. 209-276.
- Burger, D. (1973) Spore Zonation and Sedimentary History of the Neocomian, Great Artesian Basin, Queensland. Geological Society of Australia, Special Publication 4, pp. 87-118.
- Burger, D. (1980) Palynology of the Lower Cretaceous in the Surat Basin. Australian Bureau of Mineral Resources, Bulletin 189, 204 p..
- Carver, R.E. (1971) Procedures in Sedimentary Petrology. John Wiley and Sons Inc., New York, 653 p..
- Cherven, V.B. (1983) A Delta-Slope-Submarine Fan Model for Maestrichtian Part of Great Valley Sequence, Sacramento and San Joaquin Basins, California. American Association of Petroleum Geologists Bulletin, Volume 67, Number 5, pp. 772-816.

- Christopher, R.A. (1979) Normapolles and Triporate Pollen Assemblages from the Raritan and Magothy Formations (Upper Cretaceous) of New Jersey. Palynology, Volume 3, pp. 73-121.
- Couper, R.A. (1953) Upper Mesozoic and Cainozoic Plant Microfossils. New Zealand Geological Survey, Paleontological Bulletin 32, pp. 1-77.
- Couper, R.A. (1958) British Mesozoic Microspores and Pollen Grains: A Systematic and Stratigraphic Study. Palaeontographica Abt. B, Volume 103, pp. 75-179.
- Cox, L.R., Newell, N.D., Boyd, D.W., Branson, C.C., Raymond, C., Chavan, A., Coogan, A.H., Coulette, D., Fleming, C.A., Haas, F., Hertlein, L.G., Kauffman, A., Nyra, A., LaRocque, A., McAlester, A.L., Moore, R.C., Nuttall, C.P., Perkins, B.F., Puri, H.S., Smith, L.A., Soot-Ryen, T., Stenzel, H.B., Trueman, R.E., Turner, R.D. and Weir, J. (1969) Bivalvia. <u>In Treatise on</u> Invertebrate Paleontology, Part N, Volumes 1-3. <u>Edited</u> by R.C. Moore and C. Teichert. The Geological Society of America, Inc. and The University of Kansas, 1224 p...
- Croxton, C.A. (1980) <u>Aquilapollenites</u> from the Late Cretaceous - Paleocene (?) of Central West Greenland. Grønlands Geologogiske Undersøgelse, Volume 101, pp. 5-27.
- Cwynar, L., Burden, E.T. and McAndrews, J. (1979) An Inexpensive Sieving Method for Concentrating Pollen and Spores from Fine-Grained Sediments. Canadian Journal of Earth Sciences, Volume 16, Number 5, pp. 1115-1120.
- Davis, P. N. (1963) Palynology and Stratigraphy of the Lower Cretaceous Rocks of Northern Wyoming. Unpublished Ph.D Thesis, University of Oklahoma, U.S.A., 199 p.
- de Jersey, N.J. (1962) Triassic Spores and Pollen Grains from Ipswich Coalfield. Queensland Department of Mines, Publication 307, 18 p.
- Dettmann, M.E. (1963) Mesozoic Microfloras from Southeastern Australia. Proceedings of the Royal Society of Victoria, Volume 77, Part 1, pp. 1-138.
- Dettmann, M.E. (1973) Angiospermous Poller from Albian to Turonian Sediments of Eastern Australia. Special Publication 4, Geological Society of Australia, pp. 3-34.

- Doerenkamp, A., Jardine, S. and Moreau, P. (1976) Cretaceous and Tertiary Palynomorph Assemblages from Banks Island and Adjacent Areas (N.W.T.). Bulletin of Canadian Petroleum Geology, Volume 24, Number 3, pp. 372-417.
- Drugg, W.S. (1967) Falynology of the Upper Moreno Formation (Late Cretaceous-Paleocene) Escarpado Canyon, California. Palaeontographica Abt. B, Volume 120, pp. 1-71.
- Ekdale, A.A., Bromley, R.G. and Pemberton, S.G. (1984) Ichnology: The Use of Trace Fossils in Sedimentology and Stratigraphy. Society of Economic Geologists and Mineralogists, Short Course Number 15, 317 p.
- Elsik, W.C. (1968) Palynology of a Paleocene Lignite, Milam County, Texas. I. Morphology and Taxonomy. Pollen et Spores, Volume 10, Number 2, pp. 253-314.
- Elsik, W.C. and Jansonius, J. (1974) New Genera of Paleogene Fungal Spores. Canadian Journal of Botany, Volume 52, pp. 953-958.
- Farabee, M.J., and Canright, J.E. (1986) Stratigraphic Palynology of the Lower Part of the Lance Formation (Maestrichtian) of Wyoming. Palaeontogaphica Abt. B, Volume 199, pp. 1-89.
- Felix, C.J. and Burbridge, P.P. (1973) A Maestrichtian Age Microflora from Arctic Canada. Geoscience and Man, Volume 7, pp. 1-30.
- Fensome, R.A. (1983) Miospores from the Jurassic-Cretaceous Boundary Beds, Aklavik Range, Northwest Territories, Canada: Incorporating Taxonomic Reviews of Several Groups of Mid-to Late Mesozic Miospores. Unpublished Ph.D Thesis, University of Saskatchewan, Saskatoon, 762 p.
- Filatoff, J. (1975) Jurassic Palynology of the Perth Basin, Western Australia. Palaeontographica Abt. B, Volume 154, pp. 1-30.
- Frederikson, N.O., Ager. T.A., and Edwards, L.E. (1988) Palynology of Maastrichtan and Paleocene Rocks, Lower Colville River region, North Slope of Alaska. Canadian Journal of Earth Sciences, Volume 25, pp. 512-527.

- Grenier, H.R. (1963) Malloch Dome and Vicinity, Ellef Ringnes Island. In Geology of the North-Central part of the Arctic Archipelago, Northwest Territories (Operation Franklin). Geological Survey of Canada Memoir 320, pp. 563-570.
- Groot, J.J. and Fenney, J.S. (1960) Plant Microfossils and Age of Nonmarine Cretaceous Sediments of Maryland and Delaware. Micropaleontology, Volume 6, Number 2, pp. 225-236.
- Groot, J.J., Penny, J.S. and Groot, C.R. (1961) Plant Microfossils and Age of the Raritan, Tuscaloosa, and Magothy Formations of the Eastern United States. Palaeontographica Abt. B, Volume 108, pp. 121-140.
- Groot, J.J. and Groot, C.R. (1962) Some Plant Microfossils from the Brightseat Formation (Paleocene) of Maryland. Palaeontographica Abt. B, Volume 111, pp. 161-171.
- Hedberg, H.D. (1976) International Stratigraphic Guide. John Wiley and Sons Inc., New York, 200 p..
- Hedlund, R.W. and Norris, G. (1968) Spores and Pollen Grains from Fredericksburgian (Albian) Strata, Marshall County, Oklahoma. Pollen et Spores, Volume 24, pp. 119-155.
- Herngreen, G.F.W. and Chlonova, A.F. (1981) Cretaceous Microfloral Provinces. Pollen et Spores, Volume 23, Number 3-4, pp. 441-555.
- Hills, L.V. and Wallace, S. (1969) <u>Paraalnipollenites</u>, A New Form Genus from Uppermost Cretaceous and Paleocene Rocks of Arctic Canada and Russia. Geological Survey of Canada, Bulletin 182, pp. 139-145.
- Holmes, A. (1965) Principles of Physical Geology. Thomas Nelson and Sons Ltd., London, England, 1288 p..
- Hopkins, W.S. (1969) Palynology of the Eccene Kitsilano Formation, Southwest British Columbia. Canadian Journal of Botany, Volume 47, pp. 1101-1131.
- Hopkins, W.S. (1971) Palynology of the Lower Cretaceous Isachsen Formation on Melville Island, District of Franklin. Geological Survey of Cane 14, Bulletin 197, pp. 109-133.

- Hopkins, W.S. (1974) Some Spores and Pollen from the Christopher Formation (Albian) of Ellef and Amund Ringnes Island, and Northwestern Melville Island, Canadian Arctic Archipelago. Geological Survey of Canada, Paper 73-12, 39 p.
- Hopkins, W.S. Jr. and Balkvill, H.R. (1973) Description, Palynology and Paleoecology of the Hassel Formation (Cretaceous) on Eastern Ellef Ringnes Island, District of Franklin. Geological Survey of Canada, Paper 72-37, 31 p.-
- Ioannides, N.S. (1986) Dinoflagellate Cysts from Upper Cretaceous - Lower Tertiary Sections, Bylot and Devon Islands, Arctic Archipelago. Geological Survey of Canada, Bulletin 371, 97 p.
- Jackson, G.D. (1969) Reconnaissance of North-Central Baffin Island. In Report of Activities, Part A, Geological Survey of Canada, Faper 69-1, pp. 171-176.
- Jackson, G.D., Davidson, A., and Morgan, W.C. (1975) Geology of the Pond Inlet Map-Area, District of Franklin. Geological Survey of Canada, Paper 74-25, 33 p..
- Jackson, G.D. and Davidson, A. (1975) Bylot Island Map-Area, District of Franklin. Geological Survey of Canada, Paper 74-29, 12 p..
- Jackson, G.D. and Iannelli, T.R. (1981) Rift-Related Cyclic Sedimentation in the Neohelikian Borden Basin, Northarn Baffin Island. In Proterozoic Basins of Canada. <u>Edited</u> <u>by</u> F.H.A. Campbell. Geological Survey of Canada, Paper 81-10, pp. 269-102.
- Jansonius, J. and Hills, L.V. (1976) Genera File of Fossil Spores. Special Publication - Department of Geology, University of Calgary.
- Jansonius, J. (1976) Paleogene Fungal Spores and Fruiting Bodies of the Canadian Arctic. Geoscience and Man, Volume 15, pp. 129-132.
- Jarzen, D.M. (1982) Palynology of Dinosaur Provincial Park (Campanian) Alberta. Syllogeus, Number 38, 69 p..
- Jeletzky, J.A. (1971) Marine Cretaceous Biotic Provinces and Paleogeography of Western and Arctic Canada: Illustrated by a Detailed Survey of Ammonites. Geological Survey of Canada, Paper 71-22, 92 p..

- Jerzykiewicz, T. and Sweet, A.R. (1986) The Cretaceous -Tertiary Boundary in the Central Alberta Foothills. I: Stratigraphy. Canadian Journal of Earth Sciences, Volume 23, pp. 1356-1374.
- Jutard G., and Plauchut, B.P. (1973) Cretaceous and Tertiary Stratigraphy Northern Banks Island. <u>In</u> Proceedings of the Symposium on the Canadian Arctic. <u>Edited by</u> J.D. Aitlen and D.J. Glass. Geological Association of Canada, Saskatoon, pp. 203-219.
- Kalgutkar, R.M. and Sweet, A.R. (1988) Morphology, Taxonomy and Phylogeny of the Fossil Fungal Genus <u>Pesavis</u> from Northwestern Canada. Geological Survey of Canada, Bulletin 379, pp. 117-133.
- Kedeves, M. (1980) Palynological Investigations of Sediments on the Lower Fish Clay, Denmark, II. Acta Mineralogica Petrographica Szeged. Volume 24, Number 2, pp. 355-376.
- Kerr, J.W. (1980) Structural Framework of Lancaster Aulacogen, Arctic Canada. Geological Survey of Canada, Bulletin 319, 24 p.
- Krutzsch, W. (1959) Einige neue Formgattungen und Arten von Sporen und Pollen aus der mitteleuropaischen Oberkreide und dem Tertiar. Palaeontographica Abt. B, Volume 105, pp. 125-157.
- Krutzsch, W. (1969) Taxonomie Syncolp(or)ater Und Morphologisch Benachbarter Pollengattungen Und -Arten (Sporae Dispersae) Aus Der Oberkreide Und Dem Tertiaer. Teil I: Syncolp(or)ate Und Syncolp(or)atoide Pollenforman. Pollen et Spores, Volume 11, Number 2, pp. 397-424.
- Langille, A.B. (1987) Sedimentology and Palynology of Cretaceous and Tertiary Strata, Southwest Baffin Island, Northwest Territories, Canada. Unpublished M.Sc. Thesis, Memorial University of Newfoundland. 163 p.
- Leffingwell, H.A. (1971) Palynology of the Lance (Late Cretacecus) and Fort Union (Paleocene) Formations of the Type Lance area, Wyoming. In Symposium on Palynology of the Late Cretaceous and Early Tertiary. Edited by R.M. Kosanke and A.T. Cross. Geological Society of America, Special Paper 127, pp. 1-64.
- Lentin, J.K. and Williams, G.L. (1975) Fossil Dinoflagellates: Index to Genera and Species (Supplement 1). Canadian Journal of Botany, Volume 53, pp. 2147-2157.

- Lentin, J.K. and Williams, G.L. (1976) A Monograph of Fossil Peridinioid Dinoflagellate Cysts. Bedford Irstitute of Oceanography, Dartmouth, Rep. Ser., Bir. 75-16, 237 p..
- Lentin, J.K. and Williams, G.L. (1977) Fossil Dinoflagellates: Index to Genera and Species, 1977 Edition. Bedford Institute of Oceanography, Report Series BI-R-81-12, 209 p.
- Low, A.P. (1906) The Cruise of the Neptune, 1903-1904. Government Printing Bureau, Ottawa, Canada, 355 p..
- Martin, H.A. and Rouse, G.E. (1966) Palynology of Late Tertiary Sedinents from Queen Charlotte Islands, British Columbia. Canadian Journal of Botany, Volume 44, pp. 171-208.
- May, F.E. (1980) Dinoflagellate cysts of the Gymodiniaceae, Peridiniaceae and Gonyaulacaceae from the Upper Cretaceous Monmouth Group, Atlantic Highlands, New Jersey. Palaeontographia Abt. B, Volume 172, pp. 10-116.
- McBride, E.F. (1963) A Classification of Common Sandstones. Journal of Sedimentary Petrology, Volume 33, pp. 664-669.
- McGregor, D.C. (1965) Triassic, Jurassic, and Lower Cretaceous Spores and Pollen of Arctic Canada. Geological Survey of Canada, Paper 64-55, 32 p...
- McIntyre, D.J. (1974) Palynology of an Upper Cretaceous Section, Horton River, District of Mackenzie, N.W.T.. Geological Survey of Canada, Paper 74-14, 57 p..
- McIntyre, D.J. (1975) Morphologic Changes in <u>Deflandrea</u> from a Campanian Section, District of MacKenzie, N.W.T., Canada. Geoscience and Man, Volume 11, pp. 61-76.
- McIntyre, D.J. (1989) Paleocene Palynoflora from Northern Somerset Island, District of Franklin, N.W.T. In Current Research, Part G, Geological Survey of Canada, Paper 89-1G, pp. 191-197.
- McPherson, J.G., Shanmugam, G. and Moiola, R.J. (1987) Fan-Deltas and Braid-Deltas: Varieties of Coarse Grained Deltas. Geological Society of America Bulletin, Volume 99, Number 3, pp. 331-340.

- Médus, J., Boch, A., Parron, C., Lauverjat, J. and Triat, J.M. (1980) Turonian Normapolles from Portugal and Southern France: Correlations. Review of Palaeobotany and Palynology, Volume 31, pp. 105-153.
- Meneley, R.A., Henao, D. and Merritt, R.K. (1975) The Northwest Margin of the Sverdrup Basin. In Canada-Continental Margins and Offshore Petroleum Exploration. Canadian Society of Petroleum Geologists, Memoir 4, pp. 531-544.
- Miall, A.D. (1975) Post-Paleozoic Geology of Banks, Prince Patrick and Eglinton Islands, Arctic Archipelago. In Canada-Continental Margins and Offshore Petroleum Exploration. Canadian Society of Petroleum Geologists, Memoir 4, pp. 557-587.
- Miall, A.D. (1977) A Review of the Braided-River Depositional Environment. Earth-Science Reviews, Volume 13, pp. 1-62.
- Miall, A.D. (1978) Lithofacies Types and Vertical Profile Models in Braided River Deposits: A Summary. In Fluvial Sedimentology. <u>Edited by</u> A.D. Miall. Canadian Society of Petroleum Geologists, Memoir 5, pp. 537-604.
- Miall, A.D. (1979) Mesozoic-Tertiary Geology of Banks Island, Arctic Canada: The history of an Unstable Craton Margin. Geological Survey of Canada, Memoir 387, 235 p.
- Miall, A.D. (1980) Cyclicity and the Facies Model Concept in Fluvial Deposits. Bulletin of Canadian Petroleum Geology, Volume 28, Number 1, pp. 59-80.
- Miall, A.D. (1986) The Eureka Sound Group (Upper Cretaceous-Oligocene), Canadian Arctic Islands. Bulletin of Canadian Petroleum Geology, Volume 34, pp. 240-270.
- Miall, A.D. (1988) The Eureka Sound Group: Alternative Interpretations of the Stratigraphy and Paleogeographic Evolution - Discussion. Geological Survey of Canada, Paper 88-10, pp. 143-147.
- Miall, A.D., Balkwill, H.R., and Hopkins Jr, W.S. (1980) Cretaceous and Tertiary Sediments of Eclipse Trough, Bylot Island Area, Arctic Canada, and their Regional Setting. Geological Survey of Canada, Paper 79-23, 20 p..

- Mikhelis, A.A. (1981) Normapolles Follen in Cretaceous/Falaeogene Boundary Deposits of the Priazov'ye (Azov Sea Area). Review of Palaeobotany and Palynology, Volume 35, pp. 209-229.
- Mutti, E. and Ricci Lucchi, F. (1972) Turbidites of the Northern Apennines: Introduction to Facies Analysis (English Translation by T.H. Nilsen, 1978). International Geology Review, Volume 20, pp. 125-166.
- Nelson, N.H. and Nilson, T. (1974) Depositional Trends of Modern and Ancient Deep-Sea Fans. In Modern and Ancient Geosynciinal Sedimentation. <u>Edited By</u> R.H. Dott and R.H. Shaver. Special Publication of Society of Economic Paleontologists and Mineralogists, Volume 19, pp. 69-91.

Sec.

- Nichols, D.J. and Ott, H.L. (1978) Biostratigraphy and Evolution of the <u>Monipites-Caryapollenites</u> Lineage in the Early Tertiary in the Wind River Basin, Wyoming. Palynology, Volume 2, pp. 93-112.
- Norris, G. (1967) Spores and Pollen from the Lower Colorado Group (Albian-? Cenomanian) of Central Alberta. Palaeontographica Abt. B, Volume 120, pp. 72-115.
- Norris, G. (1986) Systematic and Stratigraphic Palynology of Eccene to Pliocene Strata in the Imperial Nuktak C-22 Well, MacKenzLe Delta Region, District of MacKenzle, N.W.T. Geological Survey of Canada, Bulletin 140, 89 p...
- Norris, G., Jarzen, D.M. and Awai-Thorne, B.V. (1975) Evolution of the Cretaceous Terrestrial Palynoflora in Western Canada. The Geological Association of Canada, Special Paper 13, pp. 333-364.
- North American Commission on Stratigraphic Nomenclature (1983)) North American Stratigraphic Code. American Association of Petroleum Geologists, Volume 67, No. 5, pp. 841-876.
- Norton, N.J. and Hall, J.W. (1969) Palynology of the Upper Cretaceous and Lover Tertiary in the Type Locality of the Hell Creek Formation, Montana, U.S.A. Palaeontographica Abt. B, Volume 125. pp. 1-64.
- Oltz, D.F. (1969) Numerical analyses of Palynological Data from Cretaceous and Early Tertiary Sediments in East Central Montana, U.S.A. Palaeontographica Abt. B, Volume 128, pp. 90-166.

- Pacltová, B. (1981) The Evolution and Distribution of Normapolles Pollen During the Cenophytic. Review of Palaeobotany and Palynology, Volume 35, pp. 175-208.
- Piel, K.M. (1971) Palynology of Oligocene Sediments from Central British Columbia. Canadian Journal of Botany, Volume 49, pp. 1885-1920.
- Pierce, R.L. (1961) Lower Upper Cretaceous Plant Microfossils from Minnesota. Minnesota Geological Survey, Bulletin 42, 86 p..
- Playford, G. (1971) Palynology of Lower Cretaceous (Swan River) Strata of Saskatchewan and Manitoba. Palaeontology, Volume 14, Part 4, pp. 533-565.
- Pocock, S.A.J. (1962) Microfloral Analysis and Age Determination of Strata at the Jurassic-Cretaceous "Noundary in the Western Canada Plains. Palaeontographica Abt. B. Volume 111, pp. 1-95.
- Potonié, R. (1956) Synopsis der Gattungen der Sporae dispersae. I. Teil: Sporites. Beihefte zum Geologischen Jahrbuch, 23, 103 p..
- Potonié, R. (1958) Synopsis der Gattungen der Sporae dispersae. II. Teil: Sporites (Nachtrage), Saccites, Aletes, Praecolpates, Polyplicates, Monocolpates. Beihefte zum Geologischen Jahrbuch, 31, 114 p..
- Potonié, R. (1960) Synopsis der Gattungen der Sporae dispersae. III. Teil: Nachtrage Sporites, Fortsetung Pollenites mit Generalregister zu Teil I - III. Beihefte zum Geologischen Jahrbuch, 39, 189 p..
- Potonié, R. and Kremp, G. (1954) Die Gattungen der Palaozoischen Sporae und ihre Stratigraphie. Geologisches Jahrbuch, 69, pp. 111-194.
- Putnam, P.E., and Oliver, T.A. (1980) Stratigraphic Traps in Channel Sandstones in the Upper Mannville (Albian) of East-Central Alberta. Bulletin of Canadian Petroleum Geology, Volume 28, Number 4, pp. 489-508.
- Reading, H.G. (1978) Sedimentary Environments and Facies. Elsevier, New York, 569 p..
- Reineck, H.E. and Singh, I.B. (1973) Depositional Environments. Springer-Verlag, Berlin-Heidelberg, 439 p..

- Ricci-Lucchi, F. (1975) Depositional Cycles in Two Turbidite Formations of the Northern Apennines (Italy). Journal of Sedimentary Petrology, Volume 45, pp 3-43.
- Ricketts, B.D. and Sweet, A.R. (1986) Stratigraphy, Sedimentology and Palynology of the Kootenay-Blaimore Transition in Southwestern Alberta and Southeastern British Columbia. Geological Survey of Canada, Paper 84-15, 41 p..
- Rouse, G.E. (1957) The Application of a New Nomenclatural Approach to Upper Cretaceous Plant Microfossils from Western Canada. Canadian Journal of Botany, Volume 35, pp. 349-375.
- Rouse, G.E. (1977) Paleogene Palynomorph Ranges in Western and Northern Canada. American Association of Stratigraphic Palynologists, Contribution Series 5A, pp. 48-65.
- Rouse, G.E., Hopkins, W.S. and Piel, K.M. (1971) Palymology of some Late Cretaceous and Early Tertiary Deposits in British Columbia and adjacent Alberta. <u>In Symposium on Palymology of the Late Cretaceous and Early Tertiary. Edited by R.M. Kosanke and A.T. Cross. Geological Society of America. Special Pager 127. pp. 213-246.</u>
- Rouse, G.E., and Srivastava, S.K. (1972) Palynological Zonation of Cretaceous and Early Tertiary Rocks of the Bonnet Plume Formation. Northeastern Yukon, Canada. Canadian Journal of Earth Sciences, Volume 9, pp. 1163-1179.
- Samoilovitch, S. R. and Mchedlishvili, N.D. (1961) Pollen and Spores of Western Siberia. Trudy. Vsesoyuznyi Nauchno-Issledovatel'skii Geologorazvedochnyi Institut, Leningrad, Volume 177, 659 p. (Unpublished English Translation).
- Samoilovitch, S.R. (1967) Tentative Botanico-Geographical Subdivision of North Asia in the Late Cretaceous Time. Review of Palaeobotany and Palynology, Volume 2, pp. 127-139.
- Schumacker-Lambry, J. (1978) Palynologie du Landenien inferieur (Paleocene) a Gelinden-Overbroek / Belgique. Relations entre les microfossils et le sediment. Univ. Liege, 157 p..
- Scott, L. (1976) Palynology of the Lower Cretaceous Deposits from the Algoa Basin (Republic of South Africa). Pollen et Spores, Volume 18, Number 4, pp. 563-609.

- Shanmugam, G. and Moiola, R.J. (1988) Submarine Fans: Characteristics, Models, Classification, and Reservoir Potential. Earth Science Reviews, Volume 24, pp. 383-428.
- Shaozhi, M. and Norris, G. (1988) Late Cretaceous Early Tertiary Dinoflagellates and Acritarchs from the Kashi Area, Tarim Basin, Xinjiang Province, China. Life Sciences Contributions, Number 150, Royal Ontario Museum, 93 p.
- Singh, C. (1964) Microflora of the Lower Cretaceous Mannville Group, East-Central Alberta. Research Council of Alberta, Bulletin 15, 239 p..
- Singh, C. (1971) Lower Cretaceous Microfloras of the Peace River Area, Northwestern Alberta. Research Council of Alberta, Bulletin 28, 542 p..
- Singh, C. (1983) Cenomanian Microfloras of the Peace River Area, Northwestern Alberta. Alberta Research Council, Bulletin 44, 322 p.
- Souther, J.G. (1963) Geological Traverse across Axel Heiberg Island from Buchanon Lake to Strand Fiord. In Geology of the North-Central part of the Arctic Archipelago, Northwest Territories (Operation Franklin). Geological Survey of Canada, Memoir 320, pp. 426-448.
- Srivastava, S.K. (1966) Upper Cretaceous Microflora (Maestrichtian) from Scollard, Alberta, Canada. Pollen et Spores, Volume 8, Number 3, pp. 497-552.
- Srivastava, S.K. (1967) Palynology of Late Cretaceous Mammal Beds, Scollard, Alberta (Canada). Palaeogrography, Palaeoclimatology, Palaeoecology, Volume 3, pp. 133-150.
- Srivastava, S.K. (1969) Assorted Angiosperm Pollen from the Edmonton Formation (Maestrichtian), Alberta, Canada. Canadian Journal of Botany, Volume 47, pp. 975-989.
- Srivastava, S.K. (1971) Monolete Spores from the Edmonton Formation (Maastrichtian), Alberta (Canada). Review of Palaecobstany and Palynology, Volume 11, pp. 251-265.

9

Srivastava, S.K. (1972) Systematic Description of Spores from the Edmonton Formation (Maestrichtian), Alberta, Canada. Falaeontographica Abt. B. Volume 139, pp. 1-36.

- Srivastava, S.K. (1975) Microspores from the Fredericksburg Group (Albian) of the Southern United States. Paleobiologie Continentale, Volume 6, Number 2, pp. 1-119.
- Srivastava, S.K. (1981) Stratigraphic Ranges of Selected Spores and Pollen from the Fredericksburg Group (Albian) of the Southern United States. Palynology, Volume 5, pp. 1-28.
- Stanley, E.A. (1965) Upper Cretaceous and Paleocene Plant Microfossils and Paleocene Dinoflagellates and Hystrichosphaerids from Northwestern South Dakota. Bulletin of American Paleontology, Volume 49, Number 222, 384 p.
- Stockmarr, J. (1971) Tablets with Spores used in Absolute Pollen Analysis. Pollen et Spores, Volume 13, pp. 615-621.
- Stone, F.J. (1973) Palynology of the Almond Formation (Upper Cretaceous), Rock Springs Uplift, Wyoming. Bulletin of American Paleontology, Volume 64, Number 278, 135 p.
- Stott, D.F. (1969) Ellef Ringnes Island, Canadian Arctic Archipelago. Geological Survey of Canada, Paper 68-16, 44 p..
- Stover, L.E. and Evans, P.R. (1973) Upper Cretaceous-Eocene Spore - Pollen Zonation, Offshore Gippsland Basin, Southeastern Australia. Geological Society of Australia, Special Publication 4, pp. 55-72.
- Sweet, A.R. (1978) Palynology of the Lower Part, Type Section, Tent Island Formation, Yukon Territory. <u>In</u> Current Research, Part B, Geological Survey of Canada, Paper 78-1B, pp. 31-37.
- Sweet, A.R. (1986) The Cretaceous Tertiary Boundary in the Central Alberta Foothills. II: Missopre and Pollen Taxonomy. Canadian Journal of Earth Sciences, Volume 23, pp. 1375-1388.
- Sweet, A.R. and McIntyre, D.J. (1988) Late Turonian Marine and Nonmarine Palynomorphs from the Cardium Formation, North-Central Alberta Foothills, Canada. In Sequences, Stratigraphy, Sedimentology: Surface and Subsurface. <u>Edited by</u> D.P. James and D.A. Leckie. Canadian Society of Petroleum Geologists, Memoir 15, pp. 499-516.

- Sweet, A.R., Ricketts, B.D., Cameron, A.R. and Norris, D.K. (1989) An Integrated Analysis of the Brackett Coal Basin, Northwest Territories. <u>In Current Research</u>, Part G, Geological Survey of Canada, Paper 89-1G, pp. 85-69.
- Svift, D.J.P., Hudelson, P.M., Brenner, R.L. and Thomgson, P. (1987) Shelt Construction in a Foreland Basin: Storm Beds, Shelf Sandbodies, and Shelf-Slope Depositional Sequences in the Upper Cretaceous Mesaverde Group, Book Cliffs, Utah. Sedimentology, Volume 34, pp. 423-457.
- Thomson, P.W. and Pflug, H. (1953) Pollen und Sporen des Mitteleuropäishen Tertiärs. Palaeontographica Abt. B, Volume 94, pp. 1-138.
- Thorsteinsson, R. and Tozer, E.T. (1963) Mesozoic and Tertiary Stratigraphy. In Geology of the North-Central part of the Arctic Archipelago, Northwest Territories (Operation Franklin). Geological Survey of Canada, Memoir 320, pp. 74-95.
- Ting, W.S. (1968) Fossil Pollen Grains of Coniferales from Early Tertiary of Idaho, Nevada and Colorado. Pollen et Spores, Volume 10, Number 3, pp. 557-598.
- Truswell, E.M. (1983) Recycled Cretaceous and Tertiary Pollen and Spores in Antarctic Marine Sediments: A Catalogue. Palaeontographica Abt. B, Volume 186, pp. 121-174.
- Tschudy, B.D. (1969) Species of Aquilapollenites and Fibulapollis from two Upper Cretaceous Localities in Alaska. United States Geological Survey, Professional Paper 643-A, 17 p..
- Tschudy, B.D. (1973) Palynology of the Upper Campanian (Cretaceous), Judith River Formation, North-Central Montana. United States Geological Survey, Professional Paper 770, 42 p..
- Tschudy, R.H. (1975) Normapolles Pollen from the Mississippi Embayment. United States Geological Survey, Professional Paper 865, 42 p.
- von der Brelie, G. (1964) Eine Unterkretazische Mikroflora aus dem nördlichen Sauerland. Fortschritte in der Geologie von Rheinland und Westfalen, Volume 12, pp. 117-168.

- Walker, R.G. (1967) Turbidite Sedimentary Structures and their Relationship to Proximal and Distal Depositional Environments. Journal of Sedimentary Petrology, Volume 37, pp. 25-43.
- Walker, R.G. (1975) Generalized Facies Models for Resedimented Conglomerates of Turbidite Association. Bulletin of the Geological Society of America, Volume 86, pp. 737-748.
- Walker, R.G. (1977) Deposition of Upper Mesozoic Resedimented Conglomerates and Associated Turbidites in Southwestern Oregon. Bulletin of the Geological Society of America, Volume 88, pp. 273-825.
- Walker, R.G. and Mutti, E. (1973) Turbidite Facies and Facies Associations. In Turbidites and Deep Water Sedimentation. Society of Economic Paleontologists and Mineralogists Pacific Section, Short Course, Anaheim, pp. 119-157.
- Walker, R.G. and Cant, D.J. (1984) Sandy Fluvial Systems. In Facies Models (Second Edition). Edited by R.G. Walker. Geoscience Canada, Reprint Series 1, pp. 71-90.
- Watts, K.F. (1987) Triassic Carbonate Submarine Fans along the Arabian Platform Margin, Sumeini Group, Oman. Sedimentology, Volume 34, pp. 43-71.
- Weyland, H. and Krieger, W. (1953) Die Sporen und Pollen der Aachener Kreide und ihre Bedeutung für die Charakterisierung des Mittleren Senons. Palaeentographica Abt. B. Volume 95, pp. 6-29.
- Wiggins, W.G. (1976) Fossil Oculata Pollen from Alaska. Geoscience and Man, Volume 15, pp. 51-76.
- Williams, G.L. and Brideaux, W.W. (1975) Palynologic Analysis of Upper Mesozoic and Cenozoic Rocks of the Grand Banks, Atlantic Continental Margin. Geological Survey of Canada, Bulletin 236, 163 p..
- Williams, G.L. and Bujak, J.P. (1977) Cenozoic Palynostratigraphy of Offshore Eastern Canada. In Contributions of Stratigraphic Palynology, Volume 1, Cenozoic Palynology, Edited py W.E. Elsik. American Association of Stratigraphic Palynologists, Contributions Series, No. 5a, pp. 14-47.
Williams, G.L. and Bujak, J.P. (1985) Mesozoic and Cenozoic Dinoflagellates. In Plankton Stratigraphy. <u>Edited by</u> H.M. Boli, J.B. Saunders and K. Perch-Nielson. Cambridge University Press, Cambridge, pp. 847-964.

- Williams, P.F. and Rust, B.R. (1969) Sedimentology of a Braided River. Journal of Sedimentary Geology, Volume 39, pp. 649-679.
- Wilson, M.A. (1978) Palynology of Three Sections Across the Uppermost Cretaceous/Paleocene Boundary in the Yukon Territory and District of Mackenzie, Canada. Palaeontographica Abt. B, Volume 166, p. 99-183.
- Wilson, L.R. and Webster, R. (1946) Plant Microfossils from the Fort Union Coal of Montana. American Journal of Botany, Volume 33, Number 4, pp. 271-278.
- Wingate, F.H. (1980) Plant Microfossils from the Denton Shale Member of the Bokchito Formation (Lower Cretaceous, Albian) in Southern Oklahoma. Oklahoma Geological Survey, Bulletin 130, 89 p..
- Wingate, F.H. (1983) Palynology and Age of the Elko Formation (Eocene) near Elko, Nevada. Palynology, Volume 7, pp. 92-131.
- Wodehouse, R.P. (1933) Tertiary Pollen-II. The Oil Shales of the Eccene Green River Formation. Bulletin of the Torrey Botanical Club, Volume 60, pp. 479-524.
- Wodehouse, R.P. (1959) Follen Grains. Their Structure, Identification and Significance in Science and Medicine. Hafner Publishing Co., New York, 574 p..
- Wolfe, J.A. (1976) Stratigraphic Distribution of Some Pollen Types from the Campanian and Lower Maestrichtian Rocks (Upper Cretaceous) of the Middle Atlantic States. United States Geological Survey, Special Paper 977, 18 p..
- Young, F.G., Rahmani, R.A. and Pemberton, S.G. (1983) Bioturbation Structures in Clastic Rocks. In Use of Sedimentary Structures for Recognition of Clastic Environments. <u>Edited by M.S. Shawa</u>. Canadian Society of Petroleum Geologist, pp. 41-52.

PLATES

All specimens, except those in plate 1, are illustrated in inturference contrast unless otherwise stated. In the explanation of figures, the species name is followed by a palynology processing number, field sample number, microscope coordinates, magnification and view. Plate 1 specimens (Figures 4, 5 and 6) were identified by D.A. Russell of the National Museum of Natural Science. The aforementioned specimens are housed in collections of the Paleobiology Division, National Museum of Natural Science, Ottawa, and have been assigned the acronym (NMC).

Plate 1

Figure 1. Trace fossils, <u>Chondrites</u> ?; Byam Martin Formation, Twosnout Creek.

Figure 2. Burrows; Byam Martin Formation, South Coast. *Note: Knife is 20cm in length.

Figure 3. Trace fossils, <u>Thalassinoides</u>; Sermilik Formation, Lithotype 2.

Figure 4. Marine lizard vertebrae, Mosasauridae,

(NMC - 40769); Sermilik Formation, Lithotype 2.

Figure 5. Shark Teeth, cf. Scapanorhynchus sp.,

(NMC - 40758); Sermilik Formation, Lithotype 2.

Figure 6. Dinosaur bone (3rd left metatarsal), <u>Hadrosauridae</u>, (NMC - 40757); Sermilik Formation, Lithotype 1.



- Figure 1. <u>Annulispora</u> sp. 1; 420(2), TCG-151, 98.9 19.6, 650X, Mid-focus.
- Figure 2. <u>Cinqutriletes pocockii</u> (Burger) Burden; 399(3), SC-22, 89.9 10.2, 650X, Proximal.

Figure 3. <u>Densoisporites velatus</u> (Weyland and Kreiger) Krasnova; 394(1), SC-33, 105 21, 650X, Proximal.

- Figure 4. <u>Antulsporitos</u> <u>distaverrucosus</u> (Brenner) Archangelsky and Gamerro; 401(1), SC-23, 103.1 15.7, 650X, Distal.
- Figure 5. <u>Cirratriradites teter</u> Norris; 394(4), SC-33, 96.1 12.6, 520X, Proximal.
- Figure 6. Appendicisporites bifurcatus Singh; 399(3),

SC-22, 85 14, 650X, Equatorial, Plane polarized light.

Figure 7. Appendicisporites erdtmanii Pocock; 420(2),

TCG-151, 87.3 15.7, 520X, Distal.

- Figure 8. <u>Camarozonosporites ambigens</u> (Fradkina) Playford; 382(2), SC-30, 89.2 8.4, 650X, Proximal.
- Figure 9. <u>Hamulatisporis amplus</u> stanley; 396(2), TC-105, 102.2 8.7, 650X, Distal.
- Figure 10. <u>Hamulatisporis</u> sp. cf. <u>H. loeblichii</u>; 3⁷5(1), TCG-156, 92 12.1, 416X, Distal.
- Figure 11. <u>Gleicheniidites senonicus</u> Ross; 399(1), SC-22, 94.5 13, 650X, Proximal.
- Figure 12. <u>Gleicheniidites</u> sp. cf. <u>G. circinidites</u>; 399(1), SC-22, 77 4, 650X, Proximal.

- Figure 14. <u>Ornamentifera baculata</u> Singh 421(1), TC-92, 81.8 15.2, 650X, Mid-focus.
- Figure 15. <u>Tappanispora reticulata</u> (Singh) Srivastava; 374(2), TC-85, 84.1 14, 650X, Distal.
- Figure 16. <u>Lvcopodiacidites canaliculatus</u> Singh; 374(1), TC-85, 86.9 15.5, 650X, Distal.
- Figure 17. Lycopodiacidites sp. 1; 371(2), SC-31, 87 19.2, 650X, Proximal.
- Figure 18. <u>Cicatricosisporites australiensis</u> (Cookson) Potonié, 382(1), SC-30, 111.3 11.1, 650X, Equatorial.



Figure 1. Cicatricosisporites sp. 1; 415(1), SC-207,

99.2 12.5, 520X, Oblique.

- Figure 2. <u>Radialisporis radiatus</u> (Krutzsch) Krutzsch; 373(2), TC-98, 99.8 13.1, 650X, Distal.
- Figure 3. <u>Retitriletes</u> <u>austroclavatidites</u> (Cockson) Doring, Krutzsch, Mai, and Schultz; 416(2), TC-114, 92.1 8, 1040X, Distal.
- Figure 4. <u>Foveotriletes</u> <u>subtriangularis</u> Brenner; 420(1), TCG-151, 82 19, 650X, Proximal.
- Figure 5. <u>Stereisporites</u> antiguasporites (Wilson and Webster); 415(1), SC-207, 83.5 13.2, 1040X, Proximal.
- Figure 6. Stereisporites regium (Drozhastchich) Drugg;

371(1), SC-31, 110.9 18.2, 1638X, Proximal.

- Figure 7. <u>Cingulatisporites</u> <u>dakotaensis</u> Stanley; 408(2) SC-115, 89 20.5, 650X, Distal.
- Figure 8. <u>Deltoidospora hallii</u> Miner; 381(4), SC-26, 79.1 13.2, 650X, Proximal.
- Figure 9. <u>Cibotiumspora juncta</u> (Kara-Murza) Singh; 414(2), SC-301, 88.1 14, 1300X, Proximal.
- Figure 10. <u>Cyathidites minor</u> Couper; 370(1), SC-24, 106 10, 1040X, Proximal.
- Figure 11. <u>Gemmatriletes clavatus</u> Brenner; 373(1), TC-98, 101 12, 650X, Oblique.
- Figure 12. <u>Osmundacidites wellmanii</u> Couper; 415(4), SC-207, 96.2 9.1, 650X, Proximal.

- Figure 13. <u>Baculatisporites comaumensis</u> (Cookson) Potonié; 401(3), SC-23, 71 16.9, 650X, Distal.
- Figure 14. <u>Laevigatosporites</u> <u>haardti</u> (Potonié and Venitz) Thomson and Pflug; 394(1), SC-33, 90 8.2, 650X, Equatorial.
- Figure 15. Laevigatosporites sp. cf. L. ovatus; 415(1), SC-207, 102.1 4.1, 1040X, Polar.
- Figure 16. <u>Hazaria sheoparii</u> Srivastava; 408(2), TC-115, 98.1 15.5, 650X, Equatorial.
- Figure 17. Taxodiaceaepollenites hiatus (Potonié) Kremp; 399(3), SC-22, 68.5 18.3, 650X, Mid-focus.
- Figure 18. <u>Taxodiaceaepollenites vacuipites</u> (Wodehouse) Wingate; 374(1), TC-85, 75.5 15, 650X, Mid-focus.



- Figure 1. <u>Podocarpidites multesimus</u> (Bolkovitina) Pocock; 370(2), SC-24, 104.6 10.8, 650X, Proximal.
- Figure 2. <u>Podocarpidites</u> sp. 1; 415(2), SC-207, 83.5 9.2, 416X, Proximal.
- Figure 3. <u>Pityosporites constrictus</u> Singh; 381(4), SC-26, 79.5 12.2, 650X. Obligue.
- Figure 4. <u>Pityosporites alatipollenites</u> (Rouse) Singh; 372(2), SC-36, 98 7, 650X, Equatorial.
- Figure 5. <u>Pityosporites</u> sp. 1; 382(1), SC-30, 83.9 13, 650X, Equatorial.
- Figure 6. <u>Pityosporites</u> sp. 2; 381(2), SC-26, 103.5 5.2, 650X, Distal.
- Figure 7. <u>Pityosporites</u> sp. 3; 394(1), SC-33, 100 10.2, 650X, Equatorial.
- Figure 8. <u>Alisporites bilateralis</u> Rouse; 415(3), SC-207, 108.4 16.2, 1040X, Distal.
- Figure 9. <u>Alisporites grandis</u> (Cookson) Dettmann; 402(1), SC-201, 78.3 7, 416X, Distal.
- Figure 10. <u>Cedripites canadensis</u> Pocock; 382(2), SC-30, 77.8 18.2, 650X, Equatorial.
- Figure 11. Pristinuspollenites sp. cf. P. inchoatus; 415(1), SC-207, 94.2 4.1, 416X, Oblique.
- Figure 12. Lilipollis sp. 1; 370(2), SC-24, 105.9 19.5, 650X, Oblique.

SC-31, 105.9 21.3, 1300X, Equatorial.

- Figure 14. Cycadopites fragilis Singh; 371(1), SC-31, 102.2 19.8, 650X, Proximal.
- Figure 15. <u>Monosulcites</u> sp. 1; 414(2), SC-301, 87 7.8, 650X, Proximal.

Warman and



Figure 1. Entylissa nitidus Balme; 399(2), SC-22,

98.6 16.6, 650X, Oblique.

- Figure 2. <u>Tricolpites</u> sp. 1; 407(2), SC-209, 81.9 12.1, 650X, Equatorial.
- Figure 3. <u>Rousea georgensis</u> (Brenner) Dettmann; 394(3), SC-33, 81.1 13.5, 1040X, Polar.
- Figure 4. <u>Cupuliferoidaepollenites parvulus</u> (Groot and Penney) Dettmann; 374(2), TC-85, 74.3 9.7, 650X, Polar.
- Figure 5. <u>Fraxinoipollenites variabilis</u> Stanley; 401(3), SC-23, 91 12, 1040X. Equatorial.
- Figure 6. <u>Tubifloridites</u> <u>lilliei</u> (Couper) Farabee and Canright; 394(3), SC-33, 83.7 6.8, 1040X, Polar.
- Figure 7. <u>Porosipollis porosus</u> Krutzsch; 372 (1), SC-36, 89 7.9, 1040X, Polar.
- Figure 8. <u>Caprifoliipites longus</u> Stanley; 382(1), SC-30, 88.5 15.5, 1300X. Eguatorial.
- Figure 9. <u>Cranwellia</u> sp. cf. <u>C.</u> <u>striata</u>; 402(2), SC-201, 98.6 16, 1300X, Equatorial.
- Figure 10. <u>Pseudoplicapollis</u> longiannulata Christopher; 401(3), SC-23, 80.9 5.1, 1300X, Polar.
- Figure 11. <u>Pseudoplicapollis</u> sp. G of Christopher, 1979; 402(1), SC-201, 89.9 17.3, 1300X. Polar.
- Figure 12. <u>Pseudoplicapollis</u> sp. G of Christopher, 1979; 401(2), SC-23, 82.9 8.1, 1300X, Polar.

- Figure 13. <u>Carpinipites ancipites</u> (Wodehouse) Srivastava; 416(2), TC-114, 94.2 10, 1300X, Polar.
- Figure 14. <u>Paraalnipollenites alterniporus</u> (Simpson) Srivastava; 408(2), TC-115, 111.1 19, 1638X, Polar.
- Figure 15. <u>Momipites wyomingensis</u> Nichols and Ott; 415(2), SC-207, 98.2 12.4, 1040X, Polar.
- Figure 16. <u>Triatriopollenites rurensis</u> Pflug and Thomson; 404(2), TC-113, 78.3 7.1, 65uX, Polar.
- Figure 17. <u>Trudopollis variabilis</u> Tschudy; 408(2), TC-115, 93 10.2, 1638X, Polar.
- Figure 12. <u>Trudopollis conrector</u> Pflug; 412(1), TC-107, 94 13.2. 1300X. Polar.



- Figure 1. <u>Megatropollis</u> sp. 1; 394(1), SC-33, 76.9 19.6, 1300X, Polar.
- Figure 2. <u>Neotriangulipollis</u> sp. 1 of Azema 1981; 371(11), SC-31, 104 10.3, 1300X, Polar.
- Figure 3. <u>Extratriporopollenites</u> sp. 2 of McIntyre 1974; 402(1), SC-201, 86.9 12.6, 1040X, Polar.
- Figure 4. <u>Polyvestibulopollenites trinus</u> (Stanley) Norris; 408(1), TC-115, 82.8 14.9, 1040X, Polar.
- Figure 5. <u>Polyatriopollenites</u> stellatus (Potonié) Pflug; 401(1), SC-23, 87.5 21, 650X, Polar.
- Figure 6. <u>Tetraporites</u> sp. 1 of McIntyre 1974; 414(1), SC-301, 68 12.5, 650X, Polar.
- Figure 7. Tetraporites sp. 2; 408(1), TC-115, 82 12.4, 650X, Polar.
- Figure 8. <u>Wodehouseia</u> <u>gracile</u> (Samoilovitch) Pokrovaskaya; 370(1), SC-24, 92.3 9.5, 650X, Mid-focus.
- Figure 9. Wodehouseia sp. 1; 415(3), SC-207, 83.3 8, 650X, Mid-focus.
- Figure 10. <u>Singularia aculeata</u> Samoilovitch; 415(1), SC-207, 76 13.8, 1300X, Mid-focus.
- Figure 11. <u>Mancicorpus trapeziforme</u> Mtchedlishvili; 402(2), SC-201, 105.9 13.2, 650X, Oblique.
- Figure 12. <u>Aquilapollenites</u> scabridus Tschudy; 414(2), SC-301, 94.9 12.8, 650X, Equatorial.

Figure 13. Aquilapollenites sp. 1; 402(1), SC-201,

91.2 12.6, 1300X, Equatorial.

- Figure 14. Unknown Specimen 1; 411(1), TC-109, 82 10, 650X, Proximal.
- Figure 15. Unknown Specimen 2; 403(1), TC-115, 86.3 12.4, 650X, Proximal.

Figure 16. <u>Pesavis parva</u> Kalgutkar and Sweet; 416(1), TC-114, 79.2 5.1, 1300X, Mid-focus.

- Figure 17. <u>Ceratiopsis diebelii</u> (Alberti) Vozzhennikova; 402(1), SC-201, 105.2 19, 520X, Oblique.
- Figure 18. <u>Palaeoperidinium kozlowskii</u> (Gorka) Davey; 394(3), SC-33, 87 16.6, 416X, Mid-focus.

A THE ADDRESS TO A THE A THE A



Appendices A₁ - A₄ Stratigraphic Sections

Four complete stratigraphic sections are included for late Cretaceous strata on Bylot Island (Appendices $\lambda_1 - \lambda_2$). Locations for all palynomorph and sedimentology samples are specified on the appropriate sections. Section C, Appendix λ_3 , is occasionally informally referred to as the "Twosnout Creek Gorge (TGG)" section. As well, note that Sections C and D are vertically exaggerated.

Grain size scale: m = mudstone, s = siltstone, sandstone: vf = very fine, f = fine, m = medium, c = coarse, vc = very coarse.









-

-=

ົ

21

--- conformable contact





APPENDIX B.

. PETROLOGY

Methods:

Seventeen samples of sandstones from the Sermilik, Byam Martin and Bylot Island Formations were thin-sectioned for petrographic analysis (Appendices B_{χ} - B_{η}). Unconsolidated beds were rare, comprising less than 10% of strata in examined sections. For this reason, all samples collected were lithified, allowing for the accurate determination of matrix, cement, porosity, roundness, sorting and original texture.

A minimum of 300 point counts per thin section supplied data on the mineralogical composition. This number ensures that for the major constituents, the volume of the observed constituents is between 2.5-6.0% of the actual volume, with a confidence level of 95.4% (Carver, 1971). Sandstone samples were classified according to %CBride (1963). This system is designed to be purely descriptive, with end member constituents being grouped to indicate gross provenance.

Framework grains counted were classified as follows: (1) Quartz - monocrystalline and polycrystalline; (2) Feldspar; and (3) Rock fragments - volcanic, plutonic, metamorphic and sedimentary. Other grains counted included micas (muscovite and biotice), opaques, glauconite and miscellaneous minerals.

Appendix B,

Formation	Sp	Qtz	F	Rf
Hassel	83	96.5	3.1	2.4
Byam Martin	3 91	91.9 95.4	7.7 4.0	0.4 0.4
Bylot Island	93 95 99 111 120	95.3 90.8 89.2 94.8 96.5	4.3 8.8 10.4 4.4 3.1	0.4 0.4 0.4 0.4 0.4
Sermilik	149 153 1 2 11 806 905 51 52	71.0 76.7 91.8 81.3 94.5 87.8 91.6 89.9 79.6	16.7 14.2 5.4 16.6 3.4 10.3 7.1 9.3 15.9	12.3 9.1 2.8 2.1 1.9 1.3 0.8 4.5

Petrology - Detrital Mode Percentages

Sp = sample, Qtz = quartz, F = feldspar, Rf = rock fragments

Appendix 83	 Pointcount Percentages 	number of detrital grains)
	48.	λq
	Petrolo	percent

Formation	sp	do	u o	¥	14	51	E X	200	1	15	d	Ē
Hossel	83	0.3	96.3	2.3	0.6					5	-	0.3
Byan Martin	5	15.0	73.0	4.6	2.6					2.3	1.0	1.3
Svam Martin	16		92.6	1.3	2.6	5		×	1.6	1.6		0.3
Bylot Island	53	0.3	67.6	1.3	2.0		÷		1.6	24.3	2.0	6.0
Bylot Island	56	1.0	77.6	5.0	2.6	ï	x	÷	2.0	10.0	1.0	1.6
Bylot Island	66	1.3	79.0	7.6	1.6				2.3	6.6	0.6	1.0
Bylot Island	111	1.3	91.3	3.0	1.3	0.3	0.3	÷	0.6	1.0	0.6	0.3
Sviot Island	120	1.0	91.6	2.0	1.0		0.3		1.0	1.0	1.6	0.3
Sermilik	671	15.0	53.0	10.6	5.3	11.0	ł	0.0	2.0	0.6	0.6	0.3
Sermilik	153	12.0	61.6	9.6	4.0	8.0	0.6	×	1.6	H	1.0	1.6
Sermilik	-	5.6	84.3	4.4	0.6	2.6	•	•	1.3	•	5.0	0.6
Sermilik	2	3.0	73.6	3.6	12.0	2.0	3		2.0		3.6	ŝ
Sermilik	:	1.3	91.0	3.0	0.3	2.0	•		0.6	2	1.3	0.3
Sermilik	806	1.6	83.0	8.6	1.3	1.6			0.6	ŝ	2.3	6.0
Serailik	506	9.9	76.0	5.0	1.3	1.0	•		1.3		10.6	0.6
Serailik	15	0.6	83.0	2.0	6.6	0.6	x		1.6	1.0	1.6	2.9
Sermilik	52	8.6	69.3	5.6	10.0	2.0	9	ъ	1.6	0.3	1.6	1.0

Sp = Sample, Gp = Polycrystalline quartz, Gm = Monocrystalline quartz, K = Potassium feldspar, Pl = Plagiociase, 1g = 1gneous rock fragments, Mm = Metamorphic rock fragments Sed = Sedimentary rock fragments, Bi = Blotite, Gi = Glauconite, Op = Opaque Brains Mi = Miscellaneous minerals.

T = Irace

Appendix C,

Twosnout Creek

Field Number	Processing Number	Section
TC-114	416	A
TC-115	408	A
TC-113	404	A
TC-116	395	A
TC-109	411	A
TC-107	412	A
TC-105	396	A
TC-102	484	A
TC-720	406	A
TC-98	373	A
TC-92	421	A
TC-89	418	A
TC-85	374	A

TWOSNOUT CREEK GORGE

TCG-156	375	с
TCG-151	420	C
TCG-142	380	с

SOUTH COAST

SC-209	407	В
SC-207	415	В
SC-203	385	В
SC-201	402	В
SC-301	414	D
SC-36	372	В
SC-35	378	B
SC-33	394	В
SC-31	371	В
SC-26	381	B
SC-30	382	В
SC-24	370	В
SC-23	401	В
SC-22	399	В

na janijani zako deseta na v na bitelo ela dana di la bitelo ela dana dana dana dana dana dana dana da			
	na 111210	10	
LexU0.5045at		50	SPI ZONE
Adut Island Reporting		2.3	
lane 18 besidies		101111111111111111111111111111111111111	PHI ZONE
iye fith bracks		1.2 1.2	GAU ZONE
HIRRIT OLD			
		5.0	13% 20KE
wist him henetia		5.0	IFAI ZONE
		12	
iya Mia lanca		1111	IGAU ZOHE
SHERE'S LINES			
Hist is all describings a		50 50 50	(37) 201E

APPENDIX C2

BARE DUAT FOR E2.FEE TROOP STATA







