HEREDITARY, CONTINUOUS, HOMOTOPY, ISOTOPY, PRODUCTIVE AND EXPANSIVE TOPOLOGICAL PROPERTIES

CENTRE FOR NEWFOUNDLAND STUDIES

TOTAL OF 10 PAGES ONLY MAY BE XEROXED

(Without Author's Permission)

MARGARET MARY MOORE







HEREDITARY, CONTINUOUS, HOMOTOPY, ISOTOPY,
PRODUCTIVE AND EXPANSIVE TOPOLOGICAL PROPERTIES

@ Margaret Mary Moore, B.A.(Ed.), B.A.(Hons.)

A Thesis submitted in partial fulfillment of the requirements for the degree of Master of Science

Department of Mathematics and Statistics

Memorial University of Newfoundland

November 1988

Permission has been granted to the National Library of Canada to microfilm this thesis and to lend or sell copies of the film.

The author (copyright owner) has reserved other publication rights, and neither the themis nor extensive extracts from it may be printed or otherwise reproduced without his/her written permission.

L'autorisation a sté accordée à la Bibliothèque, nationale du Canada de microfilmer cette thèse et de prêter ou de vendre des exemplaires du film.

L'auteur (titulaire du droit d'auteur) se réserve les autres droits de publication, n' la thèse in i de long extraits de celle-ci ne doivent être imprimés ou autrement reproduits ans son autorisation écrite.

ISBN 0-315-50442-0

ABSTRACT

Several separation axioms, connectedness and compactness as well as a number of other general topological properties are studied to determine whether or not they are invariant with respect to heredity, closed heredity, open heredity, continuity, open continuity, closed continuity, divisibility, retractions, projections, homotopy and isotopy equivalences, as well as finite, countable, and arbitrary products, and also with, respect to contractions and expansions of the topology of a space. Of the resulting four hundred eighty questions, all but seven tan be answered. The resulting answers, found and discussed in the paper, are also tabulated in several tables.

ACRNOWLEDGMENTS

and the second of the second second second second

I would like to express my gratitude to my supervisor, Dr. S. Thomeier, for his invaluable guidance, encouragement thoroughness and assistance in the preparation of this thesis.

I would also like to acknowledge the encouragement and moral support of Mr. A. B. Fekete and Dr. S. P. Singh and the kindness of Dr. P. Booth during the time of my graduate studies.

I wish to thank the departmental secretaries for their kindness and help.

Finally, I pay tribute to my husband, son and daughter for their great patience and the opportunity to leave home in order to become a student again, and to my father who, during sickness and death, taught me the value of persistence.

TABLE OF CONTENTS

	Introduction				. 4: 1
4:					
	Chapter 0				2 1
					9 1
	DEFINITIONS OF PROPERTI	ES \			. 5
. ,		4			1.
. /					1
	Chapter 1				
	HEREDITARY, ISOTOPY AND	HOMOTOPY	PROPERTIES	5 .	11
			. 1		
	Chapter 2				
				_	
	OPEN, ISOTOPY AND HOMOT	OPY PROPER	TIES \		22
. 1	24			• 11	
	Chapter 3				
,					
	PRODUCTIVE AND EXPANSIV	E PROPERTI	ES		. 33
	Chapter 4				
	APPENDIX ON SOME BASIC	THEOREMS A	ND CORNER	REXAMPLES	26
	APPENDIX ON SOME BASIC	THEUREMS A	AND COUNTER	REXAMPLES	3,6
	· ·				
	Bibliography			4.	58
	ninital and a series				
		,			

LIST OF TABLES

			1		
Table I	3				. 0
TABLE OF PR	OPERTIES-	. 5			4
1,			. •		
m-11- 17			2.5		
Table II				٠.	190
HEREDITARY,	номотору А	ND ISOTOPY	PROPERTI	ES .	21
Table III		F	0.3		2.0
	" a ft		. 1		
CONTINUOUS	PROPERTIES	6		2.7	31
	5 /			~ *	*
Table fy .	. / ``			E	
	1,75	* A	· 1	1.0	
OPEN, ISOTO	PY AND HOMO	TOPY PROPER	TIES		32
		5 8 9		. 1	
Table V			W 10 W		\neg
		2 0 0	7		
PRODUCTIVE	AND EXPANSIV	E PROPERTI	BS .		. 35
	. 1 10		2	7.	
Table VI	A 7				
COUNTEREXAM	ntne non we	-HEREDITAR	v nhanen	MTDE	. 54
COUNTEREXAN	PLES FOR NO	-nakabitak	I PROPER		. 34
_Table VII		3 N N		*	
COUNTEREXAM	PLES FOR NO	-continuop	S PROPER	TIES	5.5
Table VIII	1975	N. 1			
COUNTEREXAN	PLES FOR NO	-CONTRACTI	VE AND N	ON-HOMOTOP	Y
18			E E (4)		
PROPERTIES			4.5		56
)	. g an	*			
Table IX	usi, 14			. 1	
		· \	n		
COUNTEREXAN	PLES FOR NO	-PRODUCTIV	E AND NO	N-EXPANSIV	• •
PROPERTIES	10.00				57

INTRODUCTION

hat the transfer of the state of the same of the same of the state of the same of the same

this paper is the result of a study done on thirty topological properties in order to compile a table of invariances and non-invariances for all those properties with respect to homotopy and isdeepy equivalences and certain constructions such as subspaces, continuous functions, product spaces and expansions or the topology of a space. The resulting four hundred eighty questions are a studied in detail and the answers to all but sever of these questions are given in this paper and are compiled in Table 1 at the end of this introduction.

On the basis of results by S.T. Hu (28, 1961) and D. Gottlieb (23, 1964), we are able to conclude that most of those thirty properties are not homotopy properties but the majority are isotopy properties.

In order to fix our terminology, a definition of each of the thirty topological properties considered is given in Chapter 0.

In chapter 1 the following types of properties for topological spaces are defined, hereditary, closed hereditary, open hereditary, retractive, projective, homotopy and isotopy topological properties. Two theorems

from Hu, Gottlieb's theorem on closed hereditary and isotopy properties and the proof that path connectedness is a homotopy property are given and we show that, for any topological property, P, the following implications hold:

Hereditary *> retractive *> projective*

Hereditary *> closed hereditary and open hereditary

Hereditary *> isotopy

harden and department of the state of the state of the

Closed hereditary > not isotopy > not homotopy

not closed hereditary

Chapter 2 gives the definitions of continuous, open continuous, closed continuous, divisible, contractive and open topological properties as well as Gottlieb's mesults on open, homotopy and isotopy topological properties and we show that the following implications, for any topological property, P, hold:

Continuous divisible retractive projective
Divisible peen continuous and closed continuous
Continuous contractive

Open hereditary and open continuous copen
Open and not hereditary not isotopy not homotopy
Homotopy and open continuous not open hereditary.

In Chapter 3 the invariance under finite, countable,

and arbitrary products and under expansions of the topology of a space are considered, and the following implications hold:

Arbitrarily productive > countably productive

For completeness, Chapter 4 contains the proofs of some basic statements which were found in the literature without proof. Several counterexamples are also given, as well as four tables tabulating those counterexamples.

Table I .

	"-" indicates that the			pe:			is					nt.				
		P	ro	pe:	rt	y	is	no						ıt.		
		T			T	YP	e c	f	11	nv.	ar:	lar	nce	-		
	-1.		Г	Γ	Γ								1.		T	T
	ı	1											П	Arbitrarily productive	Countably productive	2
			hereditary	,			1		. 8	Closed continuous			П	npo	Countably productiv	3
			t t	hereditary	ı				continuous	nu			Н	Pr	0 :	릶
		+	ed	#	١.			П	nu	ti			П	H	ad i	잌
		12	er	ed	nB		V.	Ve	t 1	on	1 v		П	=	M '	90
		t a		er	9	P	#	t I	0	O	t	ρY	N	B	1 .	44
		둫	ed		42	18	ac	e-		ed	ra	t t	8	t	ta t	n s
	Property	Hereditary	Closed	Open	Continuous	pivisible	Retractive	Projective	Open	0.8	Contractive	Tomoropy	Isotopy	P	8 7	Expansiv
	Property	He	5	0 p	00	10	Re	Pr	9	CI	8	ů	13	Ar	0 3	김ᄶ
1	To	+	+	+	=	-	+	+	-	-	-		+	+	+14	
2	Ti .	Ŧ	#	+	Ξ	~	+	+	Ξ	+	-	=			+ +	
3	72	Ŧ	+	+	=	=	+	+	-	-	-	=	+	+	+ +	1:
5	Taya	+	+	+	ļ=.	=	+	+	=	-	=/	=	+		+ +	
6	Ta Ta Va	+	1	+	E	=	+	+	-	-	=	-	+	#	+ +	
7	T.	芒	÷	-	E	E	+	÷	-	-	Ξ	Ξ	1		-1:	
8	Ts à	+	+	+	=	-	+	+	-	+	-	-	+	=	-1-	
9	Regular	+	+	+	=	-	+	+	-	-	-	-	+		+ +	
10	Completely regular	Ŧ	+	+	Ξ	-	+	+	-	-	-	7	+		+ +	Ŀ
11.	Normal .	ᆂ	+	=	Ξ.	=	+	+	-	+	-	-	-	=	- -	4:
3	Completely normal	±	+	+	=	-	+	+	-	+	-	-	+	⇟	+ +	+=
14	Path 'connected	믚	H	-	÷	÷	÷	-	7	÷	÷	-	+		##	
15	Locally connected	恄	-	+	÷	+	+	+	+	+	-	÷	-	-	-17	+
16	Totally disconnected	+	+	+	-	-	+	+	-		-	-	+	+	+ +	
17	Compact	Ε	+	=	+	+	+	+	+	+	+	Ξ	Ξ	÷Ι	+ +	Ε3
18	Lindelöf	1-	+	-	+	t	+	+	+	+	+	-	-	=		1-
19	Locally compact	ᆂ	+	=	=	7	+	+	+	=	-	-	=	=	- +	4=
20	Countably compact 'Paracompact	+=	+	-	+	+	+	÷	+	+	+	=	-	-	=	+=
22	Separable	+=	-	+	Ŧ	+	+	-	7	Ť	+	=	=		7	1
23	Second countable	+	+	+	Ė	÷	+	+	+	-	÷	=	+		+ +	
24	First countable	1+	+	+	4	-	+	+	+	-	-	-	+	-	+ +	-
25	Discrete '	1+	+	+	Ξ	+	+	+	+	+	Ξ	Ξ	+		-1+	
26	Indiscrete	+	+	+	Ŧ	+	+	+	+	+	+	-	+		+ +	
27	Metrizable	<u> +</u>	+	+	-	-	+	+	=	=	=	=	+	=	+ +	1=
	Pixed point	1-	-	-	ы	-	+	+	_	=	=	-	=1	ΞĿ	-1-	1=
29	Contractible	_	-	-		-	+	+				+1	+	+	+ +	

CHAPTER O

Property of the property of th

DEFINITIONS OF PROPERTIES

In this chapter we collect the definitions of basic properties for the convenience of the reader, and to fix our terminology.

0.11 Definition

0.2: Definition:

A topological space (X,T) is a Ti-space if for any a,b,e,X,

Hopen sets U,ye,T, s.t. a,s,U, b,s,V and a,s,V, b,s,U

0.3: Definition:

A topological space (X,T) is a <u>Ta_space</u> if for any a,bex,

dopen sets U,VET s.t. aeU, beV and UnV = \$.

0.4: Definition:

, A topological space (X,T) is a <u>Tak-space</u> if for any a,b∈X, ∃ open sets U,V∈T s.t. a∈U, b∈V and Ū∩V̄ = Ø.

0.5: Definition:

A topological space (X,T) is a T3-space if A is a closed

set in X and b \in X s.t. b \notin A; then \exists open sets \exists U, V \in T s.t. A \subset U, b \in V and \exists \cap V = \not s.

0.6: Definition:

A topological space (X,T) is a $\underline{Ty(x_1-x_2)ace}$ if A is a closed set in X and $b\in X$ s.t. $b\notin A$, then \exists a continuous function f(X,T)=(0,1) s.t. f(A)=0 and $f(\{b\})=1$.

0.7: Definition:

A topological space $\chi(X,T)$ is a $\frac{T_{+}-space}{t}$ if λ,B are disjoint closed sets in X, then Ξ open sets $U,V \in T$ s.t. $\lambda \subset U$, $B \subset V$, and $U \cap V = A$.

0.8: Definition:

A topological space (X,T)' is a $T_{S-BPACO}$ if X and B are separated sets in X (i.e. $\overline{X}\cap B=X\cap \overline{B}=\emptyset$), then \overline{X} U.V.C. T s.t. $X\subset U$, $B\subset V$, and $U\cap V=\emptyset$.

0.9: Definition:

A topological space is a regular space if it is To and T3.

0.10: Definition:

A topological space is a completely regular space if it is τ_{o} and $\tau_{aV_{o}}$.

0.11: Definition:

A topological space is a normal space if it is T and Tu.

.0.12: Definition:

A topological space is a completely normal space if it is $\tau_{_1}$ and $\tau_{_2}$.

0.13: Definition:

A topological space X is a <u>connected space</u> if and only if it has no nontrivial separation, i.e. if U,V are non-empty open sets in X s.t. $U \cup V = X$, then $U \cap V = X$.

0.14: Definition:

A topological space X is a path connected space if it has only one path component, i.e. X is path-connected \iff for any two points a, b \(X, \extstyle \) a path p: \(I \ightarrow \chi \text{x} \) s.t. p(0)=a and p(1)=b.

0.15: Definition:

A topological space X is a <u>locally connected space</u> if for vevery point peX and for every neighbourhood U of p, 3 a connected neighbourhood VCU of p.

0.16: Definition:

A topological space X is a totally disconnected space if for each pair of points p,qeX, A a disconnection UUV of

with peU and qeV.(U U V is a disconnection of X \iff U \neq β , V \neq β , U \cap V = β and U \cup V = X.)

0.17: Definition:

A topological space X is a <u>compact space</u> if and only if every open cover of X has a finite subcover.

0.18: Definition:

A topological space X is a <u>Lindelor space</u> if and only if every open cover of X has a countable subcover.

0.19: Definition:

A topological space X is a <u>locally compact space</u> if for every point pg X, p has at least one compact neighbourhood in X.

. 0.20: Definition:

A topological space X is a <u>countably compact space</u> if and only if each countable open cover of X has a finite subcover.

0.21: Definition:

A topological space X is a <u>paracompact space</u> if and only if it is T₂ and every open cover of X has a locally finite ... open refinement, i.e. a refinement which is a locally finite open cover of X. 0.22: Definition:

A topological space X is a <u>separable space</u> if and only if it contains a countable subset which is dense in X.

0.23: Definition:

A topological space X is a second countable space if and only if its topology has a countable basis.

0.24: Definition:

A topological space X is a <u>first countable space</u> if and only if it has a countable local basis at each of its points.

0.25; Definition:

A topological space X is a <u>discrete space</u> if every subset of X is open.

0.26: Definition:

A topological space X is an <u>indiscrete space</u> if the only open sets are \$\text{g}\$ and \$\times\$.

0.27: Definition:

A topological space X is a <u>metrizable space</u> if and only if \exists a metric $d:X^2 \longrightarrow R$ which induces the topology of X.

0.28: Definition:

A topological space X has the <u>fixed point property</u> if every continuous function $f:X \to X$ is s.t. f(x)=x for at least $x \in X$.

0.29: Definition:

A topological space X is a contractible space if it has the same homotopy type as a singleton point.

0.30: Definition:

A topological space X is a <u>locally contractible space</u> if for every point peX and for every nexableourhood U of peX 3 a neighbourhood VCU of p which is contractible in U.

CHAPTER

making and a water and a state of

HEREDITARY, ISOTOPY AND HOMOTOPY PROPERTIES

This chapter will present theorems by Ru'(28) and Gottlieb (23) which allow us to conclude that sixteen of the thirty properties defined in Chapter 0 are hereditary properties, and hence are isotopy properties, but are not homotopy properties. Seven other properties, but are not homotopy properties. Seven other properties, but are not homotopy properties. Counterexamples, to show that these seven properties are not hereditary properties, will be given in Chapter 4. Three further properties are homotopy properties, and hence are isotopy properties, but are not closed hereditary, and thus not hereditary properties. The remaining four properties, for which a conclusion with respect to hereditary, homotopy, and isotopy properties cannot be dimny from the theorems presented in this chapter, will be considered in chapters, 2 and 4.

1.1: Definition:

Property P of topological spaces is a hereditary property if and only it P is inherited by every subspace of a space which has P.

1.2: Definition:

Property P of topological spaces is a closed hereditary

property if and only if P is inherited by every closed subspace of a space which has P. \checkmark

1.3: Definition:

Property P of topological spaces is an open hereditary property if and only if P is inherited by every open subspace of a space which has P.

1.4: Definition:

A map fix — Y from a given space X into a given space Y is sid to be a homotopy equivalence if and only if there exists a map giv — X from Y into X such that the composed maps gofix — > X and fogiv — Y are homotopic to the identity maps on X and Y respectively. That is, there exist homotopies HiXXI — > X and KiYXI — Y, where I is the closed unit interval, such that

H(x,0) = 1x(x), ∀'x∈X

 $H(x,1) = gf(x), \forall x \in X$

 $K(y,0) = 1_{y}(y), \forall y \in Y$

K(y,1) = fg(y), ∀ y∈ Y

In this case, the map $g:Y \longrightarrow X$ is also a homotopy equivalence.

1.5: Definition:

Property P of topological spaces is a homotopy property if and only if P is inherited by every space that is

homotopically equivalent to a space which has P.

to prevent the second of the s

1.6: Definition:

A map is a continuous function.

147: Definition:

If $f: X \longrightarrow Y$ is an injective map from a space X into x space Y which defines a homeomorphism from X onto f(X), then f is an imbedding of X into Y.

1.8: Definition:

A homotopy h. i. X - Y, (t & 1), is said to be an isotopy if, for each t & I, h. is an imbedding. Two imbeddings f, g: X - Y are said to be isotopic if there exists an isotopy h. X - Y, (t & I), such that h. f = f and h. f = g. An imbedding f: X - Y is said to be an isotopy equivalence if there exists an imbedding g: Y - X such that the composity imbeddings of and f of gare isotopic to the identity imbeddings on X and Y respectively. Two topological spaces X and Y are said to be isotopically equivalent it symbol, X - Y if there exists an isotopy equivalence f: X - Y.

The relation \(\times \) among topological spaces is obviously an equivalence relation. (28, page 168).

1.9: Definition:

Property P of topological spaces is an isotopy property if

and only if P is inherited by every space that is isotopically equivalent to a space which has P.

....

Hence the following implication holds trivially:

P is a homotopy property > P is an isotopy property.

1.10: Definition:

If λ is a subspace of the topological space X and $r: X \longrightarrow \lambda$ is a continuous onto function such that $roi = 1_A$, where i is the inclusion map $i: \lambda \longrightarrow X$, then λ is called a retraction X and the function X is called a retraction of X onto λ .

1.11: Definition:

Property P of topological spaces is a <u>retractive property</u>
if and only if P is inherited by every retract of a space
which has P.

1.12: Definition:

Property P of topological spaces is a <u>projective property</u>
if and only if P is preserved by every projection of an
arbitrary product space which has P.

Since every factor space, X, where is from some arbitrary index set A (not necessarily finite or countably), obtained from a projection map, p, from a

topological space X onto the factor space, is a retract of X, i.e. p o i = 1, , where i is the inclusion map i: $X_i \longrightarrow X$, and every retract, open subspace, and closed subspace is a subspace of X, the following implications for any topological property hold trivially:

Hereditary ⇒ retractive → projective
Hereditary → open hereditary
Hereditary → closed hereditary

1.13: Theorem (Ha's Theorem (28, page 170))

If P is a closed hereditary topological property such that
every singleton apace has P and that there exists a space X
which does not have P, then P is not a homotopy property.

Proof:

Let X be a space which does not have P. Consider the cone, CX, over X which is the quotient space obtained by identifying the top, X x {0}, of the cylinder, X x I, to a single point, v, called the vertex of CX. X is homeomorphic with the bottom, X x {1}, of CX since the inclusion map, i: X → X x {1}, is continuous and bijective, the projection, p: X x {1}, is continuous and p = 1 X x {1}, is a closed subspace of CX since X is a closed subspace of X, {1}, is a closed subspace of X is a closed subspace of X is a closed step in closed. Since P is a closed thereditary property which X does not have, it follows that CX cannot have P. Consider the inclusion map, j: v → CX, and the

projection map, q: CX \longrightarrow v, which is a constant map. Claim: j is a homotopy equivalence, that is, j o q \cong 1, and q o j \cong 1, . Now q o j = 1, \Longrightarrow q o j \cong 1, and j o q = constant map, c: CX \Longrightarrow v \cong CX. Claim: there exists a homotopy, h_i: CX \Longrightarrow CX, such that h_i connects h_o = c with h_i = 1_{CX}. Consider h_i (x,s) = (x,st), \forall s, t \in I. Now h_o (x,s) = c(x,s) = (x,s) = v And h_i (x,s) = 1_C(x,s) = (x,s), \forall s \in I. So CX and the singleton space; $\{v\}$, have the same homotopy type. Hence CX is contractible. Since the singleton space, $\{v\}$, has P and CX does not have P, P is not a homotopy property.

A contrapositive statement of Hu's theorem can be used in order to conclude that homotopy properties are not closed hereditary properties.

1.14: Theorem:(Corollary to Mu's Theorem)

If P is a homotopy topological property such that every singleton space has P and that there exists a space, X, which does not have P, then P is not a closed hereditary property and hence not a hereditary property.

1.15: Theorem:

Hereditary properties are isotopy properties.

Proof: (28, page 171)

Let P be any hereditary topological property of spaces.

Assume that f: X Y is an isotopy equivalence and that
the space, X, has the property P. It suffices to prove that
Y also has P. By definition of an isotopy equivalence,
there exists an imbedding, g: Y X, such that the
computed imbeddings, g o f and f o g, are isotopic to the
identity imbeddings on X and Y respectively. The image, g(Y),
is a subspace of X. Since P is hereditary, this implies that
g(Y) has P. As an imbedding, g is a homeomorphism of Y onto
g(Y). Since P is a topological property and g(Y) has P, it
follows that Y also has P.

Since the singleton space has all the properties defined in Chapter 0 and for each property there exist spaces for which that property falls, then Hu's theorem can be applied to the properties in Chapter 0 whenever necessary.

1.16 Theorem:

"Being ath connected" is a homotopy property.

Proof:

Let $f: X \longrightarrow Y$ be a homotopy equivalence with homotopy inverse $g: Y \longrightarrow X$ and let X be path connected. Let $Y_g: Y_1 \subseteq Y$. We shall show that Ξ a path $\alpha: I \longrightarrow Y$ s.t. $\alpha: Y \subseteq Y_g$ and $\alpha: Y_g: Y_g$. Let $X: Y \times I \longrightarrow Y$ be a homotopy

from f o g to 1_y , 1_x , x_y , y, y = f o g(y) and $x(y,1) = \frac{1}{2}(y), \forall y \in Y. \text{ Let } x \longrightarrow x \text{ be a path a.t. for any } x_0, x_1 \in X, p(0) = x_0 = g(y_0) \text{ and } p(1) = x_1 = g(y_1).$

$$y_{a} = \frac{1}{\sqrt{y_{a}}} \frac{1}{\sqrt{y_{a$$

Define \ll (t) = $\begin{cases} K(y_0, 1-3t), & 0 \le t \le 1/3 \\ f & 0 & p(3t-1), 1/3 \le t \le 2/3 \end{cases}$

 $(K(Y_1, 3t-2), 2/3 \le t \le 1$

od (t) is well defined:

Sings K(y,t), $0 \le t \le 1$, and f o p are continuous maps an ∞ is continuous since ∞ (o) = y_0 and ∞ (1) = y_1 , ∞ is indeed a path in Y connecting y_0 with y_1 , as required.

Hence Y is path connected and 'being path connected' is a homotopy property and hence not a closed hereditary property and not a hereditary property.

1.17. Theorem:(23, page 562)

Let P be a cataged hereditary topological property which
holds on some space X and on X x I, where I is the closed
unit interval. Let S be a subspace of X which does not have
P. Then P is not an isotopy property.

Proof i

Let Y = \{(x,t)/\ 0 \leq t \neq 1, or x \neq 3\} be a subspace of X x x.
Y does not possess P since S x\{\bar{\chi}\} = \{(x,1)/\ x \neq 3\} is closed
in Y and S x\{\bar{\chi}\} is homeomorphic to S which does not have P.
Y is indicpically equivalent to X x I since the maps

i: Y \rightarrow X x I s.t. i(x,t) = (x,t) and

 $j: X \times I \longrightarrow Y$ s.t. j(x,t) = (x,t/2)

are imbeddings since i is the inclusion map of Y into X, $j(X \times I) = \frac{1}{2}(x,t)/0 \le t \le \frac{1}{2}, x \in Y \subset X^2$ is the inclusion map of X x I into Y and inclusion maps are imbeddings. The maps $h_{S^1} \times X \times I \longrightarrow X \times I$ s.t. $h_{S^1}(x,t) = \frac{1}{2}(x,(1+s)t/2)$ and $h_{S^1} \times Y \longrightarrow Y$ s.t. $h_{S^1}(x,t) = \frac{1}{2}(x,(1+s)t/2)$ wre

isotopies since ho = i o j and h = 1 xxI,

 $\kappa_0=j$ of i and $\kappa_1=i_V$, and i,j are imbeddings. Since X x i has P and Y does not have P, P is not an isotopy property.

Since the closed unit interval is T₄, normal, compact, Lindelöf, locally compact, countably compact and paracompact, the Tychonoff plank, T, has all of those properties and contains a subspace which does not have any of those properties, except for local compactness, and T x I has all of those properties, and since I x I is locally compact but the rationals are not, the above theorem can be used to conclude that none of those properties are isotopy properties.

Table II indicates which properties of Chapter O, that can be derived from these theorems, are or are not hereditary, closed hereditary, open hereditary, retractive, projective, homotopy and isotopy properties. The "x" in each line indicates that the proof of invariance of that property is either given in this paper or referenced in the literature. The "+" indicates that the invariance is implied by the "x" on the same line. The "-" indicates that, as a result of Hu's and Gottlieb's theorems, non-invariance is implied by the "x" in the same line.

. Table II

ERREDITARY, HOMOTORY AND ISOTORY PROPERTIES

CI	osed hereditary & not	her	e	11 t	ar	Y	*	no	t isotopy
не	reditary > closed here	dit	aı	Y	*	nc	t	ho	motopy
	reditary - open heredi								T.
He	reditary > retractive	> r	r	16	ct	iv	e		
He	reditary => isotopy				25	Ho	mc	to	py shisotopy
Ho	motopy > not closed he	red	111		v	->	ne	t	hereditary >
	"x" indicates proof	1	1		1	-	<u> </u>	-	
of	invariance is either	1.							
71	ven in this paper or		7						
	und in the literature.		12	-					Reference to
	"+" indicates the		1#	1 2					Kararanca to
14	variance is implied		P	4					
		M	H	hereditary	0	0			proof of "x".
" X	" in the same line.	Hereditary	e		Retractive	rojective			
3.0	"-" indicates the	4	-	6	4	4	2	M	
	n-invariance is	끃	8	٩	2	0	S	8	CN stands for
	plied by "x" in the	0	8	п	H	-	0	4	completely
sa	me line.	1 4	Closed	Open	4	ដ	8	Isotopy	normal.
	Property	m			24	D.	m	H	
1	To .	×	+		+	+	-	+	4.1
2	"Ti	x	+	+	+	+	Σ.	+	44, p.79
3	To	×	+	+	+	+	=	+	27, p.55
-4	T240	×	+	+	+	+	=	+	4.2
- 5	T3.	×	+	+	+	+	=	+	30, p.78
6	T5%	×	+	+	+	+	Ξ	4	30 / p.78
7	T#	†=-	×	-	-	-	=	-	36, p.78
8	T ₅	† ∓	+	+	+	+	=	+	24, p.75(CN=TA
9	Régular	x-		+	+	+	=	+	30, p.78
10	Completely regular	Î.		+	+	+	=	+	30, p.78
11	Normal	<u>-</u> -	×	ا -		-	=	÷	30, p.78
12	Completely normal	×		+	+	+	=	+	4.3
13	Connected	12-	-	 - -	Ξ-	-	×	+	
14	Path connected	+							44, p.229 .
15	Locally connected	ļ=-	÷.			-	X.	+	1.16
				×.		-			4.4
16	Totally disconnected	×.		±.	+	<u>+</u>	=.	+	4.5
	Compact		×.			-	=	Ξ.	27, p.62
18	Lindelöf	!	x		_	1	=	=	30, p.78
19	Locally compact	1_	x	-			=	=	27, p.66
20	Countably compact	1_	x				-	-	30, p.158.
21	Paracompact	1_	x				-	-	56, p.148
22	Separable :			x					41, p.258
23	Second countable	x		+		+	-	+	30, p.78
24	First countable .	x	+	+	+	+	-	+	14, p.59
25	Discrete	×	+	+	+	+	-	Ŧ	4.6
26	Indiscrete	×	+	+	+	+	=	+	4.7
27	Metrizable	×	+	+	+	+	=	7	27, p.98
28	Fixed point	Ť	-	-			-1	-1	
29	Contractible	1=	=	-	-	-	x	Ŧ	27, p.53

CHAPTER 2

OPEN, ISOTOPY AND HOMOTOPY PROPERTIES

This chapter will present theorems by Gottlieb(23)
which allow us to conclude that "being locally connected"
and "being separable" are not isotopy properties and hence,
not homotopy properties. Furthermore, "being locally
contractible" is not a homotopy property. Also,
connectedness, path connectedness and contractibility,
which were found to be not closed hereditary properties,
are not open hereditary properties either.

2.1: Definition:

Property P of topological spaces is a <u>continuous property</u>
if and only if P is preserved by every onto map.

2.2: Definition:

Property P of topological spaces is an <u>open continuous</u>

<u>property</u> if and only if P is preserved by every open, onto

2.3: Definition:

Property P of topological spaces is a <u>closed continuous</u>

<u>property</u> if and only if P is preserved by every closed,
onto map.

2.4: Definition:

A map f: X - YY from a topological space X onto a topological space Y is a <u>quotient map</u> if and only if O C Y is open in X whenever f (0) C X is open in X.

2.5: Definition:

Property P of topological spaces is a <u>divisible property</u> if and only if P is preserved by every quotient map.

2.6: Definition:

Property P of topological spaces is a <u>contractive property</u> if and only if whenever T', T are topologies on a space X, T'C T and (X,T) has P, then (X,T') also has P.

2.7: Definition:

Property P of topological spaces is an <u>open property</u> if and only if P is preserved by every open map and inherited by every open subspace of a space which has P.

2.8: Theorem:

Every retraction is a quotient map.

Proof:

Let $r\colon X\longrightarrow A$ be a retraction from a topological space X onto a retract A of X. Now r is continuous and onto M need to show that a subset $0\subset A$ is open in A whenever $r^{-1}(0)\subset X$ is open in X. Assume that $r^{-1}(0)\subset X$ is open in X.

Define a relation \sim on X s.t. \forall x,y \in X, $x \sim y \Leftrightarrow$ r(x) = r(y). Clearly, \sim is an equivalence relation. Let p: $X \longrightarrow X/\sim$ be the quotient map from X onto the topological space X/\sim , where X/\sim is given the quotient topology by p defined by p(x) = [x], where $[x] = \{y \in X/r(y) = r(x)\}$.

Then the map $q: X/\sim \longrightarrow \lambda$, defined by q([x]) = r(x) for $[x] \in X/\sim$, is well defined because $q \circ p(x) = r(x)$. Since r is onto, then q is onto. We shall show that q is 1-1. Assume that r(a) = r(b) where $r(a), r(b) \in \lambda$. Then by definition of q, q([a]) = q([b]).

Therefore [a] = $\begin{cases} y \in \mathbb{X}/ r(y) = r(a) \end{cases}$ = $\begin{cases} y \in \mathbb{X}/ r(y) = r(b) \end{cases}$ = [b]

Hence q is 1-1. Since q is 1-1 and onto, q is bijective and q o $q^1 = 1_A$. Now, since p is a quotient map, $r^1(0) = p^1(0^i)$ C X is open in X inplies that $0^i = q^1(0)$ C X/ α is open in X/ α . Since q is bijective and continuous it is open, and $q(0^i) = q - q^{-1}(0) = 0$ C λ is open in λ . Therefore r is a quotient map and every retraction is a quotient map.

2.9: Theorem:

If X is a non-empty set and T^I/T are topologies on X s.t. $T^I \subset T$, then the function $f_X^I: (X,T) \longrightarrow (X,T^I)$ is a continuous function.

Proof: Since $T' \subset T$, an open set $0 \subset T'$ is open in T. Hence, $1 \stackrel{!}{\times} (0) \subset (X,T)$ is open in (X,T) whenever $0 \subset (X,T')$ is open in (X,T'). Therefore i, is a continuous function.

2.10: Theorem: Every open, onto map is a quotient map.

Proof: (55 , page 103)

2.11: Theorem:

Every closed, onto map is a quotient map.

Proof: (55 , page 103)

Since projective maps are retractions and since retractions, open, onto maps and closed, onto maps are quotient maps, and quotient maps are continuous functions, and since, whenever a topological space is contractive, a continuous function can be defined from the space with topology T to the space with a coarser topology T, the following implications for any topological property, P, hold trivially:

Continuous >> divisible -> retractive >> projective
Continuous -> open continuous
Continuous -> closed continuous
Continuous -> contractive

Divisible => open continuous and closed continuous

2.12: Theoremi(Gottlieb's Theorem (23, page 564))
Any open property P is not an isotopy property if there
exists a space X and a subspace S such that X x I possesses
P but S does not.

Proof:

Let S and X be chosen as in the hypothesis. Consider the subspace Y of X x I such that

Y = $\frac{5}{2}(x,t)$ 0 \leq t \leq 3/4 or x \in 8 $\frac{3}{2}$.

Now S x (3/4, 1] is open relative to Y and Yould have P if Y had P. But S x (3/4, 1] is also a product space and since S is the image of an open projection map from S x (3/4, 1], we see that S would have P if S x (3/4, 1] had P. Since S $\frac{3}{2}$ does not have P, S x $\frac{3}{2}(3/4, 1]$ does not have P and hence, since-P is open hereditary, Y does not possess P. The fact that Y is isotopically equivalent to X x I can be shown by taking the same maps i,j,h, and k, as we take in theorem 1.17. Thus the Shorem is proven.

2.13: Theorem: (23, page 565)

Let P be an open property such that X has P implies X x I has P. Then P is an isotopy property if and only if P is hereditary.

If P is hereditary, it is an isotopy property by theorem 1.15. If P is not hereditary, there exists a space X enjoying P with a subspace S not possessing P. Since X x has P, all the conditions of the preceding theorem are satisfied and so P is not an isotopy property.

2.14: Theorem: (23, page 564) Open properties are not homotopy propert:

Proof:

Let P be an open property. The singleton space { v} must have P, for there exists a space X with P and the constant map c: X -> {v} is open. Let S be a space without P. S x J, where, J is the open unit interval, cannot have P since S = W (S x J), where W is the projection of S x J onto S. Since T is an open map, S without P implies that S x J does not possess P. Consider C(S), the cone over S. S x J is homeomorphic to an open subset of C(S), so C(S) does not possess P since P is open. Since C(S) is homotopically equivalent to 3v2, P cannot be a homotopy property.

Gottlieb used the method of proof of the above theorem to arrive at a more general result.

2.15: Theorem: (23, page 564) Let P be a property such that (i) the singleton space 5v? has P, (ii) P is an open hereditary property, and (iii) such that there exists a space S where S x J does not possess P, where J is the open unit interval. Then P is not a homotopy property.

Proof

Let P be an open hereditary property such that the singleton space {v} has P. Let S be a space where S x J does not possess P. S x J is homeomorphic to an open subset of C(S), the cone over S. Hence C(S) does not possess P. Since C(S) is homotopically equivalent to 2v2, and 2v3 has P. P is not a homotopy property.

Since X = Comb space (41, page 187) is not locally connected, it is not locally contractible. Then S x J is not locally contractible and the above theorem can be used to prove that 'being locally contractible' is not a homotopy property.

A contrapositive statement to theorem 2.15 will be used in order to conclude that hosotopy properties are not open hereditary properties and is stated as a corollary.

2.16: Theorem:(Corollary to theorem 2.15)

Let P be a homotopy property such that the singleton space $\frac{2}{3}$ has P and such that there exists a space S where S x J does not possess P. Then P is not an open hereditary.

property.

Remark:

The assumption that S x J does not possess P is, in fact, necessary, as the following example shows:

The singleton space [v], the open unit interval J and the cone C([v]) = I, the closed unit interval, are locally contractible. [v] x J is an open subspace of C([v]) and [v] x J is locally contractible. "Being locally contractible is an open hereditary property. Hence C([v]) is locally contractible. C([v]) and [v] are homotopically equivalent. Therefore "being locally contractible" would be a homotopy property, which is not so.

Table III indicates which properties of Chapter 0 are continuous, divisible, retractive, projective, open continuous, closed continuous, and/or contractive properties. The "x" and/or "*" in a line indicate that the proof of invariance of that property is either given in this paper or referenced in the literature and a reference to the proof is given in the last column of the table. The "+" indicates that the invariance is implied by the "x" in the same line by one of the implications stated at the top of the table.

Table IV indicates that, by using theorem 2.12, two of

the thirty properties defined in Chapter 0 are not isotopy properties and hence not homotopy properties. Also three of the properties are not open hereditary and hence not . hereditary properties. The table also indicates that, by using theorem 2.16, three of the properties are not open properties. The table also indicates which of the properties are open continuous properties and which are open properties. The "x" and/or "*" in a line indicate that the proof of invariance of that property is either given in this paper or referenced in the literature and a reference to the proof is given in the last column of the table. The "+" and/or "0" in a line indicates that the invariance is implied by "x" in the same line by one of the implications stated at the top of the table. The "#" in a line indicates that the invariance is either implied by "x" and "*" in the same line or implied by "+" and "*" in the same line by one of the implications stated at the top of the table. (Note: "@" in a line does not imply anything whereas "+" in a line may help to imply "#" in the same line') The "-" in a line indicates that, as a result of theorems 2.12, 2.13, 2.14, 2.15 and 2.16, the non invariance is implied by "#" or

Table III

	CONTINUC	ous	'P	RO	PB	RT	IB	s	1
Co	ntinuous => divisible =>	re	tr	ac	t1	ve	-	P	rojective
Co	ntinuous > open continu	100	18	&	Cl	08	ed	. C	ontinuous
	ntinuous > contractive								and the second
Di	visible > open continu	101	18	&	c1	08	ed	_ c	ontinuous . ;
	ب الله الله الله الله الله الله الله الل						a		
* 7	" & "*" indicate that	1							8 N V
th	e proof of invariance							1	Reference to
18	either given in this	П		1			Closed continuous		
pa	per or found in the	П				10	ē		proof of
11	terature.	2				B	17		Table State
						continuous	12	Contractive	"x"and"*".
	" indicates that the	Continuous	9	Retractive	Projective	4	0	1	
	oof of invariance is	ğ	Divisible	4	4	0	1.	Ü	T2 indicates
10	plied by "x".	1 n	81	a	0	0	P	E H	that
		4	ゼ	1	2	1 1	8	4	reference to
5		ō	5	ě	Ĭ	Open	H	0	proof is in
	Property	0	-	-	_	0	0	Lo	Table II.
1	To			×	+				T2
2	T ₁	Ι.		×	+	1	* .		x-T2 *-41,p19
. 3	T2			x	+	L	L	L.	T2
4	T21/2			x	+			L	T2
5	TS			X.	+	_	·	_	T2
6	T51/2 1	Ш	_	X.	+_	_		L	T2
7	T#:		_	×.	+		*	L.	x-4.9 *-41,p24
8	T5			X.	+		*		x-T2 *-24,p81
9	Regular	Ŀ	_	X.	÷.				T2/
10	Completely regular	Ш	_	×.	+	_			T2 .
1.1	Normal	L		X.	+		*.	ш	x-4.10 *-41,p24
12	Completely normal	_	_	×.	+	_	*	_	x-T2 *-4.11
13	Connected	x		+	+	+	+		27,p78
14	Path connected	×		+	+	+	+	±.	27,p85
15	Locally connected	-	x	+	+	+	+.	Н	56,p200
16	Totally disconnected.	Н	-	x	+	-	-	-	T2
17	Compact	×		+	+_	+	+		27,p62
18	Lindelöf	×	+	+	+	+	+	<u>+</u>	24, p48
19	Locally compact	-	+	×	+	+	-	-	x-3,p19 *-56,p13
20	Countably compact	×	+	+.		7	+	-	30,p158
21	Paracompact	-		×	+	-		+	x-4.12 *-15,p16
22	Separable	X.	+	<u>+</u>	÷-	+	+		24,p48
24	Second countable	\vdash	-	x		-	-		x-T2 *-15,p17 x-T2 *-41,p25
25	First countable Discrete	-	-	<u>×</u> .	+	-	+		
26	Indiscrete	-	×.	+	+				31,p128 4.13
27		x	۲.	÷-	+	7	ᅫ		T2
28	Metrizable Fixed point	-				-	-		29,p20
29	Contractible	-		×	+	-	-4	-+	29,p25
30	Locally contractible	-		Ě-	7-		-	-+	
30	LOCALLY CONTRACTIBLE		-	<u>تح.</u>	Ľ		_	_1	29,p26

Table IV

OPEN, ISOTOPY AND HOMOTOPY PROPERTIES

Op	en hereditary & open c	on	ti	nuc	u	3 1	> <	pen
Op	en > not homotopy							opy ⇒ isotopy
Ho	motopy => not open	0	per	n l	ne i	e	111	ary pot homotopy
								ot open continuous
	en and not hereditary		no	t :	Isc	oto	P	
	& *** indicate proof of		1	Г	Г	Г	F	Reference to
inva	riance is either given	1		ı		ì	1	
in t	his paper or found in		1			ı		proof of
	literature.	1	P4			1	-	
"+"8	"@"indicate invariance	1	tary	continuous	1	1		"x" and "*".
is i	mplied by "x".	1	4	1 a			1	1 -
	indicates invariance	l M	hered	1 5	l			
is i	mplied by "x"&"*" or	1	9	4		1		T2&T3 indicate
++ 5	***.	4	9	. 0		10	M	that reference
	indicates that non-	Hereditary	ч			Homotopy	Isotopy	to proof is in
inva	riance is implied by	0	8	- 4	H	9	#	Table II or
-#"	or "x".	9	Open	Open	Open	Ö	8	Table III.
	Property	1 111	0	0	l٥		1	
1	To	x	+	7		-	9	72
2	Ti	x	+			-	0	T2
3	Ta	×	+			-	0	T2
4	T2/2	×	+	Γ.		=	9	T2
5	T3		+			=	e	T2
. 6	T 31/2	×	+	-	-	=	9.	T2
-7	T +	-		_		1		· · · · · · · · · · · · · · · · · · ·
8	To	x	+			-	9	Т2
9.	Regular	×	+	-	_	-	0	T2
10	Completely regular	x	+		-	-	9	T2
11	Normal ·	-				_		
12	Completely normal	×	+	-		-,	9	Т2
13	Connected	-	-	٠	-	×		*-T3 x-T2
14	Path connected	-	-	•	=	x	0	*-T3 x-T2
15	Locally connected ·	_	x	٠	#	-	=	x-T2 *-T3
16	Totally disconnected	×	+			-	0	T2
17	Compact	-		٠	-		-	тз .
18	Lindelöf			٠				Т3
19	Locally compact	-		٠		_		Т3
20	Countably,compact			•	-	_		T3
21	Paracompact	1		-	-	-	-	
22	Separable	_	×	•	#	=	-	x-T2 *-T3
23	Second countable	×	+		#	=		x-T2 · *-T3
24	First countable		÷			-		x-T2 *-T3
.25	Discrete .	×	+		#	-	0	x-T,2 *-T3
26	Indiscrete	×		•	*		0	x-T2 *-T3
27	Metrizable		+	-			9	T2
	Fixed point	Ĥ	H	-	-	-	Ť	
	Contractible	-	-		-1	x		T2 *
	Locally contractible	-	×		-	-	-	T2

CHAPTER 3

PRODUCTIVE AND ETPANSIVE PROPERTIES

In this chapter we will conclude that fourteen of the properties defined in Chapter 0 are arbitrarily productive properties, eighteen are countably productive properties and twenty-one are finitely productive properties. Also six of the properties are expansive properties, 1. e. are preserved under refinements of the topological structure of a space.

3.1: Definition: +

Property P of topological spaces is an <u>arhitrarily productive property</u> is and only if P is inherited by an uncountable product of spaces all of which have P.

3.2: Definition:

Property P of topological spaces is a <u>countably productive</u>
property if and only if P is inherited by a countableproduct of spaces all of which have P.

3.3: Definition:

Property P of topological spaces is a <u>finitely productive</u>

<u>property</u> if and only if P is inherited by a finite product

of spaces all of which have P.

3.4: Definition:

Property P of topological spaces is a <u>expansive property</u> if and only if whenever T', T are topologies on a space X, T' and (X, T') has P, then (X, T) also has P.

Since a finite product can be viewed as a special case of a countable product and a countable product can be viewed as a special case of an arbitrary product, the following implications for any topological property, P, hold trivially:

Arbitrarily productive >> countably productive ->> finitely productive

Table V indicates which properties of Chapter 0 are arbitrarily productive, countably productive, finitely productive and/or expansive properties. The "x" and the """ in a line indicate that the proof of invariance of that property is either given in this paper or referenced in the literature. The "+" in a line indicates that the invariance is implied by the "x" in the same line.

Table V

PRODUCTIVE AND EXPANSIVE PROPERTIES

Ar	bitrarily productive ⇒					y productive
th is pa	" & "*" indicate that e proof of invariance either given in this per or found in the terature.	productive	Countably productive	productive		Reference to
	". indicates that the oof of invariance is		prod	rodu		proof of
	plied by "x".	Arbitrarily	tably	tely p	Expansive	
	Property	Arbit	Count	Finitely	Expar	
1]	To	x	+	+	•	x-30,p86 *-49,p14
2	TI	x			٠	x-4.14 *-49,p14
3	Ta	×	+	+	*	x-27,p56 *-49,p14
4	T 2 1/2	×	+	+	.*	x-4.15 *-49,p14
5	Т3	×	+	+		30,p86
6	T 31/2		+	+		30,p86-
7	T#	T		-		
8	Tf	1	Ι	1	Γ.	
9	Regular	×	+	+		24,p80
10	Completely regular		+.		-	30,p86
11	Normal	1	1	1	-	
12	Completely normal	1	-	1	-	
	Connected	1×	+	+		27,p79
	Path connected	×		+	-	27,p85
	Locally connected	7=-	1	×	-	41,p186
16	Totally disconnected	×	+	+		x-56,p210 *-4.19
17	Compact	1×	+	+	-	27,p64
	Lindelöf	†=-	-	1		
19	Locally compact	1-	-	×	-	30,p86 or p165
20	Countably compact	1	-	1		
21	Paracompact	 	1	1	-	
22	Separable	1:-	×	+	-	41,p258
23	Second countable	1	×	+		30,p86
24	First countable *	1	÷	+	-	24,p46
25	Discrete	 	 -	×	-	x-4.16 *-4.20
26	Indiscrete	Tx-	+	1	-	4.17
27	Metrizable ,	†	×		-	27,p98
28	Fixed point	+	 - -	ا - -	-	EZY
29	Contractible	+ ×	1=-	1-	-	27,p50
	Locally contractible	1 <u>~</u> _	17	1	L	211930

CHAPTER 4

APPENDIX ON SOME BASIC THEOREMS AND COUNTEREXAMPLES

For completeness, this chapter contains the proofs of a few basic facts for which we could find no explicit proof in the literature. A number of counterexamples are also given, in order to complete the four tables of counterexamples found at the end of the chapter.

4.1: Theorem:

Every subspace of a To-space is a To-space.

Proof:

Let $\{Y, T_y\}$ be a subspace of the T_0 -space $\{X, T\}$. Let $a, b \in Y \subset X$ s.t. $a \neq b$. Since $\{X, T\}$ is a T_0 -space, \exists an open set $U \in T$ s.t. either: $a \in U$, $b \notin U$ or: $a \notin U$, $b \in U$.

By the definition of subspace, Y O U is a Ty-open set.

Hence, either: $a \in Y$, $a \in U \Rightarrow a \in Y \cap U$ $b \in Y$, $b \notin U \Rightarrow b \notin Y \cap U$

or: a ∈ Y, a ¢ U ⇒ a ¢.Y∩U

; PEN PERSON

Therefore (Y,T,) is a To-space

4.2: Theorem:

Every subspace of a T2/2-space is a T2/2-space

Let (Y, T_Y) be a subspace of the T_A(x-space (X, T)).

Let a, b∈ Y ⊂ X s.t. s ≠ b. Since (X, T) is a T_A(x-space, ∃

open sets U, V ∈ T s.t. s ∈ U, b ∈ V and Ū ∩ V̄ = \$.

By the definition of subspace, Y ∩ U and Y ∩ V are T_Y-open,

sets. Hence, a ∈ Y, a ∈ U ⇒ a ∈ Y ∩ U

b ∈ Y, b ∈ V ⇒ b ∈ Y ∩ V

and $(\widehat{Y} \cap \overline{U}) \cap (\widehat{Y} \cap \overline{V}) \subset (\overline{Y} \cap \overline{U}) \cap (\overline{Y} \cap \overline{V}) = \overline{Y} \cap (\overline{U} \cap \overline{V})$ = $\overline{Y} \cap Y$

Thus, $(Y \cap U) \cap (Y \cap V) = \emptyset$. Therefore (Y, T_V) is a T_{2V_0} -space.

4.3: Theorem:

Every subspace of a completely normal ce is a completely normal space.

Proof:

Since a completely normal space is a Tg-space that is Tt, every subspace of a Tg-space is a Tg-space and every subspace of a Tt-space is a Tt-space, then every subspace of a completely normal space is a completely normal space.

4.4: Theorem:

Every open subspace of a locally connected space is a locally connected space.

Let (X,T) be a locally connected space. Let (Y,T_Y) be an open subspace of (X,T). Let $p\in Y$ be arbitrary. Since Y is a neighbourhood of p in X and (X,T) is a locally connected space, \exists a neighbourhood Y of p which is connected and $Y\subset Y$. Therefore (Y,T_Y) is a locally connected space.

4.5: Theorem:

Every subspace of a totally disconnected space is a totally disconnected space.

Proof:

Let (X,Y) be a totally disconnected space. Let (Y,Ty) be a subspace of (X,T). Let a,b G Y C X s.t. a ≠ b.

Since (X,T) is a totally disconnected, ∃ a disconnection

ULIV of X with a G U and b G V.

.By the definition of subspace, Y \bigcap U and Y \bigcap V are Ty-open sets.

Hence, UUV = X ⇒ (Y∩U) ∪ (Y∩V) = Y∩(UUV).

= Y ()

UN = | (X ∩ U) ∩ (X ∩ V) = X ∩ (U ∩ V) = X ∩

 $a \in Y$, $a \in U \Rightarrow a \in Y \cap U \Rightarrow Y \cap U \neq \emptyset$ $b \in Y$, $b \in V \Rightarrow b \in X \cap V \Rightarrow Y \cap V \neq \emptyset$

Therefore $(Y \cap U) \cup (Y \cap V)$ is a disconnection of Y with a $\in Y \cap U$ and $b \in Y \cap V$. Therefore (Y, T_V) is a totally

disconnected space.

4.6: Theorem:

Every subspace of a discrete space is a discrete space.

Proof:

Let (X,T) be a discrete space. Let (Y,T_y) be a subspace of (X,T). Since (X,T) is discrete, every subset of X is an open set and T = set of all subsets of X.

= A: A is a subset of x .

By the definition of subspace, Y \(\Omega\), \(\daggered\) A, \(\daggered\) A \(\daggered\) Ence, \(\tag{T}\), \(\daggered\) \(\daggered\) A is a subset of \(\daggered\).

- set of all subsets of Y.

Therefore (Y,Ty) is discrete.

4.7: Theorem:

Every subspace of an indiscrete space is an indiscrete space.

Proof:

Let (X,T) be an indiscrete space. Let (Y,T_y) be a subspace of (X,T). Since (X,T) is indiscrete, T = {\(\delta_1\)X\}.

By the definition of subspace, Y \(\textit{D}\)\ \delta and Y \(\pi\) \(\text{X}\) are the only Ty-open sets. But \(\delta\)\(\pi\)\(\delta\) and Y \(\pi\)\(\delta\).

Hence Ty = {\(\delta_1\)X\}. Therefore (Y,T_y) is indiscrete.

4.8: Theorem:

Every open subspace of a locally contractible space is a locally contractible space.

Proof:

Let (Y,T_Y) be an open subspace of the locally contractible space (X,T) and let $X\in Y$ be any point. Since Y is a neighbourhood of X in X and (X,T) is locally contractible, \exists a neighbourhood U of X in X such that $U\subset Y$ and U is contractible in Y. Therefore (Y,T_Y) is locally contractible.

4.9: Theorem:

Every retract of a T_{ϕ} -space is a T_{ϕ} -space (Note: In our definition of T_{ϕ} -spaces, T_{ϕ} -spaces are not assumed to be T_{ϕ} .)

Proof:

Let X be a T_q -space. Let $T:X \longrightarrow \lambda$ be a retraction from X onto the retract λ of X. Let $M,N \subset \lambda \subset X$ be disjoint, closed sets. Since X is a T_q -space, \exists open sets $U,V \subset X$ s.b. $U \cap V = \beta$ and $M \subset U$, $N \subset V$. Since λ is a retract of X, $\lambda \cap U$ and $\lambda \cap V$ are disjoint, open sets in λ and $M \subset \lambda$, $M \subset U \Longrightarrow M \subset \lambda \cap U$, $M \subset \lambda$, $M \subset V \Longrightarrow M \subset \lambda \cap V$. Hence λ is a T_u -space.

4.10: Theorem:

"Being a normal space" is a retractive property.

Since a normal space is T_{ij} and T_{ij} and since "being T_{ij} " and "being T_{ij} " are retractive properties, then "being a normal space" is a retractive property.

4.11: Theorem:

"Being a completely normal space" is a closed continuous property.

Proof:

Since a completely normal space is T₅ and T₁ and since
"being T₅" and "being T₁" are closed continuous properties,
then "being a completely normal space" is a closed
continuous property.

4.12: Theorem:

Every retract of a paracompact space is a paracompact space.

Proof:

Since a paracompact space is a T2-space and "being paracompact" is a closed hereditary property and since every retract of a T2-space is closed/(29, page 18), then every retract of a paracompact space is a paracompact space

4.13: Theorem:

"Being an indiscrete space" is a continuous property.

Clearly, a function from an indiscrete space X onto a topological space Y can only be continuous if Y is also indiscrete, therefore "being an indiscrete space" is a continuous property.

4.14 Theorem:

The arbitrary product of T_1 -spaces is a T_1 -space.

Proof:

Let $\{x_i:i\in I\}$ be a collection of T_i -spaces. Let $X=\overline{\mathbb{T}^1}$ X_i be the product space.

Let $p = \langle a_i : i \in I \rangle$ and $q = \langle b_i : i \in I \rangle$ be distinct points in X. Then p and q must differ in at least one coordinate space, say X_{i_0} , i.e. $a_{j_0} \neq b_{j_0}$.

By hypothesis, X_{j0} is a T_{j} -space, hence there exist open subsets U and V of X_{j0} such that $a_{j0} \in U$, $b_{j0} \in V$ and

 $a_j \notin V$, $b_j \notin U$.
By the definition of the product space, the projection

 $\prod_{j_0}^{l}: x \longrightarrow x_{j_0}$ is continuous. Accordingly, $\prod_{j_0}^{l}[U]$ and $\prod_{j_0}^{l}[Y]$ are open sets of X and $P \in \prod_{j_0}^{l}[U]$, $q \in \prod_{j_0}^{l}[U]$. Hence X is also a T_j -space.

4.15: Theorem:

The arbitrary product of T21/2-spaces is a T21/2-space.

Let {X; il & I} be a collection of Tall-spaces.

Fet X = T X be the product space.

let p = <a; i e I > and q = <b; i e I > be distinct notate in X.

Then p and q must differ in at least one coordinate space, say $X_{i_0}^*$, i.e. $a_{i_0}^*$, $b_{i_0}^*$.

by hypothesis, $X_{|j|}$ is a $T_{|j|/2}$ -space, hence there exist disjoint open sets U and V in $X_{|j|}$ such that $u_{|j|} \in V$, $v_{|j|/2} \in V$ and $\overline{U} \cap \overline{V} = \emptyset$.

By the definition of the product/space, the projection $\prod_{i=1}^n x \longrightarrow x_{i_0}^* \text{ is continuous.}$

Accordingly, $\prod_{i=1}^{n} \{U\}$ and $\prod_{i=1}^{n} \{V\}$ are disjoint open sets in X such that $p \in \prod_{i=1}^{n} \{U\}$, $q \in \prod_{i=1}^{n} \{V\}$ and

$$\frac{\Pi_{ij}^{*}(v)}{\Pi_{ij}^{*}(v)} \cap \frac{\Pi_{ij}^{*}(v)}{\Pi_{ij}^{*}(v)} \subset \frac{\Pi_{ij}^{*}(v)}{\Pi_{ij}^{*}(v)}$$

$$= \frac{\Pi_{ij}^{*}(v)}{\Pi_{ij}^{*}(v)} \cap \frac{\Pi_{ij}^{*}(v)}{\Pi_{ij}^{*}(v)}$$

Hence $\overline{\Pi_{j_0}^i[U]} \cap \overline{\Pi_{j_0}^i[V]} = \beta$.
Therefore X is also a T_{2y_0} -space.

4.16: Theorem:

The finite product of discrete spaces is a discrete space.

Proof:

Without loss of generality, let X and Y be discrete spaces. Let $Z = X \times Y = \frac{1}{2}(x,y) / X \in X$, $y \in Y_0^2$ be the product space. By the definition of product space, the projections $p\colon z\longrightarrow x\text{ and }q\colon z\longrightarrow Y\text{ are continuous. Let }\{x_i^2\}\text{ and }\{y_i^2\}$ be any singleton open sets in X and Y respectively. Then $p^{-1}(\{x_i^2\}) = \frac{2}{2}(x_i^2,y_n^2) \dots (x_i^2,y_i^2) \dots \frac{2}{2}\text{ are open sets in Z and } q^{-1}(\{y_i^2\}) \cap q^{-1}(\{y_i^2\}) = \frac{2}{2}(x_i^2,y_i^2) \text{ is any open singleton set in Z. Hence Z is a discrete space.}$

4.17: Theorem:

The arbitrary product of indiscrete spaces is an indiscrete space.

Proof:

Let $\{x_i^*, i \in I_i^2\}$ be a collection of indiscrete spaces. Let $x = \prod_i x_i^*$ be the product space. By the definition of product space, the projection $\prod_{i=1}^{n} x_i \longrightarrow x_i^*$ is an open map. Hence $\beta = \prod_{i=1}^{n} (\beta)$ and $x = \prod_{i=1}^{n} (x_i)$ are open sets of x. $U \subset X$ is open in X implies that $\prod_{i=1}^{n} (U) \subset X_i^*$ is, open in X_i^* and $\prod_{i=1}^{n} (U) = \beta$ or X_i^* .

Therefore U must be p or X and X is an indiscrete space.

4.18: Theorem:

"Being locally contractible" is a finitely productive property.

Proof: (Based on proof of theorem 10.35 in 41, page 185) Let $\{x_i^2\}$, $i=1,2,\ldots,n$, be a finite family of locally contractible topological spaces. Let \mathcal{B}_i be a contractible basis of X_i .

Then $\{B_i \times B_2 \times ... \times B_n / B_i \in \mathcal{B}_i$, i N is a contractible basis of the product space.

4.19: Theorem:

Total disconnectedness is an expansive property.

Let X be a non-empty set and let T^I/T be topologies on X such that $T^I\subset T$ and (X,T^I) is totally disconnected. Since the identity map $I_{X^I}(X,T^I)\longrightarrow (X,T)$ is a closed map, then any set, A, that is closed in T is also closed in T. Also if $A\subset T^I$ is closed in T^I , then X-A is open in T^I and also in T since I_X is also an open map. Hence, (X,T) is a totally disconnected space.

4.20: Theorem:

"Being distrete" is an expansive property.

Proof:

Let X be a set and T', T be topologies on X such that $T' \subset T$ and (X, T') is discrete. Then T' = T and hence (X, T) is also discrete. Thus 'belong discrete' is an expansive property.

4.21: Counterexample:

The fixed point property is not closed hereditary.

Let X = [0,1] and Y = 101 U 11.

Y with the discrete subspace topology is a closed subspace of X. It is well known by the Brouwer fixed point theorem that X has the fixed point property. Y does not have the fixed point property, since the continuous function $f: Y \longrightarrow Y$ s.t. f(0) = 1 and f(1) = 0 is such that for any $y \in Y$, $f(y) \neq y$.

4.22: Counterexample;

The fixed point property is not open acceditary.

Let X = [0,1] and Y = (0,1).

Y is an open subspace of X. X has the fixed point property. Y does not have the fixed point property, since the continuous function f, $Y \longrightarrow Y$ s.t. f(y) = y/2 is such that for any $y \in Y$, $f(y) \neq y$.

4.23: Counterexample:

"Being locally contractible" is not closed hereditary.

Let X = [0,1] and Y = the Cantor set(49, page 57).

X/is locally contractible (26, page 191).

Y is a closed subspace of X and Y is not locally connected since it is totally separated. Hence Y is not locally

4:24: Counterexample:
Completely normal, T₅ and T_{2/2} are not open continuous
topological properties.

Let X be the union of the lines y=0 and y=1 in \mathbb{R}^2 with the usual topology. Let Y be the quotient space of X obtained by identifying each point (x,0), for $x\neq 0$, with the corresponding point (x,1). The resulting projection map $p(X) \to Y$ is continuous and open, but p(0,0) and p(0,1) are distinct points of Y-which do not have disjoint neighbourhoods. Hence Y is not T_2 and thus not $T_{2/2}$, T_3 and completely normal but X possesses all of those properties. (56, page 88, example 13.9(b))

4.25: Counterexample:

Total disconnectedness is not an open continuoum
topological property.

The rationals, Q, with the usual topology is a totally disconnected topological space (41, page 177). The set $\{0,1\}$ with the indiscrete topology is not a totally disconnected topological space. Let $a/b \in Q$ be in lowest terms where $b \neq Q$ and define a/b to be odd if a is an odd integer and even if a is an even integer. Let $f: Q \longrightarrow \{0,1\}$ be defined such that

 $f(a/b) = \begin{cases} 0, & \text{if } a/b \text{ is even} \\ 1, & \text{if } a/b \text{ is odd} \end{cases}$

Then f is continuous since $\{0,1\}$ is indiscrete and f is open since the image of any open set in 0 has to be the set $\{0,1\}$ or \emptyset .

4.26: Counterexample:

T 21/2 is not a closed continuous topological property.

(56, page 87, example 13.9(a)) is an example of a closed continuous image of a $T_{2}y_{2}$ -space which is not a $T_{2}y_{3}$ -space.

4.27: Counterexample:

The fixed point property and "being contractible" are not closed continuous properties and hence not divisible properties.

Let X = [0,1] and Y = S'.

It is well known that [0,1] is contractible and has the fixed point property and S^I is not contractible and does not have the fixed point property. The map $f\colon X\longrightarrow Y$ s.t.

 $f(x) = (\cos 2\pi x, \sin 2\pi x)$ is a closed map (37, page 101).

4.28: Counterexample:

If property P is a \mathbf{T}_0 , \mathbf{T}_1 , \mathbf{T}_2 , $\mathbf{T}_{A_{1}N_2}$, regular, completely regular, normal, completely normal, totally disconnected, discrete or metrizable topological property, then P is not a contractive property.

Let X be a finite set with at least two points and T^{\prime} , T be

the indiscrete and discrete topologies on X respectively. $T' \subset T$. Then, by (49, pages 170-171), (X,T) has all of the above properties and (X,T') has none of those.

4.29: Counterexample:

If property P is a T3, T3, T4, T5 or peracompact topological property, then P is not property property

Let X be an uncountable set . Let T be the discrete topology on X. Let T be the countable complement appology on X. T \subset T. Then, by (49, pages 170-171), (X,T) has all of the above properties and (X,T because of those.

4.30: Counterexample:

The fixed point property is not a contractive or expansive topological property.

[0,1] with the usual topology has the fixed point property but if it is given the discrete or the indiscrete topology it does not have the fixed point property for let [0,1] — 1[0,1] be the function such that

$$f(x) = \begin{cases} x + \frac{1}{2}, & 0 \le x \le \frac{1}{2} \\ x - \frac{1}{2}, & \frac{1}{2} < x < 1 \end{cases}$$

Then f is continuous if the domain is discrete or the range is indiscrete and $f(x) \neq x$ for any $x \in [0,1]$.

4.31: Counterexample:

"Being locally contractible" is not a contractive topological property.

The set of rationals, Q, with the usual topology is not locally connected and hence, not locally contractible. Q with the discrete topology is locally contractible.

4.32: Counterexample:

The fixed point property is not an isotopy property and hence not a homotopy property.

[0,1] is isotopic to (0,1).(28, page 191; proposition 3.3)
[0,1] has the fixed point property and (0,1) does not.

4.33: Counterexample:

"Being locally contractible" is not an isotopy property.

(22, page 44)

Let X be the set of all points in the plane composing the square $\frac{1}{2}(x,y)$ / $0 \le x \le 1$, $0 \le y \le 1$.

Let $Y = X \cup \{(x,y) / 1 \le y \le 2, x \text{ is rational and } 0 \le x \le 1\}$. X is locally contractible and Y is not locally contractible

as any picture will show. X is isotopically equivalent to Y:

 $f: X \longrightarrow Y: (x,y) \longrightarrow (x,y)$

g: Y → X: (x,y) → (x,y/2)

 $h_{t}: X \longrightarrow X: (x,y) \longrightarrow (x,y/(2-t))$

 $k_{\perp}: Y \longrightarrow Y: (x,y) \longrightarrow (x,y/(2-t))$

H: X x I \longrightarrow X: (x,y,t) \longrightarrow h_t(x,y) = (x,y/(2-t))

 $K: Y \times I \longrightarrow Y: (x,y,t) \longrightarrow k_{t}(x,y) = (x,y/(2-t))$

Now all functions defined above are continuous since $P: \mathbb{R}^{A'} \times I \longrightarrow \mathbb{R}^{A} \times I: (x,y,t) \longrightarrow (x,y/(2-t)) \text{ is continuous since } \Pi_i \text{ o P and } \Pi_{A} \text{ o P are both continuous where } \Pi_i \text{ and } \Pi_{A} \text{ are the projections of } \mathbb{R}^{A} \longrightarrow \mathbb{R}. \text{ Then } f, g, h_f, \text{ and } k_f \text{ are imbeddings since they are clearly 1-1 and ogen on their images. By inspection fog = k_and l_y = k_i, g of = h_0 and l_X = h_i, hence X and Y are homotopically equivalent.}$

4.34: Counterexample:

"Being locally contractible" is not a countably productive topological property.

The Cantor set is not locally contractible since it is not locally connected and it is the contable product of locally contractible spaces. $\{0,2\}^N$ where $\{0,2\}$ is given the discrete topology, is homeomorphic to the Cantor set)

4.35; Counterexample:

Being contractible is not an expansive topological property.

The set, R, of reals, with the usual topology is contractible but with the discrete topology is not

connected hence, not contractible.

4.36: Counterexample:
"Being locally contractible" is not an expansive
topological property.

The set, R, of reals, with the usual topology, is locally contractible but with the indiscrete rational(irrational) extension of R, which is finer than the usual topology on R, is not locally connected and hence is not locally contractible.

Table VI indicates the properties which are not hereditary and not closed hereditary and /or not open hereditary properties. The "=" in a line indicates that a counterexample is either given in this paper or referenced in the literature. The "-" in a line indicates the moninvariance is implied by the "=" in the same line, due to the contrapositive statements of the following:

Hereditary => chosed hereditary & open hereditary.

Table VII indicates the properties which are not continuous, not divisible and not open continuous and/or not closed continuous properties. The "-" in a line indicates that a counterexample is either given in this paper or referenced in the literature. The "-" in a line indicates the non-invariance is implied by the "-" in the

same line due to the contrapositive statements of the following:

Continuous > divisible

popen continuous & closed continuous.

Table VIII indicates the properties which are not continuous and not contractive properties as well as those which are not isotopy and not homotopy properties. The """ in a line indicates that a conterexample is either given. In this paper or referenced in the literature. The "" in a line indicates the non-invariance is implied by the "" in the same line, due to the contrapositive statements of the following:

Continuous => contractive.

Homotopy => isotopy

Table IX indicates the properties which are not arbitrarily productive, not countably productive and/or not finitely productive properties as well as those which are not expansive properties. The "e" and/or "" in a line indicate that a counterexample is either given in this paper or referenced in the literature. The "-" in a line indicates that the non-invariance is implied by the "" in the same line due, the contrapositive statement of the following: Arbitrarily productive countably productive.

Table V

COUNTEREXAMPLES FOR NON-HEREDITARY PROPERTIES

No	ot closed hereditary.⇒ ot open hereditary ⇒ no	noi t l	he.	he re	reditary ditary
gi	" indicates that a punterexample is either even in this paper or				Reference to "=".
f	ound in the literature.		A		N stands for
3			8	2	normal.
	" indicates that the		7	t a	
	n-invariance of the	1	9	Ξ	CH stands for
	operty is implied by	13	e	ě	closed hereditary.
""		12	_	9	OH stands for
100	A	2	9	1".	open hereditary.
1	Property	Hereditary	0.0	Open hereditary	ben neredrury.
	,	He H	2	0	L
1	To				
2	T1 /1-	1		L.	
3	T2.	1	L	<u> </u>	
5	T21/2	-	-	-	
6	T 3/2	+-	-	-	
7	T#	-	-	=	41,p253 (Tu is N)
8	Te	1	-	1	
9	Regular				
10	Completely regular		Ŀ	_	
11	Normal	=	-	=	41,p253 (N is T+)
13	Completely normal	+	-	-	
14	Path connected	╁┼	-	 −-	
15	Locally connected	+-	-	-	41,p185
16	Totally disconnected	1		†=	
17	Compact	Ε.		Ξ	41,p271-273
18	Lindelöf	Ξ		Ξ	41,p271-273
19	Locally compact	-		=	41,p280
21	Countably compact	1=	-	=	41,p271-273
21	Paracompact Separable	+=-	-	-	15,p162 30,p50
23	Second countable	 	H	-	301830
24	First countable	t	Н	-	
25	Discrete	\Box			· · · · · · · · · · · · · · · · · · ·
26	Indiscrete				
27	Metrizable				7
28	Fixed point	1=	=	=	CH-4.21 OH-4.22
30	Contractible	1-1	-	-	<u> </u>
30	Locally contractible	1-1	-	L_	4.23

Table VII

COUNTEREXAMPLES FOR NON-CONTINUOUS PROPERTIES

No	t open continuous > no	t i	41	vi	si	ble > not continuous
	" indicates that a	П				
	unterexample is either	П		'		-
	ven in this paper or	П			89	Reference to "="
	und in the literature.	Ш		_	5	Kererence co
	and in our literacule.	1		B	ed continuous	
	· indicates that the	1		9	2	OC stands for
	n-invariance of the			5	t	open continuous.
	operty is implied by	ē		4	ō	open continuous.
		18	7 P	6	0	CC stands for-
		15	겊	U	Pe	closed
		4	7	n	osed	continuous.
100	Property	Continuous	Divisible	1 2	Close	
		0	A		o	
1	To .	Ξ	=	Ξ	Ξ	OC & CC-41,p198
2	T1	ļ=.	-	=	L_	41,p198
3	T2	=	-	=	=	
4	T21/2	ļ=.	=	=		OC-4.24 CC-4.26
5	T3	=	=	7	=	OC-41,p208 CC-41,
6	T 3/2.	7	1=	Ž	LΞ	OC-41,p243 CC-41,
7	T#	ξ=.	2		-	41,p254
8	T.5	 =	E	=	-	4.24
	Regular	 = -				OC-41,p208 CC-41,p
10	Completely regular	1=	=		1=	OC-41,p243 CC-41,
12	Normal	1=	1=	=	-	41,p254
13	Completely normal	+=-	+=	ᄩ	-	4.24
14	Path connected	╌	╁	├		
15	Locally connected	+	-	-	-	ļ
16	Totally disconnected	+-	+	=		4.25
17	Compact	ŧ÷.	=	F	-	7.23
18	Lindelöf	-	+	-	-	
19	Locally compact	+-	+	-	-	56,p133 ·
20	Countably compact	-	+-	-	+=	
	Paracompact	+	+=	,=	-	56,p149
22	Separable	+	+-	-	-	
23	Second countable	-	1=	-	-	34,p104
24	First countable	=	+=	-		34,p104
25	Discrete	1-	t-	+-	-	
	Indiscrete	1-	 -	-	-	
27	Metrizable	1=	1=		-	OC-50,p697 CC-50,
28	Fixed point '	1=	=	-	-	CC-4.2
29	Contractible	1=	1=	-	-	CC-4.2
30	Locally contractible	1-	t-	1		

Table VIII

COUNTEREXAMPLES FOR NON-CONTRACTIVE AND NON-HOMOTOPY

PROPERTIES

N	ot contractive > not co	ont	in	u o	u.s	
	ot isotopy - not homoto				•	
.,	" & "#" indicate	1	т	Т	т-	T
	hat a counterexample	1	١,	1	1	/
	either given in		1	"		× ×
	his paper or found	1				11
	the literature.		1		١.	
11	the literature.	1	1		1	Reference to
-	· · · · · · · · · · · · · · · · · · ·	1				
	-"&"*" indicate that	1	0			"=" and "#".
	ne non-invariance of	Continuous	Contractiv	1		
ti	ne property is	10	4	1		,
	aplied by "="&"#"	l a	1 0	Homotopy	Isotopy	
re	espectively.	17	1	1	0	
1	S 0 0	l a	ä	ă	00	
1	Property	18	20	100	8	
L		Ľ	Ľ	[_	Γ.	
1	To	1-	=	L		4.28
2	Ti.	1-	=	_	_	4.28
3	Ta	. =.	=	<u> </u>	_	4.28
4	Tate	4=	Ξ		-	4.28
5	T3	+=	=	├-	-	4.29
6	T 3½	4=	=		ļ	4.29
7	T# C	1=	=	L_	L	4.29
8	TS	+=	=	_		4.29
9	Regular	-	=	_	L.	4.28
10	Completely regular	4=	=	_	L.	4.28
11	Normal	-	=	_	_	4.28
12	Completely normal	1-	=	_		4.28
13	Connected	1	_	_	_	
14	Path connected	1		_		
15	Locally connected	1-	=			. 41,p185
16	Totally disconnected	1-	=			4.28
.17	Compact	\perp				, ,
18	Linde18f	\perp		`		
19	Locally compact	1-1	=			41,p280
20	Countably compact	1				
21	Paracompact	1-	12			4.29
22	Separable					
23	Second countable	1-	=			41,p273
24	First countable	ΙΞI	=			41,p273
25	Discrete	1-1	=			4.28
26	Indiscrete	\square				
27	Metrizable	Œ	-			4.28
28	Fixed point	Œ	=	٠	#	=-4.30 #-4.32
29	Contractible	T = T				
30	Locally contractible	1-1	=	*	#	=-4.31 #-4.33

COUNTEREXAMPLES FOR NON-PRODUCTIVE AND NON-EXPANSIVE

PROPERTIES /

Not	finitely productive					tably productive trarily productive
tha eit pap lit	and "= indicate t a counterexample is her given in this er or found in the erature. indicates that the -invariance of the porty is implied by	9		Finitely productive		Reference to
11	To .	+~	٥	1	-	~~~~~
2.	Ti	+	-	Н	-	
3	T 2	+-		Н	-	
4	T 21/2	+	-	-	-	
5	T3	+-		-	*	49,p88-89.
6	T 31/2	+-		-	#	49,p88-89
7	TH	1=	=	=	#	=-41,p254 #-49,p88-89
8	To	+=	-		#	=-24,p80 #-49,p88-89
9	Regular	+-			#	49,p88-89
10	Completely regular	+			#	49,p88-89
11	Normal	1=	-	=	#	=-41,p254 #-49,p88-89
12	Completely normal	1=	-	=	#	=-24,p80 #-49,p88-89
13	Connected	T			#	.49,p41-43
14	Path connected	T			#	49,p41-43
15	Locally connected	I-				=-41,p186 #-41,p185
16	Totally disconnected	L				
17	Compact	L			#	41,p271-273
18	Lindelöf	ΙΞ	-	=		→ & #-41,p271-273
19	Locally compact	ΙΞ	=		#	=-49,p121 #-41,p280
20	Countably compact	I-	Ξ		*	=-30,p192 #-41,p271-3
21	Paracompact	1-	=	=	*	=-24,p69 #-49,p88-89
22		1=		_	#	=-49,p123 #-41,p271-3
23	Second countable	1=	_	1_	*	=-49,p123 #-41,p271-3
24		=	_	-	*	=-49,p123 #-41,p271-3
25		1=	=	1		49,p121
26	Indiscrete	+	-	-		49,p41-42
27	Metrizable	=	_	1		=-30,p100 #-49,p88-89
28	Fixed point	ļ=.	=	=		=-12,p977 #-4.30
29	Contractible	-	-			#-4.035
30	Locally contractible	1=	=	1	#	=-4.34 #-4.36

BIBLIOGRAPHY

- Armstrong, M.A., Basic Topology, Springer-Verlag, New York, 1983.
- 2. Bing, R.H., The elusive fixed point property., Amer. Math. Month., 76 (1969), 119-132.
- Borsuk, K., Theory of Retracts, Polish Scientific Publishers, Warsaw, 1966.
- Bourbaki, N., Elements of Mathematics: General Topology, Part 1., Addison-Wesley, Reading, Mass., 1966.
- Bourbaki, N., Elements of Mathematics: General Topology, Part 2., Addison-Wesley, Reading, Mass., 1966.
- 6. Bredon, G. E., Some examples for the fixed point property., Pac. J. Math., 38 (1971), 571-575.
- Brown, R., Elements of Modern Topology., McGraw-Hill, New York, 1968.
- Brown, R. F., The fixed point property and cartesian products., Amer. Math. Month., 89 (1982), 654-678.
- 9. Cech. E., Topological Spaces., Revised by F. Zdenek & M. Kafetov., John Wiley & Sons, London, 1966.
- Chinn, W.G. and Steenrod, N.E., First Concepts of Topology: The Geometry of Mappings of Segments, Curves, Circles, and Disks., Random House, Toronto, 1966.
- Christenson, C. O. and Voxman, W. L., Aspects of Topology., Marcel Dekker, New York, 1977.
- Connell, E.H., Properties of fixed point spaces., Proc. Amer. Math. Soc., 10 (1959), 974-979.
- Conover, R.A. A First Course in Topology: An Introduction to Mathematical Thinking, Milliams & Wilking, Baltimore, 1975.
- Cullen, H. F., Introduction to General Topology., D. C. Heath, Boston, 1968.
- 15. Dugundji, J., Topology., Allyn & Bacon, Boston, 1966.
- Fadell, E., Recent results in the fixed point theory of continuous maps., Bull. Amer. Hath. Soc., 76 (1970), 10-29.

- Fairchild, W. W. and Tulcea, C. I., Topology., W. B. Saunders, Philadelphia, 1971.
- Franz, W., Algebraic Topology., Frederick Ungar, New York, 1968.
- Franz, W. General Topology., Frederick Ungar, New York, 1965.
- Gamelin, T.W. and Greene, R.E., Introduction to Topology., Saunders College, New York, 1983.
- Gemignani, M. C., Elementary Topology., Addison-Wesley, Reading, Mass., 1967.
- 22. Gottlieb, D. H., Homotopy and isotopy properties Dissertation, UCLA, 1962.
- 23. Gottlieb, D.H., Homotopy and isotopy properties of topological spaces., Can. J. of Math., 16 (1964), 561-571.
- 24. Greever, J., Theory and Examples of Point-Set Topology., Brooks/Cole, Belmont, Ca., 1967.
- 25. Harris, D., Structures in Topology., American Mathematical Society, Providence, R. I., 1971.
- Hocking, J. G. and Young, G. S., Topology., Addison-Wesley, Reading, Mass., 1961.
- Hu, S. T., Elements of General Topology., Holden-Day, San Francisco, 1964.
- Hu, S. T., Homotopy and isotopy properties of topological spaces., Can. J. Math., 13 (1961), 137-176.
- 29. Hu, S. T., Theory of Retracts., Wayne State University Press, Detroit, 1965.
- Husain, T., Topology and Maps., Plenum Press, New York, 1977.
- 31. Joshi, K. D., Introduction to General Topology., Halsted Press, New York, 1983.
- Kahn, D. W., Topology: An Introduction to the Point-Set and Algebraic Areas., Williams & Wilkins, Baltimore, 1975.
- Kasriel, R. H., Undergraduate Topology., W. B. Saunders, Philadelphia, 1971.

- Relley, J., General Topology., Van Nostrand, New York, 1955.
- Lopez, W., An example in the fixed point theory of polyhedra., Bull'. Amer. Math. Soc., 73 (1967), 922-924.
- Mansfield, M. J., Introduction to Topology., Van Nostrand, Princeton, N. J., 1963.
- McCarty, G., Topology: An Introduction with Application to Topological Groups., McGraw-Hill, New York, 1967.
- 38. Mendelson, B., Introduction to Topology, 2nd. Edition.,
- Michael, E., Local properties of topological spaces., Duke Nath. J., 21 (1954), 163-171.
- 40. Munkres, J., Topology: A First Course., Prentice-Hall, Englewood Cliffs, N. J., 1975.
- 41. Murdeshwar, M. G., General Topology., Halsted Press, New York, 1983.
- 42. Pervin, W. J., Foundations of General Topology., Academic Press, New York, 1964.
- Schubert, H., Topology. Translated by S. Moran., Allyn & Bacon, Boston, 1968.
- 44. Schurle, A. W., Topics in Topology., Elsevier North Holland, New York, 1979.
- Simmońs, G. F., Topology and Modern Analysis., McGrav-Hill, New York, 1963.
- Singer, I. M. and Thorpe, J. A., Lecture Notes on Elementary Topology and Geometry., Scott, Foresman, Glenview, 111., 4967
- Sorgenfrey, R. H., Oh the topological product of paracompact spaces. Bull. Amer. Hath. Soc., 53 (1947), 631-632.
- 48. Spanier, E. H., Algebraic Topology., Springer-Verlag,
- Steen, L. A. and Seebach, J. A., Counterexamples in Topology., Holt, Rinehart & Winston, New York, 1970.
- Stone, A. H., Metrizability of decomposition spaces., Proc. Amer. Math. Soc., 7 (1956), 690-700.

- Stone, A. H., Paracompactness and product spaces.
 Bull. Amer. Math. Soc., 54 (1948), 977-982.
- Strother, W. L., On an open question concerning fixed points., Proc. Amer. Math. Soc., 4 (1953), 988-993.
- Sutherland, W. A., Introduction to Metric and Topological Spaces., Clarendon Press, Oxford, 1975.
- 54. Wall, C. T. C., A Geometric Introduction to Topology., Addison-Wesley, Reading, Mass., 1972.
- 55. Wilansky, A., Topology for Analysis., Ginn, Waltham, Mass., 1970.
- 56. Willard, S., General Topology., Addison-Wesley, Reading, Mass., 1970.
- Whyburn, G. and Duda, E., Dynamic Topology., Springer-Verlag, New York, 1979.







