

POSSIBLE RECENT RANGE EXPANSION OF *ALCATAENIA LONGICERVICA* (EUCESTODA: DILEPIDIDAE) PARASITIC IN MURRES *URIA* SPP. (ALCIDAE) INTO THE NORTH ATLANTIC

SABIR B. MUZAFFAR¹, ERIC P. HOBERG² & IAN L. JONES¹

¹*Department of Biology, Memorial University of Newfoundland, St. John's, Newfoundland and Labrador, A1B 3X9, Canada (v35sbm@mun.ca)*

²*US National Parasite Collection & Animal Parasitic Disease Laboratory, USDA, Agricultural Research Service, BARC East No. 1180, 10300 Baltimore Avenue, Beltsville, Maryland, 20705, USA*

Received 17 October 2005, accepted 12 January 2006

A wide range of helminths has been recorded from murre *Uria* spp. (Threlfall 1971, reviewed by Hoberg 1984a, Muzaffar & Jones 2004). Of the cyclophyllidean cestodes that parasitize murre, the genus *Alcataenia* Spasskaya 1971 (Dilepididae) is represented by ten species, eight of which are restricted and specific to the auks; the other two species occur in gulls (Laridae). The intermediate hosts of *Alcataenia* are euphausiid crustaceans such as species of *Thysanoessa* (Shimazu 1975), which also form an important dietary component of breeding and wintering murre, particularly Thick-billed Murre *Uria lomvia* (Gaston & Noble 1985, Birkhead & Nettleship 1987, Elliot *et al.* 1990, Rowe *et al.* 2000, Ainley *et al.* 2002). *Alcataenia armillaris* Rudolphi 1810 and *A. meinertzhageni* Baer 1956 are restricted to the murre and have been recorded from different locations in the Holarctic (Threlfall 1971; Hoberg 1984a, 1986). *Alcataenia longicervica* Hoberg 1984 was described from murre in the North Pacific basin and has been regarded as endemic to the region (Hoberg 1986) because relatively extensive surveys of murre from the North Atlantic and Arctic basin have previously failed to find the species (e.g. Markov 1937, 1942, Belopolskaya 1952, Threlfall 1971, Galkin *et al.* 1994).

We report a new geographic record of *A. longicervica* in wintering Thick-billed Murre and Common Murre *Uria aalge*. Thick-billed Murre and Common Murre were collected by hunters in February 2005 from Harbor Breton (47.467°N, 55.800°W) and St. Mary's Bay (46.933°N, 53.583°W), Newfoundland. Each bird was dissected, and the gastrointestinal tracts were frozen. These samples were later analyzed under a dissecting microscope to collect helminths. Helminth specimens were preserved in 70% ethanol and subsequently stained in Semichon acid carmine stain and mounted entire for identification. The armed rostellum was dissected from some specimens and mounted separately to facilitate accurate determination of form, number, distribution and dimensions of hooks. Identifications were based on descriptions or re-descriptions (Hoberg 1984b) and direct comparison to holotype, paratype or known voucher specimens representing species of *Alcataenia* held in the US National Parasite Collection (USNPC), Beltsville, Maryland, USA.

Seven Thick-billed Murre yielded 39 specimens of *A. longicervica* and 40 specimens of *A. armillaris*. In addition, five Common Murre yielded eight *A. longicervica* and 24 *A. armillaris*. No specimens of *A. meinertzhageni* were found. Infections with *Alcataenia* are generally associated with lesions, because the tapeworm is deeply imbedded in the host's intestinal mucosa, making it susceptible to secondary infections (Threlfall 1971, Hoberg 1984b). Specimens

of *A. longicervica* have been deposited in the collections of the USNPC under accession numbers 97386–97388.

The appearance of *A. longicervica* in the Northwest Atlantic can be explained in two ways, involving either inadequate earlier sampling or a recent range expansion and establishment in the North Atlantic among populations of *Uria* spp.

Under the first scenario, *A. longicervica* has been continuously present across the Holarctic as a component of the helminth fauna of murre established in the Pleistocene (e.g. Hoberg 1986), but was not previously detected in the Atlantic basin. Studies by Belopolskaya (1952) in the Barents Sea and Threlfall (1971) in the Northwestern Atlantic examined large numbers of murre and other alcids and reported the common occurrence of *A. armillaris* and, to a much lesser extent, *A. meinertzhageni*. Other surveys from the North Atlantic and Arctic basin, although involving smaller sample sizes for either species of murre, also apparently did not document dilepidid cestodes other than *A. armillaris* or *A. meinertzhageni* (e.g. Krabbe 1869; Markov 1937, 1941; Baer 1956, 1962; Galkin *et al.* 1994; Galaktionov 1996).

Alcataenia longicervica was described based on an extensive collection of specimens in *U. lomvia* and *U. aalge* from the eastern Bering Sea (Hoberg 1984b, 1986) and, because of its similarity to *A. armillaris*, may have escaped recognition in the North Atlantic in the course of earlier surveys. We believe that scenario to be unlikely, because unequivocal diagnostic characters serve to distinguish *A. longicervica* from its congeners in murre. An elongated neck region, the structure and dimensions of the rostellar sac, and attributes of the genital organs in mature cestodes clearly distinguish this cestode. The specimens of Belopolskaya (1952) or Threlfall (1971) are no longer available because they are unrepresented in museum collections. However, all cestodes remaining from collections by Krabbe (1869) and Baer (1956, 1962) were examined during a re-description and found to be consistent with *A. armillaris* (see Hoberg 1984b). Although such circumstantial evidence does not fully resolve the issue of the historical distribution of *A. longicervica*, it also does not refute an alternative hypothesis.

Alternatively, assuming that *A. longicervica* was formerly absent in the northwest Atlantic, the current finding could be considered in light of the distributional patterns of recognized intermediate hosts for species of *Alcataenia*. Shimazu (1975) found cysticeroids of *A. armillaris* in the euphausiid *Thysanoessa inermis*, and it

is suggested that in general, species of *Thysanoessa* or perhaps *Euphausia* are intermediate hosts for other *Alcataenia*. Changing patterns of distribution for species and populations of euphausiids may be implicated in the putative range expansion and geographic colonization for *A. longicervica* across the Arctic basin into the North Atlantic.

In the scenario that we propose, increasingly pronounced eastward currents such as the Transpolar Drift (Dyke *et al.* 1997) may have facilitated the advective movement of infected euphausiids (specifically species of *Thysanoessa*) leading to geographic colonization of high-latitude seas adjacent to the Canadian Arctic and western Greenland. Because globally significant colonies of Thick-billed Murres occur in these regions (Gaston & Hipfner 2000, CAFF 2004) this exposure via infected *Thysanoessa* most likely facilitated the establishment of populations of *A. longicervica* in Thick-billed Murres. Infected Thick-billed Murres could then have further disseminated the parasite through populations of *Thysanoessa* in the Labrador Sea and to the Grand Banks in Newfoundland waters, subsequently exposing Common Murres, distributed further south and co-occurring with Thick-billed Murres in those regions (Gaston & Jones 1998, Gaston & Hipfner 2000, Ainley *et al.* 2002, CAFF 2004). Long-term changes are also evident in the distributions of many marine invertebrates, including euphausiids and calanoid copepods (Beaugrand *et al.* 2002, Barnard *et al.* 2004), and such shifts in distribution could contribute to the apparent spread of parasitic organisms among marine vertebrates (Hoberg 1996, 2005).

A test of our hypothesis could be achieved through phylogeographic comparisons based on mitochondrial DNA derived from populations of *Alcataenia* in the North Atlantic and North Pacific basins (e.g. Avise 2000). In this case, a contemporary range expansion would be consistent with the absence of geographic partitioning between populations of *A. longicervica* across the Holarctic and would argue against a longer history for this species in the western North Atlantic.

ACKNOWLEDGMENTS

We thank Greg Robertson [Canadian Wildlife Service (CWS)] and Johanne Dussureault (Atlantic Cooperative Wildlife Ecology Research Network, CWS) for providing the bird specimens used in this study. Funding was provided by the Atlantic Cooperative Wildlife Ecology Research Network, CWS/NSERC (Natural Sciences and Engineering Research Council of Canada) Industrial Chair and NSERC equipment grants to ILJ.

REFERENCES

- AINLEY, D.G., NETTLESHIP, D.N., CARTER, H.R. & STOREY, A.E. 2002. Common Murre. In: Poole A. & Gill F. (Eds). The birds of North America. No. 666. Philadelphia, PA, & Washington, DC: The Academy of Natural Sciences & The American Ornithologists' Union.
- AVISE, J.C. 2000. Phylogeography: the history and formation of species. Cambridge: Harvard University Press.
- BAER, J.G. 1956. Parasitic helminths collected in west Greenland. *Meddeleser om Grønland* 124: 1–55.
- BAER, J.G. 1962. Cestoda. Volume II, Part 12. The zoology of Iceland. Reykjavik: Ejnar Munksgaard. pp. 1–63.
- BARNARD, R., BATTEN, S.D., BEAUGRAND, G., BUCKLAND, C., CONWAY, D.V.P., EDWARDS, M., FINLAYSON, J., GREGORY, L.W., HALLIDAY, N.C., JOHN, A.W.G., JOHNS, D.G., JOHNSON, A.D., JONAS, T.D., LINDLEY, J.A., NYMAN, J., PRITCHARD, P., REID, P.C., RICHARDSON, A.J., SAXBY, R.E., SIDEY, J., SMITH, M.A., STEVENS, D.P., TAYLOR, C.M., TRANTER, P.R.G., WALNE, A.W., WOOTTON, M., WOTTON, C.O.M. & WRIGHT, J.C. 2004. Continuous plankton records: plankton atlas of the North Atlantic Ocean (1958–1999). II. Biogeographical charts. *Marine Ecology Progress Series* (Suppl): 11–75.
- BEAUGRAND, G., REID, P.C., IBÁÑEZ, F., LINDLEY, J.A. & EDWARDS, M. 2002. Reorganization of North Atlantic copepod biodiversity and climate. *Science* 296: 1692–1694.
- BELOPOLSKAYA, M.M. 1952. Parazitofauna morskich vodoplavaiushchikh ptits. Leningrad Universitet Uchenie Zapiski Annals. *Seria Biologicheskikh Nauk* 141: 127–180.
- BIRKHEAD, T.R. & NETTLESHIP, D.N. 1987. Ecological relationships between Common Murres, *Uria aalge*, and Thick-billed Murres, *Uria lomvia*, at the Gannet Islands, Labrador. III. Feeding ecology of the young. *Canadian Journal of Zoology* 65: 1638–1649.
- CONSERVATION OF ARCTIC FLORA AND FAUNA (CAFF). 2004. Distribution of murre outside the breeding season. Circumpolar murre banding program, North Atlantic Region. [Technical Report No. 13.] Akureyri, Iceland: CAFF International Secretariat.
- DYKE, A.S., ENGLAND, J., REIMNITZ, E. & JETTE, H. 1997. Changes in driftwood delivery to the Canadian Arctic archipelago: the hypothesis of postglacial oscillations of the transpolar drift. *Arctic* 50: 1–8.
- ELLIOT, R.D., RYAN, P.C. & LIDSTER, W.W. 1990. The winter diet of Thick-billed Murres in coastal Newfoundland waters. *Studies in Avian Biology* 14: 125–138.
- GALKIN, A.K., GALAKTIONOV, K.V., MARASAEV, S.F. & PROKOFEV, V.V. 1994. Tsestody ryboiadnykh ptits O. Kharlov i Semli Frantsa-Iosifa. *Parazitologiya* 28: 373–383.
- GASTON, A.J. & NOBLE, D.N. 1985. The diet of Thick-billed Murres (*Uria lomvia*) in west Hudson Strait and northeast Hudson Bay. *Canadian Journal of Zoology* 63: 1148–1160.
- GASTON, A.J. & JONES, I.L. 1998. The auks. Oxford: Oxford University Press.
- GASTON, A.J. & HIPFNER, J.M. 2000. Thick-billed Murre. In: Poole A. & Gill F. (Eds). The birds of North America. No. 497. Philadelphia, PA, & Washington, DC: The Academy of Natural Sciences & The American Ornithologists' Union.
- HOBERG, E.P. 1984a. Systematics, zoogeography and ecology platyhelminth parasites of the seabird family Alcidae (Charadriiformes: suborder Alcae) [PhD dissertation]. Seattle: University of Washington. 324 pp.
- HOBERG, E.P. 1984b. *Alcataenia longicervica* sp. n. from murre, *Uria lomvia* (Linnaeus) and *Uria aalge* (Pontoppidan) in the North Pacific basin, with redescription of *Alcataenia armillaris* (Rudolphi, 1810) and *Alcataenia meinertzhageni* (Baer, 1956) (Cestoda: Dilepididae). *Canadian Journal of Zoology* 62: 2044–2052.
- HOBERG, E.P. 1986. Evolution and historical biogeography of a parasite–host assemblage, *Alcataenia* spp. (Cyclophyllidea: Dilepididae) in Alcidae (Charadriiformes). *Canadian Journal of Zoology* 64: 2576–2589.

- HOBERG, E.P. 1996. Faunal diversity among avian parasite assemblages: the interaction of history, ecology and biogeography in marine systems. *Bulletin of the Scandinavian Society for Parasitology* 6: 65–89
- HOBERG, E.P. 2005. Marine birds and their helminths. In: Rohde, K. (Ed). *Marine parasitology*. Collingwood, Australia: CSIRO Publishing. pp. 414–420.
- KRABBE, H. 1869. Bidrag til Kundskab om Fugelenes Baendelorme. Kongelige Danske Videnskabernes Selskabs Skrifter. Naturvidenskabelig og Matematisk Afdeling 8: 251–368.
- MARKOV, G.S. 1937. Changes with age in the parasite fauna of the Novaya Zemlya Murre (*Uria lomvia lomvia*) [Russian]. *Transactions of the Leningrad Natural Society* 66: 456–466.
- MARKOV, G.S. 1941. Parasitic worms of Bezymiannaya Bay (Novaia Zemlia). *Reports USSR Academy of Sciences* 30: 579–582.
- MUZAFFAR, S.B. & JONES, I.L. 2004. Parasites and diseases of the auks (Alcidae) of the world and their ecology—a review. *Marine Ornithology* 32: 121–146.
- ROWE, S., JONES, I.L., CHARDINE, J.W., ELLIOT, R.D. & VEITCH, B.G. 2000. Evidence for recent changes in the winter diet of murre (*Uria* spp.) in coastal Newfoundland waters. *Canadian Journal of Zoology* 78: 495–500.
- SHIMAZU, T. 1975. Some cestode and acanthocephalan larvae from euphausiid crustaceans collected in the northern North Pacific Ocean [Japanese]. *Bulletin of the Japanese Society of Science and Fisheries* 41: 813–821.
- THRELFALL, W. 1971. Helminth parasites of alcids in the northwest North Atlantic. *Canadian Journal of Zoology* 49: 461–466.
-