

PARASITES AND DISEASES OF THE AUKS (ALCIDAE) OF THE WORLD AND THEIR ECOLOGY—A REVIEW

SABIR BIN MUZAFFAR & IAN L. JONES

*Department of Biology, Memorial University of Newfoundland, St John's, Newfoundland, A1B 3X9, Canada
(v35sbm@mun.ca)*

Received 1 March 2004, accepted 1 June 2004

SUMMARY

MUZAFFAR, S.B. & JONES, I.L. 2004. Parasites and diseases of the auks (Alcidae) of the world and their ecology—a review. *Marine Ornithology* 32: 121–146.

We reviewed all the organisms that are known to parasitize auks (Aves: Alcidae). Of the 23 extant auk species, parasites have been described from 19 species; no published information was found on parasites of Xantus's Murrelet *Synthliboramphus hypoleucus*, Craveri's Murrelet *S. craveri*, the Japanese Murrelet *S. wumizusume* or the Long-billed Murrelet *Brachyramphus perdix*. Our survey identified 184 taxa parasitic or pathogenic on auks. Endoparasitic microorganisms included 21 viruses, 13 bacteria, 3 dinoflagellates, 6 protozoa and 3 fungi. Other endoparasitic organisms included 57 platyhelminth (34 digenean, 23 cestode), 9 acanthocephalan and 22 nematode taxa. Ectoparasites (all arthropods) included 2 pentastomid, 14 acari and 30 insect taxa. We reviewed published information on the effects that these parasites have on the biology of auks. Most studies did not investigate relationships between the ecology, the breeding condition or the physiologic state of the birds and the presence of parasites. Even though episodes of mass mortality of seabirds are periodically recorded, few of those episodes have been exclusively linked to parasitic infestations. Viral isolates from auks have been recorded from several breeding colonies, but their epizootiologic consequences are unknown. Bacterial isolates, of which the most noteworthy species is the Lyme disease-causing agent *Borrelia burgdorferi* (sensu lato), have been recorded in auks and their tick ectoparasite *Ixodes uriae*. A few life cycles of digeneans and cestodes recorded from auks have been determined. Growing evidence indicates prey switching by auk species alters their endoparasitic fauna. Ectoparasitic organisms often play a role in transmitting endoparasites and may affect the reproductive success of their hosts. Some links have been clearly established; many others are only implied and urgently need close attention. The role of parasitism in host population dynamics and as reservoirs is discussed in the context of seabird ecology as a whole.

Key words: Charadriiformes, Alcidae, ectoparasite, endoparasite, disease

INTRODUCTION

Parasites play a very important part in the lives of their hosts (Rothschild & Clay 1957, Dogiel 1964). Nevertheless, relative to the enormous diversity in the multitude of parasites that live in or on wild birds (Loye & Zuk 1991, Fowler 1993, Clayton & Moore 1997), little information is available on the intricate life cycles of those parasites. Information on parasitic acari, for example, is limited to conspicuous taxa that are readily captured and isolated from their hosts. Less-evident parasitic organisms, whether internal or external, are usually overlooked, and their complex interactions and potential influence on host life cycles are largely unknown. Parasites may incur costs for their avian hosts and are often regarded as an important drawback to colonial breeding (Brown & Brown 1986).

Nest ectoparasites may reduce breeding success in Cliff Swallows *Hirundo pyrrhonota* (Brown & Brown 1986, 1996). Ectoparasites have been shown to cause nest desertion and chick mortality in a range of seabirds (Feare 1976; King *et al.* 1977a, 1977b; Duffy 1983, 1991). There is also clear evidence of the impact of ectoparasites on seabird reproductive success and population dynamics (Boulinier & Danchin 1996, Gauthier-Clerc *et al.* 1998, Bergström *et al.* 1999, Boulinier *et al.* 2001, Mangin *et al.* 2003). Outbreaks of disease likely related to parasitic infestations have

rarely been studied in detail (Petermann *et al.* 1989, Debacker *et al.* 1997), and the influence of diseases on seabird populations remains unclear (Gauthier-Clerc *et al.* 2003).

The general lack of information on seabird parasites applies equally to the Alcidae (auks or alcids). Although behavior, ecology and physiology are well known (reviewed by Gaston & Jones 1998), relatively few studies have documented the incidence of parasites and their influence on their hosts (Eveleigh 1974, Eveleigh & Threlfall 1976, Fitzpatrick & Threlfall 1977, Choe & Kim 1987, Hoberg 1996, Muzaffar 2000).

Individual seabirds in breeding colonies may expose one another to a multitude of viruses and bacteria (e.g. Oprandy *et al.* 1988, Gylfe *et al.* 1999). Many seabirds, including the auks, breed during summer in large colonies (Gaston & Jones 1998) that also serve as ideal habitats for ectoparasites (e.g. Eveleigh & Threlfall 1975, Clifford 1979). Many parasitic species have become highly specialized in synchronizing their life cycles with their hosts' breeding phenology (e.g. Eveleigh & Threlfall 1975, Clifford 1979, Lvov 1980, Hoberg 1996, Gylfe *et al.* 1999, McCoy & Tirard 2002). However, most auk species spend the greater part of the year dispersed at open sea where transmission and survival of parasites represents a major challenge. Nevertheless, foraging behavior and diet facilitate the transmission of a variety of endoparasites through

marine invertebrate and vertebrate species, although those parasites, too, are better transmitted during the breeding season because of the abundance of marine littoral intermediate hosts (Hoberg 1996).

The objectives of our study were

- to present a review of the diversity of diseases and parasites of alcids as reported in the literature, and
- to highlight key studies documenting the ecology of host–parasite interactions and their relationship to the marine environment.

TAXONOMY

The literature exhibits some taxonomic uncertainty, particularly for the helminth fauna of alcids. Our account is based on a search of the literature for the period 1899–2003 (including review papers), and it clumps organisms into easily recognizable taxonomic groups. However, given that systematics and taxonomy vary among the groups, the taxonomic groupings are not consistent. Bacteria, for example, form a Kingdom, whereas the Eucestoda represents a Class. We attempt to identify genera and species wherever possible. We also attempt to classify organisms in tables, and we provide notes explaining taxonomic ambiguity wherever necessary.

PARASITES AND DISEASES OF THE AUKS

Viruses

Ticks of the family Argasidae and Ixodidae are widely distributed and are often associated with auks (Clifford 1979). The life cycles of seabird ticks tend to be closely attuned to seabird breeding seasons [e.g. McCoy & Tirard 2002 (see the later discussion of ectoparasites)]. Generally, *Ixodes uriae* (Ixodidae) serve as the primary vectors in more northern latitudes; *Ornithodoros* spp. (Argasidae) play the same role in more temperate southern locations (Clifford 1979, Lvov 1980).

I. uriae have been collected from seabird colonies in Russia (Lvov *et al.* 1979, Lvov 1980) and from the west and east coasts of North America (Yunker 1975, Main *et al.* 1973, Oprandy *et al.* 1988). Viral isolation techniques from argasid and ixodid (mostly *I. uriae*) ticks alike revealed viruses representing five families of RNA viruses (mostly arthropod-borne viruses or arboviruses) and two families of DNA viruses (Clifford 1979, Lvov *et al.* 1979, Lvov 1980, Oprandy *et al.* 1988; Table 1). The presence of virus-specific antibodies (which reflect exposure and an immune response to viruses) in the auk and other seabird hosts showed that viremia may develop in the hosts and that prevalence of particular viruses varied considerably between years (Lvov *et al.* 1979, Oprandy 1988).

Viruses such as Tyuleniy virus have produced disease symptoms in experimentally infected Thick-billed Murres *Uria lomvia*, leading to lesions in the central nervous system that can sometimes be fatal (Lvov *et al.* 1979, Lvov 1980). On the other hand, infections with Paramushir, Great Island and Bauline viruses produced no signs of disease in the hosts (Main *et al.* 1973, 1976; Lvov *et al.* 1979; Oprandy *et al.* 1988).

Enzootic foci centered on colonies of alcids and larids suggest that the viruses are transmitted and maintained among the hosts and their tick vectors (Clifford 1979, Lvov 1980, Oprandy *et al.* 1988).

The existence of active foci of viruses in the low Arctic latitudes is governed by adaptations of the viruses and the tick vectors to environmental extremes (Lvov 1980). Murres in the low Arctic, for instance, may have tick prevalences of 10%–45%, increasing to as high as 100% in chicks (Lvov *et al.* 1979, Lvov 1980). With infestations of as many as 1000 ticks per bird, a means exists for build-up and active exchange of viruses in the host population.

Viruses, along with their tick hosts, overwinter within the rocks or the substrate of the auk colony site at depths of up to 40 cm, where temperatures are adequate for their continued survival. Viruses have been known to remain active in ticks for more than two years (Lvov *et al.* 1979). Viremic birds may also represent a source of infection to a colony during the breeding season, but evidence to suggest that the viruses may overwinter in their seabird hosts is lacking (Lvov *et al.* 1979, Lvov 1980).

Bacterial infections

A wide range of bacterial infections occur in free-ranging birds. Such infections are regarded as important causes of morbidity and mortality (Davis *et al.* 1971, Steele & Galton 1971).

Escherichia coli is generally part of the natural intestinal flora of birds, and it may cause disease in conjunction with other infections (e.g. Morishita *et al.* 1999). Infections with *E. coli* were recorded in 14% of the murres (*Uria* spp.) collected in a German study (Petermann *et al.* 1989).

Pasteurella multocida, the causative agent of the contagious avian cholera may occur as both a chronic and an acute infection in free-ranging birds, often leading to death (Friend & Franson 1999). It has also been recorded from murres (Petermann *et al.* 1989).

An earlier study, also in Germany, prompted by an outbreak of *Salmonella paratyphi* B infections in humans, revealed heavily contaminated seawater in seabird feeding areas (Steiniger & Hahn 1953). Subsequently, isolates of the bacterium from droppings of Razorbill *Alca torda*, Common Murre *U. aalge* and gull (*Larus* spp.) were reported, suggesting that foraging areas may be important in the spread of bacterial infections.

A long-term study on beached Common Murres on the Belgian coast revealed salmonella and other pathogenic infections, although it was not clear whether the infections caused mortality (Debacker *et al.* 1997). Many *Salmonella* serotypes have been isolated from gulls, and the feeding habits of gulls in contaminated areas are believed to possibly serve as a source of infection and a means of spread of the pathogen (Monaghan *et al.* 1985). Gulls have also been implicated as reservoirs of the bacterium, because adults may harbor the disease agent without apparent ill effects (Friend & Franson 1999). Many of these bacterial infections are probably widespread in the Alcidae, but potentially go unnoticed because of chronic levels of infection.

The causative agent of Lyme disease, *Borrelia burgdorferi* (sensu lato), is transmitted primarily by ticks in the *I. ricinus-persulcatus* complex (Lane *et al.* 1991, Baranton *et al.* 1992). The transmission cycle of *B. burgdorferi* in nature is very complex. It spans both hemispheres and involves *Ixodes* ticks and a wide range of birds and mammals. *B. garinii*—one of the genomic species of *B. burgdorferi* (sensu lato)—has been isolated mostly from *I. uriae* ticks and from their auk (for example, Atlantic Puffins *Fratercula*

TABLE 1
Microorganisms^a of auks and vectors

Microorganism	Host species ^b	Vectors	Locality	References
RNA viruses				
Raviviridae, <i>Flavivirus</i> Group B Tyuleniy	<i>Uria aalge</i> , <i>U. lomvia</i> , <i>Fratercula cirrhata</i>	<i>Ixodes uriae</i>	Tyuleniy I., Sakhalin I., Commodore Is., Magadan region, and Murmansk, Russia; OR, USA	Lvov <i>et al.</i> 1975, Lvov <i>et al.</i> 1979, Lvov 1980
Reoviridae, <i>Orbivirus</i> , Kemerovo (KEM) Bauline	<i>Alca torda</i> , <i>U. aalge</i> , <i>F. arctica</i>	<i>Ixodes uriae</i>	Great I., NL, Canada	Oprandy <i>et al.</i> 1988, Main 1978, Main <i>et al.</i> 1973
Great Island	<i>A. torda</i> , <i>U. aalge</i> , <i>F. arctica</i>	<i>Ixodes uriae</i>	Great I., NL, Canada	Oprandy <i>et al.</i> 1988, Main 1978, Main <i>et al.</i> 1973
Okhotskiy	<i>U. aalge</i>	<i>Ixodes uriae</i>	Tyuleniy I., Commodore Is., Magadan region, Murmansk, and SE Chukotka, Russia	Lvov <i>et al.</i> 1975, Lvov <i>et al.</i> 1979, Lvov 1980
Arbroath	<i>F. arctica</i>	<i>Ixodes uriae</i>	Arbroath I., Scotland	Moss & Nuttall 1985
Unnamed		<i>Ixodes uriae</i> , <i>Ornithodoros maritimus</i>	Great Saltee I., Ireland	Nuttall <i>et al.</i> 1984
Unnamed		<i>Ixodes uriae</i>	Isle of May, Scotland	Spence <i>et al.</i> 1985
Bunyaviridae, <i>Nairovirus</i> , Sakhalin (SAK) Sakhalin	<i>U. aalge</i> , <i>F. cirrhata</i>	<i>Ixodes uriae</i>	Tyuleniy I., Sakhalin I., Iona I., Commodore Is., Magadan region, Murmansk, and SE Chukotka, Russia; Gull I., AK, USA; OR, USA	Lvov <i>et al.</i> 1975, Lvov 1980
Avalon	<i>F. arctica</i>	<i>Ixodes uriae</i>	Great I., NL, Canada; Sakhalin I., Russia	Main <i>et al.</i> 1976
Bunyaviridae, <i>Nairovirus</i>, Hughes (HUG)				
Farallon	<i>U. aalge</i>	<i>Ornithodoros</i> sp. NR <i>denmarki</i>	Northwest USA	Clifford 1979
Soldado		<i>O. maritimus</i>		Hoogstraal <i>et al.</i> 1976
Unnamed		<i>O. maritimus</i>	Great Saltee I., Ireland	Nuttall <i>et al.</i> 1984
Bunyaviridae, <i>Phlebovirus</i>, Uukuniemi (UUK)				
Zaliv Terpeniya	Both <i>Uria</i> spp.	<i>Ixodes uriae</i>	Tyuleniy I., Commodore Is., Magadan region, Murmansk coast, and SE coast of Chukotka, Russia	Lvov <i>et al.</i> 1975, Lvov <i>et al.</i> 1979, Lvov 1980
Paramushir		<i>Ixodes uriae</i>	Tyuleniy I., Commodore Is., Russia	Lvov 1980
Unnamed	<i>F. arctica</i>	<i>Ixodes uriae</i>	Arbroath Island, Scotland	Moss & Nuttall 1985
Unnamed		<i>Ixodes uriae</i> , <i>Ixodes rothschildi</i>	Great Saltee I., Ireland	Nuttall <i>et al.</i> 1984
Paramyxoviridae				
<i>Paramyxovirus</i> New Castle	<i>U. aalge</i>		German Bight	Petermann <i>et al.</i> 1989
Orthomyxoviridae, <i>Influenzavirus A</i>				
Influenza	<i>U. aalge</i>		Sakhalin I., Russia	Sazonov <i>et al.</i> 1977

TABLE 1 (continued)

Microorganism	Host species ^b	Vectors	Locality	References
DNA viruses				
Adenoviridae				
<i>Adenovirus</i>	<i>U. aalge</i>		German Bight	Petermann <i>et al.</i> 1989
Poxviridae, <i>Avipoxvirus</i>				
Avian pox	<i>U. aalge</i>			Enstipp & Grémillet 2002
Bacteria				
Proteobacteria, Enterobacteriaceae				
Proteobacteria, Enterobacteriaceae <i>Escherichia coli</i>	<i>U. aalge</i>		German Bight	Petermann <i>et al.</i> 1989
<i>Salmonella enteritidis</i>	<i>U. aalge</i>		Belgian coast	Debacker <i>et al.</i> 1997
<i>S. paratyphi B</i>	<i>U. aalge</i> , <i>A. torda</i>		Swedish coast	Steineger & Hahn 1953
<i>S. typhi</i>	<i>A. torda</i>		Swedish coast	Steineger & Hahn 1953
<i>Yersinia intermedia</i>	<i>U. aalge</i>		German Bight	Petermann <i>et al.</i> 1989
<i>Y. pseudotuberculosis</i>	<i>U. aalge</i>		No locality	Lvov & Iljicev 1979
Pasteurellaceae				
<i>Pasteurella multocida</i>	<i>U. lomvia</i> , <i>U. aalge</i> , <i>F. cirrhata</i>		German Bight	Petermann <i>et al.</i> 1989
<i>Actinobacillus</i> sp.	<i>U. aalge</i>		German Bight	Petermann <i>et al.</i> 1989
Campylobacteraceae				
<i>Campylobacter jejuni</i>	<i>F. arctica</i>		No locality	Hubálek 1994
Mollicutes, Mycoplasmataceae				
<i>Mycoplasma gallisepticum</i>	<i>U. aalge</i>		German Bight	Petermann <i>et al.</i> 1989
Firmicutes, Clostridiaceae				
<i>Clostridium perfringens</i>	<i>U. aalge</i>		German Bight	Petermann <i>et al.</i> 1989
Streptococcaceae				
<i>Streptococcus</i> sp.	<i>U. aalge</i>		German Bight	Petermann <i>et al.</i> 1989
Spirochaetes, Spirochaetaceae				
<i>Borrelia garinii</i>	<i>U. aalge</i> , <i>A. torda</i>	<i>Ixodes uriae</i>	Widespread: Egg I., AK, USA; Flatey I., Iceland; Nolsoy, Faeroe Is.; Bonden I., Sweden	Olsen <i>et al.</i> 1993, Olsen <i>et al.</i> 1995, Jaenson <i>et al.</i> 1994, Gylfe <i>et al.</i> 1999

TABLE 1 (continued)

Microorganism	Host species ^b	Vectors	Locality	References
Dinoflagellata				
Gonyaulacaceae				
<i>Protogonyaulax tamarensis</i>	<i>A. torda</i> , <i>U. aalge</i> , <i>F. arctica</i>	Mussels and other marine shellfish	Northumberland and north Durham coasts, UK	Coulson <i>et al.</i> 1968, Armstrong <i>et al.</i> 1978
<i>Gonyaulax catenella</i>	<i>U. aalge</i> , <i>F. cirrhata</i>		WA, USA	McKernan & Scheffer 1942
Dinophyceae				
<i>Dinophysis</i> spp.	<i>U. aalge</i>		NE coast of England	Shumway <i>et al.</i> 2003
Apicomplexa				
Plasmodiidae				
<i>Plasmodium relictum</i>	<i>U. aalge</i>	Biting Diptera	No locality	Rhodhain & Adrienne 1952 in Bennett <i>et al.</i> 1982
<i>Plasmodium</i> sp.	<i>U. aalge</i>	Biting Diptera	No locality	Rewell 1948 in Bennett <i>et al.</i> 1982
Eimeriidae				
<i>Eimeria fraterculae</i>	<i>F. arctica</i>		Great I., NL, Canada	Leighton & Gajadhar 1986
<i>Eimeria</i> sp.	<i>U. aalge</i>		German Bight	Petermann <i>et al.</i> 1989
Sarcocystidae				
<i>Sarcocystis</i> spp.	<i>U. aalge</i>		Triangle I., BC, Canada	Enstipp & Grémillet 2002
Microsporidia				
	<i>F. corniculata</i>		Captive birds	Tocidlowski <i>et al.</i> 1997
Fungi				
Ascomycota, Onygenaceae				
<i>Blastomyces</i> sp.	<i>U. aalge</i>		German Bight	Petermann <i>et al.</i> 1989
Trichocomaceae				
<i>Aspergillus fumigatus</i>	<i>F. cirrhata</i> , <i>Cepphus columba</i> , <i>Cerorhinca monocerata</i> , <i>U. aalge</i>		Coast of OR, USA; AK, USA; Triangle I., BC, Canada	Monroe <i>et al.</i> 1994, Enstipp & Grémillet 2002
<i>Aspergillus</i> spp.	<i>U. aalge</i>		German Bight	Petermann <i>et al.</i> 1989, Jauniaux & Coignoul 1994

^a Viruses are classified as RNA or DNA viruses and are assigned to families and genera. Each genus is further classified into serogroups with many different viruses, each bearing a name. Some viruses representing a serogroup remain unnamed. Other microorganisms are classified into phyla/divisions, families, genera and species.

^b Empty spaces in this column indicate that the microorganism has been isolated only from the vector or vectors, but in areas that alcid may occupy during the breeding season.

arctica and Razorbills) and other seabird hosts (Olsen *et al.* 1993, Olsen *et al.* 1995, Gylfe *et al.* 1999). Seabirds at their breeding colonies and this tick species are thought to play an important role in the maintenance of *B. garinii* in enzootic foci in the Northern Hemisphere (Olsen *et al.* 1995, Gylfe *et al.* 1999). The bacterium is widespread in North America and Europe. Atlantic Puffins have been implicated as the most likely reservoir host in the Faeroe Islands (Gylfe *et al.* 1999).

I. uriae feeds over a period of 4–10 days (Barton *et al.* 1996)—a period that may allow the ticks to disperse between colonies (over small spatial scales) with their hosts [for example, Black-legged Kittiwakes *Rissa tridactyla* (Danchin 1992, Boulinier *et al.* 2001, Gasparini *et al.* 2001)]. Such dispersal may facilitate the spread of *B. garinii* between colonies that are not separated by large distances. Dispersal of *B. garinii* over much larger spatial scales has been implied (Olsen *et al.* 1995), but seems unlikely based on the existence of distinct molecular strains in the two hemispheres (Lane *et al.* 1991, Bunikis *et al.* 1996). Pathogenicity in the bird hosts has not been reported, although the disease represents an important human health hazard (Olsen *et al.* 1993).

Dinoflagellates

Paralytic shellfish poisoning (PSP) is a widely-occurring illness that is a concern in temperate regions and is often responsible for the deaths of many seabirds (Okaichi *et al.* 1989, Shumway *et al.* 2003). The causative agents of PSP are dinoflagellates belonging to various genera that are toxic to animals higher in the food chain. Mortality of alcids has been recorded, although small percentages of local populations were affected (Coulson *et al.* 1968, Armstrong *et al.* 1978; Table 1). There is evidence to indicate that reduced feeding activity, reproductive failure and impaired motor functioning may also be caused by toxic algae (Shumway *et al.* 2003).

Many dinoflagellates can cause PSP, but the only species reported to have caused mortality in alcids is *Protogonyaulax tamarensis* (Coulson *et al.* 1968, Armstrong *et al.* 1978). The actual incidences and impacts of toxic algal blooms are probably not adequately reported (Shumway *et al.* 2003).

Apicomplexa and Microsporidia (Protozoa)

Plasmodium spp. are widely-occurring blood parasites that cause avian malaria (Davis *et al.* 1971). The alcids have been searched extensively for this “protozoan” and other blood parasites, but only Common Murres have so far shown evidence of infection (Table 1). The lack of adequate vectors for the transmission of the parasite may be a reason for the very few instances of *Plasmodium* in auks.

Another protozoan, *Eimeria fraterculae*, recorded from Atlantic Puffins, causes renal coccidiosis that produces a range of morphologic abnormalities in the kidneys (Leighton & Gajadhar 1986). Little is known about the prevalence and host specificity of *Eimeria fraterculae* and its occurrence in other alcids.

The protozoan *Sarcocystis* has caused mortality in wild-caught Common Murre chicks (Enstipp & Grémillet 2002). Mortality of wild-caught captive Horned Puffin *F. corniculata* chicks has also been attributed to infections caused by unidentified Microsporidia (Tocidlowski *et al.* 1997). Nothing is otherwise known about the protozoan parasites of alcids in the wild.

Fungi

Wild-caught Tufted Puffins *F. cirrhata* may harbor *Aspergillus fumigatus*, and alcids and penguins are generally believed to be very susceptible to aspergillosis in captivity (Monroe *et al.* 1994; Table 1). Alcid species reported to be infected in captivity are Pigeon Guillemots *Cephus columba*, Rhinoceros Auklets *Cerorhinca monocerata* and Common Murres, with the chicks being more susceptible to infection than the adults (Stoskopf 1993). In the wild, Common Murres may be infected with *Aspergillus* spp. (Petermann *et al.* 1989), although fungal diseases are generally less common in alcids than are infections with other parasitic groups (e.g. Threlfall 1971, Petermann *et al.* 1989, Stoskopf 1993, Debacker *et al.* 1997).

Platyhelminthes (Digenea and Eucestoda)

Platyhelminths are the most important component of the endoparasite fauna of auks. They comprise two major subgroups: the digeneans (flukes; Table 2) and the cestodes (tapeworms; Table 3). Although the digeneans seemingly represent the most species-rich parasitic group occurring in the alcids (Table 2), many of the records are incidental, thereby inflating the true species richness (Hoberg 1996).

Digeneans

Digeneans of vertebrates generally have a life cycle involving one or two intermediate hosts (Dogiel 1964). The eggs released in the environment may be eaten by the first intermediate host (an invertebrate). Alternatively, the eggs may hatch into a larval stage that enters the first intermediate host either as food or by penetration. The transformation into the subsequent developmental stage of the digenean takes place within the first intermediate host. Entry into the second intermediate host (usually an invertebrate or a vertebrate) takes place when the first intermediate host is eaten by the second. Further development in the second intermediate host results in the formation of the final infective stage of the digenean, which enters the final avian host and transforms into the adult digenean, the eggs of which are evicted with the feces. Across the spectrum of digeneans, a certain degree of variation occurs within this general life cycle.

Adult digeneans recorded in alcids occur in the intestine, gall bladder and kidneys, and usually have one or two intermediate hosts (e.g. Dubois 1953, Bykhovskaia–Pavlovskaja 1962, Hoberg 1981, Hoberg 1984e, Nolsø 2002). *Gymnophallus deliciosus*, for example, occurs mainly in the gall bladders of Atlantic Puffins as adults (Bykhovskaia–Pavlovskaja 1962, Nolsø 2002) and usually has bivalve mollusks as first and second intermediate hosts. Some *Gymnophallus* spp. may also use polychaetes as intermediate hosts. Digeneans such as *Cryptocotyle lingua* utilize snails and fish as their first and second intermediate hosts respectively (Stunkard 1930).

Occurrence of digeneans is relatively rare in auks and other pelagic seabirds and tends to be associated with foraging patterns and prey diversity (Hoberg 1996). *Peusdogymnophallus* is the only digenean that is unique to the auks, being recorded from the puffins and auklets (Hoberg 1981). Hoberg (1984e) considered most digeneans other than *Renicola* spp. and *Pseudogymnophallus alcae* to be incidental. Large numbers of digeneans of other genera (such as *Gymnophallus*) recorded from the alcids suggest that many may be common in auks, but are underreported (Nolsø 2002).

TABLE 2
Platyhelminthes Digenea^a parasites of auks and possible intermediate hosts

Species	Host species	Intermediate host(s)	Locality	References
Digenea Strigeidae				
<i>Cotylurus cornutus</i>	<i>Alca torda</i>	Freshwater mollusks	Europe, N. America	Dubois 1953
<i>Cotylurus pileatus</i>	<i>A. torda</i> , <i>Uria aalge</i>	Freshwater mollusks	Eurasia	Dubois 1953
<i>Ichthyocotylurus erraticus</i>	<i>Fratercula corniculata</i>	Freshwater mollusks	Ugaiushak I., AK, USA	Dubois 1981, Hoberg 1984e
<i>Ichthyocotylurus platycephalus</i>	<i>A. torda</i> , <i>U. aalge</i> , <i>Cepphus grylle</i>	Freshwater mollusks	England; Griefswald, Germany	Dawes 1946, Dubois 1953, Dubois 1978
<i>Apatemon minor</i>	<i>C. grylle</i>		Barents Sea	Bykhovskaia–Pavlovskaja 1962
<i>Apatemon somateriae</i>	<i>F. cirrhata</i>	Mollusks	Kodiak I., AK, USA	Dubois 1981, Hoberg 1984e
Echinostomatidae				
<i>Himasthla</i> sp.	<i>F. corniculata</i>	Lamellibranchs and gastropods	Ugaiushak I., AK, USA	Hoberg 1984e
<i>Echinostoma</i> sp.	<i>C. grylle</i>		England	Nicoll 1923
<i>Stephanoprora denticulata</i>	<i>A. torda</i>	Mollusk and fish species	England	Nicoll 1923, Dawes 1946
<i>Stephanoprora spinosa</i>	<i>C. grylle</i>		No locality	von Linstow 1878 in Hoberg 1984e
Psilostomatidae				
<i>Sphaeridiotrema globulus</i>	<i>A. torda</i>	Gastropods	Europe, N. America	Dawes 1946
Microphallidae				
<i>Maritrema afanassjewi</i>	<i>F. cirrhata</i>		N. Pacific	Skrjabin & Mamaev 1968
<i>Pseudospelotrema uriae</i>	<i>C. carbo</i>		Rimsky–Korsakov Is., Vladivostok, Russia	Alekseev & Smetanina 1970
<i>Pseudospelotrema japonicum</i>	<i>C. columba</i>		Komandorski Is., Russia	Yamaguti 1939
<i>Spelotrema arenaria</i>	<i>F. arctica</i>	Amphipods and gastropods	East Murmansk, Russia	Yamaguti 1975
<i>Liliatrema skrjabini</i>	<i>C. carbo</i>		Kuril I., Russia	Bykhovskaia–Pavlovskaja 1962
<i>Liliatrema sobolevi</i>	<i>F. cirrhata</i>		Sea of Okhotsk	Skrjabin & Mamaev 1968, Belogurov <i>et al.</i> 1968
<i>Microphallus pirum</i>	<i>C. columba</i> , <i>F. cirrhata</i>		St Matthew I., AK, USA	Skrjabin & Mamaev 1968, Hoberg 1984e
<i>Microphallus</i> sp.	<i>C. columba</i>		Gray's Canyon, WA, USA	Hoberg 1984e
Diplostomatidae				
<i>Diplostomum mahonae</i>	<i>U. aalge</i>	Gastropod and fish species	Europe	Dubois 1953
<i>Diplostomum spathaceum</i>	<i>A. torda</i> , both <i>Uria</i> spp.	Gastropod and fish species	Europe; W Aleutian Is., AK, USA	Dubois 1953, Hoberg 1984e

TABLE 2 (continued)

Species	Host species	Intermediate host(s)	Locality	References
Opisthorchiidae				
<i>Metorchis xanthosomus</i>	<i>A. torda</i>	Mollusk and fish species	England	Nicoll 1923, Dawes 1946
Gymnophallidae				
<i>Gymnophallus deliciosus</i>	<i>Aethia pusilla</i> , <i>F. arctica</i>	Polychaetes, mollusks	Barents Sea, Kuril I., Rimsky–Korsakov Is., Vladivostok, Russia	Beloposkaya 1952, Bykhovskaia–Pavlovskaja 1962, Alekseev & Smetanina 1970
<i>Gymnophallus</i> sp.	<i>F. arctica</i>	Polychaetes, mollusks	Nolsoy, Faeroe Is.	Nolsø 2002
<i>Pseudogymnophallus alcae</i>	<i>A. torda</i> , <i>F. cirrhata</i> , <i>F. corniculata</i> , <i>Cyclorhynchus</i> <i>psittacula</i> , <i>Aethia cristatella</i> , <i>Aethia pusilla</i>	Polychaetes, mollusks	St Mary's I., QC, Canada; Cape Thompson, St Lawrence I., St Paul I., St Matthew I., W Aleutian Is., Ugaiushak I., USA	Hoberg 1981, Hoberg 1984e
Schistosomatidae				
<i>Ornithobilharzia lari</i>	<i>U. aalge</i>	Marine prosobranchs	Witless Bay, NL, Canada	Threlfall 1971
Prosthogonimidae				
<i>Prosthogonimus ovatus</i>	<i>C. grylle</i>	Freshwater snails, dragonfly larvae	England	Dawes 1946
Heterophyidae				
<i>Cryptocotyle lingua</i>	Both <i>Uria</i> spp., <i>C. carbo</i> , <i>C. grille</i> , <i>F. arctica</i> , <i>A. torda</i>	Gastropod and fish species	Widespread: Arctic, N. Atlantic, N. Pacific	Dawes 1946, Bykhovskaia– Pavlovskaja 1962, Skrjabin & Mamaev 1968, Alekseev & Smetanina 1970, Threlfall 1971
<i>Cryptocotyle concavum</i>	<i>A. torda</i>		England	Dawes 1946
<i>Cryptocotyle</i> sp.	<i>U. aalge</i>		Sequim Bay, Washington	Hoberg 1984e
<i>Galactosomum humbargari</i>	<i>F. corniculata</i> , <i>C. monocerata</i>	Fish species	Protection I., WA, and Ugaiushak I., AK, USA	Hoberg 1984e
<i>Galactosomum</i> sp.	<i>Brachyramphus</i> <i>marmoratus</i>		Point Roberts, WA, USA	Hoberg 1984e

TABLE 2 (continued)

Species	Host species	Intermediate host(s)	Locality	References
Renicolidae				
<i>Renicola keimahuri</i>	<i>C. carbo</i> , <i>F. cirrhata</i>	Mollusks	Sea of Okhotsk, Barents Sea	Yamaguti 1939, Belogurov <i>et al.</i> 1968
<i>Renicola quinta</i>	<i>Synthliboramphus antiquus</i> , <i>C. carbo</i>		Kuril I., Russia	Bykhovskaia–Pavlovskaja 1962
<i>Renicola umigarasu</i>	Both <i>Uria</i> spp., <i>S. antiquus</i> , <i>C. carbo</i> , <i>Aethia pusilla</i> , <i>F. cirrhata</i>		Kuril I., Rimsky–Korsakov Is., Vladivostok, Russia	Yamaguti 1939, Bykhovskaia–Pavlovskaja 1962, Skrjabin & Mamaev 1968, Alekseev & Smetanina 1970
<i>Renicola vladica</i>	<i>F. corniculata</i> , <i>F. cirrhata</i> , <i>S. antiquus</i>		Sea of Okhotsk, Kuril I., Rimsky–Korsakov Is., Russia	Bykhovskaia–Pavlovskaja 1962, Skrjabin & Mamaev 1968, Belogurov <i>et al.</i> 1968
<i>Renicola</i> sp.	<i>Cyclorhynchus psittacula</i>		Primore, N. Pacific	Oshmarin 1963 in Hoberg 1984e
^a Digeneans or flukes.				

TABLE 3
Platyhelminthes: Eucestoda^a of auks and possible intermediate hosts

Species	Host species	Intermediate host(s)	Locality	References
Eucestoda				
Pseudophyllidea, Diphylobothriidae				
<i>Diphylobothrium dendriticum</i>	<i>Uria lomvia</i>		Cape Thompson, AK, USA	Swartz 1960 in Hoberg 1984e
<i>Diphylobothrium</i> sp.	<i>Fratercula cirrhata</i> , <i>Cephus monocerata</i> , <i>U. aalge</i>	Copepoda and fish species	Kodiak I., western Aleutian I., Alaska	Hoberg 1984e
<i>Schistocephalus solidus</i>	<i>Alca torda</i> , <i>U. aalge</i> , <i>C. grylle</i>	<i>Gasterosteus aculeatus</i>	No locality	Joyeux & Baer 1936
Tetrabothriidea, Tetrabothriidae				
<i>Tetrabothrius cylindraceus</i>	Both <i>Uria</i> spp.	Unknown: cephalopods, crustaceans, teleosts	Greenland; Funk I. and Witless Bay, NL, Canada; Labrador Sea	Ditlevsen 1917, Römer & Schaudinn 1918, Threlfall 1971
<i>Tetrabothrius erostris</i>	<i>U. aalge</i> , <i>C. grille</i> , <i>C. columba</i>	Unknown: cephalopods, crustaceans, teleosts	Greenland; Funk I. and Witless Bay, NL, Canada; Labrador Sea	Ditlevsen 1917, Römer & Schaudinn 1918, Threlfall 1971

TABLE 3 (continued)

Species	Host species	Intermediate host(s)	Locality	References
<i>Tetrabothis jagerskioldi</i>	<i>A. torda</i> , both <i>Uria</i> spp., <i>Brachyramphus marmoratus</i> , <i>Synthliboramphus antiquus</i> , all <i>Cepphus</i> spp., all <i>Fratercula</i> spp.	Unknown: cephalopods, crustaceans, teleosts	Greenland; Funk I. and Witless Bay, NL, Canada; Labrador Sea; Barents Sea	Joyeux & Baer 1936; Baer 1956; Threlfall 1971; Temirova & Skjarbin 1978; Smetanina 1979; Belopolskaya 1952, 1963
<i>Tetrabothis macrocephalum</i>	<i>U. aalge</i> , <i>B. marmoratus</i>		Sakhalin I., Russia	Yamaguti 1959
<i>Tetrabothis</i> spp.	Both <i>Uria</i> spp., <i>Alle alle</i> , <i>S. antiquus</i> , <i>B. marmoratus</i> , <i>Ptychoramphus aleuticus</i> , <i>Aethia cristatella</i> , <i>Aethia pusilla</i> , <i>C. columba</i> , <i>C. grille</i> , <i>F. cirrhata</i> , <i>F. corniculata</i> , <i>C. monocerata</i> , <i>Cyclorhynchus psittacula</i>	Unknown: cephalopods, crustaceans, teleosts	Arctic, N. Atlantic, N. Pacific	Threlfall 1971, Hoberg 1984d, Hoberg 1984e
Cyclophyllidea				
Dilepididae				
<i>Paricterotaenia</i> sp.	<i>U. lomvia</i>		Cape Thompson, Alaska	Swartz 1960 in Hoberg 1984e
<i>Lateriporus</i> sp.	<i>U. lomvia</i>		Cape Thompson, AK, USA	Swartz 1960 in Hoberg 1984e
<i>Alcataenia armillarlis</i>	Both <i>Uria</i> spp., <i>A. torda</i> , <i>F. corniculata</i> , <i>C. monocerata</i> , <i>Aethia pusilla</i> , <i>C. columba</i>	Euphausiid crustaceans	Widespread: Arctic, N. Atlantic, N. Pacific	Ditlevsen 1917, Römer & Schaudinn 1918, Hoberg 1979, Baer 1956, Markov 1937, Hoberg 1984a, Threlfall 1971
<i>Alcataenia atlantiensis</i>	<i>A. torda</i>		N. Pacific	Hoberg 1991
<i>Alcataenia campylacantha</i>	All <i>Cepphus</i> spp., both <i>Uria</i> spp.	Euphausiid crustaceans	Widespread: Arctic, N. Atlantic, N. Pacific	Ditlevsen 1917, Römer & Schaudinn 1918, Joyeux & Baer 1936, Threlfall 1971, Hoberg 1984d, Hoberg 1984e
<i>Alcataenia cerorhincae</i>	<i>C. monocerata</i>	Euphausiid crustaceans	Destruction & Protection Is. and Grays Harbor, WA, USA; Forrester I., W Aleutian Is., AK, USA	Hoberg 1984b

TABLE 3 (continued)

Species	Host species	Intermediate host(s)	Locality	References
<i>Alcataenia fraterculae</i>	<i>U. aalge</i> , <i>F. cirrhata</i> , <i>F. corniculata</i> , <i>Aethia cristatella</i> , <i>C. monocerata</i>	Euphausiid crustaceans	Buldir I., Ugaiushak I., Kodiak I., Amchitka I., St Paul I., St Matthew I., St Lawrence I., Cape Thompson, AK, USA	Hoberg 1984b
<i>Alcataenia larina</i>	<i>Aethia pygmaea</i> , <i>Aethia cristatella</i> , <i>U. aalge</i>	Euphausiid crustaceans	Widespread: Arctic, N. Atlantic, N. Pacific	Smetanina 1979, Belopolskaya 1952, Hoberg 1984b, Hoberg, 1984e
<i>Alcataenia longicervica</i>	Both <i>Uria</i> spp., <i>F. corniculata</i> , <i>F. cirrhata</i>	Euphausiid crustaceans	St Matthew I., Buldir I., W Aleutian Is., Ugaiushak I., Kodiak I., Semidi Is., St Paul I., St Lawrence I., Cape Thompson, AK, USA; Humboldt Bay, CA, USA	Hoberg 1984a
<i>Alcataenia meinertzhageni</i>	<i>C. grille</i> , both <i>Uria</i> spp.	Euphausiid crustaceans	Greenland; Peter the Great Bay, Russia; Buldir I., St Matthew I., AK, USA; Sequim Bay, WA, USA	Baer 1956, Threlfall 1971, Smetanina 1979, Hoberg 1984a
<i>Alcataenia micracantha</i>	<i>U. aalge</i> , <i>C. grylle</i>	Euphausiid crustaceans	Barents Sea; NW Atlantic	Baylis 1919, Threlfall 1971, Hoberg 1984a
<i>Alcataenia pygmaeus</i>	<i>Aethia pygmaea</i>	Euphausiid crustaceans	Buldir I., AK, USA	Hoberg 1984c
<i>Alcataenia</i> spp.	<i>B. brevirostris</i> , both <i>Uria</i> spp., <i>Alle alle</i> , <i>C. monocerata</i> , <i>Aethia cristatella</i> , <i>F. cirrhata</i> , <i>P. aleuticus</i>	Euphausiid crustaceans	N. Atlantic, N. Pacific	Threlfall 1971, Hoberg 1984d, Hoberg 1984e
<i>Anomotaenia</i> cf. <i>clavigera</i>	<i>P. aleuticus</i>		Grays Harbour, WA, USA	Hoberg 1984e
<i>Microsomacanthus</i> cf. <i>ductilis</i>	<i>B. marmoratus</i> , <i>C. columba</i>		Amchitka I., St Lawrence I., AK, USA	Hoberg 1984e
<i>Choanotaenia stercorarii</i>	<i>F. arctica</i>		Barents Sea	Belopolskaya 1952
Hymenolepididae				
<i>Aploparaksis</i> cf. <i>hirsutus</i>	<i>F. cirrhata</i>		Nunivak I., AK, USA	Hoberg 1984e
<i>Diorchis pelagicus</i>	<i>Aethia pygmaea</i> , <i>Aethia cristatella</i>	Crustaceans	Buldir I., AK, USA	Hoberg 1982, Hoberg 1984e
^a Cestodes or tapeworms.				

Feeding specialization, geographic distribution patterns of first and second intermediate hosts and the mobility of certain developmental stages (such as cercariae) have been postulated to limit the interaction between the digeneans and their auk hosts to oceanic islands (Hoberg 1996). Transmission and prevalence of digeneans in pelagic habitats are restricted, because the most frequently used intermediate hosts generally occur in the littoral zone, making islands the enzootic focus of infection (Bustnes & Galaktionov 1999). Many of the digeneans strongly “shape” the populations of their intermediate hosts by selectively causing mortality (Lauckner 1987, Davies & Knowles 2001). Changes in populations of intermediate hosts, in turn, may influence seabird feeding patterns and hence transmission of the digeneans between seabirds and their intermediate hosts (Bustnes & Galaktionov 1999, Bustnes *et al.* 2000).

Because of their more diverse feeding habits, gulls of various species harbor far greater digenean loads than do auks (e.g. Threlfall 1968, 1971). The parasites common to both gulls and auks are more typically found in auks breeding in colonies that have large gull populations (Belopolskaya 1952, Galaktionov 1995). Those findings suggest that gulls may be critical in forming a link between auks and their digenean fauna (Galaktionov 1995).

Cestodes

The life cycles of cestodes are similar to those of digeneans, in that the cestodes also utilize one or more intermediate hosts, with development to a subsequent stage taking place in each intermediate host (Dogiel 1964). Cestodes from three orders are known to occur in the intestines of alcids (Table 3). The genus *Alcataenia* has received particular attention from taxonomic, biogeographic and evolutionary perspectives (Hoberg 1984a, 1984b, 1984c, 1984d, 1986a). However, details of the prevalence, intensity and life cycles of the various species in the genus remain largely unknown.

It is evident that many *Alcataenia* species are host-specific; cestodes found in an incidental host fail to develop to maturity (Hoberg 1984a, 1984b). Cysticeroids (a developmental stage occurring in one of the intermediate hosts) of *Alcataenia armillaris* have been reported from *Thysanoessa inermis* [Crustacea: Euphausiidae (Shimazu 1975)], which forms an important component of the diet of some alcids (Gaston & Jones 1998). Prevalence is high among various alcid hosts in areas where these prey are infected with cysticeroids (Hoberg 1984c, Hoberg 1984e).

A study of similar platyhelminths of Greater Shearwaters *Puffinus gravis* suggests that platyhelminths of pelagic seabirds may be acquired before the birds migrate to northern latitudes in summer, depending on the consumption of infected invertebrates in warmer waters (Hoberg & Ryan 1989). Long-term patterns in the distribution of cestodes of auks may also be linked to the abundance of gulls, fish prey and foraging ecology (Galaktionov 1995).

Digenean and cestode infections in alcids are assumed to be low in pathogenicity, although evidence is lacking. One reason for the assumption could be the apparent low rate of platyhelminth infections in the alcids relative to other seabirds. Threlfall (1971), for example, investigated the prevalence and abundance of several platyhelminth genera from alcids in Newfoundland and concluded that alcids are relatively less infected by platyhelminths than are

gulls in the same area. Hoberg (1979) found a similar higher prevalence and abundance of platyhelminths in gulls than in alcids. The apparent absence of negative impact on host fitness suggests that digenean parasite burdens may be below some threshold value that causes declines in host fitness (Hoberg & Ryan 1989).

Acanthocephala

Acanthocephalans recorded from alcids (Table 4) are generally rare and are usually considered incidental (Threlfall 1971; Hoberg 1984e, 1986b). Some species of the genus *Corynosoma* have been found in seabirds in the Antarctic, but the feeding behavior of hosts and oceanographic factors may be responsible for restricting the range of species (Hoberg 1986b). Life cycles of few *Corynosoma* spp. have been determined. The intermediate hosts utilized by Antarctic *Corynosoma* include nearshore fish and amphipods. That genus does not use euphausiids as intermediate hosts, reducing the chances of infection in auks. Furthermore, acanthocephalans of the genus are primarily parasites of marine mammals, making auk infections very rare (Hoberg 1984e).

Nematodes (roundworms)

Nematodes representing four orders—most of which have life cycles involving more than one host—have been collected and described from auks (Table 4). *Eufilaria lari*, isolated from murre, for example, use ceratopogonid flies as vectors (Sonin 1966). *Eulimdana* spp. are believed to be transmitted by amblyceran louse vectors (Bartlett *et al.* 1989, Bartlett 1993; also see “Phthiraptera” later in this subsection). However, most nematodes of auks use aquatic invertebrates and fish species as intermediate hosts (Norris & Overstreet 1976, Anderson 2000). A wide spectrum of invertebrates and many fish species have been reported to carry larval *Contracaecum* (Norris & Overstreet 1976). *Contracaecum* spp. *Seuratia shipleyi* and *Stegophorus* spp. are widespread (Hoberg & Ryan 1989), but have been reported from only a few auk species (Table 4). That rare occurrence could again be attributable to the dietary specializations in auks as compared with gulls and other seabirds (Hoberg 1996, Gaston & Jones 1998).

Nematodes that occur in the alimentary tract of the definitive host may in many cases cause lesions, ulcerations and tumors (Nagasawa *et al.* 1998b, Anderson 2000). Experimentally infected Ring-billed Gulls *Larus delawarensis* showed that larval *Cosmocephalus obvelatus* larvae initially invaded the proventriculus of the host; but, after moulting into the fourth larval stage, they were found across the entire length of the esophagus. Transmission to young seabird chicks may occur when the chicks consume larvae regurgitated by the adults during feeding. *Cosmocephalus obvelatus*, in conjunction with other endoparasites (for example, the cestode *Tetrabothrius*) may have negative impacts on gull hosts (Bosch *et al.* 2000). Whether similar negative effects occur in auks infected with those endoparasites is not clear (Nagasawa *et al.* 1998b).

ARTHROPODA

Pentastomida

The Pentastomida represents a class of specialized crustaceans that are strictly parasitic (Bakke 1972; Table 5). *Reighardia sterna* is surmised to have a life cycle involving a single fish intermediate host (Threlfall 1971). Hoberg (1987) suggested that transmission of this pentastome occurs during the nesting season in sub-Antarctic gulls and that similar factors may control distribution patterns in

TABLE 4
Acanthocephala and Nematoda reported from the auks and their possible intermediate hosts

Species	Host species	Intermediate host(s)/vectors	Locality	References
Acanthocephala				
Polymorphidae				
<i>Polymorphus magnus</i>	<i>F. corniculata</i>	Amphipods	Komandorski I., Russia	Tsimbaliuk 1965
<i>Polymorphus minutus</i>	<i>Alle alle</i> , <i>U. lomvia</i> , <i>C. carbo</i> , <i>C. grille</i> , <i>F. cirrhata</i>		N. Pacific	Skrjabin & Mamaev 1968
<i>Polymorphus phippii</i>	<i>F. corniculata</i>		Barents Sea	Skrjabin & Ryzhikov 1973
<i>Corynosoma semerme</i>	<i>U. lomvia</i>	Crustacean fish	Komandorski I., Russia	Tsimbaliuk 1965
<i>Corynosoma strumosum</i>	<i>F. cirrhata</i> , <i>U. aalge</i> , <i>C. columba</i>	Crustacean fish	Komandorski I., Russia	Tsimbaliuk 1965
<i>Corynosoma villosum</i>	<i>U. aalge</i>		Komandorski I., Russia	Tsimbaliuk 1965
<i>Corynosoma</i> sp.	Both <i>Uria</i> spp., <i>F.</i> <i>cirrhata</i> , <i>C. columba</i>		St Matthew I., Kodiak I., AK, USA	Hoberg 1984e
<i>Bulbosoma</i> sp.	Both <i>Uria</i> spp.			Hoberg 1984e
Echinorhynchidae				
<i>Echinorhynchus alcae</i>	<i>A. torda</i>		Greenland	Wesenburg–Lund 1917
Nematoda				
Acuardiidae				
<i>Stegophorus stellae-polaris</i>	Both <i>Uria</i> spp., <i>C. carbo</i> , <i>C. columba</i> , <i>Alle alle</i> , <i>A. torda</i> , <i>F. cirrhata</i> , <i>S. antiquus</i> , <i>Cyclorhynchus</i> <i>psittacula</i> , <i>Aethia</i> <i>pygmaea</i>		Widespread: N. Atlantic, N. Pacific	Schmidt 1964, Alekseev & Smetanina 1968, Threlfall 1971, Barus <i>et al.</i> 1978, Nagasawa <i>et al.</i> 1998b
<i>Stegophorus stercorarii</i>	<i>A. torda</i> , <i>U. lomvia</i> , <i>C. grylle</i> , <i>Aethia</i> <i>pusilla</i> , <i>Aethia</i> <i>crisatella</i> , <i>F. cirrhata</i>		Wrangel I. and Rimsky–Korsakov Is., Russia	Alekseev & Smetanina 1968, Leonov & Shvetsova 1970, Barus <i>et al.</i> 1978
<i>Cosmocephalus imperialis</i>	<i>U. aalge</i>	Amphipods	Sakhalin I., Russia	Morishita 1930
<i>Cosmocephalus obvelatus</i>	<i>A. torda</i> , both <i>Uria</i> spp., <i>C. grille</i> , <i>C. carbo</i> , <i>C. monocerata</i>	Amphipods	N. Atlantic, N. Pacific	Cram 1927, Barus <i>et al.</i> 1978, Threlfall 1971
<i>Paracuararia adunca</i>	<i>A. torda</i>	Amphipods	No locality	Anderson & Wong 1982

TABLE 4 (continued)

Species	Host species	Intermediate host(s)/vectors	Locality	References
<i>Paracuaria tridentata</i>	<i>U. aalge</i> , <i>Cyclorhynchus psittacula</i> , <i>Aethia pygmaea</i> , <i>C. columba</i>	Amphipods	Komandorski I., Russia	Gibson 1968, Tsimbaliuk & Belogurov 1964, Barus <i>et al.</i> 1978
<i>Skrjabinocerca prima</i>	<i>U. aalge</i> , <i>F. cirrhata</i>		N. Pacific	Skrjabin & Mamaev 1968
<i>Streptocara crassicauda</i>	<i>A. torda</i> , <i>U. lomvia</i> , <i>C. grylle</i> , <i>C. columba</i> , <i>Aethia cristatella</i> , <i>F. corniculata</i> , <i>F. cirrhata</i>	Amphipods and fish species	Widespread: Arctic, N. Atlantic, N. Pacific	Cram 1927, Belopolskaya 1952, Gibson 1968, Threlfall 1971, Leonov & Shvetsova 1970
<i>Streptocara</i> sp.	<i>F. arctica</i>		Barents Sea; Nolsoy, Faeroe Is.	Belopolskaya 1952, Nolsø 2002
<i>Seuratia shipleyi</i>	<i>F. arctica</i> , <i>F. corniculata</i>	Crustaceans	Widespread: N. Atlantic, N. Pacific	Threlfall 1971, Barus <i>et al.</i> 1978
<i>Seuratia</i> sp.	<i>F. arctica</i> , <i>F. corniculata</i>	Crustaceans	Nolsoy, Faeroe Is.	Swartz 1960 in Hoberg 1984e, Nolsø 2002
Ascaridae				
<i>Ascaris spiculigera</i>	<i>A. torda</i> , <i>U. aalge</i>		Greenland	Römer & Schaudinn 1918
Heterakidae				
<i>Heterakis kurilensis</i>	<i>Aethia cristatella</i>		No locality	Barus <i>et al.</i> 1978
Diectophymidae				
<i>Eustrongylides ignotus</i>	<i>U. aalge</i>		Sakhalin I., Russia	Morishita 1930
<i>Eustrongylides mergorum</i>	<i>A. torda</i> , both <i>Uria</i> spp., <i>C. grylle</i>	Invertebrates, fish species	Novaya Zemlaya, Russia; Greenland; NW Atlantic	Markov 1941, Threlfall 1971, Barus <i>et al.</i> 1978
<i>Eustrongylides tubifex</i>	<i>A. torda</i> , <i>U. aalge</i>		No locality	Barus <i>et al.</i> 1978
Anisakidae				
<i>Anisakis</i> sp.	<i>U. aalge</i> , <i>F. cirrhata</i>	Crustaceans, cephalopods and anadromous/marine teleosts	Rimsky-Korsakov Is., Russia; Greenland; Funk I. and Witless Bay, NL, Canada; Labrador Sea	Alekseev & Smetanina 1968, Threlfall 1971, Barus <i>et al.</i> 1978
<i>Contracaecum osculatum rudolphi</i>	<i>A. torda</i> , both <i>Uria</i> spp., <i>S. antiquus</i> , <i>C. carbo</i> , <i>C. grille</i> , <i>F. cirrhata</i> , <i>F. corniculata</i>		Widespread: Arctic, N. Atlantic, N. Pacific	Threlfall 1971, Belopolskaya 1952, Alekseev & Smetanina 1968, Leonov & Shvetsova 1970, Belogurov <i>et al.</i> 1968
<i>Contracaecum spiculigerum</i>	<i>A. torda</i> , both <i>Uria</i> spp., <i>C. carbo</i> , <i>F. cirrhata</i>	Wide range of invertebrates and fish species	Greenland; Newfoundland; Labrador Sea; N. Pacific	Skjarbin & Mamaev 1968, Threlfall 1971
<i>Contracaecum variegatum</i>	<i>A. torda</i> , both <i>Uria</i> spp., <i>F. cirrhata</i>	Wide range of invertebrates and fish specie	Bering Sea	Nagasawa <i>et al.</i> 1998a

TABLE 4 (continued)

Species	Host species	Intermediate host(s)/vectors	Locality	References
<i>Contracecum</i> sp.	<i>U. aalge</i> , <i>F. arctica</i>	Wide range of invertebrates and fish species	N. Atlantic, N. Pacific	Threlfall 1971, Nagasawa <i>et al.</i> 1998a, Nolsø 2002
Onchocercidae				
<i>Eulimdana lari</i>	Both <i>Uria</i> spp., <i>C. grylle</i>	Louse species	Wrangel I.	Sonin 1966, Leonov & Shvetsova 1970
<i>Eufilaria lari</i>	Both <i>Uria</i> spp., <i>C. grylle</i>	Ceratopogonid flies	Barents Sea; Komandorski I., Russia	Belopolskaya 1952, Tsimbaliuk & Belogurov 1964
Filariidae				
<i>Capillaria contorta</i>	<i>U. lomvia</i> , <i>C. grylle</i>		Greenland	Römer & Schaudinn 1918, Barus <i>et al.</i> 1978
<i>Capillaria</i> sp.	<i>F. arctica</i>		Barents Sea	Belopolskaya 1952

TABLE 5
Arthropod parasites of auks

Species	Host species	Locality	References
Crustacea			
Pentastomida Reighardiidae			
<i>Reighardia lomviae</i>	<i>Uria aalge</i> , <i>Fratercula arctica</i>	Norway; Faeroe Is.	Dyck 1975, Bakke 1978
<i>Reighardia sterna</i>	<i>F. arctica</i> , <i>U. aalge</i>	Greenland; Newfoundland; Labrador Sea	Threlfall 1971
<i>Reighardia</i> sp.	Both <i>Uria</i> spp., <i>Synthliboramphus antiquus</i> , <i>F. cirrhata</i> , <i>F. arctica</i> , <i>Aethia cristatella</i>	Nolsoy, Faeroe Is.; Kodiak I., St Lawrence I., Ugaiushak, AK, USA; Humboldt Bay, CA, USA; Sea of Okhotsk	Hoberg 1984e, Nolsø 2002
Acari			
Metastigmata Ixodidae			
<i>Ixodes (Ceratiixodes) uriae</i>	<i>Alca torda</i> , all <i>Fratercula</i> spp., both <i>Uria</i> spp., <i>Ptychoramphus aleuticus</i> , <i>Cyclorhynchus psittacula</i> , <i>Cephus monocerata</i>	Arctic, N. Atlantic, N. Pacific	Eveleigh & Threlfall 1975, Hoberg & Wehle 1982, Choe & Kim 1987, Morbey 1996, Muzaffar 2000
<i>Ixodes signatus</i>	Both <i>Uria</i> spp., <i>F. cirrhata</i>	Buldir I., Pribilof Is., AK, USA	Hoberg & Wehle 1982, Choe & Kim 1987
Argasidae			
<i>Ornithodoros capensis</i>	<i>U. aalge</i>	Pacific	Clifford 1979
<i>Ornithodoros</i> sp. near <i>denmarki</i>	<i>U. aalge</i>	Oregon coast	Clifford 1979
<i>Ornithodoros maritimus</i>	<i>U. aalge</i> , <i>A. torda</i>	N. Atlantic	Hoogstraal <i>et al.</i> 1976
Hypoderidae			
<i>Thalassornectes (Alcicletes) aukletae</i>	<i>Aethia cristatella</i> , <i>C. psittacula</i>	N. Pacific	Pence & Hoberg 1991

TABLE 5 (continued)

Species	Host species	Locality	References
Mesostigmata			
<i>Rhinonyssus caledonicus</i>	Both <i>Uria</i> spp.	Pribilof Is., AK, USA	Choe & Kim 1987
Prostigmata			
<i>Neotrombicula</i> sp.	Both <i>Uria</i> spp.	Pribilof Is., AK, USA	Choe & Kim 1987
Astigmata Alloptidae			
<i>Alloptes conurus</i>	Both <i>Uria</i> spp.	Pribilof Is., AK, USA	Choe & Kim 1991
<i>Alloptes crassipes</i>	<i>A. torda</i> , <i>F. arctica</i> , both <i>Uria</i> spp., <i>C. grylle</i>	N. Atlantic	Canestrini & Kramer 1899, Römer & Schaudinn 1918
<i>Alloptes fraterculae</i>	<i>F. arctica</i>	Barents Sea	Belopolskaya 1952
<i>Alloptes phaethontis</i>	<i>F. arctica</i>	Greenland	Römer & Schaudinn 1918
<i>Alloptes</i> spp.	<i>A. torda</i> , <i>F. arctica</i> , both <i>Uria</i> spp.	Gannet I. and Gull I., NL, Canada	Eveleigh 1974, Muzaffar 2000
Laminosioptidae			
<i>Calamioptes</i> sp.	Both <i>Uria</i> spp.	Pribilof Is., AK, USA	Choe & Kim 1987
Dermanyssidae			
<i>Dermanyssus gallinae</i>	<i>F. arctica</i>	N. Atlantic	Dobson 1952
Insecta			
Phthiraptera Amblycera Menoponidae			
<i>Austromenopon nigropleurum</i>	<i>A. torda</i> , <i>Alle alle</i> , <i>F. arctica</i> , <i>S. antiquus</i>	Gannet I. and Gull I., NL, Canada	Wheeler & Threlfall 1989, Eveleigh & Threlfall 1976, Eveleigh & Amano 1977, Muzaffar 2000
<i>Austromenopon phippii</i>	<i>U. lomvia</i>	Gull I., NL, Canada	Eveleigh & Threlfall 1974
<i>Austromenopon uriae</i>	<i>U. aalge</i>	Gannet I. and Gull I., NL, Canada	Timmermann 1954, Eveleigh & Threlfall 1976, Eveleigh & Amano 1977, Muzaffar 2000
<i>Austromenopon</i> sp.	<i>U. aalge</i>	N. Atlantic	Wheeler & Threlfall 1989
Ischnocera Philopteridae			
<i>Quadriceps aetherea</i>	<i>Alle alle</i> , <i>C. grylle</i> , <i>Aethia pusilla</i>	Gull I., NL, Canada	Eveleigh & Threlfall 1976, Eveleigh & Amano 1977, Timmerman 1974
<i>Quadriceps aethereus</i>	<i>Aethia pusilla</i>		Emerson 1972
<i>Quadriceps alcae</i>	<i>A. torda</i>	Gannet I. and Gull I., NL, Canada	Eveleigh & Threlfall 1976, Eveleigh & Amano 1977, Muzaffar 2000
<i>Quadriceps antiqua</i>	<i>S. antiquus</i>	Langara I., British Columbia; Petrel I., Alaska	Timmermann 1974, Hoberg & Wehle 1982
<i>Quadriceps helgovauki</i>	<i>F. arctica</i>	Gannet I. and Gull I., NL, Canada	Eveleigh & Threlfall 1976, Eveleigh & Amano 1977, Muzaffar 2000
<i>Quadriceps maritimus</i>	<i>U. lomvia</i> , <i>P. aleuticus</i>		Emerson 1972, Wheeler & Threlfall 1989
<i>Quadriceps obliquus</i>	Both <i>Uria</i> spp.	Gannet I. and Gull I., NL, Canada	Emerson 1972, Eveleigh & Threlfall 1976, Muzaffar 2000

TABLE 5 (continued)

Species	Host species	Locality	References
<i>Quadriceps</i> sp.	<i>Brachyramphus marmoratus</i> , <i>S. antiquus</i> , <i>F. corniculata</i>	Ugaiushak I. and Kodiak I., AK, USA	Hoberg & Wehle 1982, Wheeler & Threlfall 1989
<i>Saemundssonina acutipecta</i>	<i>C. monocerata</i>	OR and CA, USA	Price <i>et al.</i> 2003
<i>Saemundssonina boschi</i>	<i>Aethia pusilla</i>	St Lawrence I., St Paul I. and Buldir I., AK, USA	Price <i>et al.</i> 2003
<i>Saemundssonina calva</i>	Both <i>Uria</i> spp.	Gannet Is. and Gull Is., NL, Canada; Greenland; Faeroe Is.; AK and ME, USA	Eveleigh & Threlfall 1976, Eveleigh & Amano 1977, Muzaffar 2000, Price <i>et al.</i> 2003
<i>Saemundssonina celidoxa</i>	<i>A. torda</i>	Gannet Is. and Gull I., NL, Canada; England	Eveleigh & Threlfall 1976, Muzaffar 2000, Price <i>et al.</i> 2003
<i>Saemundssonina fraterculae</i>	<i>F. arctica</i> , <i>F. corniculata</i> , <i>F. cirrhata</i>	Gannet Is. and Gull I., NL, Canada; Buldir I. and Kodiak I., AK, USA; Faeroe Is.	Eveleigh & Threlfall 1976, Eveleigh & Amano 1977, Hoberg & Wehle 1982, Muzaffar 2000, Price <i>et al.</i> 2003
<i>Saemundssonina grylle</i>	<i>C. grylle</i> , <i>C. columba</i>	Labrador; Gull I., NL, Canada; Faeroe Is.	Eveleigh & Threlfall 1976, Eveleigh & Amano 1977, Price <i>et al.</i> 2003
<i>Saemundssonina insolita</i>	<i>P. aleuticus</i>	BC, Canada; CA, USA	Price <i>et al.</i> 2003
<i>Saemundssonina merguli</i>	<i>Alle alle</i>	Gull I., NL, Canada; Faeroe Is.; FL, USA	Eveleigh & Threlfall 1976, Eveleigh & Amano 1977, Price <i>et al.</i> 2003
<i>Saemundssonina montereyi</i>	<i>S. antiquus</i> , <i>B. marmoratus</i> , <i>P. aleuticus</i>	Bay of Monterey, CA, and Petrel I., AK, USA	Kellogg 1896, Hoberg & Wehle 1982, Price <i>et al.</i> 2003
<i>Saemundssonina wumisuzume</i>	<i>Aethia cristatella</i> , <i>Aethia pygmaea</i>	AK, USA	Price <i>et al.</i> 2003
<i>Saemundssonina</i> sp.	<i>F. cirrhata</i> , <i>B. marmoratus</i>	Buldir I. and Kodiak I., AK, USA	Hoberg & Wehle 1982
Diptera Hippoboscidae	<i>Icosta americana</i>	<i>U. lomvia</i>	Wheeler & Threlfall 1989
Muscomorpha			
Unknown sp.	<i>A. torda</i> , <i>F. arctica</i>	Gannet I., NL, Canada	Muzaffar 2000
Siphonaptera Pulicidae			
<i>Actenopsylla suavis</i>	<i>P. aleuticus</i> , <i>F. cirrhata</i>	N. Pacific	Holland 1984
<i>Spilopsyllus cuniculi</i>	<i>F. arctica</i>	W coast of England; Coronados I., CA, USA	Rothschild & Clay 1957, Guiguen <i>et al.</i> 1983
<i>Ornithopsylla laetitiae</i>	<i>F. arctica</i>	Scilly I. and Skomer & Skokoholm Is., Wales; Ireland's Eye, Great Skellig, Ireland	Rothschild & Clay 1957, Guiguen <i>et al.</i> 1983
Ceratophyllidae			
<i>Ceratophyllus vagabundus</i>	<i>U. lomvia</i>	Capes Lisburne and Thompson, AK, USA	Hoberg & Wehle 1982
<i>Ceratophyllus gallinae</i>	<i>F. arctica</i>	England	Rothschild & Clay 1957, Guiguen <i>et al.</i> 1983
<i>Ceratophyllus garei</i>	<i>F. arctica</i>	England	Rothschild & Clay 1957, Guiguen <i>et al.</i> 1983
<i>Ceratophyllus borealis</i>	<i>F. arctica</i>	England	Rothschild & Clay 1957, Guiguen <i>et al.</i> 1983
<i>Orchopeas leucopus</i>	<i>U. lomvia</i> , <i>F. arctica</i>	Gannet I., NL, Canada	Muzaffar 2000

the Northern and Southern Hemispheres. *Reighardia lomviae* has been recorded from murre (Dyck 1975), and the genus is believed to be the only pentastome to parasitize avian hosts, particularly gulls of the Northern Hemisphere (Bakke 1972). Pathogenicity in auks has not been studied, although the high load recorded from a murre and the hypothesized consumption of lung capillaries (Dyck 1975) may indicate potential negative impacts on the host.

Acari (ticks and mites)

I. uriae is known to parasitize more than 50 species of seabirds, including many alcid species (Eveleigh & Threlfall 1975, Mehl & Traavik 1983, Danchin 1992, Jaenson *et al.* 1994, Barton *et al.* 1996, Morbey 1996, McCoy *et al.* 1999, Bergström *et al.* 1999, Muzaffar 2000, Frenot *et al.* 2001, McCoy & Tirard 2002). *I. uriae* is distributed in the subpolar and temperate regions of both hemispheres (Mehl & Traavik 1983, Olsen *et al.* 1993; Table 5). It is regarded as a generalist parasite feeding on a variety of seabirds, although recent studies indicate genetic differences within species that have a preference for specific hosts (Boulinier *et al.* 2001, McCoy *et al.* 2001, McCoy *et al.* 2002, McCoy *et al.* 2003). The life cycle of *I. uriae* involves reproductive strategies to overcome environmental extremes (McCoy & Tirard 2002). Individuals may live from four years (Eveleigh & Threlfall 1975) to eight years in the low Arctic (Lvov 1980). Life cycles that are as short as two years have been recorded in penguin colonies in the Crozet Archipelago (Frenot *et al.* 2001).

All the life stages (larvae, nymph and adult) of *I. uriae* live in or on the substrate, with individuals getting onto the birds only for blood meals, which may last for 4–10 days (Eveleigh & Threlfall 1975, Baranton *et al.* 1992, Frenot *et al.* 2001). Ticks not only have a role as a reservoir and vector of *B. garinii* (Olsen *et al.* 1993, Olsen *et al.* 1995, Gylfe *et al.* 1999, discussed earlier); heavy tick infestations, particularly on chicks, may cause slowed growth and mortality (Morbey 1996, Gauthier–Clerc *et al.* 1998, Bergström *et al.* 1999). Alcid chicks infested with ticks have been known to have closed eyes, swollen heads and symptoms of paralysis (Harris 1984, Morbey 1996). The periodic peaks in tick density may have more important consequences in reproductive success, although such effects have not been studied in the alcids.

Feather mites are numerically the most abundant group of ectoparasites living on birds (Gaud & Atyeo 1996, Proctor 2003). Generally ranging in size from 0.3 mm to 0.7 mm, they are not eaten by the host, nor do they penetrate the host's skin. They occur on the surface of feathers, on feather barbs and within quills. Their entire life is spent on the host; accidental removal from the host results in death.

Feather mites have been given relatively little attention, primarily because they are generally considered to be benign to the host, feeding on feather fragments, desquamated skin scales and oily secretions (Krantz 1978, Gaud & Atyeo 1996, Proctor 2003). Fungal spores and diatoms may also form part of their diet (Dubinin 1951, Krantz 1978). High densities of feather mites may cause a depluming behavioral response because of skin irritation in some host species (Gaud & Atyeo 1996); however, the buildup of feather mite numbers could simply be an artifact of captivity (Proctor 2003). Individuals are often densely crowded at breeding sites, and a buildup of feather mites may be a consequence of coloniality (Muzaffar 2000).

The feather mite genus *Alloptes* (Astigmata: Alloptidae) has been recorded on auks (Table 5). Taxonomic uncertainty in the genus makes identification to the specific level difficult. *Alloptes crassipes* has been recorded from Razorbills and Atlantic Puffins by Canestrini & Kramer (1899); Belopolskaya (1952) reported specimens of *Alloptes fraterculae* from Atlantic Puffins. Choe & Kim (1991) found *Alloptes conurus* on Thick-billed Murres and Common Murres; but the species designation was only tentative, and the specimens could represent an undescribed species. Dobson (1952) reported infestations of poultry mites (*Dermanyssus gallinae*) on Atlantic Puffin chicks.

Phthiraptera (lice)

The Phthiraptera, or lice, form an important component of the ectoparasitic fauna of the auks (Eveleigh 1974, Eveleigh & Threlfall 1976, Ballard & Ring 1979, Choe & Kim 1987, Muzaffar 2000; Table 5). Lice spend their entire lives on their host (Rothschild & Clay 1957) and have received considerable attention in phylogenetic studies because of their high levels of host specificity (Smith 2001, Marshall 2002, Price *et al.* 2003). Although lice may occur in very large numbers on auks (Choe & Kim 1987), no evidence exists to imply that the lice produce a negative impact. Generally, lice have been demonstrated to cause damage to plumage and to incur energetic expenses on avian hosts (Clayton 1991, Booth *et al.* 1993).

In seabirds, such as the Black Noddy *Anous minutus*, sunning reduces *Quadraceps* louse loads, and birds with high loads are seen to “sun” even when temperatures are high (Moyer & Wagenbach 1995). Sunning therefore represents an energetically expensive means of reducing louse loads. *Quadraceps* is also common on alcids and may potentially produce similar consequences to energetic expenses.

Louse populations on alcids vary seasonally, with the highest loads occurring during the chick-rearing period. It has been suggested that chicks may have an important role in the reproduction of lice (Eveleigh & Threlfall 1976, Muzaffar 2000). Moreover, distribution patterns of the lice on alcids reflect preferences for certain microhabitats (Choe & Kim 1987) that are preened less frequently (see Clayton 1991), suggesting adaptive significance of lice to host responses.

In addition to being a source of irritation, louse genera such as *Austramenopon* act as a vector of the filarial worm *Eulimdana* in charadriiform hosts, although long-term effects of that nematode on the host life cycle have not been studied (Bartlett 1993). *Austramenopon* is also hematophagous (Eveleigh & Threlfall 1976), suggesting that high loads may reduce host condition. *Quadraceps* and *Saemundssonina* may both be potentially damaging to the feathers, with probable energetic expenses on the hosts (Clayton 1991, Moyer & Wagenbach 1995, Price *et al.* 2003).

Avian preening directly influences louse populations and distribution patterns on the hosts (Clayton 1991). Muzaffar (2000) suggested that varying densities of *Quadraceps* and *Austramenopon* on different alcid hosts may be a result of differential preening attributable to variable bill shapes, a contention supported by studies on a range of other avian hosts (Clayton & Cotgreave 1994). Although direct evidence concerning the negative attributes of the lice of alcids is lacking, circumstantial evidence indicates a relationship that is expensive to the host.

Diptera (flies)

Muzaffar (2000) found larvae of a fly species (Diptera: Muscomorpha) partially imbedded in the skin of Atlantic Puffin chicks (Table 5), suggesting that the species might feed on tissue fluid or blood, exposing the wound to secondary infections. Certain Diptera larvae are known to be hematophagous parasites of birds (Sabrosky *et al.* 1989).

Numerous genera of Calliphoridae have been recorded from a wide range of birds, but none have been recorded from seabirds (Wheeler & Threlfall 1989). The hippoboscid species *Icosta americana* has been recorded from Thick-billed Murres, although the impacts on the host are not known (Wheeler & Threlfall 1989). Dipterans are potentially significant parasites because they are capable of causing injury or infection; however, few studies have quantified their occurrence.

Siphonaptera (fleas)

In the North Atlantic region, six flea species have been recorded from Atlantic Puffins (Rothschild & Clay 1957; Table 5). Some fleas likely occur on auks accidentally in situations where their preferred hosts coexist with auks: for example, *Ornithopsylla laetitiae* (preferred host: Manx Shearwater *Puffinus puffinus*) and *Spirospyllus cuniculi* [preferred host: European rabbit *Oryctolagus cuniculus* (Rothschild & Clay 1957)]. In the North Pacific region, *Actenopsylla suavis* has been recorded from two burrow-nesting alcids: the Cassin's Auklet *Ptychoramphus aleuticus* and the Tufted Puffin (Holland 1984; Table 5). Muzaffar (2000) found *Orchopeas leucopus* specimens from Thick-billed Murres and Atlantic Puffins at the Gannet Islands in Newfoundland and Labrador, although the occurrence was likely accidental because this flea is parasitic to the Deer Mouse *Peromyscus maniculatus*, which is abundant in the archipelago. The low frequency of fleas recorded from alcids or their nesting sites indicates that fleas are less important as ectoparasites.

DISCUSSION

Our compilation of the records of parasites and diseases of alcids emphasizes the need for further parasitologic research on this group of seabirds. A diverse array of parasites and diseases has been documented. Many species of parasites (for example, digeneans and cestodes) have complex life cycles involving organisms that form fundamental components of the zooplankton and higher trophic levels in the marine food web. Nevertheless, limited information is available on the intermediate hosts and their interactions with the parasites, or on the detrimental effects of many of the parasites on seabirds in general and auks in particular.

The goal of parasitologic studies should be to integrate available information on various aspects of the biology and ecology of the host species and to derive a better understanding of their complex interactions with other organisms in the marine ecosystem. Belopolskaya (1952), Hoberg (1996), Hoberg (1997), Brooks & Hoberg (2000), and Hoberg & Adams (2000) have addressed this holistic viewpoint in the understanding of parasite communities in complex marine ecosystems. Parasites represent biologically diverse assemblages that serve as determinants of historical and contemporary biogeography, providing information on complex interactions at different trophic levels (because of their different intermediate hosts) in a predictable manner.

Feeding ecology and parasite transmission

The feeding ecology of seabirds may shape their endoparasitic fauna (Galaktionov 1995, Hoberg 1996). Belopolskaya (1952) found a rich digenean and cestode fauna in gulls and alcids from the Seven Islands archipelago in the Barents Sea in the early 1940s. A study on the same hosts from the same colonies in the early 1990s showed drastic changes in both gull and alcid endoparasites (Galaktionov 1995). A striking reduction in the digeneans and major fluctuations in the cestodes were seen in studied gulls. More drastic changes were noted in the endoparasite fauna of alcids, with a total disappearance of digeneans. The changes were attributed to a combination of

- population changes in the seabird hosts.
- variations in the important prey (fish) species (many of which served as intermediate hosts).
- reductions in invertebrate intermediate hosts.

Gulls and auks sometimes experience drastic switches in their principal prey (e.g. Birkhead & Nettleship 1987, Montevecchi & Myers 1995, Bryant *et al.* 1999, Massaro *et al.* 2000, Rowe *et al.* 2000). The impact of those switches is normally considered in terms of energetic consequences and resulting changes in productivity or overwinter survival. However, the role of changing parasite infections resulting from prey switches is rarely considered (e.g. Galaktionov 1995). Changes in diet that have occurred over the past few decades in various parts of the Northern Hemisphere could expose seabirds to a whole new suite of endoparasites with potential negative impacts.

Host population dynamics

Parasites may potentially affect seabird population dynamics by selectively reducing fitness and reproductive success (Schreiber & Burger 2002). Ticks, for instance, generally tend to have negative influences on seabird hosts (Duffy 1991, McCoy *et al.* 2002). High densities of *Ornithodoros capensis* tick have been correlated with nest desertions in several seabird species (Feare 1976; King *et al.* 1977a, 1977b; Duffy 1983). There is also evidence indicating that breeding dispersal in Black-legged Kittiwakes may be a consequence of high *I. uriae* infestations (e.g. Boulinier & Danchin 1996, Boulinier *et al.* 2001).

Tick infections can reduce nestling condition, growth rates and survival in many seabird species, with occasional incidences of mortality (Morbey 1996, Ramos *et al.* 2001, Feare & Gill 1997, Bergström *et al.* 1999, Gauthier-Clerc *et al.* 1998, Mangin *et al.* 2003). Blood hematocrit levels (an index of oxygen-carrying capacity) have been correlated with high infestations of *I. uriae* in Black-legged Kittiwakes, suggesting that ticks may represent an energetic expense to their hosts (Wanless *et al.* 1997). However, not all seabird species exhibit such a response to high tick infestations [for example, Common Murres (Wanless *et al.* 1997), King Penguins *Aptenodytes patagonicus* (Gauthier-Clerc *et al.* 2003)], making it difficult to isolate the exact consequence of tick infestations on their hosts (Gauthier-Clerc *et al.* 2003).

In endoparasitic infestations, negative consequences may be clearly evident (for example, intestinal lesions), but impacts at the population level are difficult to establish. The co-occurrence of *Tetrabothrius* spp. (cestodes) and *Cosmocephalus obvelatus* (nematode) are seemingly more important in this negative relationship between parasite loads and body condition of Yellow-legged Gulls *Larus cachinnans* (Bosch *et al.* 2000).

It is not known if these infections cause any major effects in the host populations. Cestode and nematode infections in auks may lead to moderate-to-severe intestinal lesions without apparent ill effects on the hosts (e.g. Threlfall 1971, Hoberg 1984c, Nagasawa *et al.* 1998b). In terrestrial avian host species, densities of one species of nematode, *Trichostrongylus tenuis*, can modulate population cycles in the Red Grouse *Lagopus lagopus* (Hudson & Dobson 1997, Hudson *et al.* 1998).

Tapeworm or nematode infections are rarely associated with mass mortality events in seabirds (Petermann *et al.* 1989, Hüppop 1996, Piatt & Van Pelt 1997, Gill 2004 pers. comm.) and whether they induce changes in population levels of their hosts in the long term is not known. The possibility that certain nematode or cestode infections above a threshold level may be detrimental to the seabird hosts cannot be ruled out. Population level changes are extremely difficult to document, particularly when assessing endoparasitic infestations in seabirds (Bosch *et al.* 2000, and see Schreiber & Burger 2002).

Reservoirs

Seabirds serve as reservoirs for viruses and bacteria in the wild (Lvov *et al.* 1979, Clifford 1979, Nuttall 1984, Gauthier-Clerc *et al.* 1999, Gylfe *et al.* 1999, Kerry *et al.* 2000). How these viruses circulate within or between colonies, or the role that they have as disease-causing agents remains largely unclear. Although some viruses may cause disease and mortality when seabirds are experimentally infected (e.g. Lvov *et al.* 1979), little is known about their ecology and epidemiology (Gauthier-Clerc *et al.* 2003).

The prevalence of the bacteria *B. burgdorferi* (sensu lato) in seabird colonies shows the importance of seabirds as reservoirs of human pathogens (Olsen *et al.* 1993, 1995; Gauthier-Clerc *et al.* 1999, 2003). Numerous bacterial species have either been isolated from penguins or are evident from antibodies present in blood (Kerry *et al.* 2000). Clinical signs of disease are rarely seen, and, in many cases, mortality events cannot be attributed to any particular pathogen (Clarke & Kerry 1993, Austin & Webster 1993, Kerry *et al.* 2000). Stress factors, such as starvation and low hematocrit levels, may be further aggravated by bacterial or viral infections resulting in symptomatic disease (Kerry *et al.* 2000).

Exotic diseases

Introduced viruses and bacteria constitute an important problem in seabird colonies. Such introductions may potentially cause outbreaks of infection in wild birds and serve to establish reservoirs of the infectious organisms in the wild (Gardner *et al.* 1997, Kerry *et al.* 2000). Problems of exotic diseases in seabird colonies of the Northern Hemisphere have not been reported, although the potential exists.

Emerging infectious diseases—such as the West Nile virus (WNV)—pose a serious threat to wild birds in the Western Hemisphere, although the spread of the virus is governed by the presence of the appropriate mosquito vectors, which generally do not occur in major seabird colonies (Campbell *et al.* 2002). Gulls, being associated with garbage dumps (e.g. Robertson *et al.* 2001), may be more prone to WNV infections (for example, Herring Gull *Larus argentatus* tested positive, Rappole & Hubálek 2003) and may form a link between such diseases and nearby seabird colonies.

Increased human activity in seabird colonies may enhance the risk of exotic infectious diseases becoming established in the wild. Outbreaks of poultry virus infections in the Antarctic serve as an admonishing example (Gardner *et al.* 1997). Human activity should therefore be carefully controlled, particularly with respect to recreational activity near seabird colonies. People working at seabird colonies should be particularly cautious about maintaining good personal hygiene in the field.

Taxonomy and phylogeny

Taxonomic studies of certain parasites of auks are particularly interesting for understanding the evolutionary history of host–parasite systems and for documenting coevolution and host-switching events that have culminated in current parasite fauna (Eveleigh & Amano 1977, Hoberg 1986a, Price *et al.* 2003). The phylogeny of lice occurring on auks, for instance, may not reflect the phylogeny of their hosts, making it difficult to interpret the evolutionary history of the host–parasite assemblage (Eveleigh & Amano 1977). Phylogenetic studies of parasites of seabirds in general, using molecular approaches, could lead to a more complete picture of coevolving systems in the marine environment (McCoy *et al.* 1999, Hoberg & Adams 2000, McCoy *et al.* 2003).

CONCLUSIONS

An urgent need exists to incorporate parasitologic components into modern-day ecological studies of seabirds. Seabird monitoring programs should have a more defined parasitology component that attempts to assess parasites and their harmful effects on seabirds. We believe that the importance of seabird parasitology has been underemphasized. Knowledge concerning the susceptibility of various species and populations to parasitic diseases could aid in making management decisions in auk (and other seabird) conservation. A better understanding of the marine web of life could be established through long-term studies of ecology and parasitology of seabirds.

ACKNOWLEDGEMENTS

We thank Johanne Dussureault, Mark Button, Brian Veitch, Shauna Baillie and Sherrylynn Rowe for assistance with fieldwork at the Gannet Islands; Rasul Khan, Jean Finney–Crawley and especially Eric Hoberg for useful comments on an earlier version of this manuscript; and Murray Colbo for valuable support and advice during this project. Funding was provided by Atlantic Cooperative Wildlife Ecology Research Network CWS/NSERC Industrial Chair and NSERC equipment grants to ILJ.

REFERENCES

- ALEKSEEV, V. & SMETANINA, Z. 1968. Nematody ryboyadnykh ptits ostrovov Rimskogo–Korsakova. *Gelmint. Zivotnykh Tykhogo Okeana*. pp. 97–104.
- ALEKSEEV, V. & SMETANINA, Z. 1970. Trematody ryboyadnykh ptits ostrovov Rimskogo–Korsakova. *Parazitologicheskie i zoologicheskie issledovaniia na Dalnem Vostoke*. *Dalnev. Gosudarst. Universitet. Uchenye Zapiski* 16: 96–103.
- ANDERSON, R.C. 2000. *Nematode parasites of vertebrates: their development and transmission*. Oxon: CABI Publishing.

- ANDERSON, R.C. & WONG, P.L. 1982. The transmission and development of *Paracuarria adunca* (Creplin, 1846) (Nematoda: Acuaroidea) of gulls (Laridae). *Canadian Journal of Zoology* 60: 3092–3104.
- ARMSTRONG, I.H., COULSON, J.C., HAWKEY, P. & HUDSON, M.J. 1978. Further mass seabird deaths from paralytic shellfish poisoning. *British Birds* 71: 58–68.
- AUSTIN, F.J. & WEBSTER, R.G. 1993. Evidence of ortho- and paramyxoviruses in fauna from Antarctica. *Journal of Wildlife Diseases* 29: 568–571.
- BAER, J.G. 1956. Parasitic helminths collected in west Greenland. *Meddeleser om Gronland* 124: 1–55.
- BAKKE, T.A. 1972. *Reighardia sterna* (Diesing, 1864) Ward, 1899 (Pentastomida: Cephalobaenida) from the common gull (*Larus canus* L.) in a Norwegian locality. *Norwegian Journal of Zoology* 20: 273–277.
- BAKKE, T.A. 1978. Tungenarken *Reighardia sterna* (Pentastomida) funnet i Norge. *Fauna Oslo* 31: 124–127.
- BALLARD, J.T. & RING, R.A. 1979. The ectoparasites of some marine birds from Bamfield Marine Station, British Columbia, with particular reference to Common Murres, *Uria aalge* (Pont.). *Canadian Journal of Zoology* 57: 1980–1984.
- BARANTON, G., POSTIC, D., SAINT GIRONS, I., BOERLIN, P., PIFFARETTI, J.C., ASSOUS, M. & GRIMONT, P.A.D. 1992. Delineation of *Borrelia burgdorferi* sensu stricto, *Borrelia garinii* sp. nov., and group VS461 associated with Lyme borreliosis. *International Journal of Systematic Bacteriology* 42: 378–383.
- BARTLETT, C.M. 1993. Lice (Amblycera and Ischnocera) as vectors of *Eulimdana* spp. (Nematoda: Filarioidea) in charadriiform birds and the necessity of short reproductive periods in adult worms. *Journal of Parasitology* 79: 85–91.
- BARTLETT, C.M., ANDERSON, R.C. & BUSH, A.O. 1989. Taxonomic descriptions and comments on the life history of new species of *Eulimdana* (Nematoda: Filarioidea) with skin-inhabiting microfilariae in the Charadriiformes (Aves). *Canadian Journal of Zoology* 67: 612–629.
- BARTON, T.R., HARRIS, M.P., WANLESS, S. & ELSTON, D.A. 1996. The activity period and life-cycle of the tick *Ixodes uriae* (Acari: Ixodidae) in relation to host breeding strategies. *Parasitology* 112: 571–580.
- BARUS, V., SERGEEVA, T.P., SONIN, M.D. & RZHNIKOV, K.M. 1978. Helminths of fish-eating birds of the Palaearctic Region I. Nematoda. The Hague: Academia Publishing House of the Czechoslovak Academy of Sciences.
- BAYLIS, H.A. 1919. A collection of Entozoa chiefly from birds from the Murman coast. *Annals and Magazine of Natural History* 9: 501–515.
- BELOGUROV, O.I., LEONOV, V.A. & ZUEVA, L.S. 1968. Gelmintofauna ryboiadnykh ptits (chaek i chistikov) poberezhia okhotskogo moria. *Gelminty Zhivoknykh Tikhogo Okeana*. Moskva: Akademii Nauk SSSR. pp. 105–124.
- BELOPOLSKAYA, M.M. 1952. Parazitofauna morskikh vodoplavaiushchikh ptits. *Leningradskogo Universiteta. Uchenye Zapiski* 141: 127–180.
- BENNETT, G.F., WHITEWAY, M. & WOODWORTH-LYNAS, C. 1982. A host–parasite catalogue of the avian haematozoa. St John's: Department of Biology, Memorial University of Newfoundland.
- BERGSTRÖM, S., HAEMIG, P.D. & OLSEN, B. 1999. Increased mortality of black-browed albatross chicks at a colony heavily-infested with the tick *Ixodes uriae*. *International Journal of Parasitology* 29: 1359–1361.
- BIRKHEAD, T.R. & NETTLESHIP, D.N. 1987. Ecological relationships between Common Murres, *Uria aalge*, and Thick-billed Murres, *Uria lomvia*, at the Gannet Islands, Labrador. III. Feeding ecology of the young. *Canadian Journal of Zoology* 65: 1638–1649.
- BOOTH, D.T., CLAYTON, D.H. & BLOCK, B.A. 1993. Experimental demonstration of the energetic cost of parasitism in free-ranging hosts. *Proceedings of the Royal Society of London. Series B: Biological Sciences* 253: 125–129.
- BOSCH, M., JORDI, T. & FIGUEROLA, J. 2000. A helminth community in breeding Yellow-legged Gulls (*Larus chachinnans*): pattern of association and its effect on host fitness. *Canadian Journal of Zoology* 78: 777–786.
- BOULINIER, T. & DANCHIN, E. 1996. Population trends in Kittiwakes *Rissa tridactyla* colonies in relation to tick infestation. *Ibis* 138: 326–334.
- BOULINIER, T., MCCOY, K.D. & SORCI, G. 2001. Dispersal and parasitism. In: Clobert, J., Danchin, E., Dhondt, A. & Nichols, J. (Eds). *Dispersal*. Oxford: Oxford University Press. pp. 169–179.
- BROOKS, D.R. & HOBERG, E.P. 2000. Triage for the biosphere: the need and rationale for taxonomic inventories and phylogenetic studies of parasites. *Comparative Parasitology* 67: 1–25.
- BROWN, C.R. & BROWN, M.B. 1986. Ectoparasitism as a cost of coloniality in Cliff Swallows (*Hirundo pyrrhonata*). *Ecology* 67: 1206–1218.
- BROWN, C.R. & BROWN, M.B. 1996. Coloniality in the Cliff Swallow: the effect of group size on social behavior. Chicago: University of Chicago Press. p. 566.
- BRYANT, R., JONES, I.L. & HIPFNER, J.M. 1999. Responses of Common and Thick-billed Murres to changes in prey availability at the Gannet Islands, Labrador. *Canadian Journal of Zoology* 77: 1278–1287.
- BUNIKIS, J., OLSEN, B., FINGERLE, V., BONNEDAHL, J., WILSKE, B. & BERGSTRÖM, S. 1996. Molecular polymorphism of the Lyme disease agent *Borrelia garinii* in northern Europe is influenced by a novel enzootic *Borrelia* focus in the North Atlantic. *Journal of Clinical Microbiology* 34: 364–368.
- BUSTNES, J.O. & GALAKTIONOV, K.V. 1999. Anthropogenic influences of the infestation of intertidal gastropods by seabird trematode larvae on the southern Barents Sea coast. *Marine Biology* 133: 449–453.
- BUSTNES, J.O., GALAKTIONOV, K.V. & IRWIN, S.W.B. 2000. Potential threats to littoral biodiversity: is increased parasitism a consequence of human activity? *Oikos* 90: 189–190.
- BYKHOVSKAIA-PAVLOVSKAIA, I.E. 1962. Trematody ptits fauny SSSR: ekologo-geograficheskii obzor. Moskva: Izd-vo Akademii Nauk SSSR.
- CAMPBELL, G.L., MARFIN, A.A., LACIOTTI, R.S. & GUBLER, D.J. 2002. West Nile virus. *Lancet Infectious Diseases* 2: 519–529.
- CANESTRINI, G. & KRAMER, P. 1899. Demodicidae und Sarcopitidae. *Tierreich* 7: 1–193.
- CHOE, J.C. & KIM, K.C. 1987. Community structure of arthropod ectoparasites on Alaskan seabirds. *Canadian Journal of Zoology* 65: 2998–3005.
- CHOE, J.C. & KIM, K.C. 1991. Microhabitat selection and adaptation of feather mites (Acari: Analgoidea) on murres and kittiwakes. *Canadian Journal of Zoology* 69: 817–821.

- CLARKE, J.R. & KERRY, K.R. 1993. Diseases and parasites of penguins. *Korean Journal of Polar Research* 4: 79–96.
- CLAYTON, D.H. 1991. Coevolution of avian grooming and ectoparasite avoidance. In: Loye, J.E. & Zuk, M. (Eds). *Bird–parasite interactions: ecology, evolution and behaviour*. Oxford: Oxford University Press. pp. 258–289.
- CLAYTON, D.H. & COTGREAVE, P. 1994. Relationship of bill morphology to grooming behaviour in birds. *Animal Behaviour* 47: 195–201.
- CLAYTON, D.H. & MOORE, J. 1997. *Host–parasite evolution: general principals and avian models*. Oxford: Oxford University Press.
- CLIFFORD, C.F. 1979. Tick-borne viruses of seabirds. In: Kurstak, E. (Ed). *Arctic and tropical arboviruses*. London: Academic Press. pp. 83–100.
- COULSON, J.C., POTTS, G.R., DEANS, I.R. & FRASER, S.M. 1968. Mortality of shags and other sea birds caused by paralytic shellfish poison. *Nature* 220: 23–24.
- CRAM, E.B. 1927. *Bird parasites of the nematode suborders Strongylata, Ascaridata, and Spirurata*. Washington, DC: Government Printing Office. [U.S. National Museum Bulletin 140]
- DANCHIN, E. 1992. The incidence of the tick parasite *Ixodes uriae* in Kittiwake *Rissa tridactyla* colonies in relation to the age of the colony, and a mechanism of infecting new colonies. *Ibis* 134: 134–141.
- DAVIES, M.S. & KNOWLES, A.J. 2001. Effect of trematode parasitism on the behaviour and ecology of a common marine snail (*Littorina littorea* (L.)). *Journal of Experimental Marine Biology and Ecology* 260: 155–167.
- DAVIS, J.W., ANDERSON, R.C., KARSTAD, L. & TRAINER, D.O. (Eds). 1971. *Infectious and parasitic diseases of wild birds*. Ames, IA: Iowa State University Press.
- DAWES, B. 1946. *The trematoda: with special reference to British and other European forms*. Cambridge: Cambridge University Press.
- DEBACKER, V., HOLBEEK, L., TAPIA, G., GOBERT, S., JOIRIS, C.R., JAUNIAUX, T., COIGNOUL, F. & BOUQUEGNEAU, J.M. 1997. Ecotoxicological and pathological studies of Common Guillemots *Uria aalge* beached on the Belgian coast during six successive wintering periods (1989–90 to 1994–95). *Diseases of Aquatic Organisms* 29: 159–168.
- DITLEVSEN, H. 1917. XI. Cestoder. *Meddeleser om Gronland* 23: 1119–1140.
- DOBSON, R. 1952. *The birds of the Channel Islands*. London: Staples.
- DOGIEL, V.A. 1964. *General parasitology*. London: Oliver and Boyd.
- DUBININ, V.B. 1951. Feather mites (Analgesoidea). Part I. Introduction to their study. *Fauna USSR* 6: 1–363.
- DUBOIS, G. 1953. Systématique des Strigeidae: complément de la monographie. *Mémoires de la Société neuchâteloise des sciences naturelles* 8: 2.
- DUBOIS, G. 1978. A propos des matériels originaux de *Cotylurus platycephalus* et *Cotylurus variegatus* (Creplin 1825) (Trematoda: Strigeidae). *Annales de Parasitologie Humaine et Comparée* 53: 53–62.
- DUBOIS, G., 1981. Notes helminthologiques. V: Strigeidae Railliet, Diplostomatidae Poirier et Proterodiplostomatidae Dubois. *Revue Suisse de Zoologie* 88: 227–232.
- DUFFY, D.C. 1983. The ecology of tick parasitism on densely nesting Peruvian seabirds. *Ecology* 64: 110–119.
- DUFFY, D.C. 1991. Ants, ticks and nestling seabirds: dynamic interactions. In: Loye, J.E. & Zuk, M. (Eds). *Bird–parasite interactions: ecology, evolution and behavior*. Oxford: Oxford University Press. pp. 243–257.
- DYCK, J. 1975. *Reighardia lomviae* sp. nov., a new pentastomid from guillemot. *Norwegian Journal of Zoology* 23: 97–109.
- EMERSON, K.C. 1972. Checklist of the Mallophaga of North America (North of Mexico). Part IV: Bird host list. Dugway, UT: Deseret Test Center.
- ENSTIPP, M. & GRÉMILLET, D. 2002. Experimental studies: foraging performance under controlled conditions. Interactions between the marine environment, predators, and prey: implications for sustainable Sand Eel fisheries (IMPRESS project). Strasbourg: Centre National de la Recherche Scientifique.
- EVELEIGH, E.S. 1974. A study of the ectoparasites of alcids of Newfoundland. MSc thesis, Department of Biology, Memorial University of Newfoundland, St John's, Newfoundland. 94 pp.
- EVELEIGH, E.S. & AMANO, H. 1977. A numerical taxonomic study of the Mallophagan genera *Cummingsiella* (= *Quadriceps*), *Saemundssonina* (Ischnocera: Phlopteridae), and *Austromenopon* (Amblycera: Menoponidae) from alcids (Aves: Charadriiformes) of the northwest Atlantic with reference to host–parasite relationships. *Canadian Journal of Zoology* 55: 1788–1801.
- EVELEIGH, E.S. & THRELFALL, W. 1974. A new species, and notes on a previously described species of *Austromenopon* Bedford, 1939 (Mallophaga: Amblycera) from alcids (Aves: Charadriiformes). *Proceedings of the Entomological Society of Washington* 76: 271–277.
- EVELEIGH, E.S. & THRELFALL, W. 1975. Bionomics of *Ixodes (Ceratiixodes) uriae* White 1852 on auks (Alcidae) from Newfoundland. *Canadian Journal of Zoology* 53: 82–86.
- EVELEIGH, E.S. & THRELFALL, W. 1976. Population dynamics of lice (Mallophaga) on auks (Alcidae) from Newfoundland. *Canadian Journal of Zoology* 54: 1694–1711.
- FEARE, C.J. 1976. Desertion and abnormal development in a colony of Sooty Terns *Sterna fuscata* infested by virus-infected ticks. *Ibis* 118: 112–115.
- FEARE, C.J. & GILL, E.L. 1997. The life cycle of the tick *Amblyomma loculosum* in a Sooty Tern *Sterna fuscata* colony in the Seychelles. *Journal of Zoology (London)* 241: 643–648.
- FITZPATRICK, C. & THRELFALL, W. 1977. The ectoparasites of three species of seabirds from Newfoundland, Canada. *Canadian Journal of Zoology* 55: 1205–1209.
- FOWLER, M.E. 1993. *Zoo and wild animal medicine—current therapy*. London: W.B. Saunders.
- FRENOT, Y., DE OLIVIERA, E., GAUTHIER–CLERC, M., DEUNFF, J., BELLIDO, A. & VERNON, P. 2001. Life cycles of the tick *Ixodes uriae* (White, 1852) in penguin colonies: relationship with host breeding activity. *International Journal for Parasitology* 31: 1040–1047.
- FRIEND, M. & FRANSON, C.J. 1999. *Field manual of wildlife diseases: general field procedures and diseases of birds*. Madison, WI: US Department of the Interior, US Geological Survey.

- GALAKTIONOV, K.V. 1995. Long-term changes in the helminth fauna of colonial seabirds in the Seven Islands archipelago (Barents Sea, Eastern Murman). In: Skjoldal, H.R., Hopkins, C., Erikstad, K.E. & Leinaas, K.E. (Eds.) Ecology of fjords and coastal waters. Oxford: Elsevier. pp. 489–495.
- GARDNER, H., BROUWER, S., GLEESON, L., KERRY, K. & RIDDLE, M. 1997. Poultry virus infection in Antarctic penguins. *Nature* 387: 245.
- GASPARINI, J., MCCOY, K.D., HAUSSY, C., TVERAA, T. & BOULINIER, T. 2001. Induced maternal response to the Lyme disease spirochaete *Borrelia burgdorferi* sensu lato in a colonial seabird, the kittiwake *Rissa tridactyla*. *Proceedings of the Royal Society of London B* 268: 647–650.
- GASTON, A.J. & JONES, I.L. 1998. The auks. Oxford: Oxford University Press.
- GAUD, J. & ATYEO, T. 1996. Feather mites of the world (Acarina, Astigmata): the supraspecific taxa. Parts I and II. *Musée Royal de l'Afrique Centrale, Annales, Sciences Zoologiques* 277: 1–193, 1–436.
- GAUTHIER-CLERC, M., CLERQUIN, Y. & HANDRICH, Y. 1998. Hyperinfestation by ticks *Ixodes uriae*: a possible cause of death in adult King Penguins, a long-lived seabird. *Colonial Waterbirds* 21: 229–233.
- GAUTHIER-CLERC, M., JAULHAC, B., FRENOT, Y., BACHELARD, C., MONTEIL, H., LE MAHO, Y. & HANDRICH, Y. 1999. Prevalence of *Borrelia burgdorferi* (the Lyme disease agent) antibodies in King Penguin *Aptenodytes patagonicus* in Crozet Archipelago. *Polar Biology* 22: 141–143.
- GAUTHIER-CLERC, M., MANGIN, S., LE BOHEC, C., GENDNER, J. & LE MAHO, Y. 2003. Comparison of behaviour, body mass, haematocrit level, site fidelity and survival between infested and non-infested King Penguin *Aptenodytes patagonicus* by ticks *Ixodes uriae*. *Polar Biology* 26: 379–382.
- GIBSON, G.G. 1968. Species composition of the genus *Streptocara* Ralliet *et al.*, 1912 and the occurrence of these avian nematodes (Acuariidae) on the Canadian Pacific coast. *Canadian Journal of Zoology* 46: 629–645.
- GUIGUEN, C., MONNAT, J.Y., LAUNAY, H. & BEAUCOURNU, J.C. 1983. Ectoparasites des oiseaux en Bretagne. 2. Siphonapteres. *Revue Française D'Entomologie (Nouvelle Serie)* 5: 144–146.
- GYLFE, A., OLSEN, B., STRASEVICIUS, B., MARTI RAS, N., WEIHE, P., NOPPA, L., ÖSTBERG, Y., BARANTON, G. & BERGSTRÖM, S. 1999. Isolation of Lyme disease *Borrelia* from puffins (*Fratercula arctica*) and seabird ticks (*Ixodes uriae*) on the Faeroe Islands. *Journal of Clinical Microbiology* 37: 890–896.
- HARRIS, M.P. 1984. The puffin. Calton: T&AD Poyser.
- HOBERG, E.P. 1979. Helminth parasites of marine birds (Charadriiformes: Alcidae and Laridae) occurring in the North Pacific Ocean and Gulf of Alaska. MSc thesis, University of Saskatchewan, Saskatoon, Saskatchewan. 121 pp.
- HOBERG, E.P. 1981. *Pseudogymnophallus alcae* gen. n. et sp. n. (Trematoda: Gymnophallidae) from alcids (Charadriiformes) in subarctic seas. *Proceedings of Helminthological Society of Washington* 48: 190–194.
- HOBERG, E.P. 1982. *Diorchis pelagicus* sp. nov. (Cestoda: Hymenolepididae) from the Whiskered Auklet, *Aethia pygmaea*, and the Crested Auklet, *Aethia cristatella*, in the Western Aleutian Islands, Alaska. *Canadian Journal of Zoology* 60: 2198–2201.
- HOBERG, E.P. 1984a. *Alcataenia longicervica* sp. n. from murrelets, *Uria lomvia* (Linnaeus) and *Uria aalge* (Pontoppidan) in the North Pacific basin, with redescription of *Alcataenia armillaris* (Rudolphi, 1810) and *Alcataenia meinertzhageni* (Baer, 1956) (Cestoda: Dilepididae). *Canadian Journal of Zoology* 62: 2044–2052.
- HOBERG, E.P. 1984b. *Alcataenia fraterculae* sp. n. from the Horned Puffin *Fratercula corniculata* (Naumann), *Alcataenia cerorhincae* sp. n. from the Rhinoceros Auklet, *Cerorhinca monocerata* and *Alcataenia larina pacifica* spp. n. (Cestoda: Dilepididae) in the North Pacific basin. *Annales de Parasitologie Humaine et Comparée* 59: 335–351.
- HOBERG, E.P. 1984c. *Alcataenia pygmaeus* sp. n. (Cestoda: Dilepididae) from the Whiskered Auklet, *Aethia pygmaea*, in the Western Aleutian Islands, Alaska, with a comment on the genera *Alcataenia* and *Rissotaenia*. *Proceedings of the Helminthological Society of Washington* 31: 315–319.
- HOBERG, E.P. 1984d. *Alcataenia campylacantha* (Krabbe, 1869) from Pigeon Guillemots, *Cephus columba*, Pallas and Black Guillemots, *Cephus grylle* (Linnaeus) and *Alcataenia* sp. indet. (Cestoda: Dilepididae) from Kittlitz's Murrelets, *Brachyramphus brevirostris* (Vigors), in Alaska. *Canadian Journal of Zoology* 62: 2297–2301.
- HOBERG, E.P. 1984e. Systematics, zoogeography and ecology platyhelminth parasites of the seabird family Alcidae (Charadriiformes: suborder Alcae). PhD thesis, University of Washington, Seattle, Washington. 324 pp.
- HOBERG, E.P. 1986a. Evolution and historical biogeography of a parasite–host assemblage, *Alcataenia* spp. (Cyclophyllidea: Dilepididae) in Alcidae (Charadriiformes). *Canadian Journal of Zoology* 64: 2576–2589.
- HOBERG, E.P. 1986b. Aspects of the ecology and biogeography of Acanthocephala in Antarctic seabirds. *Annales de Parasitologie Humaine et Comparée* 61: 199–214.
- HOBERG, E.P. 1987. *Reighardia sternaes* (Diesing, 1864) (Pentastomida) from seabirds in Antarctica. *Canadian Journal of Zoology* 65: 1289–1291.
- HOBERG, E.P. 1991. *Alcataenia atlantiensis* n. sp. (Dilepididae) from the razorbill (*Alcatorda* Linnaeus) in the eastern North Atlantic basin. *Systematic Parasitology* 20: 83–88.
- HOBERG, E.P. 1996. Faunal diversity among avian parasite assemblages: the interaction of history, ecology and biogeography in marine systems. *Bulletin of the Scandinavian Society of Parasitology* 6: 65–89.
- HOBERG, E.P. 1997. Phylogeny and historical reconstruction: host–parasite systems as keystones in biogeography and ecology. In: Reaka-Kulda, M.L., Wilson, D.E. & Wilson, E.O. (Eds). Biodiversity II: understanding and protecting our biological resources. Washington, DC: Joseph Henry Press. pp 243–261.
- HOBERG, E.P. & ADAMS, A. 2000. Phylogeny, history and biodiversity: understanding faunal structure and biogeography in the marine realm. *Bulletin of the Scandinavian Society of Parasitology* 10: 19–37.
- HOBERG, E.P. & RYAN, P.G. 1989. Ecology of helminth parasitism in *Puffinus gravis* (Procellariiformes) on the breeding grounds at Gough Island. *Canadian Journal of Zoology* 67: 220–225.
- HOBERG, E.P. & WEHLE, D.H.S. 1982. Host and geographic records of ectoparasites from the Alaskan seabirds (Charadriiformes: Alcidae and Laridae). *Canadian Journal of Zoology* 60: 472–475.

- HOLLAND, G.P. 1984. The fleas of Canada, Alaska and Greenland. *Memoirs of the Entomological Society of Canada* 130: 1–631.
- HOOGSTRAAL, H., CLIFFORD, C.M., KEIRANS, J.E., KAISER, M.N. & EVANS, D.E. 1976. The *Ornithodoros (Alectorobius) capensis* group (Acarina: Ixodoidea: Argasidae) of the Palearctic and Oriental Regions. *O. (A.) maritimus*: identity, marine bird hosts, virus infections, and distribution in Western Europe and Northwestern Africa. *Journal of Parasitology* 62: 799–810.
- HUBÁLEK, Z. 1994. Pathogenic microorganisms associated with free-living birds (a review). *Acta Scientiarum Naturalium Brno* 28: 1–74.
- HUDSON, P.J. & DOBSON, A.P. 1997. Host–parasite processes and demographic consequences. In: Clayton, D.L. & Moore, J. (Eds). *Host–parasite evolution: general principles and avian models*. Oxford: Oxford University Press. pp. 128–154.
- HUDSON, P.J., DOBSON, A.P. & NEWBORN, D. 1998. Prevention of population cycles by parasite removal. *Science* 282: 2256–2258.
- HÜPPOP, O. 1996. Causes and trends of the mortality of Guillemots (*Uria aalge*) ringed on the island of Helgoland, German Bight. *Die Vogelwarte* 38: 217–224.
- JAUNIAUX, T. & COIGNOUL, F. 1994. Aspergillose chez les oiseaux marins échoués a la cote belge. *Annales de Médecine Vétérinaire* 4: 59–63.
- JAENSON, T.G.T., TÄLLEKLINT, L., LUNDQUIST, L., OLSEN, B., CHIRICO, J. & MEJLON, H. 1994. Geographical distribution, host associations, and vector roles of ticks (Acari: Ixodidae, Argasidae) in Sweden. *Journal of Medical Entomology* 31: 240–256.
- JOYEUX, C.H. & BAER, J.G. 1936. Faune de France. Vol. 30. Cestodes. Paris: Paul Lechevalier et Fils.
- KELLOGG, V.L. 1896. New Mallophaga I, with special reference to a collection made from maritime birds of the Bay of Monterey, California. Ser. 2. *Proceedings of the California Academy of Sciences* 6: 31–168.
- KERRY, K., RIDDLE, M. & CLARKE, J. 2000. Diseases of Antarctic wildlife. Technical report of the Scientific Committee on Antarctic Research (SCAR) and the Council of Managers of National Antarctic Programs (COMNP). Kingston: Australian Antarctic Division.
- KING, K.A., BLANKINSHIP, D.R. & PAUL, R.T. 1977a. Ticks as a factor in the 1975 nesting failure of Texas Brown Pelicans. *Wilson Bulletin* 89: 157–158.
- KING, K.A., KEITH, J.O. & MITCHELL, G.A. 1977b. Ticks as a factor in nest desertion of California Brown Pelicans. *Condor* 79: 507–509.
- KRANTZ, G.W. 1978. A manual of acarology. 2nd ed. Corvallis: Oregon State University Book Stores.
- LANE, R.S., PIESMAN J. & BURGENDORFER W. 1991. Lyme borreliosis: relation of its causative agent to its vectors and hosts in North America and Europe. *Annual Review of Entomology* 36: 587–609.
- LAUCKNER, G. 1987. Ecological effects of larval trematode infestation on littoral marine invertebrate populations. *International Journal for Parasitology* 17: 391–398.
- LEIGHTON, F.A. & GAJADHAR, A.A. 1986. *Eimeria fraterculae* sp. n. in the kidneys of Atlantic Puffins (*Fratercula arctica*) from Newfoundland, Canada: species description and lesions. *Journal of Wildlife Diseases* 22: 520–526.
- LEONOV, V.A. & SHVETSOVA, L.S. 1970. Nematody ptits ostrova Vrangelia. Parazitologicheskie i Zoologicheskie issledovaniia na Dalnem Vostoke. *Uchenye Zapiski* 16: 96–103.
- LOYE, J.E. & ZUK, M. 1991. Bird–parasite interactions: ecology, evolution and behavior. Oxford: Oxford University Press.
- LVOV, D.K. 1980. Arboviruses in the USSR. In: Vesebjak–Hirajan, J., Porterfield, J.S. & Arslanagic, E. (Eds). *Arboviruses in the Mediterranean countries*. Stuttgart: Gustav Fischer Verlag. pp. 35–48.
- LVOV, D.K. & ILJICEV, V.D. 1979. Avian migrations and transport of infectious agents [Russian]. Moscow: Nauka Press.
- LVOV, D.K., TIMOPHEEVA, A.A., SMIRNOV, V.A., GROMASHEVSKY, V.L., SIDOROVA, G.A., NIKIFOROV, L.P., SAZONOV, A.A., ANDREEV, A.P., SKVORTZOVA, T.M., BERESINA, L.K. & ARISTOVA, V.A. 1975. Ecology of tick-borne viruses in colonies of birds in the USSR. *Medical Biology* 53: 325–30.
- LVOV, D.K., GROMASHEVSKI, V.L., SKVORTSOVA, T.M., BEREZINA, L.K., GOFMAN, Y.P., ZHDANOV, V.M., NOVOKHATSKI, A.S., KLIMENKO, S.M., SAZANOV, A.A., KHUTARETSKAYA, N.V., ARISTOVA, V.A., KONDRASHINA, N.G., FOMINA, K.B. & SARKISYAN, B.G. 1979. Arboviruses of high latitudes in the USSR. In: Kurstak, E. (Ed). *Arctic and tropical arboviruses*. London: Academic Press. pp. 21–38.
- MAIN, A.J. 1978. Mykines and Tindhölmur: two new Kemerovo group orbiviruses from *Ixodes uriae* in the Faeroe Islands. *Journal of Medical Entomology* 15: 11–14.
- MAIN, A.J., DOWNS, W.G., SHOPE, R.E. & WALLIS, R.C. 1973. Great Island and Bauline: two new Kemerovo group orbiviruses from *Ixodes uriae* in eastern Canada. *Journal of Medical Entomology* 10: 229–235.
- MAIN, A.J., DOWNS, W.G., SHOPE, R.E. & WALLIS, R.C. 1976. Avian arboviruses of the Witless Bay Seabird Sanctuary, Newfoundland, Canada. *Journal of Wildlife Diseases* 12: 182–194.
- MANGIN, S., GAUTHIER–CLERC, M., FRENOT, Y., GENDNER, J. & LE MAHO, Y. 2003. Ticks *Ixodes uriae* and the breeding performance of a colonial seabird, King Penguin *Aptenodytes patagonicus*. *Journal of Avian Biology* 30: 30–34.
- MARKOV, G.S. 1937. Changes with age in the parasite fauna of the Novaya Zemlya Murre (*Uria lomvia lomvia*) [Russian]. *Transactions of the Leningrad Natural Society* 66: 456–466.
- MARKOV, G.S. 1941. Parasitic worms of Bezymyannaya Bay (Novaya Zemlya), Republic of the USSR [Russian]. *Academy of Sciences* 30: 579–582.
- MARSHALL, I.K. 2002. Congruence and cospeciation: morphological and molecular phylogenetics of the Amblycera (Phthiraptera). PhD thesis, University of Glasgow, Glasgow, Scotland. 253 pp.
- MASSARO, M., CHARDINE, J.W. & ROBERTSON, G.J. 2000. Delayed capelin (*Mallotus villosus*) availability influences predatory behaviour of large gulls on Black-legged Kittiwakes (*Rissa tridactyla*), causing a reduction in kittiwake breeding success. *Canadian Journal of Zoology* 78: 1588–1596.
- MCCOY, K.D., BOULINIER, T., SCHJØRRING, S. & MICHALAKIS, Y. 2002. Local adaptation of the ectoparasite *Ixodes uriae* to its seabird host. *Evolutionary Ecology Research* 4: 441–456.
- MCCOY, K.D., BOULINIER, T., TIRARD, C. & MICHALAKIS, Y. 2001. Host specificity of a generalist parasite: genetic

- evidence of sympatric host races in the seabird tick *Ixodes uriae*. *Journal of Evolutionary Biology* 14: 395–405.
- MCCOY, K.D., BOULINIER, T., TIRARD, C. & MICHALAKIS, Y. 2003. Host-dependent genetic structure of parasite populations: differential dispersal of seabird tick host races. *Evolution* 57: 288–296.
- MCCOY, K.D., BOULINIER, T., CHARDINE, J., DANCHIN, E. & MICHALAKIS, Y. 1999. Dispersal and distribution of the tick *Ixodes uriae* within and among seabird host populations: the need for a population genetic approach. *Journal of Parasitology* 85: 196–202.
- MCCOY, K.D. & TIRARD, C. 2002. Reproductive strategies of the seabird tick *Ixodes uriae* (Acari: Ixodidae). *Journal of Parasitology* 88: 813–816.
- MCKERNAN, D.L. & SCHEFFER, V.B. 1942. Unusual numbers of dead birds on the Washington coast. *Condor* 44: 204–206.
- MEHL, R. & TRAAVIK, T. 1983. The tick *Ixodes uriae* (Acari, Ixodidae) in seabird colonies in Norway. *Fauna Norvegica Series B* 30: 94–107.
- MONAGHAN, D.J., SHEDDEN, C.B., ENSOR, K., FRICKER, C.R. & GIRDWOOD, R.W.A. 1985. Salmonella carriage by Herring Gulls in the Clyde area of Scotland in relation to feeding ecology. *Journal of Applied Ecology* 22: 669–680.
- MONROE, A., NOAH, P. & BROWN, S. 1994. Comparison of medical treatment regimes for aspergillosis in captive Tufted Puffins (*Lunda cirrhata*). *Penguin Conservation* 7: 1–5.
- MONTEVECCHI, W.A. & MYERS, R.A. 1995. Prey harvests of seabirds reflect pelagic fish and squid abundance on multiple spatial and temporal scales. *Marine Ecology Progress Series* 117: 1–9.
- MORBAY, Y.E. 1996. The abundance and effects of ticks (*Ixodes uriae*) on nestling Cassin's Auklets (*Ptychoramphus aleuticus*) at Triangle Island, British Columbia. *Canadian Journal of Zoology* 74: 1585–1589.
- MORISHITA, K. 1930. Two nematode parasites of the guillemot. *Japanese Journal of Zoology* 3: 61–72.
- MORISHITA, T.Y., AYE, P.P., LEY, E.C. & HARR, B.C. 1999. Survey of pathogens and blood parasites in free-living passerines. *Avian Diseases* 43: 549–552.
- MOSS, S.R. & NUTTALL, P.A. 1985. Isolation of orbiviruses and uukuviruses from puffin ticks. *Acta Virologica* 29: 158–161.
- MOYER, B.R. & WAGENBACH, G.E. 1995. Sunning by Black Noddies (*Anous minutus*) may kill chewing lice (*Quadraceps hopkinsi*). *Auk* 112: 1073–1077.
- MUZAFFAR, S.B. 2000. Ectoparasites of auks (Alcidae) at the Gannet Islands, Labrador: diversity, ecology and host-parasite interactions. MSc thesis, Department of Biology, Memorial University of Newfoundland, St John's, Newfoundland. 94 pp.
- NAGASAWA, K., BARUS, V. & OGI, H. 1998a. Descriptions of larval *Contraecum variegatum* (Rudolphi, 1809) and adult *Contraecum* sp. (Nematoda: Anisakidae) collected from seabirds of the Bering Sea. *Journal of the Yamashina Institute of Ornithology* 30: 22–30.
- NAGASAWA, K., BARUS, V. & OGI, H. 1998b. *Stegophorus stellaepolaris* (Parona, 1901) (Nematoda: Acuariidae) collected from Thick-billed Murres (*Uria lomvia*) of the Bering Sea. *Journal of the Yamashina Institute of Ornithology* 30: 31–35.
- NICOLL, W. 1923. A reference list of the trematode parasites of the British birds. *Parasitology* 15: 151–202.
- NOLSØ, A. 2002. Ecology of the endoparasites of Atlantic Puffins (*Fratercula arctica grabae*) (Alcidae: Charadriiformes) *Fóðskaparrit* 50: 131–142.
- NORRIS, D.E. & OVERSTREET, R.M. 1976. The public health implications of larval *Thynnascaris* nematodes from shellfish. *Journal of Milk and Food Technology* 39: 47–54.
- NUTTALL, P.A. 1984. Tick-borne viruses in seabird colonies. *Seabird* 7: 31–41.
- NUTTALL, P.A., KELLY, T.C., CAREY, D., MOSS, S.R. & HARRAP, K.A. 1984. Mixed infections with tick-borne viruses in a seabird colony in Eire. *Archives of Virology* 79: 35–44.
- OKAICHI, T., ANDERSON, D.M. & NEMOTO, T. 1989. Red tides: biology, environmental science and toxicology. London: Elsevier.
- OLSEN, B., DUFFY, D.C., JAENSON, T.G.T., GYLFE, A., BONNEDAHL, J. & BERGSTRÖM, S. 1995. Transhemispheric exchange of Lyme disease spirochetes by seabirds. *Journal of Clinical Microbiology* 33: 3270–3274.
- OLSEN, B., JAENSON, T.G.T., NOPPA, L., BUNIKIS, J. & BERGSTRÖM, S. 1993. A Lyme borreliosis cycle in seabirds and *Ixodes uriae* ticks. *Nature* 362: 340–342.
- OPRANDY, J.J., SCHWAN, T.G. & MAIN, A.J. 1988. Tick-borne Kemerovo group orbiviruses in a Newfoundland seabird colony. *Canadian Journal of Microbiology* 34: 782–786.
- PENCE, D.B. & HOBERG, E.P. 1991. *Thalassornectes (Alcidectes) aukletae* subgen. et sp. n. (Acari: Hypoderatiidae) from the Crested and Parakeet Auklets (Aves: Charadriiformes: Alcidae). *Journal of Medical Entomology* 28: 205–209.
- PETERMANN, V.S., GLUNDER, G. & HEFFELS-REDMANN, U. 1989. Untersuchungsbeefunde an "krnak" bzw. "Tot" gefundenen Trottelummen (*Uria aalge*), Dreizehen (*Rissa tridactyla*), Silber (*Larus argentatus*) und Lachmowen (*Larus ridibundus*) aus dem Bereich der Deutschen Bucht, 1982–1985. *Dtsch Tierartztl Wochenschr* 96: 241.
- PIATT, J.F. & VAN PELT, T.I. 1997. Mass-mortality of guillemots (*Uria aalge*) in the Gulf of Alaska in 1993. *Marine Pollution Bulletin* 34: 656–662.
- PRICE, R.D., PALMA, R.L. & CLAYTON, D.H. 2003. Taxonomic review of the genus *Saemundssonina* Timmerman (Phthiraptera: Philopteridae) from the Alcidae (Aves: Charadriiformes), with description of a new species and new host-louse records. *Proceedings of the Entomological Society of Washington* 105: 915–924.
- RAPPOLE, J.H. & HUBÁLEK, Z. 2003. Migratory birds and West Nile virus. *Journal of Applied Microbiology* (Supplement) 94: S47–S58.
- PROCTOR, H.C. 2003. Feather mites (Acari: Astigmata): ecology, behavior, and evolution. *Annual Review of Entomology* 48: 185–209.
- RAMOS, J.A., BOWLER, J., DAVIS, L., VENIS, S., QUINN, J. & MIDDLETON, C. 2001. Activity patterns and effect of ticks on growth and survival of tropical Roseate Tern nestlings. *Auk* 118: 709–716.
- ROBERTSON, G.J., FIFIELD, D., MASSARO, M. & CHARDINE, J.W. 2001. Changes in nesting-habitat use of large gulls breeding in Witless Bay, Newfoundland. *Canadian Journal of Zoology* 79: 2159–2167.
- RÖMER, F. & SCHAUDINN, F. 1918. *Fauna Arctica*. Jena: Gustav Fischer.
- ROTHSCHILD, M. & CLAY, T. 1957. *Fleas, flukes, and cuckoos*. New York: Macmillan Publishing.
- ROWE, S., JONES, I.L., CHARDINE, J.W., ELLIOT, R.D. & VEITCH, B.G. 2000. Evidence for recent changes in the winter diet of murres (*Uria* spp.) in coastal Newfoundland waters. *Canadian Journal of Zoology* 78: 495–500.

- SABROSKY, C.W., BENNETT, G.F. & WHITWORTH, T.L. 1989. Bird blow flies (*Protocalliphora*) in North America (Diptera: Calliphoridae)—with notes on the palearctic species. Washington, DC: Smithsonian Institution Press.
- SAZONOV, A.A., LVOV, D.K., WEBSTER, R.G., SOKOLOVA, T.V., BRAUDE, N.A. & PORTYANKO, N.V. 1977. Isolation of an influenza virus, similar to A/Port Chalmers/1/73 (H3N2) from a Common Murre at Sakhalin Island in USSR (Strain A/Common Murre/Sakhalin/1/74). *Archives of Virology* 53: 1–7.
- SCHMIDT, G.D. 1964. First reports of *Stegophorus stellae-polaris* (Parona, 1901) Wehr, 1934 (Nematoda: Ancyracanthidae) from North America. *Canadian Journal of Zoology* 42: 1157.
- SCHREIBER, E.A. & BURGER, J. (Eds). 2002. Biology of marine birds. New York: CRC Press.
- SHIMAZU, T. 1975. Some cestode and acanthocephalan larvae from euphausiid crustaceans collected in the northern North Pacific Ocean [Japanese]. *Bulletin of the Japanese Society of Science and Fisheries* 41: 813–821.
- SHUMWAY, S.E., ALLEN, S.M. & DEE BOERSMA, P. 2003. Marine birds and harmful algal blooms: sporadic victims or under-reported events? *Harmful Algae* 2: 1–17.
- SKRJABIN, K.I. & MAMAEV, I.U.L. 1968. Gelminty Zhivotnykh Tikhogo Okeana, Biologo-Pochvennyi Institut, Akademiia nauk SSSR. Moskva: Nauka.
- SKRJABIN, K.I. & RYZHIKOV, K.M. 1973. Gelminty ptits i akutii i sopedelnykh territorii. Nematody i akantotsefaly. Gelmintologicheskaiia Laboratoriia, Akademiia nauk SSSR. Moskva: Nauka.
- SMETANINA, Z.B. 1979. Tsestody ryboiadnykh ptits primorskogo kraia. Materialy nauchnoii konferentsii vsesoyznogo obshchestva gelmintologov Vypusk 31. Tsestody i tsestodzy. Akademii nauk SSSR. Lab. Gelmint. Moskva: Akademii Nauk USSR. pp. 120–128.
- SMITH, V.A. 2001. Avian louse phylogeny (Phthiraptera: Ischnocera): a cladistic study based on morphology. *Zoological Journal of the Linnean Society* 132: 81–144.
- SONIN, M.D. 1966. Filariata of animals and man and the diseases they cause. Part 1. Aprotoidea. [Russian]. *Osnovy Nematologii* V: 17.
- SPENCE, R.P., HARRAP, K.A. & NUTTALL, P.A. 1985. The isolation of Kemerovo group orbiviruses and Uukuniemi group viruses of the family Bunyaviridae from *Ixodes uriae* ticks from the Isle of May, Scotland. *Acta Virologica* 29: 129–36.
- STEELE, J.H. & GALTON, M.M. 1971. Salmonellosis. In: Davis, J.W., Anderson, R.C., Karstad, L. & Trainer, D.O. (Eds). Infectious and parasitic diseases of wild birds. Ames, IA: Iowa State University Press. pp. 51–58.
- STEINER, F. & HAHN, E. 1953. Ueber den Nachweis von Keimen der typhus-paratyphus-enteritis-gruppe aus vogelkot von der Stora Karlso, Sweden. *Acta Pathologica Microbiologica Scandinavica* 33: 401.
- STOSKOPF, M.K. 1993. Penguin and alcid medicine. In: Fowler, M.E. (Ed). Zoo and wild animal medicine—current therapy. London: W.B. Saunders. pp. 189–194.
- STUNKARD, H.W. 1930. The life history of *Cryptocotyle lingua* (creplin), with notes on the physiology of Metacercariae. *Journal of Morphology and Physiology* 50: 143–190.
- TEMIROVA, S. & SKJARBIN, A.S. 1978. Tetrabotriaty i mezotsestoidaty lentochnye gelminty ptits i mlekopitaiushchikh. *Osnovy Tsestologii* 9: 1–117.
- THRELFALL, W. 1968. Studies on the helminth parasites of the American Herring Gull (*Larus argentatus* Pont.) in Newfoundland. *Canadian Journal of Zoology* 46: 1119–1127.
- THRELFALL, W. 1971. Helminth parasites of alcids in the northwest North Atlantic. *Canadian Journal of Zoology* 49: 461–466.
- TIMMERMANN, G. 1954. Vorlaufige ubersicht uber das amblyseres—genus *Austromenopon* Bedford, 1939 (Mallophaga). *Bonner Zoologische Beitrage* 5: 195–206.
- TIMMERMANN, G. 1974. Grupen—revisionen bei Mallophagen. X. Die *Cummingsiella* (früher *Quadriceps*)—arten der alcidae. *Mitteilungen aus dem Hamburgischen Zoologischen Museum und Institut* 70: 165–180.
- TSIMBALIUK, A.K. 1965. Skrebni zhivotnykh ostovov beringova moria. Paraziticheskie chervi domashnikh i dikikh zhivotnykh. Vladivostok: Akademii Nauka SSSR. pp. 343–348.
- TSIMBALIUK, A.K. & BELOGUROV O.I. 1964. O faune nematod ryboiadnykh ptits ostovov beringova moria. Nauch doklady vyssheii shkoly. *Biologicheskie Nauki* 4: 7–11.
- TOCIDLOWSKI, M.E., CORNISH, T.E., LOOMIS, M.R. & STOSKOPF, M.K. 1997. Mortality in captive wild-caught Horned Puffin chicks (*Fratercula corniculata*). *Journal of Zoo and Wildlife Medicine* 28: 298–306.
- WANLESS, S., BARTON, T.R. & HARRIS, M.P. 1997. Blood hematocrit measurements of 4 species of North Atlantic seabirds in relation to levels of infestation by the tick *Ixodes uriae*. *Colonial Waterbirds* 20: 540–544.
- WESENBURG-LUND, E. 1917. XX. Acanthocephaler. *Meddeleser om Gronland* 23(Suppl): 143–156.
- WHEELER, T.A. & THRELFALL, W. 1989. Synopsis of the parasites of Canada: In: Kennedy, M.J. (Ed). Ectoparasites of birds. Edmonton: Queen's Printer.
- YAMAGUTI, S. 1939. Studies on the helminth fauna of Japan. Pt. 4. Trematodes of birds. *Japanese Journal of Zoology* 8: 2.
- YAMAGUTI, S. 1959. Systema helminthum. Vol II. The cestodes of vertebrates. New York: Interscience Publishers.
- YAMAGUTI, S. 1975. A synoptical review of life histories of digenetic trematodes of vertebrates. Tokyo: Keigaku Publishing.
- YUNKER, C.E. 1975. Tick-borne viruses associated with seabirds in North America and related islands of vertebrates. *Medical Biology* 53: 302–311.