



# A New Way of Analyzing Vowels: Comparing Formant Contours Using Smoothing Spline ANOVA




Jennifer Nycz & Paul De Decker, Department of Linguistics, New York University

## 1. Introduction

- Formant values are typically measured at 1 or more single points.
- BUT: Vowels are dynamic, time-varying acoustic events.
- Formant **trajectories** may be important for conveying/perceiving contrast (e.g. Lindblom & Studdert-Kennedy 1967, Hillenbrand & Nearey 1999) OR sociolinguistic info (Thomas 2000), and could factor into changes in progress (Majors 2005).

### Smoothing Spline Analysis of Variance (SS ANOVA)

- method for statistical comparison of curves (Gu 2002; see Davidson 2006 for use of this technique in linguistic ultrasound research)
- tells you **whether** two curves are significantly different; confidence intervals indicate **where** the differences are



Individual F1 contours and smoothing splines for *pass* and *pat*.

Smoothing splines & confidence intervals for *pass* and *pat*.


Smoothing splines are functions that best fit the data. The SS ANOVA compares values for the smoothing parameters of splines; a comparison of the smoothing parameters tells you whether curves are different (see handout for more detail!)

## 2. Procedures

- 8 speakers (6 from NJ, 1 Canadian, 1 Alabamian) produce 10 tokens each of several words containing vowels of interest: *cot/caught*, *don/dawn*; *pan/pass/pad/pat*; *pin/pen*. (each word embedded within a carrier phrase, *Say \_\_\_ very loudly*.)
- 50 point formant contours (F1, F2, F3) for each vowel extracted from sound files in Praat (Boersma & Weenink 2006).
- ANOVAs run on onset, midpoint, offset formant values for each vowel.
- Splines generated for each word and SS ANOVA run in S-PLUS.

## 3. Demonstration: Ash allophony

Allophones of [æ] may differ in nasality, lingual height/backness, and/or dynamic qualities (e.g. Labov 1994, Plichta 2002, De Decker & Nycz 2005).




C has a "nasal system" typical of many NJ speakers (Ash 2002), though examination of trajectories reveal differences between *pass* and *pad/pat*.

JD also shows pre-nasal [æ] "tensing"; the diphthongal quality of her vowel in *pan* is compared here to *pad* and *pin*.

Another nasal system, though CS's pre-nasal [æ] does not show the extreme movement of JD's.

## 4. Demonstration: Low back vowel contrast



Based on point measurements at the onset and midpoint, both KB and KR show significant F1 and F2 differences between *cot* and *caught*. However, while KR's formant trajectories for the two vowels are quite similar, KB's low back vowels are not: her *caught* formants slope upward at a greater rate than those of her *cot*.

*Future research: What role do trajectories have in low back merger? (Majors 2005)*

## 5. Conclusions & Future Research

- smoothing splines enable us to compare average formant curves, instead of inspecting individual tokens (which may or may not be representative).
- Formant trajectories may differentiate allophones that are grouped together under certain single point analyses.
- SS analyses can inform single point measurements by identifying potentially important acoustic landmarks

*Future research: To what extent are trajectory differences relevant to perception? (lots of conflicting work on this topic) How can these methods be used to analyze vowel-liquid and other difficult-to-segment sequences?*