



GEOLOGY, ORE DEPOSITS AND APPLIED
ROCK GEOCHEMISTRY OF THE BUCHANS GROUP



JOHN GEOFFREY THURLOW

000103





National Library of Canada
Collections Development Branch

Canadian Theses on
Microfiche Service

Bibliothèque nationale du Canada
Direction du développement des collections

Service des thèses canadiennes
sur microfiche

NOTICE

The quality of this microfiche is heavily dependent upon the quality of the original thesis submitted for microfilming. Every effort has been made to ensure the highest quality of reproduction possible.

If pages are missing, contact the university which granted the degree.

Some pages may have indistinct print especially if the original pages were typed with a poor typewriter ribbon or if the university sent us a poor photocopy.

Previously copyrighted materials (journal articles, published tests, etc.) are not filmed.

Reproduction in full or in part of this film is governed by the Canadian Copyright Act, R.S.C. 1970, c. C-30. Please read the authorization forms which accompany this thesis.

**THIS DISSERTATION
HAS BEEN MICROFILMED
EXACTLY AS RECEIVED**

AVIS

La qualité de cette microfiche dépend grandement de la qualité de la thèse soumise au microfilmage. Nous avons tout fait pour assurer une qualité supérieure de reproduction.

S'il manque des pages, veuillez communiquer avec l'université qui a conféré le grade.

La qualité d'impression de certaines pages peut laisser à désirer, surtout si les pages originales ont été dactylographiées à l'aide d'un ruban usé ou si l'université nous a fait parvenir une photocopie de mauvaise qualité.

Les documents qui font déjà l'objet d'un droit d'auteur (articles de revue, examens publiés, etc.) ne sont pas microfilmés.

La reproduction, même partielle, de ce microfilm est soumise à la Loi canadienne sur le droit d'auteur, SRC 1970, c. C-30. Veuillez prendre connaissance des formules d'autorisation qui accompagnent cette thèse.

**LA THÈSE A ÉTÉ
MICROFILMÉE TELLE QUE
NOUS L'AVONS REÇUE**

Ottawa, Canada
K1A 0N4

GEOLOGY, ORE DEPOSITS AND APPLIED ROCK
GEOCHEMISTRY OF THE BUCHANS GROUP, NEWFOUNDLAND

by

© John Geoffrey Thurlow, B.A.Sc., M.Sc.

A Thesis submitted in partial fulfillment
of the requirements for the degree of

Doctor of Philosophy

Department of Geology
Memorial University of Newfoundland
St. John's, Newfoundland

April, 1981

ABSTRACT

The Buchans Group is a regionally extensive, submarine, calc-alkaline volcanic suite erupted at a point in time near the Ordovician-Silurian boundary. It is subdivided into Lower and Upper Subgroups, each of which consists of interbedded mafic to felsic flows, breccias, pyroclastics and related sediments. The Lower Subgroup, which contains the economic ores, consists of four volcanic formations which have been repeated structurally by thrust faults, forming seven major volcanic sequences. The economically barren Upper Subgroup is subdivided into three major sequences.

Plutonic rocks associated with the Buchans Group include the Feeder Granodiorite, Hungry Mountain Complex and the Topsails Granite. The earliest of these, the Feeder Granodiorite, is considered to be a time-equivalent plutonic facies of the Buchans Group. Deformed plutonic facies of the Hungry Mountain Complex were tectonically emplaced upon the Buchans Group during a major episode of southeast-directed Silurian thrusting. The Siluro-Devonian Topsails Granite intrudes all older formations and, in the Buchans area, consists of two main phases: a high level alkali-feldspar phase and a coarser grained equigranular peralkaline phase. Gently dipping alkali amphibole-bearing ignimbrite forms roof pendants within the Topsails Granite. These are interpreted as extrusive

equivalents of the Granite and may be outliers of the Springdale Group.

The Buchans ores are high grade polymetallic massive sulphide deposits associated with submarine felsic volcanic rocks near the top of the Lower Buchans Subgroup. Three genetically related ore types are recognized; stockwork ore, in situ ore, and mechanically transported ore. Stockwork ore consists of networks of veinlet and disseminated base metal sulphides within highly silicified and locally chloritized host rocks. In situ ore lies stratigraphically above stockwork mineralization and is composed of banded to streaky yellow ore, black ore and barite. Transported ore consists of discrete sulphide and barite fragments in breccias which flank in situ ores and form substantial distinct orebodies. Their origin is interpreted as a result of brecciation at source, transport down-slope in paleotopographic channels by gravity-driven submarine debris flows, and deposition in depressions. The Buchans ores and their environment are similar to those of the Japanese Kuroko deposits in almost all respects.

Whole-rock chemical analyses have shown that Buchans Group volcanic rocks are similar in chemical abundances and variations to other calc-alkaline volcanic suites. There was apparently little chemical evolution in Buchans Group flows through the stratigraphic column, and

flows within the ore horizon sequences are essentially similar to their counterparts elsewhere in the Buchans Group. Pyroclastic rocks of the ore horizon are chemically similar to those at other stratigraphic horizons and show no anomaly related to mineralization with the exception of high Ba content.

Study of the unaltered pyroclastic rocks of the ore horizon has shown that there are no regional variations in element concentrations, element ratios, discriminant scores or factor scores, either along or across stratigraphy, which bear relationship to mineralization. Erratic base metal concentrations are present in host pyroclastic rocks only within 50 m of ore. Visibly altered footwall rocks have undergone chemical changes typical of rocks in most massive sulphide terranes.

Anomalously high concentrations of Ba in ore horizon pyroclastics is a regional feature which provides a means of distinguishing ore horizon pyroclastics from similar lithologies at other stratigraphic horizons. The Ba anomaly is caused by microscopic barite which was widely distributed by subaqueous pyroclastic flows which transported most ore horizon pyroclastics.

TABLE OF CONTENTS

	Page
Abstract	ii
List of Figures.....	xi
List of Tables	xv
List of Plates	xvii
 CHAPTER 1. GENERAL INFORMATION	 1
1.1. Location and Access.....	1
1.2. Topography, Vegetation, Climate and Outcrop Exposure.....	1
1.3. History of Mining.....	4
1.4. Previous Work in the Buchans Area.....	5
1.5. Previous Lithogeochemical Studies in the Vicinity of Massive Sulphide Mineralization..	10
1.6. Present Study.....	12
1.7. Acknowledgements.....	14
 CHAPTER 2. THE STRUCTURAL AND STRATIGRAPHIC SETTING OF THE BUCHANS GROUP.....	 17
2.1. Introduction.....	17
2.2. The Relationship of the Buchans Group to Struct- urally Underlying Formations.....	19
2.3. Buchans Group Correlatives.....	20
2.4. Granitoid and Related Rocks (Topsails Granite and Feeder Granodiorite).....	21
2.5. Volcanic Outliers Within the Topsails Granite (Springdale Group?).....	32

	Page
2.6. Hungry Mountain Complex	34
2.7. Contact Relations Between the Hungry Mountain Complex and the Buchans Group	41
2.7.1. Relationships Within the Hungry Mountain Complex	41
2.7.2. Relationships Within the Buchans Group ..	42
2.8. Siluro-Devonian Deformation	48
2.9. Carboniferous Rocks	50
2.10. Summary of Chapter Two	52
CHAPTER 3. GEOLOGY OF THE BUCHANS GROUP	54
3.1. Introduction	54
3.2. Stratigraphy of the Buchans Group	56
3.2.1. Stratigraphy of the Lower Buchans Subgroup	56
3.2.1.1. Footwall Basalt	56
3.2.1.1.1. Skidder Prospect	69
3.2.1.2. Footwall Arkose and Wiley's Sequence	69
3.2.1.2.1. Connel Option	76
3.2.1.3. Intermediate Footwall	77
3.2.1.4. Lucky Strike Ore Horizon Sequence	79
3.2.1.4.1. Rhyolite	83
3.2.1.4.2. Pyritic Siltstone ...	84
3.2.1.4.3. Dacitic Pyroclastics ..	87
3.2.1.4.4. Ore Horizon	98

	Page
3.2.1.5. Lake Seven Basalt.....	99
3.2.1.6. A Stratigraphic Discontinuity at the Top of the Lake Seven Basalt.....	100
3.2.1.7. Prominent Quartz Sequence..	102
3.2.1.8. Ski Hill Sequence.....	106
3.2.1.9. Oriental Intermediate Footwall	109
3.2.1.10. Oriental Ore Horizon Sequence	110
3.2.2. Stratigraphy of the Upper Buchans Subgroup	111
3.2.3. Diabase and Related Intrusives.....	119
3.3. Structure and Metamorphism of the Buchans Group	122
3.3.1. Structure of the Buchans Area.....	122
3.3.1.1. Silurian Thrust Faulting in the Buchans Group.....	122
3.3.1.2. East-West Fold Axes Related to Thrusting.....	126
3.3.1.3. Devonian Folding.....	127
3.3.2. Metamorphism of the Buchans Area...	128
CHAPTER 4. ORE DEPOSITS OF THE BUCHANS GROUP....	132
4.1. Introduction.....	132
4.2. Intermediate Footwall and Stockwork Mineraliz- ation	132
4.2.1. General Comments.....	132
4.2.2. Stratigraphy and Mineralization in the Lucky Strike Area.....	139

	Page
4.3. Lucky Strike Ore Horizon Sequence	148
4.3.1. Footwall Rocks	148
4.3.2. Lucky Strike Orebodies.....	149
4.3.3. Ore Horizon Breccias	152
4.3.4. MacLean, Rothermere, Two Level and Lucky Strike North; Transported Ores	159
4.3.5. Clementine Prospect.....	178
4.4. Oriental Ore Horizon Sequence	178
4.4.1. Oriental No. 1 Orebody	178
4.4.2. Oriental No. 2 Orebody	181
4.4.3. Sandfill Prospect	186
4.4.4. Middle Branch Prospect	189
4.4.5. Old Buchans Orebodies	196
4.5. Selected Comments	196
4.5.1. Miscellaneous Occurrences of Sulphide Clasts	196
4.5.2. Comments on Granite Boulders at Buchans	199
4.5.3. Buchans as Compared to Other Occurrences of Mechanically Transported Ores ...	201
4.6. Comparison of Buchans and Kuroko Deposits ..	201
CHAPTER 5. GENERAL PETROCHEMISTRY OF THE BUCHANS GROUP	207
5.1. Post-magmatic Chemical Changes of Buchans Group Rocks	207
5.2. General Elemental Abundances and Chemical Variations of the Buchans Group Rocks	210

	Page
5.3. Chemical Variations in Flows as a Function of Stratigraphic Position.....	234
5.3.1. Variations in Compositions of Mafic and Intermediate Flows Through the Stratigraphic Sequence	235
5.3.2. Variations in Compositions of Felsic Flows Through the Stratigraphic Sequence	236
5.3.3. Some Comments on the Chemistry of the "White Rhyolite" at Buchans.....	239
5.3.4. Summary	242
5.4. Geochemical Comparison of Flows and Cogenetic Fragmental Rocks.....	242
CHAPTER 6. GEOCHEMISTRY OF THE ORE HORIZON DACITIC TUFFS	248
6.1. Introduction.....	248
6.2. General Chemical Abundance in the Ore Horizon Dacitic Tuffs.....	248
6.3. Correlation Matrix for Ore Horizon Dacitic Tuffs,	254
6.4. R-mode Factor Analysis of the Ore Horizon Dacitic Tuffs - Major Elements.....	254
6.5. R-mode Factor Analysis of the Ore Horizon Dacitic Tuffs - Major and Trace Elements....	261
CHAPTER 7. APPLIED ASPECTS OF THE GEOCHEMISTRY OF ORE HORIZON DACITIC TUFFS.....	265
7.1. Geochemical Comparison of Ore Horizon Dacitic Tuffs to Other Dacitic Tuffs.....	265

	Page
7.2. Chemical Variations in the Ore Horizon Dacitic Tuffs as a Function of Distance to Ore	270
7.2.1. Introduction	270
7.2.2. Chemical Variations Versus Distance to Nearest Ore (Whether Transported or In Situ Ore)	271
7.2.3. Chemical Variations Versus Distance to In Situ Ore	274
7.2.4. Discriminant Scores as a Function of Distance to Ore	276
7.3. The Cause of the Ba Anomaly in the Ore Horizon Dacitic Tuffs	279
 CHAPTER 8. SUMMARY AND EXPLORATION APPLICATION...	286
8.1. Summary of Thesis	286
8.2. Exploration Applications	287
References	291
Appendix I	1-1
Appendix II	2-1
Appendix III	3-1
Appendix IV	4-1

LIST OF FIGURES

	Page
Figure 1-1 Location of Buchans with respect to tectonostratigraphic zones of Newfoundland.....	2
Figure 1-2 Buchans area drill hole location map (in back pocket)	
Figure 2-1 Generalized geological map of the Buchans area.....	18
Figure 2-2 Geological map of the Buchans area (in back pocket)	
Figure 2-3 Schematic cross section showing relationships between the Hungry Mountain Complex, Topsails Granite and Buchans Group.....	47
Figure 3-1 Schematic post-thrusting stratigraphy of the Buchans Group.....	57
Figure 3-2 Vertical section through Lucky Strike, Rothermere and MacLean Orebodies	80
Figure 4-1 Distribution of in situ and transported mineralization.....	133

	Page
Figure 4-2 Distribution of alteration zones within the Intermediate Footwall.....	138
Figure 4-3 Sketch showing bedded siltstones trans- ected by hairline fractures.....	174
Figure 4-4 Isopach map of the Oriental ore horizon sequence.....	191
Figure 4-5 Schematic cross section of the Sanfill- Middle Branch area.....	192
Figure 4-6 Paleotopographic plan of the Sandfill- Middle Branch area.....	193
Figure 4-7 Plan showing distribution of high grade sulphide clasts.....	197
Figure 5-1 Alkali: silica diagram for Buchans Group flows.....	214
Figure 5-2 AFM diagram for Buchans Group flows....	216
Figure 5-3 K_2O : SiO_2 classification of Buchans Group flows.....	217
Figure 5-4 Ti: Zr diagram for Buchans Group basalts.....	218
Figure 5-5 Ti/100: Zr: Sr/2 diagram for Buchans Group basalts.....	219

	Page
Figure 5-6 K:Rb diagram for Buchans Group flows...	220
Figure 5-7 SiO_2 histogram, Buchans Group flows....	222
Figure 5-8 Anhydrous Harker diagrams for Buchans Group flows.....	225
Figure 5-9 CaO, MgO and MnO variations in flows...	238
Figure 5-10 SiO_2 : differentiation index diagram com- paring unaltered ore horizon rhyolites to rhyolites at other stratigraphic positions.....	240
Figure 5-11 Geochemical comparison of ore horizon dacitic tuffs and rhyolite flows.....	245
Figure 6-1 Histograms for all elements determined in ore horizon dacitic tuffs.....	250
Figure 7-1 Histogram of Ba abundances in ore horizon dacitic tuffs as compared to non-ore horizon tuffs.....	266
Figure 7-2 Histogram of discriminant scores.....	268
Figure 7-3 Lateral variation of elements in ore horizon dacitic tuffs as a function of distance to nearest ore.....	272

Page

Figure 7-4	Lateral variations of elements in ore horizon dacitic tuffs as a function of distance to nearest in-situ ore.....	277
Figure 7-5	Lateral variations of discriminant scores in ore horizon dacitic tuffs as a function of distance to ore.....	280
Figure 7-6	Schematic diagram showing modes of occurrence of barite in ore horizon dacitic tuffs.....	282

		Page
Table 1-1	Summary of major element dispersions in relation to volcanogenic massive sulphide deposits.....	13
Table 3-1	Stratigraphy of the Buchans Group.....	58
Table 3-2	Buchans Group thrust correlatives - Model A.....	125
Table 3-3	Buchans Group thrust correlatives - Model B.....	125
Table 4-1	Ore deposits of the Buchans area.....	134
Table 5-1	Average chemical composition of Buchans area rocks.....	211
Table 5-2	Correlation matrix for Buchans Group flows.....	223
Table 5-3	Composition of "white rhyolite" compared to other ore horizon rhyolites.....	241
Table 6-1	Average chemical composition of ore horizon dacitic tuffs.....	249
Table 6-2	Correlation matrix for Lucky Strike ore horizon dacitic tuffs.....	255

	Page
Table 6-3 Rotated factor matrix for dacitic tuffs of the Lucky Strike ore horizon sequence - major elements.....	257
Table 6-4 Rotated factor matrix for dacitic tuffs of the Lucky Strike ore horizon sequence - major and trace elements.....	264
Table 7-1 Discriminant function data.....	269

LIST OF PLATES

		Page
Plate 2-1	Feeder Granodiorite.....	22
Plate 2-2	Core samples of Feeder Granodiorite, rhyolite, dacitic crystal-vitric tuff and volcanic conglomerate.....	24
Plate 2-3	Photomicrograph of alkali-feldspar granite.....	26
Plate 2-4	Peralkaline granite.....	28
Plate 2-5	Hornblende gabbro breccia, intrusive into the Hungry Mountain Complex.....	36
Plate 2-6	Peralkaline dyke cutting deformed Hungry Mountain Complex.....	38
Plate 2-7	Mylonite from the base of the Hungry Mountain Complex.....	40
Plate 2-8	Penetrative foliation in felsic tuffs below Hungry Mountain thrust.....	43
Plate 2-9	Carboniferous red sandstone and conglomerate.....	49
Plate 3-1	Pillow lava, Footwall Basalt.....	59
Plate 3-2	Nearly whole pillows in pillow breccia, Footwall Basalt.....	61

	Page
Plate 3-3 Highly disaggregated pillow breccia, Footwall Basalt.....	62
Plate 3-4 Graded pillow breccia, Footwall Basalt..	63
Plate 3-5 Spheroidal banded rhyolite within Footwall Basalt unit.....	65
Plate 3-6 Footwall Arkose, typical lithic arkose..	72
Plate 3-7 Hematitized ore horizon rhyolite.....	82
Plate 3-8 Pyritic siltstone, Lucky Strike ore horizon sequence.....	85
Plate 3-9 Dacitic pyroclastic, Lucky Strike ore horizon sequence.....	88
Plate 3-10 Dacitic pyroclastic from "coarse quartz" unit.....	95
Plate 3-11 Pyroclastic breccia, "coarse quartz" unit.....	96
Plate 3-12 Rhyolite, Prominent Quartz sequence....	101
Plate 3-13 Andesitic breccia, Ski Hill sequence...	107
Plate 3-14 Volcanic conglomerate of Upper Arkose..	113
Plate 3-15 Flow banded rhyolite, Upper Buchans Subgroup.....	117

	Page
Plate 4-1 Stockwork mineralization, Lucky Strike area.....	135
Plate 4-2 Green mariposite-like sericite after plagioclase.....	145
Plate 4-3 Streaky, very high grade in situ ore...	151
Plate 4-4 Typical granitoid-bearing breccia conglomerate.....	155
Plate 4-5 Granodiorite cobble in ore breccia.....	156
Plate 4-6 Low grade, transported, fragmental ore..	161
Plate 4-7 High grade transported ore.....	163
Plate 4-8' Plastically deformed ore fragments.....	164
Plate 4-9 Fragment composed of fragmental ore.....	166
Plate 4-10 Fine grained, bedded, high grade, clastic ore.....	169
Plate 4-11 Siltstone fragments with altered rims...	170
Plate 4-12 Pyritic siltstone cut by fractures with adjacent alteration.....	172
Plate 4-13 Banded Oriental No. 1 ore.....	179
Plate 4-14 Black ore pebble in drillcore sample of pyroclastic breccia.....	198

	Page
Plate 4-15 Fragmental ore from the Japanese Ainaí Kuroko deposit.....	202
Plte 4-16 Motoyama volcanic breccia, Kosaka Mine..	204
Plate 7-1 Core sample containing yellow ore clast with dark rim.....	284

CHAPTER ONE

GENERAL INFORMATION

1.1. Location and Access

The area of study is located in central Newfoundland between latitudes $48^{\circ}42'N$ and $48^{\circ}58'N$ and longitudes $56^{\circ}33'W$ and $57^{\circ}00'W$ (Fig. 1-1). The town of Buchans is accessible by 77 km of paved road which leaves the Trans Canada Highway at Badger. A number of gravel surfaced roads around the town of Buchans provide access to areas up to 4.5 km from town.

Most parts of the study area are accessible on a single day's outing on foot with transmission lines, bogs and some stream beds providing best walking conditions. All the lakes and ponds in the area are too small and shallow for float plane operation and no rivers are navigable over significant distance by canoe.

1.2. Topography, Vegetation, Climate and Outcrop Exposure

The study area is a gently rolling upland area. All parts of the area, except the extreme northwest, drain into Red Indian Lake which forms the topographic low of the area. Large areas to the north and west, underlain by granitoid rocks, form a bog-covered and boulder-strewn barren plateau. The area underlain by volcanic rocks, in




Figure 1 - 1 : Location of Buchans with respect to tectonostratigraphic zones of Newfoundland and significant post-ophiolite volcanogenic massive sulphide deposits.

general, slopes irregularly towards Red Indian Lake and is covered with poorly drained bog and shallow ponds in low-lying areas and dense spruce and fir forest on small hummocks.

In general, the inland location of the Buchans area results in a somewhat less moderated climate than the rest of Newfoundland. Field work is comfortable from May to October. Precipitation is moderate with summers being somewhat drier than spring or fall. Mosquitos, black flies and other biting insects are a problem for much of the summer and rutting moose and their hunters should be avoided during the fall.

Exposure of outcrop is, in general, very poor and would average much less than one percent over the study area. The best exposures are found in stream beds and on small abrupt hills. The intervening bog and forest normally contain a few widely scattered outcrops. Though lichens may be thick and weathering may be deep, many inland outcrop can be greatly enlarged and improved by stripping off moss cover.

During the present study, at least half of the 350,000 m of Buchans drill core was logged and scores of sections plotted. Although much of the core comes from the immediate Buchans area, a number of holes have been drilled at more distant localities. The location of drill

holes is shown on Fig. 1-2 (in pack pocket).

1.3. History of Mining

The history of the Buchans area has recently been compiled by Neary (1980).

The outcropping deposits at Old Buchans were discovered in 1905 by Matty Mitchell, a Micmac prospector employed by the Anglo Newfoundland Development Company.

Development work on the property was carried out between 1906 and 1911, but efforts were suspended due to metallurgical difficulties with the fine grained sulphide aggregate. Development began in earnest in 1926 by ASARCO as a result of their discovery of selective flotation, and production began in 1928 at a rate of 500 tons per day following the discovery of additional orebodies. In 1930 the milling capacity was increased to its present rate of 1250 tons per day.

The near-surface Oriental No. 1 and Lucky Strike Orebodies were discovered by equipotential surveys carried out in 1926 under the direction of Hans Lundberg (Lundberg 1957). (These discoveries represent the first successful application of geophysics in North America). The Rothermere and Maclean (formerly Rothermere #4) orebodies were discovered between 1947 and 1950 as a result of intensive

drilling under the direction of H.J. MacLean. E.A. Swanson became chief geologist in 1951 and was responsible for the discovery of the Oriental #2 orebody in 1953. Since that time the sub-economic Clementine and Sandfill prospects have been discovered but available tonnages are not sufficient to warrant production.

During the past twenty years virtually many methods of surface geochemistry and ground and airborne geophysics have been used in an attempt to outline new ore but with only limited success. Diamond drilling has always, and will continue to play a major role in the detailed local exploration, and over 350,000 m of drill core is stored in the core sheds in Buchans. The careful logging and re-logging of this core has made the Buchans area the most thoroughly studied local volcanic environment in the province.

1.4. Previous Geological Work in the Buchans Area

Murray (1871) described the rocks in the vicinity of Red Indian Lake in general terms and considered them to be correlative with "Silurian" rocks of the Exploits valley. He considered granites in the area to be of Laurentian (Archean) age.

Snelgrove (1928) implied the presence of two ages of granites in the Buchans area. The younger granites

cut the Buchans volcanics and he considered them to play an important role in the mineralizing process. With reference to the older granites, Snelgrove (1928, p. 1113) states that "lavas actually abut against the granites approximately five miles north of the lake (Red Indian), although the age of the intrusives is probably later than 'Laurentian' as Murray mapped them". The occurrence of Snelgrove's implied unconformity has not been mentioned by subsequent workers. Snelgrove's thin section descriptions of rocks from the mine area are the first published.

The paper by Newhouse (1931) represents the first integrated descriptive work on the regional and local geology and on the ore deposits. He considered the volcanics to be of Cambro-Ordovician age and noted the similarity of cherts of the Buchans area to those in Notre Dame Bay. Plant remains collected by Newhouse indicated that poorly indurated strata along the shores of Red Indian Lake are of Carboniferous age.

After seven years of study, George (1937) described both the regional and local geology of the Buchans area. This excellent paper includes a number of maps, photographs, photomicrographs and cross sections made available by further development of the Lucky Strike and Oriental Orebodies. Description of the local stratigraphy, structure and mineralization occupies a prominent portion of

the paper and his account of the Lucky Strike and Oriental Orebodies forms a valuable record of ores long since removed by mining. He noted the association between quartz porphyry (dacitic tuff) and ore, although he considered this unit to be of intrusive origin. George (1937) recognized the presence of two major stratigraphic horizons favourable for mineralization, and considered the Lucky Strike horizon to overlie the Oriental horizon. Lamprophyre dykes described by George are considered by later authors to be diabase.

The Buchans Staff (1955) provided the first published general descriptions of the Rothermere Orebodies as well as summaries of mining and milling practices.

Relly (1960) was responsible for the first comprehensive written report of the geology of the immediate mine area. The bulk of this work is descriptive, based on field observations and hundreds of thin and polished sections. In addition, the thesis includes the first petrochemical analyses aimed at understanding alteration related to the mineralization. Despite amassing considerable evidence in favour of a syngenetic origin for the ore, Relly considered the ore to be of epigenetic hydrothermal replacement origin.

In a paper judged best of the year by the Canadian Institute of Mining and Metallurgy, Swanson and Brown

(1962) reduced Relly's voluminous descriptive work to publishable size integrating most of the ideas of the mine staff at the time. They suggested an Ordovician age for the Buchans Group and considered it to be correlative with the Roberts Arm Group to the northeast. They considered the ore to be of epigenetic origin but Anger (1963) deduced from published descriptions that the ores were similar to Meggen and Rammelsberg and, therefore, probably syngenetic.

Williams (1967) considered rocks of the Buchans area to be of Silurian age, based on the presence of red sandstones on the southeast shore of Red Indian Lake and some lithologic similarities of the Buchans Group to the Springdale Group. Mapping by Williams (1970) at 1:250,000 represented the first published regional geological map which included the Buchans area.

Anderson (1972a) carried out regional mapping at 1:50,000 but, to date, only preliminary maps are available. Anderson (1972b) considered the Buchans Group to conformably overlie plutonic boulder conglomerate of the Exploits Group which he considered 'similar to Silurian Goldson Conglomerate', and thus implied a Silurian age for the Buchans Group.

Since the publication by Swanson and Brown (1962)

the ideas of the mine and exploration staff concerning ore genesis have changed and this has resulted in new advances in understanding the stratigraphy of the area. Entwistle and Barnes (1971) first suggested that the Oriental sequence was a fault repetition of the Lucky Strike sequence, an hypothesis supported by Walker and King (1975). The present author investigated the relationship between rock geochemistry and mineralization (Thurlow, 1973) and these findings and a geological summary were published by Thurlow, Swanson and Strong (1975) and Thurlow (1977).

In 1976, ASARCO Inc., and the Price Co. Ltd. commissioned a number of researchers to study various aspects of Buchans geology with the results published as a volume commemorative of fifty years of mining at Buchans. At the time of writing, most studies are near completion.

The coloured 1:50,000 geological map of the Buchans area accompanying this work (Fig. 2-2) was prepared to form part of a paper by Thurlow and Swanson (1980). A similar map of the Buchans sheet (12A/15) will be published shortly by Kean (1980).

In addition to the above mentioned workers, there exist several theses (e.g., Woakes, 1954; Catherall, 1960; Alcock, 1961) and a number of unpublished company reports (e.g., MacLean, 1941; Swanson, Perkins, and Higgins,

1955; Larsen, 1973).

1.5. Previous Lithogeochemical¹ Studies in the Vicinity of Massive Sulphide Mineralization

The increasing difficulty of discovering massive volcanogenic sulphide deposits has fostered abundant lithogeochemical research. Govett and Nichol (1979) have recently compiled a review of work to date. General references such as Hawkes and Webb (1962), Bradshaw et al. (1970), Boyle and Garrett (1970), Sakrison (1971), and Levinson (1974) discuss the techniques and problems encountered in economic lithogeochemical surveys and refer to the various successes and failures of numerous individual studies. Regional lithogeochemical surveys in areas of massive sulphide mineralization have met with some academic success in areas such as the Archean of the Canadian Shield (e.g., Davenport and Nichol, 1972; Descarreaux, 1973; Bennett and Rose, 1973; Larson and Webber, 1977; McConnell, 1976; Sopuck, 1977; Lavin, 1976), New Brunswick (Govett et al., 1974; Graf, 1977) and Cyprus (Govett and Pantazis, 1971). Similarly, local studies in the immediate vicinity of massive sulphide mineralization have resulted in the detection of anomalies of varying extent and intensity in such areas as the Canadian Shield (e.g., Sakrison, 1967; Spitz and Darling, 1978; Marcotte and David, 1978), Cyprus (e.g., Govett, 1972; Pantazis

1. The terms "lithogeochemistry" and "rock geochemistry" are used synonymously in this study.

and Govett, 1973), Japan (e.g., Tatsumi and Clark, 1972; Tono, 1974), the Scandinavian Caledonides (e.g., Nilsson, 1968; Gjelsvik, 1968; Nairis, 1971) and the Appalachians (e.g., Gale, 1969; Thurlow, 1973; Whitehead, 1973; Whitehead, 1973; Whitehead and Govett, 1974; Goodfellow, 1975; Govett and Goodfellow, 1975).

In spite of the apparent ability of many rock geochemical studies to be capable of guiding massive sulphide exploration, the exploration community in general, have been hesitant in applying such methods. This has been, in part, due to costs but is more a function of the relative speed and efficiency of applied geophysical, geochemical and geological techniques in unequivocally pinpointing specific targets.

Rock geochemistry is plagued by a number of other deficiencies largely related to its relative infancy. Some of these include:

- 1) a lack of universally applicable parameters to areas of unknown mineral potential,
- 2) a general lack of research into the mode of occurrence of base metals in silicate rocks and minerals,
- 3) a lack of integrated regional and local lithogeochemical surveys,

- 4) a lack of integrated stratigraphic, petrological and lithogeochemical data,
- 5) a general lack of convincing demonstration that anomalies are indeed related to mineralization (especially in regional surveys),
- 6) the presence of very subtle (and sometimes dubious) anomalies often identified only by advanced statistical techniques,
- 7) the presence of obvious alteration and/or mineralization (especially in local surveys) which requires no geochemical analysis to identify the anomalous situation.

Rock geochemical studies have been successful in characterizing the nature of chemical change in massive sulphide alteration zones; particularly the near universal removal of Na_2O and addition of MgO due to breakdown of plagioclase and chlorite formation (Table 1-1). This has proven most useful in areas where deformation and metamorphism have masked original textures. Also successful, has been the ability of rock geochemical studies to assist in local and regional correlation of mineralized horizons.

1.6. Present Study

This study stems from earlier work by the author (Thurlow, 1973) which indicated that applied rock

Table 1-1 (after Govett and Nichol, 1979)

Summary of major element dispersions in relation to
volcanogenic massive sulphide deposits

Deposit	Alteration mineralogy	Elements enriched	Elements depleted	Elements unchanged	Age
Kuroko, Japan Lambert & Sato (1974)	Mon, Ser, Chl, Kaol	K, Mg, Fe, Si	Ca		Cenozoic
Kuroko, Japan Tatsumi & Clark (1972)	Ser, Qtz, Cal	Mg, K	Na, Ca, Fe	Al	Cenozoic
Hitachi, Japan Kuroda (1961)	Cord, Anthoph.	Mg, Fe, Ba	Na, Ca, Sr		Cenozoic
Buchans, Canada Thurlow et al. (1975)	Chl, Ser, Qtz	Mg, Fe, Si	Na, Ca, K		Paleozoic
Heath Steele, Canada Wahl et al. (1975)	Chl, Ser	Mg	Na, Ca		Paleozoic
Brunswick No. 12 Canada Goodfellow (1975)	Chl, Ser, Qtz	Mg, Fe, (Mn), (K)	Na, Ca (Mn), (K)	Al	Paleozoic
Killingdal, Norway Rui (1973)	Chl, Bio, Qtz	Mg, K, Mn	Na, Ca, Si	Al, Ti, Fe (total)	Paleozoic
Skorovass, Norway Gjelsvik (1968)	Chl, Ser	Mg	Na, Ca		Paleozoic
Boliden, Sweden Nilsson (1968)	Chl, Ser, Qtz, And	Mg, K, Al	Na, Ca		Proterozoic
Mattabi, Canada Franklin et al. (1975)	Qtz, Carb, Ser, Chld, Chl, And, Gar, Kyan, Bio	Fe, Mg	Na, Ca		Archean
Millenbach, Canada Simmons et al. (1973)	Chl, Ser, Anthoph, Cord.	Mg, Fe	Na, Ca, Si		Archean
Mines de Poirier Canada Descarreaux (1973)	Chl, Ser,	Mg, K	Na, Ca	Si	Archean
Lac Dufault, Canada Sakrisson (1966)	Chl, Ser	Mg, Fe, Mn	Na, Ca	Al, Ti K, Si	Archean
East Waite, Moberg, Joutel, Poirier, Agnico-Eagle, Mattabi, Sturgeon Lake, South Bay McConnell (1976)	Qtz, Ser, Chl, Carb, Sauss, Epidote	Mg, Fe	Na, Ca	Si, Al	Archean

geochemistry may be of use in exploration at Buchans. It was realized, that in order to be properly interpreted, the geochemistry had to be expanded to a more regional scale and that a thorough understanding of the complex stratigraphy of the area was required. The purpose of this study is therefore:

- 1) to establish the relationships of the Buchans Group to surrounding lithologies,
- 2) to refine the volcanic stratigraphy of the Buchans Group
- 3) to determine the nature of the ore occurrences and to compare them with other volcanogenic sulphide deposits,
- 4) to establish the general petrochemical character of the Buchans Group, and
- 5) to determine whether rock geochemical techniques have practical exploration application at Buchans.

1.7. Acknowledgments

A study so protracted as this required persistent financial, technical and moral support from many people and organizations.

Funding from a Memorial University Fellowship, an NRC Bursary and liberal grants from Dave Strong's NRC and GSC Research Grants are all gratefully acknowledged.

Substantial support was made available through the ASARCO - Abitibi-Price Cotenancy by Dick Brown and Eric Swanson in the form of bursaries and summer research employment. The Cotenancy further sponsored a tour of the Kuroko deposits and reduced the sting at the Comptroller's Office.

Enormous technical assistance and advice was received from Jaan Vahtra, Alan Thompson, Foster Thornhill, Gerry Ford and Lloyd Warford. Gert Andrews cheerfully and patiently taught me the intricacies of the A.A. and Dave Press wrote and modified computer programs and helped in many other ways. Lloyd James made available the expertise of ASARCO's Salt Lake City lab where base metal (as sulphide) determinations were undertaken. Numerous underground tours were arranged by Ken Larson with the aid of Henry Schumacher and Juri Houtman. Mort Verbiski and Peter Brockie helped in a myriad of ways. Bev King willingly typed and re-typed the manuscript.

The author has benefited greatly from discussions with supervisors and colleagues Eric Swanson, Dave Strong, Takeo Sato, Hank Williams, Ed Perkins, Dave Barbour, Tony Pryslak, Peter Walker, Baxter Kean, Paul Dean, Don Baker and all the contributors to the Buchans Volume.

In the darkest hours of my procrastination, pep talks from Bob Stevens, Dave Strong and my wife Yvonne kept me going.

My employers, Abitibi-Price, and in particular Gerry Colborne, Hamish McGregor and Harry Rosier made available the time and facilities necessary to complete this work.

One's thesis advisor normally has tremendous input into a student's work and, in my case, Dave Strong was no exception. I have been doubly fortunate to have benefited from the experience, support, co-operation and advice of Eric Swanson without whose help, this work would not have been possible.

CHAPTER TWO
THE STRUCTURAL AND STRATIGRAPHIC SETTING OF
THE BUCHANS GROUP

2.1. Introduction

The Buchans Group and adjacent formations form part of the Dunnage Zone (Williams, 1979); one of four tectonostratigraphic sub-divisions of the Newfoundland Appalachians. Regional compilation and correlation within the Dunnage Zone by Dean (1977) and Kean et al. (1980) have shown that a pre-Caradocian "early island arc" volcanic sequence overlies Cambro-Ordovician oceanic crust. The "early arc" volcanics are in turn overlain by a widespread graptolitic Caradocian shale which is succeeded by flych and post-Caradocian volcanics, including the Buchans Group.

Rocks bounding on the Buchans Group in space and time include a large area of granitoid rocks to the east, north and west (the Topsails Granite of Baird, 1960), foliated plutonic rocks of the Hungry Mountain Complex to the north, Carboniferous rocks along Red Indian Lake (Fig. 2-2) and the thick volcano-sedimentary terrane south of Red Indian Lake (Fig. 2-1). These units have been examined in this study mainly to determine their stratigraphic relationships to the Buchans Group.




Figure 2-1. Generalized geological map of the Buchans area.

2.2. The Relationship of the Buchans Group to Structurally Underlying Formations

The relationship between the Buchans Group and underlying rock units is not known with certainty though recent mapping at 1:50,000 by Kean (1977, 1979, 1980) has elucidated many aspects.

Southeast of Red Indian Lake, volcanic and sedimentary rocks of the pre-Caradocian Victoria Lake Group are overlain by fossiliferous Caradocian black shales, slates and argillites (Kean and Jayasinghe, 1980). These are in turn overlain by a plutonic boulder-bearing greywacke-conglomerate sequence, lithologically similar to the Sansom Greywacke and Goldson Conglomerate in Notre Dame Bay (Anderson, 1972). The Buchans Group structurally overlies this sequence. Though the contact, or contact zone, is not exposed, it has been considered both a fault by Williams (1970) and a conformable contact by Anderson (1972). Through mapping and regional correlations, Kean (1977, 1980) interprets the Buchans Group as younger than the greywacke-conglomerate sequence but also in fault contact with it. The author considers this the most reasonable interpretation though uncertainties exist as to the significance of a major structural lineament traversing Red Indian Lake and a red sandstone sequence on the southeast shore of the Lake. These uncertainties cannot

be dismissed until fossil evidence of the age of the Buchans Group is obtained.

2.3. Buchans Group Correlatives

With few exceptions, geologists have considered, and still consider, the Buchans Group to be a time-equivalent regional correlative of the Roberts Arm Group of Notre Dame Bay (e.g., Swanson and Brown, 1962). There are many arguments in support of this contention despite a lack of fossil control. The Roberts Arm and Buchans Groups are both regionally extensive and lithologically similar submarine volcanic sequences. Both appear to conformably overlie a post-Caradocian flysch sequence and stratigraphically underlie the dominantly subaerial Springdale Group. Both are chemically similar mafic to felsic calc-alkaline suites (Strong, 1977). Both have undergone a similar structural history of thrust faulting and Acadian folding. Both contain a variety of base metal sulphide deposits, some of which are remarkably similar (e.g., Buchans and the Bull Road showing at Pilley's Island).

To date, the Buchans Group has proven barren of fossils diagnostic of its age. The best estimate of the age of the Buchans Group is Rb-Sr whole rock isochron of 447 ± 18 Ma, essentially on the Ordovician-Silurian

boundary (Bell and Blenkinsop, 1980). A similar age of 448 ± 7 Ma has recently been derived from the Roberts Arm volcanics (Bostock, 1978; Bostock et al., 1979).

2.4. Granitoid and Related Rocks (Topsails Granite and Feeder Granodiorite)

Granitic to gabbroic rocks underlie the area north and west of the Buchans Group (Fig. 2-1) and comprise a portion of a larger plutonic complex which underlies approximately 7200 sq. km (3000 sq. mi.) of west central Newfoundland. Brief study by the author has led to subsequent more detailed work, recently published by Taylor et al. (1980).

During the present study, the following plutonic rock types were identified (nomenclature after Streckeisen (1976), reproduced in Appendix 1).

- 1) Pink medium grained granodiorite with prominent (5-10 mm) rounded quartz (hereinafter called the "Feeder Granodiorite").
- 2) Fine grained gabbro and related hybrid granite
- 3) Pink to reddish orange to brick red, fine to medium grained, equigranular, biotite-bearing alkali-feldspar granite to alaskite (the "biotite granite" of Taylor et al., 1980)
- 4) Aegirine and/or alkali-amphibole-bearing peralkaline




Plate 2-1: Feeder Granodiorite with rounded to subhedral, cracked quartz, plagioclase (white) and altered biotite.

granite and related quartz-feldspar porphyry.

In addition, Taylor et al. (1980) have identified a metaluminous biotite-hastingsite granite which is cut by dykes of peralkaline microgranite.

Bell and Blenkinsop (in press) have established whole rock Rb/Sr isochrons as follows:

- a) Feeder Granodiorite, 480 ± 80 Ma, initial $^{87}\text{Sr}/^{86}\text{Sr} = 0.754$
- b) Peralkaline Granite, 419 ± 5 Ma, initial $^{87}\text{Sr}/^{86}\text{Sr} = 0.707$
- c) Alkali-feldspar granite, 386 ± 9 Ma, initial $^{87}\text{Sr}/^{86}\text{Sr} = 0.7067$

The Feeder Granodiorite is restricted to two small parts of the study area (Fig. 2-2). The distribution of these occurrences around the margins of the Topsails Granite suggests that the Feeder Granodiorite was once more extensive but is now largely obliterated by later granite intrusions.

The Feeder Granodiorite is distinctly more calcic than later granites and contains strongly zoned plagioclase generally in excess of two thirds of feldspars (Plate 2-1). Perhaps the most characteristic feature of these rocks is the presence of prominent (up to 1.5 cm) quartz phenocrysts generally equant and rounded or slightly



Plate 2-2: Core samples, right to left; Feeder granodiorite-like boulder from pyroclastic breccia, rhyolite, dacitic crystal-vitric tuff, volcanic conglomerate. All samples are characterized by large, rounded to subhedral, cracked quartz and are considered to be facies equivalents.

resorbed but locally with sharply defined hexagonal crystal faces (Plate 2-1). The quartz crystals commonly have poor to well developed irregular to radial cracks. The equant nature of the quartz crystals is indicative of early quartz crystallization under low pressure (i.e., high crustal level) (Tuttle and Bowen, 1958). The locally preserved hexagonal crystal outlines are characteristic of high temperature beta quartz and the radial to irregular cracks in quartz may be due to density increase and consequent volumetric decrease during transformation from beta to alpha quartz. The complex zoning of plagioclase crystals is indicative of instability of pressure and temperature in the magma chamber, a characteristic most likely to be developed at the margins of a high level intrusive body.

Textural and geochemical comparison of the Feeder Granodiorite and certain rhyolitic volcanics of the Buchans Group lead the author to conclude that these lithologies are genetically related. Massive pink to grey rhyolite flows of the Prominent Quartz Sequence contain large zoned plagioclase phenocrysts and texturally distinctive prominent (5-10 mm) rounded to hexagonal quartz phenocrysts with radial to irregular cracks (Plate 2-2). Both macroscopically and microscopically, the phenocryst mineralogy and distinctive morphology (especially of quartz) are identical in these flows to those present in the Feeder Granodiorite.




Plate 2-3: Typical alkali-feldspar granite with quartz, albite and K-feldspar. Albite is much less clouded by minute hematite than K-feldspar.

This striking textural similarity, geochemical similarities presented in Chapter Five, the occurrence of cobbles and boulders of Feeder Granodiorite-like material in the Buchans Group and the similarity of radiometric ages of the Feeder Granodiorite and Buchans Group suggest that magmas of the Feeder Granodiorite fed certain rhyolitic volcanics of the Buchans Group.

The Topsails Granite can be subdivided into peralkaline and non-peralkaline suites. The non-peralkaline suite is further divisible into the alkali-feldspar granite with associated mafic rocks and the biotite-hastingsite granite.

The alkali-feldspar granite (terminology after Streckeisen, 1973) is composed of approximately equal quantities of quartz, albite and K-feldspar with lesser altered biotite and minor hornblende. The granite is characterized by moderate to strong hematitization of feldspars imparting a deep pink to brick-red appearance in hand specimen. In all cases, K-feldspar is distinctly more clouded by minute hematite than is albite, facilitating distinction of these two minerals (Plate 2-3). Fluorite, apatite, zircon, allanite and magnetite are common accessories. Quartz and biotite contents vary and the granite locally approaches syenite and elsewhere, alaskite.



Plate 2-4: Peralkaline granite containing quartz,
perthite and black alkali amphibole.

Massive fine grained mildly alkaline gabbroic to dioritic rocks are intimately associated with the alkali-feldspar granite within the area of study. These mafic rocks are areally much more restricted than associated granites but regional magnetic and gravity data suggest that they are volumetrically more important than their surface exposure would indicate. Gabbroic bodies are surrounded and intruded by later granite except at one locality in the northwest of the area where gabbro is faulted against rhyolite of the Upper Buchans Subgroup.

The biotite-hastingsite granite crops out east of Hinds Lake and consists of K-feldspar and quartz with lesser plagioclase, hastingsite and biotite. Taylor et al. (1980) consider this granite to be comagmatic with the biotite-bearing alkali-feldspar granite.

Peralkaline granitic rocks crop out in the north-eastern portion of the area in close association with volcanic roof pendants in the granite. These are medium to coarse grained, equigranular and weather in various shades of red, reddish yellow, light purple and white. They are characterized by alkali ferromagnesian minerals (aegirine, riebeckitic-arfvedsonite and aenigmatite), perthite and quartz (Plate 2-4) i.e., are hypersolvus granites. At the contact with the volcanic roof pendants red phenocryst-rich dykes, mineralogically similar to

the peralkaline granite, with aphanitic to granophyric groundmass, cut the volcanics.

Considerable evidence exists to suggest that all granitoid rocks were intruded into, and crystallized at a high level in the crust. Mirolitic cavities are common in the alkali-feldspar granite and may be filled with fluorite, siderite and a green phyllosilicate (lepidolite?). The hypothesis that the Feeder Granodiorite feeds rhyolites of the Prominent Quartz Sequence and subsequently intrudes its own ejecta, suggests a high level of intrusion. A similar relationship between the peralkaline granite and the volcanic roof pendants in the granite is suggested in the next section. The presence of roof pendants and wide chill zones in the granites with the development of strongly porphyritic granites is further evidence of a high level of intrusion. The low grade metamorphism of the host rocks indicates that they were never buried to great depth and thus granites must have intruded into a high level. The brittle behavior of the host rocks at granite contacts and the lack of metasomatism of the host rocks at granite contacts are further evidence of a high level of intrusion. Within both the Lower and Upper Buchans Subgroups there exist boulders of the Feeder Granodiorite in density flow deposits, further suggesting a high level of intrusion.

Although no contacts are exposed between the various types of granitoids, or granitoids and the Buchans Group, a sequence of igneous intrusion can be inferred from relationships previously presented and from apophyses which maintain the characteristic mineralogy, chemistry and, in some cases, texture of their parent rocks. The Feeder Granodiorite feeds and subsequently intrudes volcanic rocks of the Lower Buchans Subgroup and is evidently the earliest exposed plutonic body in the Buchans area. It will be shown in a later section that intrusion of the Feeder Granodiorite is separated in time from later igneous activity by a period of major thrust faulting.

Relationships between the various later granitoid suites are not yet fully resolved though it is evident that all post-date the Buchans Group. Gabbroic rocks feed numerous diabase dykes which cut all portions of the Buchans Group and the Feeder Granodiorite. These dykes are petrographically and geochemically identical to the gabbros. The alkali-feldspar granite feeds porphyritic red "rhyolite" dykes which cut the Buchans Group and again, the mineralogical and geochemical character of the dykes is similar to the source plutonic body.

Dykes of alkali-feldspar granite also cut the gabbroic bodies. The contacts between the two are, in some cases, sharp but generally a diffuse zone of "hybrid

granite" is developed. Thin dykes intruding gabbro have thin hybrid zones at the contact whereas the contact between large gabbro and granite bodies is commonly marked by "hybrid" zones on the order of hundreds of meters in width. Within the Buchans Group, dykes of both "rhyolite" cutting diabase and vice versa can be observed. This suggests a temporal link between magmas of gabbroic and granitic composition. Furthermore, the gabbros exhibit mildly alkaline characteristics (see Section 5.2.) suggesting a genetic link between the gabbros and the alkali-feldspar granite.

Fine grained, spherulitic to flow-banded dykes with alkali ferromagnesian minerals, presumably apophyses from the peralkaline granite, cut the Buchans Group. Taylor et al. (1980) observe that dykes of peralkaline granite cut the hastingsite-biotite granite, in apparent conflict with the reported isotopic ages. They suggest that both criteria may be correct and that oscillatory peralkaline-peraluminous magmatism may have been operative.

2.5. Volcanic Outliers Within the Topsails Granite (Springdale Group?)

Volcanic rocks within the Topsails Granite (Fig. 2-1) were briefly examined in order to compare their gross lithologic character with volcanic rocks of the Buchans

Group. Little similarity was observed between these groups of volcanic rocks. They consist of reddish sub-aerial ignimbrite, non-welded pumice flows, breccias, pyroclastics and flow banded rhyolite, intruded by porphyritic dykes emanating from the peralkaline granite. Gently dipping ignimbrites overlying the peralkaline granite contain alkali ferromagnesian minerals suggesting an intrusive-extrusive relationship between the granite and volcanics. Non-peralkaline volcanics are reported by Taylor et al. (1980). The relationship between these and the peralkaline volcanics and granites is not clear.

Vertically cleaved, gently dipping ignimbrite on Barren Mountain occupies an area near the core of a synclinal trough and is probably an outlier of a once more extensive subaerial felsic volcanic sequence. The lithologic similarity to volcanic rocks of the Springdale Group as described by Kalliokowski (1955), Neale and Nash (1963) and Williams (1967) suggests that parts of the volcanic rocks associated with the Topsails granite are related to the Springdale Group. The major synclinal axis in the Springdale Group as mapped by Kalliokowski (1955), can be traced southward through the Topsails Granite, through the Barren Mountain outlier and into the Buchans Group.

2.6. Hungry Mountain Complex

Two belts (Western and Eastern) of deformed plutonic rocks exist approximately 5 km north and 15 km east-northeast of Buchans and occupy an area between the Topsails Granite and the Buchans Group (Figs. 2-1, 2-2). These rocks have been called the Hungry Mountain Complex (Thurlow, 1975) after a prominent hill in the area. The author considers the Western and Eastern Belts to be lateral equivalents of a once larger sheet which has subsequently been dissected into two parts by intrusion of the alkali-feldspar facies of the Topsails Granite. Mapping by Riley (1957), Anderson (1972) and Kean (1979) indicates that rocks correlative with the Western Belt of this Complex extend from the northern boundary of the present study area in a northwesterly direction along the west shore of Hinds Lake (Figs. 2-1, 2-2). Mapping by ASARCO exploration crews in the later 1940's and scattered observations by the present author indicate that the Eastern Belt extends northward at least to a point several kilometers east of Quarry.

Bell and Blenkinsop (1980) have established two Rb-Sr whole-rock isochrons of 660 ± 70 Ma and 400 ± 60 Ma from separate localities in the Western Belt.

Rocks of the Hungry Mountain Complex range in composition from gabbro to granite and are characterized

by a complex deformational history. Within the Western Belt, foliated rocks of granodioritic composition are most abundantly exposed. These are light grey to white weathering and consist of quartz, zoned plagioclase (with albite-oligoclase rims) and aligned biotite with minor K-feldspar and hornblende. Dykes of foliated granodiorite locally cut bodies of deformed diorite and hornblende gabbro. Elsewhere the foliated granodiorite contains slightly to moderately flattened, angular to rounded, resorbed hornblende quartz diorite xenoliths with diffuse margins. At one locality a hornblende-plagioclase inclusion (similar to the larger mafic portions of the Complex) contains a fabric which is not parallel to the fabric in the enclosing host rock. This is taken as evidence that some of the more mafic portions of the Complex have undergone an earlier deformational history before intrusion of the granodioritic phase.

The Western Belt of the Hungry Mountain Complex, though generally similar to the Eastern Belt, contains a greater abundance of massive medium to coarse grained gabbro. The degree of deformation is more variable ranging from locally intense to essentially undeformed. Deformed granodioritic dykes locally cut pre-deformed gabbro but in at least one locality, the opposite sequence is observed.



Plate 2-5: Hornblende gabbro breccia, intrusive into
the Hungry Mountain Complex.

The deformational fabric within the Complex is typically defined by abundant flattened quartz and aligned biotite (or chlorite after biotite) which form augen of altered plagioclase.

In more mafic phases, tabular hornblende defines the foliation. At several localities this main foliation is openly folded with interlimb angles of approximately 120 degrees. No cleavage is associated with this later folding.

A steeply dipping spaced fracture cleavage of probable Acadian age is visible locally and overprints the main foliation within the Complex.

The Hungry Mountain Complex is cut by a number of intrusives that post-date the main fabric. The earliest of these is an intrusive breccia composed of coarse hornblende gabbro in a light feldspathic matrix (Plate 2-5). Amphibole-rich portions consist of coarse hornblende (over 1 cm) with less abundant clinopyroxene, orthopyroxene and interstitial highly altered feldspars. Deuteric alteration, especially of orthopyroxene, to tremolite is common. Macroscopically a strong textural similarity to the Brighton Gabbro (Hussey, 1974) is noted.

The hornblende gabbro and the entire Complex are cut by grey to red aplite and "rhyolite" dykes emanating



Plate 2-6: Flow banded peralkaline dyke (top) cutting deformed Hungry Mountain Complex. Small circled xenolith of host rock occurs in dyke below lens cap.

from the alkali-feldspar granite, dark brownish-red, aegerine-bearing dykes from the peralkaline granite and less abundant diabase dykes related to the gabbro.

Chilled contacts against the foliated rocks of the Complex are abundantly exposed and the foliation in the latter is truncated by dyke intrusion (Plate 2-6).

Locally, angular fragments of foliated granodiorite are included in the dykes.

The contact between the Hungry Mountain Complex and the main body of the Topsails Granite is not exposed on surface but probably exists in DDH 2856 (location on Fig. 1-2). The upper 153 m of this drill hole consists of typical foliated granodiorite cut by abundant (greater than 1/3 of the volume) red "rhyolite" dykes of the alkali-feldspar Topsails Granite. At 153 m depth the foliated granodiorite is cut by a thicker body of "rhyolite" with a wide chilled margin. Near the contact, the matrix is aphanitic but this coarsens with depth until at 296 m depth (end of hole) the "rhyolite" has graded into a fine grained, slightly porphyritic granite with matrix crystals of 1-2 mm size. This is considered to be a portion of the main body of the Topsails Granite intruding the Hungry Mountain Complex. The width of the chilled margin (in excess of 140 m) indicates that the Hungry Mountain Complex was relatively "cold" at the time of the Topsails Granite intrusion.



Plate 2-7: Mylonite from the base of the Hungry Mountain Complex.

2.7. Contact Relationships Between the Hungry Mountain Complex and the Buchans Group

2.7.1. Relationship Within the Hungry Mountain Complex

The contact between the Hungry Mountain Complex and the Buchans Group is neither exposed in outcrop nor in drill core but several areas of outcrop near the contact are present. Within a few hundred meters of the contact, foliated granitoid rocks of the Complex become noticeably more deformed and acquire a protomylonitic texture. This passes gradationally towards the contact into mylonitic fabric characterized by extreme stretching of quartz resulting in a strong lineation (Plate 2-7). Quartz exhibits sutured grain boundaries, the development of sub-grain boundaries and commonly forms augen with chlorite around more resistant plagioclase. Less commonly, a planar fabric defined by flattened quartz is developed. North of Buchans the lineation trends north-northwest and plunges gently at 10 to 20 degrees. The strike of the foliation, where present, parallels a topographic break which is interpreted as the contact with the Buchans Group.

At one locality small folds are generated by the transposition of the main foliation in the Complex by the mylonitic fabric. These folds are defined by flattened quartz and biotite passing around the hinges of

isoclinal folds with attenuated limbs. They indicate that the main foliation in the Complex pre-dates the mylonitic fabric. This is further substantiated by the presence of sub-vertical foliations in the Complex which are highly discordant to the mylonite zone within several hundred meters of the mylonite.

2.7.2. Relationships Within the Buchans Group

Rocks of the Buchans Group near the contact with the Hungry Mountain Complex are exposed in the bed of Harry's River and form sporadic outcrops for a distance of about 2 km from the contact (Fig. 2-2). Within 650 m of the contact these consist of a sequence of dark green to black fine grained basaltic and andesitic tuffs. Stratigraphically below this is a sequence of subaqueous pyroclastic flows and related arkose of the Prominent Quartz Sequence (see Section 3.2.). These units strike northeast and dip to the northwest at 45 degrees.

Within 650 m of the contact with the Complex the mafic tuffs of the Buchans Group have undergone low greenschist facies metamorphism. At distances greater than 650 m from the contact, the Buchans Group exhibits its typical prehnite-pumpellyite facies of metamorphism. Moving towards the contact, chlorite decreases at the expense of actinolite and a penetrative mineral fabric at a slight angle to bedding (approximately 25°) becomes



Plate 2-8: Shallowly dipping penetrative foliation in felsic tuffs of Upper Buchans Subgroup, below Hungry Mountain thrust. Later, steeply dipping, spaced fracture cleavage is typical of all Buchans area rocks.

visible within these greenschist facies rocks. Small recumbent folds with axial planes parallel to the foliation are developed. Within 50 m of the inferred contact, hornblende appears and increases at the expense of actinolite. Within 20 m of the contact, the mineral assemblage consists of quartz, blue-green hornblende in excess of actinolite, epidote, altered plagioclase (of unknown composition) and minor chlorite and sphene, an assemblage typical of the epidote-amphibolite facies of Miyashiro (1973).

Basaltic to rhyolitic rocks of the Upper Buchans Subgroup occupy a large area to the northeast of Buchans (Fig. 2-2). These rocks contain a subhorizontal to gently northward dipping penetrative fabric (Plate 2-8) which increases gradually in intensity towards the contact with the Complex. This foliation is visible throughout a large area of the Upper Buchans Subgroup for distances up to 6 km from the contact with the Hungry Mountain Complex. This foliation is cut by a vertical spaced fracture cleavage, typical of Acadian deformation throughout the Buchans Group (Section 3.3). In contrast to this gradual lateral change in intensity of fabric, a much more rapid change in intensity of fabric and metamorphic grade is observed in vertical drill holes penetrating this sequence. For instance, DDH 2837 (location on Fig. 1-2) penetrates over 1.2 km of mafic to felsic flows, pyroclastics and arkosic sediments. From the bottom of the hole to

within 150 m of surface, the sequence is characterized by the presence of prehnite-pumpellyite-chlorite ^{or} vesicles in basalts and matrix prehnite in acid pyroclastics and arkosic sediments. No foliation is present in this sequence. However, within 150 m of surface, an initially weak but upward increasing foliation at a slight angle to bedding becomes evident. Prehnite and pumpellyite disappear and a quartz-albite-epidote-chlorite-calcite assemblage is present. The disappearance of pumpellyite above 150 m in this drill hole and the presence of typical low grade greenschist assemblage is taken to indicate that the prehnite-pumpellyite: greenschist isograd exists, with a sub-horizontal attitude, slightly below 150 m of the present erosional surface.

In view of the evidence presented (i.e., the mylonite zone in the Hungry Mountain Complex at the contact with the Buchans Group, the intensifying foliation at a low angle to bedding near the contact with the Complex and the stratigraphically inverted metamorphic zonation in the Buchans Group from sub-greenschist to epidote amphibolite facies, it is suggested that the Hungry Mountain Complex was thrust upon the Buchans Group. The thrust plane is at a moderate angle to the present erosional surface in the Harry's River area but at a very low angle in the Upper Buchans Subgroup to the east. By comparison with the width of the aureole in the Harry's

River area, it is suggested that the thrust plane existed approximately 500 m above the present erosional surface in the area of DDH 2837, i.e., had an average northward dip of approximately 10° in this area (Fig. 2-3). These variations in the present attitude of the thrust plane are attributed to a post-thrusting open Acadian folding (see Section 3.3.).

The time of thrusting can be fairly accurately determined from the relationships previously discussed. Thrusting occurred after deposition of the Buchans Group. Unfoliated "rhyolite" dykes akin to the alkali-feldspar Topsails Granite cut the mylonite zone in the Complex, indicating a pre-Topsails Granite age of thrusting. The wide chill zone in the Topsails Granite against the Complex (DDH 2856) provides further evidence that the Complex was relatively "cold" at the time of Topsails Granite intrusion. Radiometric age dates (Bell and Blenkinsop, 1980) concur with the geological data and suggest a Silurian age of thrusting.

The gently plunging north-northwest trending lineation both in the mylonite zone and in the Buchans Group is interpreted as stretching in the direction of thrusting. The thrusting direction was from north-northwest to south-southeast as indicated by facing directions of small recumbent folds in the aureole beneath




Figure 2-3. Schematic cross section showing relationships between the Hungry Mountain Complex, Topsails Granite and Buchans Group.

the thrust in the Buchans Group.

Due to the lack of systematic study of the Hungry Mountain Complex, its geological significance and pre-thrust disposition are obscure. It is evidently a package of rocks very foreign to the volcanic and plutonic evolution of the Buchans area. The remarkable similarity of gabbros in the Complex to gabbros of the Annieopsquotch igneous complex (recently interpreted as ophiolitic by Dunning and Herd, 1980), and the inversely zoned metamorphic aureole beneath the Complex, suggests that the Hungry Mountain Complex might represent part of a dismembered ophiolite suite.

2.8. Siluro-Devonian Deformation

The Buchans area has been subjected to two major structural events: 1) the pre-Topsails Granite Silurian episode of thrusting, discussed earlier, and 2) a period of Devonian broad open folding. The latter event imparting a weak northeast-striking subvertical fracture cleavage to all existing lithologies and is axial planar to a broad open syncline within the Buchans Group. The intensity of this cleavage varies considerably and is directly related to the competency of the affected rocks. In many exposures, the cleavage is not visible but it can be identified in several parts of the Buchans Group, Hungry Mountain Complex, Topsails Granite and the subaerial



Plate 2-9: Shallowly dipping red sandstone and conglomerate of presumed Carboniferous age. Upright hammer is approximately one meter length.

volcanic roof pendants on the Granite. The deformation related to this cleavage is nowhere sufficiently intense to cause visible effects of strain. The cleavage does not affect dated Carboniferous strata and is thus post-granite and pre-Carboniferous in age, i.e., Devonian.

Folds related to this deformation are broad, open and of regional extent. The axis of one major syncline within the Buchans Group can be traced through flat lying and vertically cleaved ignimbrite within the Topsails Granite and subsequently into a major syncline within the Springdale Group, a distance of approximately 100 km.

2.9. Carboniferous

Rocks of presumed Carboniferous age crop out in several small areas along Red Indian Lake and in a number of drill cores along the contact between the Footwall Arkose and the Upper Buchans Subgroup in the center of the map area. Similar rocks are reported to have been present on Buchans Island before construction of the Exploits Dam (E.A. Swanson, pers. comm. 1975).

These strata consist of flat lying to gently-dipping, poorly indurated, polymictic red conglomerate and red crossbedded sandstone (Plate 2-9). Some boulders in the conglomerate are very similar to parts of the Buchans Group, others can be confidently related to the Topsails

Granite. Similar conglomerates and sandstone are present in the banks of the Shanadithit River and underlie thin coal seams and limestone (D. Barbour, pers. comm. 1975). Plant remains collected from the latter area are of Carboniferous age (Newhouse, 1931) and miospores (Belt, 1969) reveal an assemblage typical of the Mississippian Horton Group. Palynological analysis of a single silty mudstone sample from the present study area revealed the presence of vegetal matter but no diagnostic age was determinable (J. Utting, pers. comm. 1975).

Although the contact is not exposed, there is little doubt that these Carboniferous rocks unconformably overlie the Buchans Group. Steeply dipping, deformed rocks of the Buchans Group occur within meters of gently dipping poorly indurated Carboniferous strata at several localities. Relationships suggest angular unconformity of nearly 90 degrees in these areas. The poorly indurated nature and lack of cleavage indicate that these Carboniferous rocks have not undergone Acadian or subsequent deformation. This lack of deformation provides the best evidence that the upright fracture cleavage and associated folding of the Buchans Group, Hungry Mountain Complex, Topsails Granite and Springdale Group is indeed of Acadian age.

The occurrence of pods of Carboniferous strata along the contact between the Footwall Arkose and the Little Sandy Formation is used as evidence for a pre-

Carboniferous fault contact between these lithologies.

2.10. Summary of Chapter Two

The geological history of the Buchans area and relationships between various rock units can be summarized as follows:

Middle to Late Ordovician
(Caradocian and/or later)

Deposition of greywacke and conglomerate on southeast side of Red Indian Lake

Latest Ordovician or
earliest Silurian

Extrusion of Buchans Group - Roberts Arm Group volcanics with cogentic intrusion of Feeder Granodiorite; deposition of massive sulphide ores

Silurian

Southeasterly directed thrust faulting of pre-deformed Hungry Mountain Complex upon the Buchans Group

Late Silurian to early
Devonian

Intrusion of the various phases of the Topsails Granite with extrusion of cogenetic subaerial volcanics

Late Devonian

Open folding, uplift, erosion

Early Mississippian

Formation of the Red Indian
Lake basin and deposition of
red beds

CHAPTER THREE

GEOLOGY OF THE BUCHANS GROUP

3.1. Introduction

Within this chapter an integrated account of the geology of the Buchans Group is presented. A consequence of the author's earlier work (Thurlow, 1973) was the realization that further geochemical data could be interpreted properly only if a detailed understanding of the stratigraphy, structure and modes of deposition of the Buchans Group could be gained. Within the scope of this thesis it is not possible to document all the complexities of Buchans geology and this account, although lengthy, is considered by the author to represent only a surficial summary. At the time of writing, several major structural-stratigraphic difficulties still exist and will be solved only by further detailed work with existing data and new data from ongoing diamond drilling.

The concept of volcanic cycles has been modified and the Buchans Group is subdivided into Lower and Upper Subgroups, each consisting of several sequences. The Lower Buchans Subgroup is a "volcanic cycle" consisting of voluminous basaltic volcanics overlain by arkosic rocks followed by intermediate and felsic volcanics which contain the major ore deposits. The Upper Buchans Subgroup

comprises a second "volcanic cycle" initiated with relatively minor mafic volcanism and arkose deposition overlain by voluminous, dominantly felsic volcanics. In this sense, the entire Buchans Group is a single dominantly mafic to dominantly felsic sequence composed of lesser internal increasingly felsic volcanic cycles. In detail, the nature of these cycles is different from typical Archean cycles (e.g., Goodwin, 1968) and volcanic cycles of the Horkuroko district of Japan (as described by Takahashi and Suga, p. 111-112).

The geology of the Buchans Group is extremely complex. The understanding of the stratigraphy is hampered by lack of outcrop, rapid lateral facies changes, a lack of marker beds, local subsidence, uplift, contemporaneous block faulting and diabase intrusion. Although metamorphism in the area is relatively mild, interpretation of the structure (and thus stratigraphy) is consequently hampered by a lack of bedding-cleavage relationships. Perhaps the major complicating factor is the postulated presence of major thrust faults. Without fossil control it is impossible to prove the existence of these faults. But the possibility of their presence demands the building of a stratigraphic sequence based on observed indisputably conformable contacts. Such contacts are not common as most are characterized by late diabase sills or more commonly by "innocent-looking" shears, the magnitude of

which is not possible to judge. The stratigraphy presented in the following sections is based upon structurally conformable contacts which allow no possibility of alternate interpretation. This stratigraphy is schematically shown in Fig. 3-1 and summarized in Table 3-1. The stratigraphy of the Buchans Group has not been formalized into formations, members, etc. because of the uncertainties attached to the effects of thrust faulting.

3.2. Stratigraphy of the Buchans Group

3.2.1. Stratigraphy of the Lower Buchans Subgroup

3.2.1.1. Footwall Basalt: The Footwall

Basalt is the lowermost unit of the Buchans Group and is thought to conformably overlie a greywacke-conglomerate sequence which bears lithologic similarities to the Sanson Greywacke and Goldson Conglomerate of Notre Dame Bay (Anderson, 1972a). Lithologically and stratigraphically the Basalt is considered by most workers to be correlative with basalts of the Roberts Arm Group to the northeast. The Footwall Basalt is continuous throughout the study area and crops out sporadically on hills, ridges, river beds and along the north shore of Red Indian Lake, dips northward and faces consistently northward. Basaltic rocks thought to form part of the Footwall Basalt are the host rocks for the Skidder Prospect (see next Section).




FIGURE 3-1: Schematic post-thrusting stratigraphy of the Buchans Group.

Table 3-1 Stratigraphy of the Buchans Group

Unit	Lithologies	Thick-ness	Mineralisation
Upper Buchans Subgroup	Basalt to felsic pyroclastics and breccia with lesser flows, minor volcanic sediment. Includes a basal member (Upper Arkose) of reworked conglomerate.	2000 m (?)	Little Sundry Prospect (Cu) and several areas of weakly to heavily disseminated pyrite, trace chalcopyrite.
Lower Buchans Subgroup			
Oriental Ore Member Sequence	Dacitic tuff, rhyolite breccia dikes, pyritic siltstone and wacke, pyroclastic breccia, polytuff volcanic breccia-conglomerate, "granite conglomerate".	400 m	High grade massive sulphide orebodies; Oriental #1 and #2, Old Buchans Conglomerate and possibly Old Buchans East and West. Sandfill and Middle Branch Prospects.
Oriental Intermediate	Pelagic felsic pyroclastic breccias, local altered intermediate and mafic flows, pyroclastics and breccia.	300 m	Minor ore grade stockwork mineralisation with larger areas of subeconomic epigenetic Pb-Zn-Cu.
Old Mill Sequence	Mainly andesitic pyroclastics and breccias; minor basaltic flows, dacitic flows.	1000 m	Local isolated block ore clasts.
Proterozoic Quartz Sequence	Mainly dacitic pyroclastics, pyroclastic breccias related to tuffaceous sediments, lesser rhyolite. Interbedded basaltic and andesitic units, Jasper chert.	2000 m	Local disseminated pyrite, Pb-Zn and trace Pb-Zn veining.
Basalt			
Lower Green Basalt	Basaltic pillow lava, pillow breccia and lesser pyroclastics.	350 m	Barren
Lucky Strike Ore Member Sequence	Dacitic tuff, massive rhyolite, pyritic siltstone and wacke, pyroclastic breccia, polytuff volcanic breccia-conglomerate, "granite conglomerate", local basaltic and andesitic horizons.	600 m	High grade massive sulphide orebodies; Lucky Strike, North, Two Level, Buchanan #1 and #2, Middle Branch Prospect. Subeconomic veins and isolated occurrences of sulphide clasts.
Intermediate Potash	Completely interbedded and altered mafic to felsic flows, pyroclastics and breccias, related tuffaceous pyritic siltstone and wacke.	250 m	Localised ore grade pyritic stockwork Zn-Pb-Cu mineralisation. Small, stratiform, high grade, polymetallic massive sulphides. Larger areas of subeconomic disseminated and veinlet mineralisation.
Potash Arkose and Willey's Sequence	Mainly lithic arkose; lesser arkose conglomerate, siliceous greywacke; minor siltstone, sandstone, chert. Facies equivalent Willey's Sequence is dominantly felsic pyroclastics with lesser flows and tuffaceous sediments.	2500 m	Overall pattern: very small bedded pyrite Pb-Zn occurrences; otherwise barren.
Potash Basalt	Basaltic pillow lava, pillow breccia, lesser pyroclastics, massive flows, interbedded chert. Very minor felsic pyroclastics.	2000 m (?)	Shisher Brook pyritic massive sulphide deposit (Cu-Zn).

POOR COPY
COPIE DE QUALITEE INFERIEURE



Plate 3-1: Pillow lava of the Footwall Basalt with poorly developed radial cracks and amygdule-rich bands concentrated along pillow top.

The base of the Footwall Basalt has not yet been observed north of Red Indian Lake and thus the thickness is not known with confidence. Thickness estimates are also complicated by paleotopographic irregularities on top of the Basalt and by an essentially unknown component of the structural repetition by thrust faulting. Despite these complications, the exposed thickness of the Footwall Basalt probably exceeds 2000 m within the study area. Outside the study area it may attain a total thickness of 8300 m, barring unknown repetition by folding or faulting.

The Footwall Basalt consists dominantly of basaltic pillow lavas and related pillow breccias. Pillows range from spherical to elongate and are normally from 1.0 to 1.5 m in diameter. Amygdules are common and are normally filled with calcite although various combinations of prehnite, pumpellyite, silica, chlorite and epidote also occur. Large calcite and smaller chlorite-filled amygdules commonly coexist in the same pillow. In some areas, amygdules are concentrically arranged only along the tops of pillows (Plate 3-1) and may facilitate "tops" determinations (similar pillows are described by Jones, 1969). Amygdules are normally spherical and 5 mm in diameter although in some localities abundant giant amygdules (over 3 cm) are present. If amygdale proportion and size are related to water depth as suggested by Jones (1969), then shallow water is indicated in these cases.



Plate 3-2: Nearly whole pillows in pillow breccia,
Footwall Basalt. Note concentric amygd-
ule-rich bands near pillow margins.



Plate 3-3: Highly disaggregated pillow breccia,
Footwall Basalt.



Plate 3-4: Graded pillow breccia, Footwall Basalt.

The pillow lavas are normally feldsparphyric with laths less than 2 mm long and of andesine-labradorite composition. Colourless to light brown unaltered augite phenocrysts are locally present and may attain sizes up to one centimeter, although those of 1-2 mm are most common. The groundmass consists of a fine grained intergrowth of plagioclase, clinopyroxene and much less abundant magnetite. Matrix plagioclase is variably altered to various combinations of epidote, calcite, clay minerals and sericite and is locally replaced by pumpellyite. Clinopyroxene may be altered to epidote and chlorite.

Outcrops of broken-pillow breccia (terminology after Carlisle, 1963) are, in general, poorly exposed but textural variations can be seen best in numerous polished boulders that occur in river beds cutting the basalts. Pillows occur in various stages of disruption from near whole pillows (Plate 3-2) to highly disaggregated varieties (Plate 3-3) which are locally size-graded (Plate 3-4). Isolated-pillow breccia is volumetrically much less significant than broken-pillow breccia.

Pillow lavas are commonly interbedded with chert lenses and locally contain abundant interpillow chert. Cherts occur in a variety of colours ranging from black to dark grey to green although bright red jasper chert predominates in some areas. Chert beds are normally



Plate 3-5: Spheroidal banded rhyolite within Footwall
Basalt unit.

less than one meter thick although a major exception to this generalization occurs in the Skidder Brook area. Here, in a small area of limited strike length, there occurs a sequence composed mainly of jasper chert in excess of 125 m thick. These are interbedded with less abundant green chert, red cherty mudstone, grey siltstone, greywacke and lithic arkose and a few massive basaltic flows. The Skidder Cu-Zn prospect (see next Section) is associated with similar, but thinner, jasper chert several hundred meters stratigraphically above the chert sequence at Skidder Brook.

A number of volumetrically less significant lithologies occur throughout the Footwall Basalt. These include pyroclastics (tuff, lapilli-tuff, lapillistone, agglomerate), tuff-breccias, hyaloclastic breccias, explosion breccias, blocky flow breccias and massive flows. Fine grained pyroclastics locally grade imperceptibly into greywacke and black fine grained bedded volcanic siltstone of basaltic composition.

Volcanic rocks more felsic than basalt occur at a small number of isolated localities in the Footwall Basalt. A few bodies of massive aphyric to feldsparphyric dark grey rhyolite occur northeast of Skidder Brook. These are characterized by a peculiar well developed spheroidal structure (Plate 3-5) which consists of thin

concentric colour banding which defines distinct spheroids of approximately 20 cm diameter. The banding becomes less distinct and eventually disappears both towards the center of individual spheroids and towards the massive rhyolite matrix between spheroids. The origin of this texture is not known but is suspected to be a cooling-related phenomenon.

Other types of felsic volcanic rocks within the Footwall Basalt include small lenses of strongly hematitized rhyolite breccia and thin beds of dacitic tuff. Most of these are of too small scale to be represented on the geologic map (Fig. 2-2).

Alteration of basaltic rocks of the Footwall Basalt is patchy in distribution and shows no recognizable pattern. Pervasive hematitization is present in some areas and may, or may not be combined with abundant vein and amygdale calcite. Such alteration occurred at, or shortly after, deposition as interbedded sedimentary units show no effects of alteration and many non-hematitized breccias contain strongly hematitized fragments. Epidotization is locally intense but usually occurs independently of hematitization and calcite introduction. Various combinations of silicification, pyritization and chloritization are found in some areas but are generally weak and of limited extent. Traces of

disseminated pyrite occur throughout the Footwall Basalt.

In the mine area, the distribution of the Footwall Basalt undergoes some changes that are not evident elsewhere as an anomalous inlier of the Basalt exists in the immediate Buchans area (Fig. 2-2). The rocks of this Buchans Inlier are known (through diamond drilling) to extend entirely beneath a shallow cover of later rocks (the Prominent Quartz Sequence) and to be contiguous with rocks which comprise the main portion of the Footwall Basalt south of Buchans (Fig. 2-2). The basaltic rocks of the Inlier strike east-northeast, dip consistently to the north-northwest at 30 to 50 degrees and face north-northwest, much like their counterparts to the south. Although previously thought to be an anticlinal structure (George, 1937; Relly, 1960; Swanson and Brown, 1962), these facts dictate that folding cannot be the sole cause of the presence of the Inlier.

With a few modifications, the lithologies of the Buchans Inlier are similar to those of the rest of the Footwall Basalt. Broken-pillow breccia is the most common rock type. Pillow fragments, some with chilled rims, range in size from 0.5 m down to tuff size and are normally separated by a few centimeters of fine grained matrix. Elongate fragments lie parallel to bedding with no tendency toward imbrication (cf. Carlisle,

1963, p. 64). Many fragments are highly amygdaloidal (over 50%), some so much so that they initially had densities close to that of water or may even have floated. Broken-pillow breccia horizons are locally intercalated with thin (up to a few meters) aquagene tuff beds or more rarely with isolated-pillow breccia. Although not exposed on surface, thin beds of arkose, chert and red mudstone are interbedded with basalts of the Buchans Inlier.

3.2.1.1.1. Skidder Prospect: The Skidder Cu-Zn Prospect, though not within the immediate study area, is briefly mentioned here as it characterizes the type of mineralization found at the base of the Buchans Group. It occurs as at least two massive stratiform sulphide lenses conformable within basaltic rocks of the Footwall Basalt in close association with jasper chert. Mineralogy consists mainly of pyrite with minor chalcopyrite and sphalerite and rare galena. A zone of strong silicification and disseminated mineralization exists in the stratigraphic footwall of the deposit with less intense alteration in the hangingwall.

3.2.1.2. Footwall Arkose and Wiley's Sequence: The Footwall Arkose, and the presumed time-equivalent Wiley's Sequence, form a major regional volcanic and sedimentary horizon which conformably overlies the Footwall Basalt. The term "Arkose" is derived from the abundantly

exposed pinkish, fine to coarse grained massive lithic arkose of volcanic derivation exposed in the mine area.

The Footwall Arkose occurs on surface along a strike length of more than 20 km from the area near the east end of Red Indian Lake to the Buchans mine area. It has been intersected downdip in hole BJ-44 3 km north of the northernmost outcrop. The unit strikes east to northeast, dips northward at 5° to 80° and faces consistently northward. Felsic tuffs within the Arkose form the host rocks for the Connel Option massive sulphide body (see next Section) but all economic massive sulphide deposits in the Buchans area occur stratigraphically above the Arkose.

The felsic volcanic Wiley's Sequence conformably overlies the Footwall Basalt southwest of Buchans and contains many lithologies in common with the Footwall Arkose. It is therefore considered a time-equivalent more proximal volcanic facies of the Footwall Arkose though the exact correlation in the critical area south of Buchans is tenuous. It is further considered to be a stratigraphic equivalent of the Prominent Quartz Sequence, a unit which is a structural repetition in a higher thrust slice.

The conformable nature of the contact between Footwall Arkose and the Footwall Basalt is displayed in a number of drill cores in the mine area and in outcrop in

the Buchans River due south of Oriental. Indeed the upper portion of the Footwall Basalt is characterized by a number of arkosic horizons which are indistinguishable from lithologies of the Footwall Arkose proper. In this sense, the contact between the two formations on a large scale may be termed "interbedded". The upper contact of the Arkose is bounded by a bedding plane (thrust?) fault throughout large areas of the formation, although (rarely) in the mine area the Arkose is conformably overlain by the Intermediate Footwall (e.g., DDW 885).

In the eastern portion of the study area, the Footwall Arkose attains a maximum thickness of 2500 m (barring structural repetition) from the base to the fault-bounded top.

The Arkose is exposed in over 200 drill cores in the mine area and in a few additional holes throughout the strike length of the formation. It is best exposed in the Buchans Inlier in the bed of the Buchans River.

The Footwall Arkose is a multi-lithologic volcano-sedimentary sequence. In general, the lower parts of the formation consist of interbedded green greywacke, tuffaceous wacke, resedimented conglomerate, sharpstone breccia, siltstone, mudstone and multicoloured cherts. Shales are conspicuous in their absence (or lack of exposure). Interbedded with these are pumiceous felsic lithic-crystal




Plate 3-6: Typical lithic arkose from Footwall Arkose unit composed of quartz, plagioclase and reddish rhyolitic grains. Enlargement approximately 2X.

tuffs locally containing quartz crystals larger than 1 cm, a characteristic feature of the Wiley's and Prominent Quartz Sequences.

The upper parts of the formation contain a relatively greater proportion of chert and siltstone and, especially in the mine area, lithic arkose. This rock is pinkish, fine to coarse grained and is normally massive and devoid of sedimentary structures, including bedding (Plate 3-6). It consists of varying proportions of quartz, plagioclase (oligoclase-andesine), reddish rhyolitic rock fragments and lesser K-feldspar. Crystal fragments range in size from less than 1 mm to 8 mm although those of 0.5 to 1.5 mm are most common. Sorting is normally fair to poor. A weak hematitization of feldspars and the presence of small red rhyolite fragments imparts the characteristic light pinkish colour to the arkose.

Massive arkosic rocks contain a number of types of rounded, isolated pebbles, cobbles and boulders. Pink to red rhyolites up to several centimeters in diameter are most common and may be quartz and/or feldsparphyric and flow-banded. Some varieties contain prominent (greater than 5 mm) quartz phenocrysts, a characteristic of rhyolites of the Wiley's and Prominent Quartz Sequences.

Rare occurrences of granitoid boulders are known (e.g. DDH 247). Isolated pebbles of jasper and rounded

basaltic fragments are found locally. Rounded boulders of lithic arkose which contain rhyolite pebbles are found within normal arkosic rocks at some localities in the mine area. The presence of these boulders suggests instability during arkose deposition with cannibalization of previously deposited arkose.

In considering the mode of transport and deposition of arkosic rocks of the formation, the thick, coarse grained, massive nature of the units and the presence of outsize clasts are of importance. The haphazard occurrence of these clasts within massive beds devoid of sedimentary structures rules out deposition either from suspension or from traction currents. Alternatively, the arkose displays many of the features of grain flow deposits as described by Stauffer (1967). Deposition from non-turbulent mass flows of this nature satisfactorily explains the thick, massive and homogeneous nature of the arkose, its textural immaturity, the presence of isolated boulders of variable composition, the lack of sedimentary structures including bedding, the interbedding of such divergent lithologies as basalt and arkose with relatively little basaltic detritus in the arkose and the presence of thin interbeds of siltstone and mudstone between the mass flows.

Lithic arkosic rocks of the Footwall Arkose were probably not derived from an exposed granitic terrane. The

relative lack of alteration of plagioclase feldspars, the lack of clastic biotite or hornblende and the scarcity of granite boulders dictate against this possibility.

Conversely, the presence of felsic pyroclastics interbedded with lithic arkose, of "arkose" with a locally vitric matrix and of lithic arkose with abundant pink rhyolite fragments suggests that rhyolitic pyroclastics are the probable source rock for arkosic portions of the formation. The presence of rhyolite boulders with prominent (5-10 mm) quartz and feldspar phenocrysts suggests that the stratigraphically equivalent Wiley's Sequence is the source. A single current direction on imbricated rip-up clasts indicates a westerly source and therefore supports this hypothesis.

The Wiley's Sequence forms an extensive felsic volcanic horizon southwest of Buchans, overlying the Footwall Basalt and underlying the Intermediate Footwall. Lithologies include massive porphyritic to flow banded rhyolite, dacitic to rhyolitic pyroclastics and breccias, tuffaceous sediments, lithic arkose, wacke, siltstone, mudstone, jasper chert and local basaltic horizons. Several rhyolite and pyroclastic horizons are characterized by large (5 mm to 10 mm) quartz and plagioclase phenocrysts and crystals. Also occurring is a spectacular sequence of graded subaqueous pyroclastic flow deposits.

3.2.1.2.1. Connel Option: The Connel Option (Fig. 2-2) is a small uneconomic distal massive sulphide sheet which occurs within the Footwall Arkose. It was discovered in September 1928 by trenching on an "electrical indication". The deposit is a conformable bed of chalcopyrite, sphalerite, galena and pyrite; approximately 200 m long and with a maximum width of 60 cm.

The mineralized horizon is underlain by a series of thinly interbedded, laterally extensive, multi-coloured chert, siltstone and fine grained dacitic crystal-vitric dacitic tuff and ash. A single contorted Mn-rich bed up to 30 cm thick occurs within jasper cherts approximately 90 m in the stratigraphic footwall of the deposit. In the vicinity of the mineralization this sequence is overlain by a sequence of fine grained bedded dacitic ash, 15 cm of grey chert and subsequently by massive sulphides. The deposit is overlain by up to 1.5 m of unmineralized grey-green chert.

The deposit is not underlain by an altered and sulphide impregnated stockwork. Chert and tuff in the footwall locally contain disseminated traces of Pb-Zn-Cu-pyrite mineralization and thin (on the order of 1 mm) beds of sulphide. A thin (30 cm) highly sericitic alteration of fine grained vitric dacite occurs in the footwall in a single drill core.

The massive sulphides consist of very fine grained high grade sphalerite, galena and chalcopyrite with abundant pyrite exhibiting colloform texture. Faint banding of sulphides is visible in some areas. The principle gangue minerals are pyrite and silica. No barite is present.

Unlike the major massive sulphides of the Buchans area, the Connel Option cannot be related to any known nearby volcanic center. Indeed, the thin blanket shape of the sulphides and their occurrence in a thinly bedded, laterally extensive sequence of chert, siltstone and fine grained volcanic ash suggests a lack of significant paleo-relief and implies that the deposit is distal to the volcanic source.

The source of metal for the deposit is unknown. The lack of known stockwork alteration and mineralization in the footwall suggests that a local fumarole was not the source of the metals. Either of Sato's (1972) Type 1 or Type 3 brines could satisfy the geologic constraints on the source of the ore metal. In this regard it is interesting to speculate on the ore-forming potential of the volcanic center which was responsible for deposition of the Footwall Arkose.

3.2.1.3. Intermediate Footwall: The Intermediate Footwall is a sequence of mafic to intermediate

with minor felsic volcanic and volcanoclastic rocks. Unlike the underlying regionally extensive units, the Intermediate Footwall occupies a total surface and subsurface area of approximately 130 km². The term "Intermediate" is derived from the intermediate stratigraphic position between the underlying units and the overlying ore-bearing sequence and the intermediate bulk composition as compared to the felsic ore horizon sequence and the mafic Footwall Basalt. The broadness of the term "intermediate" is also convenient because of pervasive alteration in the mine area which renders recognition of original compositions difficult. The Intermediate Footwall is especially important in terms of ore genesis in that it is host for the stockwork mineralization below the massive sulphide orebodies and itself contains at least two small stratiform massive sulphide bodies.

The Intermediate Footwall originally conformably overlay the Footwall Arkose (e.g., DDH 885). Locally, the Intermediate Footwall appears to disconformably overlies the Footwall Basalt (e.g., DDH 1038). This disconformity is marked by a mixed breccia zone several meters thick consisting of numerous fragments of basalt of the Footwall Basalt and fragments of the Intermediate Footwall. The proportion of basaltic fragments decreases rapidly with distance above the contact and the proportion of altered fragments of the Intermediate Footwall

sympathetically increases. Apart from this local disconformable relationship and the above-mentioned conformable contact in DDH 885, the basal contact of the formation is a bedding plane fault and/or is occupied by diabase or "rhyolite" sills. The upper contact of the Intermediate Footwall is also commonly marked by a bedding plane fault although pyritic siltstone of the Lucky Strike ore horizon sequence locally conformably overlies the Intermediate Footwall (e.g., DDH 435).

Thickness estimates in the mine area are complicated by both faulting of the upper and lower contact and elsewhere by paleo-erosion at the upper contact which yielded rather uneven paleotopography. Regardless of these complications the formation attains a maximum thickness of at least 250 m in the area beneath Rothermere and Lucky Strike orebodies and exceeds 1000 m thickness in areas more remote from mineralization. Further description of the nature and role of the Intermediate Footwall in the mineralizing process is given under the heading "Ore deposits of the Lucky Strike ore horizon sequence".

3.2.1.4. Lucky Strike Ore Horizon Sequence:

The paleotopography following deposition of the Intermediate Footwall was characterized by a number of topographic highs, channels and depressions on the volcanic surface (Figure 3-2). The largest of these, MacLean Depression,




Figure 3-2. Vertical cross section through Lucky Strike, Two Level, Rothermere and MacLean Orebodies. Section line is shown in plan on p. 133.

reaches a maximum width of 800 m, a length in excess of 1 km and a depth locally exceeding 200 m. Smaller depressions underlie Rothermere, Two Level and Lucky Strike Orebodies. Felsic volcanic and sedimentary units of the Lucky Strike ore horizon sequence overlie the Intermediate Footwall, filled the paleotopographic depressions and subsequently levelled the existing topography. These units form the host rocks of the high grade stratiform polymetallic sulphide bodies including Lucky Strike, Two Level, Rothermere, MacLean and the Clementine Prospect.

The Lucky Strike ore horizon sequence is confined to the Buchans area roughly between Old Buchans and Clementine, and occupies a total known surface and subsurface extent of approximately 50 sq. km. It reaches a maximum thickness locally exceeding 1000 m. The sequence is exposed on surface only in rare outcrops and in the Lucky Strike open pit but is cored top to bottom in hundreds of drill holes. The sequence conformably overlies the Intermediate Footwall (e.g., DDH 435) although the basal contact has commonly acted as the locus for low angle faulting. The upper contact of the sequence with the Ski Hill and Prominent Quartz sequences is a major low angle shear zone which has been traced for more than 25 km across the Buchans area. The Lucky Strike ore horizon sequence is conformably overlain by the Lake Seven Basalt. This




Plate 3-7: Hematitized ore horizon rhyolite with white plagioclase and somewhat smaller clear quartz. Enlargement approximately 2X.

latter contact defines the top of the Lower Buchans Subgroup.

The lithologies of the Lucky Strike ore horizon sequence are described in some detail in the following sections. In general the sequence consists of basal rhyolite, pyritic siltstone, wacke and dacitic tuffs overlain by the massive and transported orebodies with associated volcanic breccia-conglomerate which are in turn overlain by a sequence of dacitic tuffs with minor basaltic and andesitic volcanics.

3.2.1.4.1. Rhyolite: Massive rhyolite occurs at the base of the Lucky Strike ore horizon sequence in the area north (downdip) of the Lucky Strike Orebody. The rhyolite is normally structureless to autobrecciated and is only rarely interbedded with other lithologies.

The rhyolite is characterized by moderate hematitization imparting maroon to pinkish hues to the rock (Plate 3-7). Most rhyolite is feldsparphyric though aphyric varieties and those containing feldspar and small resorbed quartz phenocrysts also occur. Oligoclase-andesine phenocrysts (1 mm) are normally clouded by minute hematite and partially altered to flecks of sericite and epidote with less common pumpellyite. In more hematitized rhyolites, calcite alteration of feldspar is dominant. The groundmass is normally holocrystalline, equigranular and very fine grained though local granophyric patches exist. Accessory

zircon, sphene, apatite, magnetite and traces of altered amphibole are common.

3.2.1.4.2. Pyritic Siltstone: Pyritic siltstone and wacke flank and overlie rhyolite but more commonly directly overlie the Intermediate Footwall (e.g., DDH 435). These sediments are restricted to the immediate vicinity of mineralization and occur in troughs and depressions in the volcanic surface. Thickness is controlled by this paleotopography and locally exceeds 100 m.

At the contact, the Intermediate Footwall consists of typically altered, veined and mineralized polyolithic breccia. This is overlain by pyritic siltstone with no veins and containing numerous altered fragments of Intermediate Footwall "floating" in a matrix of siltstone. These fragments die out rapidly above the contact but small lithic fragments of the Intermediate Footwall are common throughout the remainder of the siltstone unit.

The siltstone is typically light to dark grey, siliceous, poorly sorted and massive to locally thinly bedded, but rarely cherty (Plate 3-8). Slump structures, small scale cross beds, graded beds, cut and fill structures, load casts, flame structures, intraformational siltstone breccias, and small penecontemporaneous faults are common.

The siltstones and coarser volcanic wackes are



Plate 3-8: Sample from footwall pyritic siltstone unit of Lucky Strike ore horizon sequence exhibiting ball and pillow structures, flame structures and graded bedding.

composed of quartz and plagioclase crystals, a high proportion of rock fragments, minor volcanic glass and a small proportion of detrital zircon and apatite. Quartz and plagioclase are normally broken or shattered and feldspars are commonly altered to calcite. Locally there occur very thin dacitic crystal-vitric ash horizons with strongly resorbed quartz in a matrix of altered and flattened pumice shreds and shards. These horizons are harbingers of oncoming felsic volcanic activity of the ore horizon dacitic tuff.

Rock fragments include a wide variety of silicic, chloritic, pyritic, sericitic and clay mineralized fragments derived from the Intermediate Footwall, rare jasper chert and basaltic fragments. Breccias composed entirely of siltstone (termed siltstone breccia) occur locally. In some cases it is demonstrable that the siltstone fragments were transported to their ultimate site of deposition. In other areas, zones of bedded siltstone have been severely disrupted to form very coarse and tight packed breccias that evidently developed in place. These breccias suggest a very unstable depositional environment during siltstone deposition.

The most characteristic feature of all siltstones below the ore horizon is the presence of very fine grained disseminated matrix pyrite. This pyrite is euhedral to

broken to subrounded and is normally slightly finer grained than the siltstone in which it occurs. Pyrite forms distinct beds in cross bedded siltstone and pyrite grains are graded in size in graded silt beds. These features suggest that the pyrite was in hydrodynamic equilibrium with associated silicate grains and that the pyrite is detrital. Pyrite constitutes from less than one percent to five percent of most of the siltstone, though locally there occur very thin (less than one mm) beds composed virtually entirely of pyrite. Also present throughout the siltstone are traces of broken to rounded sphalerite and barite crystals and rare specks of chalcopyrite. These sulphide and sulphate minerals are also considered to be detrital and derived from the underlying Intermediate Footwall. The presence of these sulphides and the lack of alteration in the siltstone similar to that of the Intermediate Footwall at the contact provides further evidence that some stockwork mineralization and related alteration predated the major massive sulphide mineralization.

3.2.1.4.5. Dacitic Pyroclastics: Dacitic pyroclastics form the most widespread lithology of the ore horizon sequence. They are interbedded with, and overlie the rhyolite and pyritic siltstone (e.g., DDH 2704). Beyond the limits of these units, the pyroclastics directly overlie the Intermediate Footwall (e.g., DDH 1406).



Plate 3-9: Dacitic pyroclastic composed of fragments of dacitic crystal-vitric tuff in chloritic crystal-vitric matrix. Quartz and white plagioclase are clearly visible. This is one of several common types of ore horizon dacitic tuff.

The bulk of the dacitic pyroclastics occur in the hangingwall of the major orebodies but a thin unit does occur in the footwall. This latter unit varies markedly in thickness, being thin to locally absent in the Lucky Strike area but reaching thicknesses in excess of 100 m below MacLean. The distinction between footwall and hangingwall dacitic pyroclastics is difficult in areas more remote from Lucky Strike due to a lack of alteration in the footwall, the lack of a marker bed defining the ore horizon and the lithologic similarity of footwall and hangingwall tuffs. Due to this lithological similarity, these units will be described together in this section.

Dacitic tuffs and lapilli tuffs are the most common lithology in the ore horizon. These are light to dark green and contain widely varying proportions of quartz and plagioclase crystals with essential and accessory lithic fragments set in a fine grained matrix of variably altered volcanic glass. Bedding is locally visible and is defined by diagenetic flattening of glass shards and pumice shreds or by alteration of vitric-rich and vitric-poor beds (Plate 3-9).

Quartz crystals comprise up to 30% of the tuffs (most commonly 10-20%) and range in size from microscopic to 3 mm (average 1-2 mm). Quartz is rarely euhedral,

normally being resorbed, embayed, broken or a combination of these. Halos of secondary quartz are common around quartz crystals.

Feldspar crystals generally comprise the same proportion of the dacite as quartz and exhibit the same size range and average size as quartz. Feldspars are mainly plagioclase with much less common K-feldspar. Accurate determination of feldspar composition is hindered by alteration, normal and oscillatory zoning, broken crystals and locally by lack of twinning. In general, feldspar compositions as determined by the Michel-Levy method and on rare combined Carlsbad-albite twins is andesine. Plagioclase is variably altered to a fine grained intergrowth of sericite and clay minerals or less commonly calcite, pumpellyite, prehnite or epidote.

Hornblende crystals are normally absent though locally form up to two percent of the tuffs. In general, hornblende is more common in the Oriental area than in the Lucky Strike area and is generally more abundant in more vitric-rich tuffs. Alteration of hornblende to chlorite with anomalous lilac birefringence is common.

Apatite, sphene, rutile, zircon and leucoxene are ubiquitous but minor constituents of the tuff. Traces of pyrite are microscopically visible in most samples. Traces of biotite occur rarely.

Matrix volcanic glass and its altered equivalents form from 30 to 70% of most portions of the dacitic tuffs (though all extremes from glass-poor crystal-lithic tuff to 100% vitric ash or pumice tuff exist). Alteration of glass to a fine grained sericite-clay mineral-chlorite aggregate imparts the characteristic grey to green colours of the dacite. Despite complete alteration of glass, characteristic shard shapes are commonly visible microscopically and pumice shreds and larger long tube pumice fragments are normally visible with the hand lens or naked eye.

Volcanic glass, from microscopic shards to pumice fragments several centimeters in length, is almost invariably flattened into the bedding plane imparting a psuedo-foliated appearance which is characteristic of virtually all vitric-ash pyroclastics in the Buchans area. This feature previously led the author (Thurlow, 1973; Thurlow et al., 1975) to consider that portions of the dacitic tuffs were deposited subaerially. However, such tuffs are locally interbedded with laminated mudstone and siltstone (which in some cases is graded-bedded and displays small scale cross beds and load casts). Elsewhere, vitric dacitic tuff with strongly flattened pumice is interbedded with pillow lava. On the basis of these observations the occurrence of flattened pumice and glass shards is not considered to represent collapse due to load in a glowing sheet of subaerial volcanic ejecta. Similarly flattened

1

pumice in vitric pyroclastics of demonstrable submarine origin have been described by Fiske (1963), Fiske et al. (1963), Fiske and Matsuda (1964), Muttli (1965), Fiske (1969) and in no less than five papers in Ishihara (1974) describing Miocene volcanic rocks in the vicinity of the Kuroko deposits of Japan. Fiske (1969) states that pumice can be flattened into the bedding plane to produce "a highly compact rock that, in some cases, superficially resembles welded tuff" and Ohtagaki et al. (1974) state that "in appearance, it resembles the pumice found in common welded tuffs". Ohtagaki et al. consider that flattened pumice in subaqueous sequences is a result of compaction during dewatering and diagenesis of pumice which has been depositionally and diagenetically altered to soft clay minerals. A similar mode of origin of flattened pumice in the Buchans area is considered likely.

The most common fragment types within dacitic tuffs are fragments of dacitic tuff and rhyolite grains and clasts. Fragments of dacite crystal-vitric tuff are diffuse to well defined, rounded to angular and have a composition and texture very similar to the matrix material. The colour of these fragments is normally slightly different from the matrix and serves to distinguish the fragments. Accessory lithic fragments are common especially at the base of graded subaqueous pyroclastic flows. The base of such flows consists of tightly packed

breccias with a crystal-vitric matrix and contains angular to rounded fragments of virtually every lithologic type below the dacite (i.e., basalt, jasper chert, numerous varieties of altered Intermediate Footwall, rhyolite, siltstone, granodiorite and (rarely) boulders of Feeder Granodiorite. Lithic fragments in these pyroclastic debris flows become finer grained and less abundant upward as the flow grades into crystal-vitric tuff and may be overlain by vitric ash or tuffaceous siltstone.

In the vicinity of the in situ massive sulphides the distinction between footwall and hangingwall dacite is not difficult. Near Lucky Strike and Rothermere Orebodies, the footwall dacite is normally greyer and more hydrothermally altered (to clay minerals) and may contain traces of galena, sphalerite and pyrite. However, beyond the area of significant alteration (e.g., beyond the vicinity of MacLean Orebody) the author has been unable to find any consistent macroscopic, microscopic or geochemical difference between the footwall and hangingwall dacite.

Although most portions of the dacitic tuffs are sufficiently similar such that recognition of any internal stratigraphy is difficult, there exists one distinctive type. Locally known as the "coarse-quartz dacite", this crystal-vitric tuff is characterized by quartz crystals somewhat coarser than normal (average 2-4 mm) and conformably



Plate 3-10: Typical dacitic pyroclastic from the "coarse quartz" unit. Sample contains 1-3 mm quartz, 1 mm plagioclase in matrix of black chloritic altered glass. Pinkish lithic fragments are rhyolite; dark lithics are basalt and andesite.

overlies the normal hangingwall dacite (e.g., DDH 827). It is best developed in a depositional basin which occurs between the Lucky Strike - MacLean and Clementine areas and locally exceeds 100 m in thickness. The coarse-quartz dacite is intimately associated with graded subaqueous debris flows and itself contains more lithic fragments than the normal tuff. Locally the coarse-quartz dacite is interbedded with more normal dacitic tuff but is emphasized that the first appearance of the coarse-quartz dacite occurs well in the hangingwall of the major ore deposits. This relationship is important as this dacite locally contains scattered pebbles of high grade black ore and is conformably overlain by polyolithic breccia-conglomerate which also locally contains sulphide fragments.

The coarse-quartz dacite contains a variety of lithic fragments with pinkish massive rhyolite and aphyric aphanitic andesite fragments being more common (Plate 3-10). Lithic-rich beds are commonly separated by vitric-rich beds imparting a diffuse banding to the rock.

The coarse-quartz dacite is conformably overlain (e.g., DDH's 799, 827) by a complex polyolithic breccia-conglomerate unit which, like the coarse-quartz member, occupies the depositional basin between the Lucky Strike - MacLean area and the Clementine area. This unit is characterized by an abundance of pink aphyric, to quartz and



Plate 3-11: Tight packed pyroclastic breccia from coarse-quartz member of Lucky Strike ore horizon sequence. Angular to rounded fragments are mainly rhyolite.

feldsparphyric rhyolite fragments (Plate 3-11) although a lesser proportion of basaltic and andesitic fragments (most of which are of unknown origin) are also normally present. Less abundant fragment types include cherty siltstone, pyritic siltstone, rare granodiorite boulders, pumice fragments, dark grey carbonate fragments (e.g., DDH's 629, 799) of unknown origin and sulphide fragments. Fragments are moderately to tightly packed in a lithic quartzo-feldspathic matrix with only a local significant vitric component. Breccia-conglomerate beds are normally massive although thick graded beds ranging from coarse Breccia-conglomerate to cherty siltstone are not uncommon (e.g., DDH's 2210, 629). These debris flows are interbedded with greywacke, siltstone (locally pyritic), vitric tuff and spectacular graded pumiceous pyroclastic flows (e.g., DDH's 1562, 932).

Sulphide fragments in this horizon and (less commonly) in the underlying coarse-quartz member are of interest. Sulphide clasts have been recognized macroscopically in at least 10 drill holes, i.e., DDH's 191, 1442, 2681, 2243, 2332, 2224, 2210, 1406, 2214 and 2778. Black ore clasts are most common with much less abundant yellow ore and stockwork fragments. Sulphide clasts range from angular to rounded, competent to plastic and reach a maximum size of 13 cm (although the average is

less than one cm). They are especially concentrated in the thickest portion of a basin which existed in the area between Lucky Strike and Clementine.

The source of the sulphide fragments is of interest in exploration. Since the fragments occur in a stratigraphic horizon at least 200 m (diabase removed) above the Rothermere ore horizon it is unlikely that the Lucky Strike Orebody provided the sulphide fragments in this breccia conglomerate. The relative rarity of fragments which are so abundant in the main ore horizon (i.e., granodiorite, pyritic siltstone and varieties of Intermediate Footwall) as well as the presence of basaltic, andesitic and carbonate fragments of unknown origin further suggest that the lithologies in the breccia-conglomerate are not from the Lucky Strike area. The decrease in abundance of sulphide fragments towards the Lucky Strike area further dictates against this area as a source. Some of the extraneous fragment lithologies can be matched with those in the Clementine clastic ore horizon, although the sulphide-bearing horizon lies above Clementine as well.

3.2.1.4.4. Lucky Strike Ore Horizon: Massive sulphide, transported ores and facies-equivalent breccia-conglomerate and "granite-conglomerate" overlie dacitic pyroclastics. These are described in Chapter Four.

The ores are overlain by a thick sequence of dacitic pyroclastics, as described in the previous section, and subsequently overlain by the Lake Seven Basalt.

3.2.1.5. Lake Seven Basalt: The Lucky Strike ore horizon sequence is conformably overlain by the Lake Seven Basalt (e.g., DDH 799) which subcrops west of Rothermere Orebody (Fig. 2-2). Though it forms the base of the Upper Buchans Subgroup, the Lake Seven Basalt is described in this section as it is the uppermost member of the Lucky Strike thrust block. The unit is areally restricted and consists dominantly of basaltic pillow lava, pillow breccia, pyroclastics and massive flows. Flows contain small phenocrysts of fresh augite and labradorite largely altered to calcite, sericite and epidote. The phenocrysts occur in a fine grained diabasic-textured matrix of plagioclase and clinopyroxene, the latter altered to chlorite. Locally flows are black and very fine grained and contain quenched plagioclase with swallow tail terminations and hollow cores, i.e., similar to those described by Pearce (1974).

Several features of the Lake Seven Basalt are similar to those of the Footwall Basalt and indeed, distinction of the two units on a hand specimen basis is impossible. Pink to white calcite amygdules are common in the Lake Seven Basalt and the unit has locally been affected by weak

to strong hematitization. Pyroclastic units commonly have a pink to white crystalline calcite-rich matrix (e.g., DDH 1562) and may show size grading of individual beds.

3.2.1.6. A Stratigraphic Discontinuity at the Top of the Lake Seven Basalt: The interpretation of the uppermost contact of the Lake Seven Basalt with overlying units has far-reaching effects both in terms of understanding the stratigraphy and in terms of exploration. Despite hundreds of drill penetrations, an indisputably conformable contact has not yet been observed. Normally, the contact is occupied by one of the innumerable diabase sills which cut the Buchans Group, or is a zone of poor core recovery or, in some cases, a zone of obvious shearing. The same comments apply in areas where the Lake Seven Basalt is absent and where the overlying Prominent Quartz or Ski Hill sequences lie directly on the Lucky Strike ore horizon sequence.

Two opposing interpretations are based upon the nature of this contact: 1) the contact is a zone of localized bedding plane shearing developed during regional folding and is of no regional consequence in understanding the stratigraphy, and 2) the contact represents a thrust fault superimposing older on younger lithologies. The significance of the latter hypothesis is discussed more fully in Section 3.3.1.1.




Plate 3-12: Porphyritic rhyolite of Prominent Quartz
Sequence containing large rounded to sub-
hedral quartz and smaller zoned plagioclase
phenocrysts.

3.2.1.7. Prominent Quartz Sequence: Rocks of the Prominent Quartz sequence are the lowermost units recognized above the stratigraphic discontinuity discussed in the previous section. The sequence is widely distributed to the north and west of Buchans and occupies a known surface and subsurface area of approximately 100 square kilometers. Outcrop exposures are extremely sparse but the sequence is fairly well known through drilling designed to penetrate the underlying ore horizon. The sequence is absent in areas east of Buchans but reaches thicknesses in excess of one kilometer to the west. Field, petrographic and geochemical evidence suggests that felsic volcanics of this sequence were derived from magmas of the Feeder Granodiorite. The Prominent Quartz Sequence is considered to be a thrust repetition of the Wiley's Sequence and Footwall Arkose.

The Prominent Quartz sequence consists dominantly of rhyolitic flows and pyroclastics which are commonly characterized by the presence of "prominent" (5-10 mm) quartz phenocrysts (Plate 3-12). These are normally rounded and somewhat embayed though locally well preserved hexagonal crystal outlines exist. They range in size up to a maximum of 3 cm. Irregular to radial cracks are common. Zoned plagioclase phenocrysts are normally present in approximately the same abundance as quartz but are generally smaller, reaching a maximum size of 1 cm. Oscillatory zoning is

common and even in flows plagioclase crystals are commonly broken. Plagioclase is commonly partly altered to pumpellyite, epidote, calcite and sericite.

Small (less than 1 mm) hornblende phenocrysts commonly comprise less than 5% of rhyolite flows and are normally altered to chlorite with lesser pumpellyite, epidote and calcite. Associated with these, there is normally a small proportion of magnetite which also may occur as small isolated crystals in the matrix. The matrix to flows is aphanitic and quartzofeldspathic in a wide variety of colours ranging from salmon pink to black.

Individual rhyolite flows of the Prominent Quartz sequence have been traced over areas exceeding a square kilometer. Locally thick (greater than 100 m) massive structureless rhyolite flows (e.g., DDH 2090) display no flow banding or autobrecciation and are overlain with sharp conformable contact by lithic-crystal-vitric tuff with some fragments of massive rhyolite. The sharp nature of such contacts, the lack of autobrecciated flow tops, the widespread distribution of the flows and their coarsely porphyritic nature are evidence in support of unusually fluid rhyolite magmas.

Rhyolite flows, especially those south of Lucky Strike, have locally undergone strong silicification, sericitization and pyritization but no significant

mineralization is known.

Crystal-vitric tuffaceous rocks of the Prominent Quartz sequence contain an identical phenocryst assemblage to the flows, i.e., prominent cracked quartz, zoned plagioclase and hornblende, but the matrix consists of anastomosing flattened glass shards and larger pumice shreds. Volcanic glass is commonly altered to dark green to black chlorite.

Also included in the Prominent Quartz sequence are important members of rhyolite and (related?) pumiceous pyroclastics which do not contain prominent quartz and feldspar phenocrysts but instead normally contain smaller (1 mm) quartz crystals. Particularly important are widespread, distinctive and spectacular subaqueous pyroclastic flows which are apparently confined to the northern and western portions of the Buchans Group. These are thick graded deposits ranging from coarse polyolithic breccias with fragments locally in excess of 1 m at the base (but normally less than 10 cm). These grade upward into finer grained breccias with abundant multicoloured long tube pumice and further upward into finer grained lithic-crystal-vitric tuff overlain gradationally by siltstone and cherty mudstone. Individual deposits may be tens of meters thick and the entire sequence consists of several tens of individual pyroclastic flows. Double grading such as that

described by Fiske and Matsuda (1964) is locally developed. The thickness of these deposits and the abundance of pumiceous material suggests that they were triggered by violent volcanic explosions. The coarsest facies of these units are developed in the Wiley's Sequence whereas finer grained more distal facies exist in numerous drill cores to the north (e.g., DDH's 2666, 724). This suggests northerly directed pyroclastic flow from an active (now eroded) felsic volcanic centre to the south.

A variety of tuffaceous sediments are interbedded with the volcanics of the Prominent Quartz sequence. The most unusual of these is a sequence composed dominantly of black, locally pyritic and carbonaceous mudstone which occur south of Lucky Strike (e.g., DDH's 1908, 1904). These are massive to thinly laminated and generally do not break parallel to bedding, hindering the search for macrofossils. Several samples of this material were sent for detailed microfossil analysis but proved barren (J. Utting, pers. comm., 1975).

A number of basaltic horizons are included within rocks of the Prominent Quartz sequence. These horizons are macroscopically and chemically similar to rocks of the structurally underlying Lake Seven Basalt and Footwall Basalt.

3.2.1.8. Ski Hill Sequence: The Ski Hill sequence consists of a thick pile of black to dark green breccias, pyroclastics and pillow lava of basaltic to andesitic composition. It conformably overlies the Prominent Quartz sequence (e.g., DDH's 2925, 2927, 2841) but where the underlying Prominent Quartz sequence is absent, the Ski Hill sequence rests with sheared contact directly upon the Lucky Strike ore horizon sequence. Sporadic outcrops of the Ski Hill sequence occur in an east-west trending belt extending from the Buchans River approximately 4 km west to the Lake Seven area (Fig. 2-2). The sequence occupies a known surface and subsurface area of approximately 30 square kilometers and reaches a maximum known thickness of approximately 1 kilometer.

The Ski Hill sequence consists mainly of andesitic breccias and pyroclastics interbedded with massive and pillowed flows of basaltic to andesitic composition. In general, flows of the sequence consist of plagioclase (andesine-labradorite) and augite phenocrysts set in a very fine grained diabasic-textured groundmass. Like the Footwall and Lake Seven Basalts, augite phenocrysts are normally much less altered than plagioclase phenocrysts or matrix pyroxene. Plagioclase phenocrysts are commonly zoned, aligned or feldspathic and generally partially altered to combinations of epidote, calcite, pumpellyite or clay minerals. Most flows and blocks of massive andesite in



Plate 3-13: Typical Ski Hill Sequence breccia with both elongate and spherical quartz amygdules. Elongate vesicles at top of larger fragment (near center of photograph) are filled with matrix material. Approximately true size.

breccias contain zoned amygdules filled with combinations of prehnite, quartz, pumpellyite, chlorite and calcite. Spherical, flow-stretched and amoeboid amygdules are common (Plate 3-13).

The top of the Ski Hill sequence is normally characterized by an upward increase in the abundance of exotic accessory fragments which grade upwards into poly-lithic breccias of density flow origin. These breccias contain a variety of lithic fragments ranging in composition from basalt to rhyolite and locally contain rounded granodiorite pebbles and cobbles (e.g., DDH's 885, 1003, 1038, 1185). In places, these breccias are characterized by a siliceous siltstone matrix bearing detrital pyrite, much like siltstones of the overlying Oriental ore horizon sequence. Rare occurrences of Pb-Zn beds (?) and of high grade black ore fragments in breccias (e.g., DDH's 1911, 1884) are also known in these upper portions of the Ski Hill sequence:

The distinction between rocks of the Ski Hill sequence and Footwall Basalt is locally difficult and, although some empirical generalizations can be made, there are many exceptions. Rocks of true andesitic composition are dominant in the Ski Hill sequence and rare in the Footwall Basalt. No arkosic rocks similar to the Footwall Arkose are known within the Ski Hill sequence and

the sequence is nowhere immediately overlain by arkosic rocks. Interpillow chert and chert interbeds are much more common in the Footwall Basalt than in the Ski Hill sequence. In general, calcite amygdules (especially pink calcite) are more common in the Footwall Basalt and amygdule abundance is also generally greater. Distinctive small elongate amygdules stretched during flow are common in the Ski Hill sequence and rare in the Footwall Basalt. Quartz-prehnite-pumpellyite amygdules are generally more common in the Ski Hill sequence. Pervasive hematitization is somewhat more common in the Footwall Basalt. In general, pillow lava is somewhat more common in the Footwall Basalt whereas true pyroclastics and a variety of breccia types are more common in the Ski Hill sequence. Breccias of the Ski Hill sequence are commonly polyolithic (in varieties of basalt and andesite) whereas breccias of the Footwall Basalt are most commonly monolithic. Polyolithic breccias such as those that occur at the top of the Ski Hill sequence are not known in the Footwall Basalt. Pyritic siltstone, plutonic pebbles and black ore occurrences are also not known in the Footwall Basalt.

3.2.1.9. Oriental Intermediate Footwall: A unit known as the Oriental Intermediate Footwall conformably overlies the Ski Hill Sequence and forms the host for epigenetic stringer and stockwork mineralization associated

with the Oriental #1 orebody. The Oriental Intermediate consists dominantly of an areally restricted pile of altered intermediate to felsic pyroclastics and especially pumice tuff, pumice breccia and vitric tuff.

Though this sequence is dominantly felsic, the term "intermediate" is convenient in that the more felsic end members of the sequence are quartz-poor to quartz-free in contrast to the quartz-rich pyroclastics of the Oriental ore horizon sequence. Also, as at Lucky Strike, the "intermediate" volcanics occupy a stratigraphic position intermediate between underlying mafic and overlying felsic sequences.

The Oriental Intermediate is discussed more fully in Section 4.4.

3.2.1.10. Oriental Ore Horizon Sequence: The Oriental ore horizon sequence is a significant felsic volcanic accumulation which conformably overlies the Oriental Intermediate. It forms the host rocks for the in situ Oriental #1 Orebody and the transported Oriental #2 and Old Buchans Conglomerate Orebodies as well as the subeconomic transported Sandfill and Middle Branch Prospects.

The sequence is shallowly northward-dipping and occupies a known surface and subsurface area of less than 10 square kilometers in an area north of Old Buchans and

Oriental. The sequence reaches a maximum thickness of approximately 400 meters in an area north of Oriental but thins rapidly both to the east and west. The northerly downdip extensions of the sequence have not been fully outlined by diamond drilling.

The lithologies and stratigraphy of the Oriental ore horizon sequence are remarkably similar to their counterparts at Lucky Strike. Detailed discussion of the sequence is given in Chapter Four.

3.2.2. Stratigraphy of the Upper Buchans Subgroup. The Upper Buchans Subgroup is characterized by mafic, intermediate and felsic volcanic rocks with the latter predominant. In keeping with the generally northerly dips in the area, it is generally distributed in the north part of the study area. The stratigraphy is less well known than that of the Lower Buchans Subgroup largely because of the lack of economic incentive to drill these rocks systematically. Information is largely derived as a consequence of drilling of the underlying Oriental ore horizon sequence. The understanding of the stratigraphy is further hampered by significant thrust faults, the effects of which are not yet fully understood.

There was no recognizable break in volcanism-sedimentation between Lower and Upper Subgroups but the

boundary marks the end of the Oriental ore horizon felsic volcanic sequence and the upper stratigraphic limit of known economic mineralization. It also marks the onset of renewed mafic volcanism on a widespread scale and concurrent influx of arkosic density flows containing detritus extraneous to the intermediate Buchans area. As such, the boundary marks the beginning of a second volcanic cycle, not totally dissimilar to that of the Lower Buchans Subgroup.

The Oriental ore horizon sequence is conformably overlain by two separate units which interfinger in complex fashion at their margins. The Sandy Lake Basalt conformably overlies the Oriental ore horizon sequence in the west (e.g., DDH's 2853, 1582) whereas the Upper Arkose forms the base of the Upper Buchans Subgroup to the east (conformable contacts in DDH's 1683 and 1872, among others).

The Sandy Lake Basalt consists of basaltic pillow lava, pillow breccia and pyroclastics all characterized by abundant calcite amygdules, labradorite and/or augite phenocrysts and locally intense diagenetic calcite-hematite introduction. As such, this basaltic unit is virtually indistinguishable from basalts of the Lower Buchans Subgroup.

On the other hand, the Upper Arkose is among one of the more unique and interesting units in the Buchans area. The Upper Arkose occupies and levels the remnants



Plate 3-14: Volcanic conglomerate of Upper Arkose.

Rounded cobbles are varieties of rhyolite,
some with large quartz phenocrysts.

of a deep paleotopographic trough in the underlying Oriental ore horizon sequence. Towards the western margin of Arkose deposition, the Upper Arkose conformably overlies (e.g., DDH's 1582, 1758) and interfingers with the Sandy Lake Basalt. The Upper Arkose reaches a maximum unshered thickness in excess of 100 m and locally attains a total thickness in excess of 300 m.

The Upper Arkose is composed almost entirely of lithic arkose and arkosic conglomerate. The conglomerate is polymictic and contains boulders of massive rhyolite and dacitic crystal-vitric tuff, both with prominent (5-10 mm) quartz phenocrysts and pink rhyolite with or without small quartz and feldspar phenocrysts (Plate 3-14). Less common are calcite amygdaloidal basalt (very similar to those of the Sandy Lake Basalt), arkosic conglomerate boulders, and minor granitoid boulders. Rely (1960, p. 82, 97) reported an occurrence of a sulphide boulder in the Upper Arkose in the Oriental east pit though none have been noted by the author, E.A. Swanson or other workers. Most boulders are well rounded with moderate sphericity and vary from tightly to loosely packed. The matrix is normally immature and poorly sorted, consisting of small lithic fragments and quartz and feldspar crystals. The conglomerates are normally massive and devoid of all sedimentary structures, though local thick graded beds exist and rare inverse to normally graded beds occur (e.g., DDH 1492). Near the base

of the conglomerate in the Oriental area, there exists at least one horizon of dacitic crystal-vitric tuff macroscopically, microscopically and geochemically identical to that of the underlying Oriental ore horizon sequence (e.g., DDH 1606). The presence of this interbedding at the upper contact of the Lower Buchans Subgroup provides strong evidence that there was no depositional break at this juncture.

Rocks of this arkosic conglomerate horizon and those of the Footwall Arkose Formation are virtually indistinguishable. Only the stratigraphic position is diagnostic in identifying these lithologies. However, in general the Upper Arkose normally contains a greater abundance of conglomerate and a greater variety of boulder lithologies. Felsic volcanic boulders with prominent quartz (5-10 mm), granitoid boulders and basaltic boulders are relatively more common in the Upper Arkose than in the Footwall Arkose.

The Upper Arkose is considered to have been deposited by the same mechanism as the Footwall Arkose (i.e., density flow) and evidence supporting this hypothesis is similar to that quoted for the Footwall Arkose. Indeed, their textural and lithologic similarity suggests that the same volcanic source may have supplied detritus in both units. The vast difference in lithologies between those

of the Footwall-Upper Arkose and those of the ore horizon sequence suggests that the volcanic sources for these sequences were spatially separated. The occurrence of ore horizon dacitic crystal-vitric tuff identical to that of the Lucky Strike Formation interbedded with the Upper Arkose probably represents an interfingering of volcanic materials from these two sources and further represents the last volcanic gasp from the source which produced the Oriental ore horizon sequence. Evidently volcanic activity shifted back and forth between these two centers with very little temporal overlap.

The Upper Arkose is conformably overlain by the Little Sandy sequence (e.g., DDH's 1606, 1683). This dominantly volcanic unit is the uppermost sequence of the Buchans Group and, although widely distributed to the north and east of Buchans (Fig. 2-2), is probably the most poorly known in the area. The sequence forms the host rocks for the Little Sandy copper mineralization several kilometers east of Buchans.

The base of the Formation in the east is everywhere fault-bounded, juxtaposing rocks of the Little Sandy sequence and Footwall Arkose. The contact is marked by shearing, fault gouge, diabase dykes or by discontinuous linear patches of poorly indurated red sandstone and conglomerate containing abundant boulders of peralkaline rhyolite




Plate 3-15: Rhyolite of the Little Sandy sequence, Upper Buchans Subgroup. Note contorted flow banding.

and related lithologies (e.g., DDH BJ-24). MacLean (1943) considered (with some reservations) these lithologies to be interbedded with the Buchans Group but they are herein interpreted as remnants of presumed Carboniferous age deposited unconformably upon the Buchans Group along the paleo-eroded fault contact.

The upper contact of the Little Sandy sequence is marked either by a thrust contact juxtaposing the Hungry Mountain Complex and the Little Sandy sequence or by intrusive contact with the alkali-feldspar phase of the Topsails Granite. In the poorly known northwestern portion of the study area, the Hungry Mountain Complex may originally have been absent and lithologies thought to be correlative with the Springdale Group may lie unconformably upon the Buchans Group (this relationship is present between the Springdale Group and Roberts Arm Groups in Notre Dame Bay, c.f. Strong, 1973).

The Little Sandy sequence consists of abundant fine grained andesitic to rhyolite pyroclastics with less abundant volcanogenic sediments and basaltic flows and volcaniclastics.

Intermediate and felsic volcanics are much more common than mafic varieties. Massive rhyolite flows with wildly contorted flow banding (Plate 3-15), pumice flows, pyritic tuff, rhyolitic autobreccias and thick units of

bedded andesitic tuff are the most common lithologies. Locally there occurs a distinctive dacitic crystal-vitric tuff indistinguishable from that of the ore horizon sequences (e.g., DDH 1606).

Strong alteration is characteristic of large portions of felsic sequences yielding typically rubbly, white to rusty weathering outcrops. Alteration consists of varying degrees of silicification, chloritization, sericitization, clay mineral alteration and ubiquitous pyritization. Interbedded breccias locally contain fragments which display alteration rims similar to those in the ore horizon sequences. The Little Sandy copper prospect occurs in such a sequence of altered intermediate to rhyolitic rocks as thin, reticulating, chalcopyrite-bearing chloritic quartz veins and open space filling between breccia fragments. The deposit, though epigenetic in detail, is stratiform on a large scale and considered to be a product of near surface or surface fumarolic activity.

3.2.3. Diabase and Related Intrusives: All units in the Buchans area are cut by fine to coarse grained, dark green to black diabase dykes and sills. These bodies range in size from small dykelets to large intrusives locally in excess of 700 m in thickness. They consist of porphyritic to sub-ophitic labradorite and augite with traces of opaques, sphene, leucoxene and green amphibole

(hornblende?) normally replaced by chlorite. Quench textures in pyroxene and plagioclase are common and chlorite locally replaces matrix glass. Grain size varies sympathetically with diabase thickness and may exceed 1 cm in thicker intrusions. Chlorite-calcite-quartz-filled amygdules are common, indicating a shallow depth of emplacement. Locally in thicker sills, amygdule size and proportion are related to stratigraphic height. Chilled margins are common and cumulus layering is noticeable in thicker dykes and sills. Macroscopically visible contact metamorphism is restricted to the margins of thicker intrusions.

The author's previous work (Thurlow, 1973) determined the presence of two chemically distinct diabase types, i.e., a relatively more abundant "high titanium diabase" characterized by brownish to purplish to pinkish (titaniferous?) augite and "low titanium diabase" which exhibits many features in common with the Buchans Group volcanics. These two diabase types are easily separable on many geochemical plots, e.g., Fig. 5-4. Petrographic examination of diabase and gabbro related to the alkali-feldspar phase of the Topsails Granite indicates that these are mineralogically and texturally indistinguishable from the high titanium diabase and a temporal and genetic link of these mildly alkaline intrusives is suggested. In contrast, the low titanium diabase is geochemically identical to the

Buchans Group volcanics and thus may represent feeders to basaltic rocks of the Buchans Group. Locally, difficulty exists in separating sills from massive flows in drill core.

Diabase occurs commonly as sills associated with gouge zones at important geological contacts. Some of these contacts can nowhere be demonstrated to be conformable despite hundreds of penetrations, raising the possibility of low-angle thrust faulting, a hypothesis that finds support in a number of geological contexts. Consequently, diabase distribution may be controlled to some degree by the presence of earlier-formed thrust faults. Some sills have themselves undergone shearing which, according to this model, must have occurred as rejuvenations along old fault zones, probably during regional flexural folding. Detailed study of the distribution of high and low titanium diabase dykes has not been undertaken but, according to the present model, only high titanium diabase could be present as thrust fault-plane fillings.

Dykes of felsite and granite also cut the Buchans Group, commonly in association with diabase. Mutual cross-cutting relations of felsite and diabase have been observed at several localities. Granitic dykes are macroscopically, microscopically and geochemically identical to the alkali-feldspar phase of the Topsails Granite and are almost certainly apophyses from this body.

3.3. Structure and Metamorphism of the Buchans Area

3.3.1. Structure of the Buchana Area: The structure of the Buchans area, though relatively simple on a regional scale, is complex on the detailed scale used in exploration. The Buchans area has been subjected to two structural events of regional consequence, i.e., a period of Silurian thrust faulting and a later period of Devonian (?) folding.

3.3.1.1. Silurian Thrust Faulting in the Buchans

Group: The evidence supporting an important regional event of Silurian thrust faulting has been presented in Section 2.7. This southeastward directed event caused the present superposition of the Hungry Mountain Complex upon the Buchans Group. As far as the Hungry Mountain Complex is concerned, there seems to be general acceptance of this hypothesis. However, the suggestion that similar major faults occur within the Buchans Group has been the subject of debate. The author is of the opinion that such faults do exist though, as for opinions to the contrary, no proof can be offered. The possibility and implications of such faulting are outlined in this section.

The relative stratigraphic positions of the Lucky Strike and Oriental orebodies and their original positions within the stratigraphy are at the crux of the thrusting problem. In the past, various workers have interpreted the

Lucky Strike ore horizon sequence to either overlie the Oriental ore horizon sequence (e.g., George, 1937), or to be overlain by the Oriental sequence (e.g., Swanson and Brown, 1962; Thurlow, Swanson and Strong, 1975) or to be stratigraphically equivalent to the Oriental sequence (e.g., Relly, 1960). According to a thrust hypothesis, the Oriental sequence now structurally overlies the Lucky Strike sequence but represents a thrust repetition of sequences that were originally stratigraphic equivalents.

The thrust hypothesis is based upon two undeniable facts; 1) that the Oriental ore horizon sequence represents a remarkable lithologic and stratigraphic repetition of the Lucky Strike ore horizon sequence, and 2) that no indisputably conformable contact has been observed between the blocks of rock containing the Lucky Strike and Oriental orebodies.

The remarkable stratigraphic repetition of lithologies formed the basis for the concept of volcanic cycles at Buchans. This same repetition led Entwistle and Barnes (1971) to suggest that the Oriental sequence is a faulted repetition of the Lucky Strike sequence. Walker and King (1975) provided the first hard evidence in support of this concept by demonstrating a lack of stratigraphic conformity between the Ski Hill and Lucky Strike ore horizon sequences. The possibility of such

faults has demanded the construction of a stratigraphic column based on indisputably conformable contacts. This stratigraphy has been outlined in previous sections and summarized in Fig. 3-1.

The suggestion that Lucky Strike and Oriental orebodies originally lay at the same stratigraphic horizon implies that certain rock sequences are also stratigraphic correlatives. These are outlined in Table 3-2. Without exception, these implied correlations are extremely plausible and provide further circumstantial evidence of thrust faulting.

A second possible stratigraphic reconstruction is presented in Table 3-3. This model implies not only that Lucky Strike and Oriental are correlative but also that the Footwall and Upper Arkose are correlative. Whether or not this reconstruction is valid, hinges on the interpretation of contact relations in DDH 885. It is beyond the scope of this thesis to elaborate fully on this point. This second model has far-reaching implications both in terms of regional and local stratigraphy and in terms of exploration.

The hypothesis that thrust faulting is an important element in Buchans geology is supported also by the following points:

Table 3-2: Buchans Group Thrust Correlatives - Model A

Lucky Strike Fault Block

Oriental Fault Block

Little Sandy Sequence

~~~~~ fault ~~~~~

Lake Seven Basalt

Sandy Lake Basalt  
and Upper Arkose

Lucky Strike Ore Horizon  
Sequence

Oriental Horizon  
Sequence

Intermediate Footwall

Ski Hill Sequence  
and Oriental Inter-  
mediate

Footwall Arkose and Wiley's  
Sequence

Prominent Quartz  
Sequence

~~~~~ fault ~~~~~

Footwall Basalt

Table 3-3: Buchans Group Thrust Correlatives - Model B

Block A

Block B

Block C

Little Sandy Sequence

~~~~~ fault ~~~~~

Footwall Arkose

Upper Arkose

~~~~~ fault ~~~~~

Footwall Basalt

Lake Seven Basalt

Sandy Lake Basalt

Victoria Lake
Volcanics

Lucky Strike Sequence—Oriental Sequence

Intermediate Footwall—Ski Hill and Oriental
Intermediate

Wiley's Sequence — Prominent Quartz
Sequence

~~~~~ fault ~~~~~



- 1) the anomalously rapid pinching out of some units along sheared contacts,
- 2) the occurrence of minor shears repeating lithologies, especially in the Clementine area, and areas north of MagLean Orebody, and
- 3) the occurrence of apparent Footwall rocks overlying the West orebody.

The suggestion that thrust faults are not a significant factor in Buchans geology can also be forcefully argued. The bedding plane shears visible at many important contacts can be interpreted as bedding plane slip developed during regional folding.

The fact that demonstrably similar lithologies represent definite stratigraphic repetitions is a further argument in favour of cyclic volcanism and the lack of any need to invoke thrust faulting. The prominent quartz-bearing Upper Arkose is stratigraphically hundreds of meters above the Prominent Quartz sequence and represents a remarkable stratigraphic repetition of similar lithologies. Other such examples can be cited; e.g., dacitic crystal-vitric tuffs of the Oriental and Little Sandy sequences.

#### 3.3.1.2. East-West Fold Axes Related to Thrusting:

In several parts of the Buchans Group there is some evidence which implies an early period of small scale

folding. Open warps with east-west trending axes of several kilometers length have become evident through recent diamond drilling and stratigraphic re-interpretations. Because of generally shallow dips, these folds have given rise to unusual map distributions of some sequences, especially in the area west of Lucky Strike (Fig. 2-2).

Apart from limited areal extent, there exists some instances where these folds have only limited vertical extent and are confined to units overlying a postulated thrust fault. It is noteworthy also that the axes of these small folds are perpendicular to the proposed direction of Silurian thrusting and that the folds are subsequently "cross-folded" by broad Devonian folding (see next section). In light of these facts it is postulated that these small warps with east-west axes are genetically related to the episode of Silurian thrusting.

3.3.1.3. Devonian Folding: All rocks of the Buchans area, except Carboniferous strata, have been affected by open Devonian folding. This event produced a weak, subvertical, spaced, fracture cleavage in most lithologies, the strength of which varies according with location. Many of the more competent lithologies do not bear this cleavage.

The cleavage generally strikes northeast and

is axial planar to a broad synclinal structure whose axis can be traced from the Buchans Group, through the Topsails Granite and into the Springdale Group (see Section 2.3.). Northwest dipping and facing sequences on the southeast side of Red Indian Lake appear to constitute a deeper structural level exposed on the south limb of this same broad structure.

Flattening related to this deformation is negligible to not detectable throughout the study area. Fragments within pyroclastics, breccias and conglomerates throughout the Buchans Group show no visible effects of strain. Beyond the study area to the southwest, deformation increases in intensity within the Footwall Basalt to the point where noticeably flattened pillows occur.

3.3.2. Metamorphism of the Buchans Area: The metamorphism of the Buchans Group is a complex topic, a full discussion of which is beyond the scope of this thesis. The Buchans Group has evidently been subjected to burial metamorphism, local dynamothermal metamorphism during thrusting, a possible subsequent episode of re-equilibration of burial metamorphic facies and subsequent regional metamorphism during open folding. Whereas the details of these events must await future study, the general metamorphic environment is discussed below such that the effects of metamorphic alteration on rock chemistry can be reasonably

evaluated.

Greenschist facies rocks occur only in the lowermost exposed portions of the Footwall Basalt in the extreme southwest of the study area. Here, foliated and slightly flattened pillow lavas contain chlorite, albite, actinolite, calcite and epidote, mineralogy characteristic of the greenschist facies. This rapid southwestward increase in tectonic flattening of pillows and of metamorphic grade continues in a regular fashion beyond the study area.

Prehnite-pumpellyite facies metamorphism, (Coombs, 1960) is by far the most widespread metamorphic grade achieved by the Buchans Group. Metamorphic alteration is heterogeneous on outcrop scale with pyroclastics displaying considerably more extensive mineralogical change than that of associated massive basaltic flows and blocks in breccias. Within the latter lithologies calcic plagioclase phenocrysts are commonly partially altered to combinations of sericite, calcite, epidote and pumpellyite. Augite phenocrysts are normally fresher than plagioclase though incipient alteration to epidote, chlorite and calcite along fractures is common. Amygdules are filled with various combinations of quartz, chlorite, calcite, epidote, prehnite and pumpellyite, all of which may occur in a single thin section. Mafic pyroclastic rocks contain all of the above-mentioned metamorphic minerals in greater abundance than in flows, with the addition of minor authigenic

albite and sphene. Within felsic pyroclastics, illite, and mixed-layer illite-montmorillonite, and montmorillonite-chlorite commonly replace vitric material (Henley and Thornley, 1980).

Monomineralic metadomains as described by Jolly and Smith (1972) and Jolly (1974) are poorly developed in the Buchans Group. Numerous thin veinlets of probable synmetamorphic age transect the volcanic rocks in several areas but these are normally polymineralic (combinations of quartz, epidote, calcite, prehnite) and do not show gradational transitions into albite basalts as do metadomains. Nevertheless, these fractures probably acted as important fluid channelways during metamorphism.

Locally, near the top of the stratigraphic sequence, zeolite-like minerals fill amygdules in massive basaltic flows. Other amygdules in the same rock are filled with epidote, calcite, prehnite and pumpellyite and plagioclase phenocrysts display partial alteration to pumpellyite. The occurrence of zeolite (?) in these prehnite-pumpellyite facies rocks attests to the sluggish reaction rates so typical of low grade metamorphic assemblages and concurs with the observation by Jolly and Smith (1974) that the stratigraphic range of secondary void filling minerals is commonly somewhat greater than the range in which they replace rock forming minerals.

The presence of the zeolite implies that the stratigraphic top of the Buchans Group existed near the boundary between zeolite and prehnite-pumpellyite metamorphic facies. The lack of pervasive low grade metamorphic alteration (such as that described by Smith, 1968) and the lack of equilibrium even on thin section scale (as evidenced by the above-mentioned zeolites) further implies that the duration of metamorphism of the Buchans Group was insufficient to allow reactions to proceed to completion.

Henley and Thornley (1980) have compared the metamorphic mineral assemblages of the Buchans Group to thoroughly studied areas elsewhere, and concluded that metamorphic temperatures were in the  $260^{\circ} - 300^{\circ} \text{C}$  range.

## CHAPTER 4

### ORE DEPOSITS OF THE BUCHANS GROUP

#### 4.1. Introduction

The ore deposits of the Buchans Group and their host rocks are described in this chapter. Three genetically related ore types occur at Buchans, i.e., 1) stockwork ore, 2) in situ ore, and 3) mechanically transported ore. All three types of mineralization are known to occur in two distinctly separate horizons, the Lucky Strike ore horizon sequence and the Oriental ore horizon sequence.

In terms of economic importance, in situ and transported ores are of approximately equal significance and together have provided about 98% of production. Large tonnages of subeconomic stockwork mineralization are present but only local zones attain ore grade.

The distribution of the various ores is shown in Fig. 4-1. Tonnages, grades and other features of the Buchans ore zones are summarized in Table 4-1.

#### 4.2. Intermediate Footwall and Stockwork Mineralization

4.2.1. General Comments: The Intermediate Footwall is a thick, localized volcanic sequence which, beyond the area of stockwork mineralization and alteration, consists




Figure 4-1: Distribution of in situ (solid black) and transported fragmental (stippled) mineralization at Buchans.



POOR COPY  
COPIE DE QUALITEE INFERIEURE

| Orebody                           | Gross Tonnage | Grade (Mill Heads) |                 |      |       |       | MINERAL PERCENTAGES BY WEIGHT <sup>2</sup> |      |      |      |      |      | Sulfide Plus Barite |     |
|-----------------------------------|---------------|--------------------|-----------------|------|-------|-------|--------------------------------------------|------|------|------|------|------|---------------------|-----|
|                                   |               | Au <sup>3</sup>    | Ag <sup>3</sup> | Cu   | Pb    | Zn    | Fe                                         | Cp   | Ga   | Si   | Py   | Ba   |                     |     |
| Lucky Strike Ore Horizon Sequence |               |                    |                 |      |       |       |                                            |      |      |      |      |      |                     |     |
| In Situ Ores                      |               |                    |                 |      |       |       |                                            |      |      |      |      |      |                     |     |
| Lucky Strike Main                 | 4,752,373     | .050               | 3.29            | 1.67 | 8.42  | 18.36 | 9.3(8.0)                                   | 4.8  | 9.7  | 27.4 | 14.0 | 26   | 81.9                |     |
| Lucky Strike - New Year           | 803,112       | .041               | 3.23            | 1.40 | 9.75  | 18.76 | 5.5(4.7)                                   | 4.0  | 11.3 | 28.0 | 7.4  | 26   | 76.7                |     |
| Sub-total In Situ                 | 5,555,485     | .049               | 3.28            | 1.63 | 8.61  | 18.42 | (7.5)                                      |      |      |      |      |      |                     |     |
| Transported Ores                  |               |                    |                 |      |       |       |                                            |      |      |      |      |      |                     |     |
| Lucky Strike - North              | 620,510       | .049               | 3.58            | 0.46 | 4.54  | 8.20  | 1.8(1.5)                                   | 1.3  | 5.2  | 12.2 | 2.4  | 55   | 76.1                |     |
| Two Level                         | 328,596       | .047               | 3.64            | 0.50 | 4.56  | 8.02  | 2.5(2.2)                                   | 1.4  | 5.3  | 12.0 | 3.8  | 55   | 77.5                |     |
|                                   | 949,106       | .048               | 3.60            | 0.47 | 4.55  | 8.14  | (1.7)                                      |      |      |      |      |      |                     |     |
| Rothermere #1                     | 3,402,000     | .033               | 3.94            | 1.15 | 7.73  | 12.68 | 4.3(3.8)                                   | 3.3  | 8.9  | 18.9 | 6.0  | 31   | 68.1                |     |
| Rothermere #2                     | 195,615       | .034               | 3.59            | 1.44 | 7.65  | 13.73 | 6.6(5.7)                                   | 4.2  | 8.8  | 20.5 | 9.5  | 30   | 73.0                |     |
|                                   | 3,597,615     | .033               | 3.92            | 1.16 | 7.72  | 12.74 | (3.9)                                      |      |      |      |      |      |                     |     |
| MacLean                           | 3,653,000     | .028               | 3.84            | 1.13 | 7.46  | 13.50 | 4.3(3.8)                                   | 3.3  | 8.6  | 20.1 | 6.0  | 24   | 62.0                |     |
| Sub-Total Transported             | 8,199,721     |                    |                 |      |       |       |                                            |      |      |      |      |      |                     |     |
| Oriental Ore Horizon Sequence     |               |                    |                 |      |       |       |                                            |      |      |      |      |      |                     |     |
| In Situ Ores                      |               |                    |                 |      |       |       |                                            |      |      |      |      |      |                     |     |
| Oriental #1                       | 2,738,664     | .061               | 3.93            | 1.71 | 8.47  | 15.80 | 6.0(5.1)                                   | 4.9  | 9.8  | 23.6 | 7.7  | 24   | 70.0                |     |
| Old Buchans East                  | 133,353       | .067               | 4.55            | 1.65 | 7.57  | 14.27 | 10.0(8.5)                                  | 4.8  | 8.7  | 21.2 | 15.1 | 24   | 73.8                |     |
| Old Buchans West                  | 19,907        | .045               | 3.00            | 1.70 | 10.40 | 16.80 | 6.0(5.1)                                   | 4.9  | 12.0 | 25.0 | 7.7  | 24   | 73.6                |     |
| Sub-Total In Situ                 | 2,891,924     | .061               | 3.95            | 1.70 | 8.44  | 15.73 | (5.2)                                      |      |      |      |      |      |                     |     |
| Transported Ores                  |               |                    |                 |      |       |       |                                            |      |      |      |      |      |                     |     |
| Oriental #2                       | 928,863       | .046               | 6.15            | 0.76 | 6.20  | 9.41  | 3.2(2.8)                                   | 2.2  | 7.2  | 14.0 | 4.6  | 30   | 58.0                |     |
| Old Buchans Conglomerate          | 72,763        | .045               | 3.70            | 0.76 | 5.88  | 9.47  | 2.6(2.3)                                   | 2.2  | 6.8  | 14.1 | 3.5  | 31   | 57.6                |     |
| Sub-Total Transported             | 1,001,626     | .046               | 5.97            | 0.76 | 6.18  | 9.47  | (2.7)                                      |      |      |      |      |      |                     |     |
| Total In Situ                     | 8,447,409     | 48%                | .053            | 3.51 | 1.66  | 8.55  | 17.50                                      | 6.74 | 4.8  | 9.8  | 26.0 | 14.4 | 25                  | 80% |
| Total Transported                 | 9,201,346     | 52%                | .034            | 4.08 | 1.04  | 7.12  | 12.20                                      | 3.51 | 3.0  | 8.3  | 18.2 | 7.5  | 30                  | 67% |
| Total Gross Tonnage               | 17,648,756    | 100%               | .043            | 3.81 | 1.33  | 7.81  | 14.74                                      | 5.06 | 3.8  | 9.0  | 22.0 | 8.2  | 28                  | 73% |
| Total Milled To December 31/78    | 17,426,000    |                    | .043            | 3.69 | 1.34  | 7.60  | 14.62                                      |      |      |      |      |      |                     |     |

1. Fe in sulphide assays from drill holes. Assay in brackets based on 15% dilution to conform more closely with mill assays.
2. Percentages of major ore-forming minerals (chalcopyrite, galena, sphalerite, pyrite, barite) as calculated from grades.
3. Au and Ag in oz./ton.

Table 4-1: Ore deposits of the Buchans Area




Plate 4-1: Stockwork mineralization, Lucky Strike area.  
Top; chalcopyrite-pyrite veinlets cutting sulphide-impregnated chloritized host. Bottom; silicified and chloritized stockwork.

mainly of andesitic pyroclastics and breccias. However, within 1 km of Lucky Strike, the unit gradually changes character into a complex mound of pyroclastics and breccias ranging from basaltic to rhyolitic composition. These rocks have undergone progressive alteration and mineralization which increases in intensity towards the Lucky Strike area and obscures the original character of the host rocks. The Intermediate Footwall is the host for several zones of stockwork mineralization and ore and at least two small syngenetic massive sulphides.

Strong stockwork mineralization consists of a network of quartz-sulphide veins and veinlets cutting strongly altered and sulphide-impregnated host rocks (Plate 4-1). The largest concentration of stockwork mineralization in the Buchans area occurs at Lucky Strike. Fine to coarse grained euhedral pyrite is the most abundant sulphide within the stockwork with much lesser quantities of chalcopyrite, sphalerite and galena. Quartz, chlorite, Barite and calcite are the dominant gangue minerals. Compared to the overlying massive sulphide, stockwork mineralization is more pyrite-rich and, in terms of metal ratios, contains proportionally more Cu. Only small zones of stockwork contain economic quantities of base metal sulphides.

In general, the development of the Intermediate Footwall in the Lucky Strike area is one of continuous

volcanism with concurrent mineralization and alteration. Within the overall alteration envelope, internal centers of hydrothermal activity migrate both areally and stratigraphically with time, resulting in several partially overlapping alteration zones. The major period of stockwork mineralization which fed Lucky Strike entirely post-dates this earlier mineralization and alteration history.

The strongly altered and mineralized zone is shaped like a thin flat-topped mushroom with most of the underlying stem cut off by faults. The zone of strongest alteration and mineralization is crudely concordant with stratigraphy and is approximately 600 m wide and up to 100 m thick. The overall alteration zone covers an area of several square kilometers but rarely exceeds 100 m thickness, and is thus more blanket-shaped. This distribution pattern of stockwork mineralization and alteration is unlike many massive sulphide deposits worldwide (c.f. Sangster, 1972).

In plan view, three roughly concentric zones consisting of varying degrees of alteration and sulphide mineralization can be outlined (Fig. 4-2). Peripheral zones of the alteration halo are characterized by a clay mineral-pyrite assemblage, the intensity of which increases inward towards zones of stronger alteration. Henley and Thornley (1980) have identified illite as the most abundant clay




Figure 4-2. Distribution of alteration zones within the Intermediate Footwall.

mineral with lesser montmorillonite and mixed layer clays. This peripheral alteration zone passes gradually into a zone of strong clay mineral-pyrite alteration with local silicification, chloritization and minor base metal mineralization (Fig. 4-2). This in turn passes gradually into a zone of strong silicification and chloritization containing consistently in excess of 1% base metals. This latter zone is confined to areas immediately adjacent to in situ mineralization and contains all the significant stockwork occurrences (Fig. 4-2). This generalized alteration pattern is presently one of the major criteria used in guiding exploration for deeply buried in situ deposits.

Kowalik et al. (1980) have recently completed a stable isotope study of the stockwork zone. They conclude that: 1) sea water was the dominant fluid involved in the alteration and mineralizing process, 2) sea water sulphate was the source of sulphur in the deposits, 3) temperatures in the stockwork zone range from 140° C to 370° C, with temperatures increasing both laterally and vertically towards Lucky Strike.

4.2.2. Stratigraphy and Mineralization in the Lucky Strike Area: The early history of the Intermediate Footwall has largely been obliterated by subsequent explosion and alteration. The lowermost member of the formation above the fault base consists of tightly packed, coarse polyolithic

(in varieties of intermediate composition) breccias, locally pumiceous breccias interbedded with minor amygdaloidal flows and related breccias of probable basaltic to andesitic composition. Some fragments in these breccias display altered rims, a feature which is common at higher stratigraphic levels in the pile and there can be shown to be caused by alteration along fractures with subsequent explosion (see Section 4.3.). Thus the earliest consolidated rocks of the formation were fractured, altered and exploded to form poly lithic breccias. These breccias were subsequently strongly and pervasively altered. This alteration varies in intensity, composition and style and consists of varying degrees of silicification, sericitization, chloritization, pyritization, clay mineralization and less common carbonate alteration. Alteration occurs both as veins and as pervasive alteration away from veins such that the original breccia texture is locally obscured. Introduction of vein material has locally caused brecciation of the host rock. For the most part, sulphide mineralization associated with this alteration is pyrite, though minor sphalerite, galena, chalcopryite and less abundant barite are common. No massive sulphide mineralization is known to be related to this episode of alteration and mineralization.

Overlying the earlier mineralized and altered portion of the Intermediate Footwall Formation is a second unit which is characterized by the presence of abundant



altered fragments derived from the underlying unit. This unit is the most extensive unit of the altered Intermediate Footwall. Polyolithic breccias are the most abundant constituent of this unit and contain numerous varieties of strongly altered fragments of original basaltic to rhyolitic composition. Evidence that these breccias are in part derived from the previously mineralized and altered breccias includes 1) the presence of strongly altered fragments with variable degrees and types of alteration, locally in a much less altered matrix, 2) the presence of fragments composed of coarse euhedral pyrite (interpreted as vein pyrite from the underlying unit), 3) the presence of numerous fragments composed entirely of black chlorite and euhedral pyrite, similar to some pervasively altered material of the underlying unit, 4) some altered fragments contain pyrite-sphalerite-galena-chalcopyrite mineralization in thin fractures and veins which are terminated at fragment boundaries.

A number of other fragment types are present in the breccias including pink rhyolite fragments, locally with altered rims. The polyolithic breccias commonly contain a visible vitric component in the matrix and normally contain varying proportions of angular to flattened long tube pumice. All variations exist between vitric polyolithic breccias (pyroclastic breccias) and breccias composed almost entirely of tightly packed pumice fragments



and wispy shreds (pumice breccias). Recognizable vitric ash horizons have locally escaped pervasive alteration and obliteration of primary texture. The breccias are normally moderately to well packed and poorly sorted, with average fragment size usually less than 5 cm. Breccia beds locally display a crude size grading over a few meters. The presence of abundant vitric material in the breccias suggests that the brecciation of the previously altered unit was accomplished by explosion related to violent frothy eruption. The products of eruption and exploded lithic fragments were subsequently redeposited as homogenized subaqueous density flows.

This second member of the Intermediate Footwall suffered locally intense alteration and stockwork mineralization which fed two small stratiform massive sulphide bodies; the Rothermere Footwall Orebody and the West Orebody. Tonnages and grades of these deposits are given in Table 4-1. The Rothermere Footwall Orebody is a small, thick, massive polymetallic lens concordant with bedding and some 15 m below the upper contact of the formation. The footwall to the deposit consists of strongly altered polyolithic breccia locally containing vein sulphide clasts. The alteration varies from strong sericitization, pyritization and clay mineralization although the most strongly altered and veined portions consist of virtually 100% chlorite and disseminated and vein euhedral pyrite and traces

of base metal sulphide. Elsewhere, intense silicification is dominant, emanating from milky, pyritic, fine grained quartz veins and cutting the above-mentioned earlier alterations. Alteration evidently continued after the major period of sulphide deposition, as hangingwall breccias are moderately altered and contain minor disseminated sulphide and local sulphide veins. The deposit itself is lens-shaped, high grade and polymetallic. Ore mineralogy, in order of abundance, consists of sphalerite, galena, chalcopryrite, pyrite and tetrahedrite. Dark grey to white barite occurs throughout the deposit but, towards the top, becomes increasingly common and forms a massive mainly baritic cap. Apart from the obvious barite zoning, no base metal zonation is evident within the deposit. A unique feature of the sulphide portion of the deposit is the occurrence of numerous rounded chalcopryrite-pyrite aggregates with diffuse outlines. These may represent sulphide flocculates which settled into the accumulating sulphide mud.

The West Orebodies, a group of small sulphide deposits, occur within the Intermediate Footwall approximately 150 to 250 m southwest of Lucky Strike Orebody. Their stratigraphic position with respect to the Rothermere Footwall Orebody is uncertain. The footwall of the deposits consists of strongly altered and-pyritized breccias and tuffs, locally of probable original dacitic composition. Minor

pyritic and base metal-rich veins cut this altered unit. The contact with the ore zone is characterized by a rapid increase (normally over less than one meter) of disseminated and vein sulphide until a sulphide mass consisting dominantly of coarse euhedral pyrite with finer grained chalcopyrite, galena and sphalerite is attained. This sulphide mineralization is associated with strong calcite alteration of completely obliterated host rock. Commonly there occur zones of very fine grained massive high grade Pb-Zn ore which display considerable textural similarity to syngenetic ores in the major massive sulphide deposits. These massive sections are locally cut by coarse grained pyritic stringers. The deposit is one of the few areas in the Buchans camp where stockwork ore and related syngenetic massive sulphide can be seen to merge. The average base metal content of the entire deposit (Table 4-1) displays characteristics of both stockwork and syngenetic sulphide. The hangingwall contact of the deposit is normally sharp and consists of strongly altered tuffs and breccias probably of original dacitic composition. Pyritic veins, similar to that in the stockwork portion of the deposit, locally cut hangingwall rocks.

The Engine House Orebody, the largest deposit of stockwork ore, occurs approximately 200 m southeast of Lucky Strike. Tonnage and grade figures are given in Table 4-1. The deposit occurs entirely within the Intermediate



Plate 4-2: Green mariposite-like sericite after plagioclase; a typical feature of proximal areas of the clay mineral alteration envelope.

Footwall and no connection is known between this strong stockwork mineralization and Lucky Strike Orebody. Both the footwall and hangingwall contacts of the deposit are marked by a rapid gradation from normally altered and mineralized Intermediate Footwall tuffs and breccias into pervasively altered and mineralized ore. The deposit consists dominantly of coarse euhedral vein and disseminated pyrite with local enriched zones of sphalerite, galena, chalcopyrite and tetrahedrite in a pervasively silicified and/or chloritized host (Plate 4-1). On the whole, silicic alteration is dominant, although local zones of 100% chlorite-pyrite with minor base metal are present. Barite is of minor importance and is generally more closely associated with silicic than chloritic alteration. Calcite is minor but ubiquitous throughout the deposit.

The character of the immediate host rock for the Engine House deposit has been largely obliterated by pervasive alteration and mineralization. However, local occurrences of somewhat less obscured host are known. Most commonly, these occur as fragments of greyish aphanitic altered flow (?) of compositions originally ranging from basalt to rhyolite. Relict plagioclase in these flows is commonly altered to waxy green illite or locally to a bright green mariposite-like mineral (Plate 4-2). Samples of the latter material submitted for X-ray diffraction are apparently sericite and show no chromium enrichment (R. Henley, pers.

comm., 1976). Larger fragments of flow are locally brecciated by intrusion of quartz-sulphide veins producing breccia aggregates consisting of fragments with altered rims.

The second unit of the Intermediate Footwall and the related small massive sulphide orebodies are conformably overlain by another sequence of altered polyolithic breccias and pumiceous tuffs. These are interbedded with fine grained, pyritic, light grey, tuffaceous, siliceous siltstone and wacke, in part derived from reworking of previously deposited breccias. Detrital sulphides (especially pyrite) are characteristic of the unit. The unit displays a gradual facies change from dominantly breccia with minor intercalated sandy tuff and tuffaceous siltstone in the Lucky Strike area, to dominantly tuffaceous pyritic siltstone in the area 500 m east of Lucky Strike (e.g., DDH 2520). The breccias of this unit are similar to underlying breccias and will not be discussed further. The interbedded and facies-equivalent tuffaceous siltstone and siliceous wacke are generally light grey and normally massive with bedding faint to absent. These tuffs are locally vitric and may grade into subaqueous pumice flows similar to those described by Yamada (1973). Graded beds ranging from fine grained breccia to pyritic siltstone occur locally, as does small scale cross bedding. The unit locally contains a few isolated pebbles of strongly altered material derived from the underlying breccias, small (2 mm) pyritic pebbles and

reddish rhyolite pebbles. Minor silica-pyrite-carbonate veining is common and the entire unit has been slightly, but pervasively silicified.

#### 4.3. Lucky Strike Ore Horizon Sequence

4.3.1. Footwall Rocks: Rhyolite with flanking and overlying pyritic siltstone and dacitic tuff form the base of the Lucky Strike ore horizon sequence (see Chapter Three) and overlie the earlier altered and mineralized Intermediate Footwall. The stockwork which fed Lucky Strike Orebody post-dates the earlier alteration in the Intermediate Footwall and the basal rocks of the ore horizon sequence. Alteration related to this mineralization consists of strong silicification and chloritization, similar to the previous alteration assemblages. Pyritic and baritic base metal-bearing veins cut the altered and sulphide impregnated host rocks. Henley and Thornley (1980) have identified the Black chlorite of the stockwork as clinochlore-penninite, a variety more magnesian than ambient Buchans Group chlorites.

Lucky Strike Orebody overlies the stockwork though the critical contact zone is poorly known due to shearing and removal of this material from the Lucky Strike glory hole.

4.3.2. Lucky Strike Orebodies: The Lucky Strike Orebodies consist of a number of contiguous ore zones which together form the largest single body of massive sulphide within the Buchans area. They were discovered in suboutcrop in 1926 as a result of equi-potential prospecting by Hans Lundberg. From 1928, when mining began, until 1957 the mine produced 6,175,995 tons of ore with an average grade of 26%  $\text{BaSO}_4$ , 17.40% Zn, 8.20% Pb, 1.51% Cu, 3.31 oz/ton Ag and 0.049 oz/ton Au. The average mineral percentages by weight, as derived from the grade, are given in Table 4-1. It consists both of in situ massive sulphide and mechanically transported sulphides intimately associated with stockwork mineralization in the underlying Intermediate Footwall. Information regarding the ore itself and contact relations with surrounding rocks is scarce due to the early practice of assaying of entire drill cores. Nevertheless, valuable information can be gained from the Lucky Strike open pit and from excellent early descriptions of the ore (e.g., George, 1937).

The ore overlies either altered dacitic crystal-vitric tuff or strongly altered and mineralized stockwork of the Intermediate Footwall. Contacts are generally not cored due both to shearing and to generally poor core recovery of most early drilling. Locally, conformable contacts of ore on dacitic tuff or of ore on Intermediate Footwall are visible. The deposit is conformably overlain



by dacitic crystal-vitric tuff or by diabase sills .  
(possibly filling fault planes).

In cross section, the deposit has a thick lens shape (Fig. 3-2), a characteristic peculiar to Lucky Strike and in keeping with its largely in situ origin. It consists of a zoned aggregate of sphalerite and barite with lesser galena, chalcopyrite and pyrite with minor tetrahedrite and bornite. Secondary smithsonite, native silver, and Zn, Cu, Pb and Fe sulphates occurred locally. George (1937) noted the presence of mineralogical zoning within the deposit and the following description is derived partly from his work. The Main Orebody consisted of approximately 90% Pb-Zn ore (black ore) and 10% Cu-Zn ore (yellow ore) with the latter confined mainly to the base of the deposit. In general, Cu-Zn ore contained more pyrite, chalcopyrite and sphalerite and less barite and galena than Pb-Zn ore. Gold and silver were concentrated towards the hangingwall (i.e., in the Pb-Zn ore). Although no ferruginous chert bed is known in the hangingwall of the deposit, it is locally overlain by red (hematitized) barite containing minor pyrite and commonly also by hematitized dacitic crystal-vitric tuffs. This zoning bears a remarkable similarity to that of many of the Kuroko deposits.

Ore textures, as preserved in a few drill cores and in exposures in the Lucky Strike Pit, indicate that the



Plate 4-3: Typical in situ wispy and streaky very high grade ore. Yellow areas are chalcopyrite, light grey are fine grained galena-sphalerite and dark grey to black is barite.

bulk of the deposit consisted of fine grained massive structureless ore and of "streaky ore" (Plate 4-3). These latter ores consist of thin, discontinuous wispy yellow streaks of chalcopyrite-pyrite-rich ore within more abundant black galena-sphalerite-rich ores. Elliston (1963) has described and illustrated similar textures within mudstones. He calls the process responsible for this texture "fluidal wisping" and gives evidence to suggest that it forms as a result of lateral flow in unconsolidated watery plastic muds. In view of evidence presented in subsequent paragraphs, a similar origin for streaky ore at Buchans is plausible.

Well developed compositional banding was not common at Lucky Strike but, according to Relly (1960), it existed locally. Portions of Lucky Strike Orebody consist dominantly of discrete sulphide and sulphate fragments associated with the breccia-conglomerate and "granite conglomerate". The occurrence of these ores is discussed in detail in the following sections.

4.3.3. Ore Horizon Breccias: The ore horizon of many worldwide massive volcanogenic sulphide deposits is commonly marked by a weakly mineralized cherty tuffaceous exhalite horizon (e.g., Sangster, 1972; Ridler, 1973; Sangster and Scott, 1976). In contrast the Lucky Strike and Oriental ore horizons are characterized by a complex series of beds composed of coarse polyolithic rudaceous rocks

with associated volcanic arenites and wackes. Within the rudaceous fraction all gradations exist from volcanic breccia to volcanic conglomerate with breccia-conglomerate being most abundant (terminology after Fisher, 1960). Because of the lithologic similarity of these rocks in the Lucky Strike and Oriental ore horizon sequences, they are described together in this section.

The horizons occupy an area in excess of 3 sq. km in the mine area and are intimately associated with the transported massive sulphide orebodies. Thickness varies sympathetically with underlying paleotopography from zero on paleotopographic highs to over 100 m in paleotopographic lows. Within the unit there is a common gradation from underlying volcanic breccia and breccia-conglomerate to overlying granitoid-bearing volcanic breccia-conglomerate (hereinafter referred to as "granite conglomerate"). This sequence is best displayed in the Oriental area, particularly in the vicinity of Oriental No. 2 Orebody.

The breccia-conglomerate beds overlie either pyritic siltstone or dacitic crystal-vitric tuff. They consist of a poly lithic assemblage of rounded to angular fragments ranging from silt size to fragments over two meters in diameter. Scores of locally derived fragment types can be identified and can be grouped into eight basic lithologic categories, i.e.,

- 1) various altered lithologies of the Intermediate Footwall including stockwork and vein sulphide fragments,
- 2) pyritic siltstone and wacke fragments,
- 3) varieties of massive rhyolite fragments,
- 4) varieties of ore horizon dacitic tuff as fragments,
- 5) fragments of basalt, andesite, jasper chert, etc., possibly derived from the Footwall Basalt,
- 6) rare fragments of volcanic arenites and wackes derived from the breccia-conglomerate itself
- 7) varieties of rounded granitoid pebbles, cobbles, and boulders, and
- 8) fragments of barite and massive sulphide ore (black ore and yellow ore).

The proportion of these various fragment types varies dramatically within individual beds and between separate beds such that the breccia-conglomerate is locally composed almost entirely of one lithologic type. The transported orebodies (e.g., MacLean) are an example of breccias composed mainly of massive sulphide fragments and are one of several facies within the unit. However, more commonly the breccia-conglomerate is an intimate mixture of numerous fragment types with those of types one to four (above) being most common. It is anomalous, and perhaps significant, that fragments of all lithologies underlying the breccia-conglomerate occur within the unit except fragments that can be confidently related to the Footwall





Plate 4-4: Typical polyolithic granitoid-bearing breccia conglomerate. Both photos from hangingwall of Old Buchans Conglomerate Orebody.





Plate 4-5: Granodiorite cobble in ore breccia; MacLean Extension deposit, 20-4 drift. Cobble is texturally identical to the Feeder Granodiorite.

Arkose.

The "granite conglomerate" (Plate 4-4) conformably and gradationally overlies the breccia-conglomerate and is especially well developed in the Oriental area. The "contact" between the two is normally characterized by a gradual stratigraphic increase in plutonic boulders until a breccia-conglomerate composed of approximately 20% (average) granitoid boulders is achieved. In contrast to other lithologies, granitoid boulders are invariably well rounded with moderate to high sphericity. Numerous granitoid types exist, ranging in composition from (rare) diorite to quartz monzonite but pink to grey varieties of granodioritic composition normally predominate. Granodiorite boulders range in size from up to a maximum of 5.8 m (Swanson and Brown, 1962) but those of 25 cm and less are most common. Granitoid boulders are commonly the largest fragment type within the "granite conglomerate". Many granitoid boulders are texturally identical to the Feeder Granodiorite (Plate 4-5).

Textures and structures of the "granite conglomerate" and breccia-conglomerate are similar and will be discussed together below. The basal contact of the breccia-conglomerate with the underlying siltstone or dacite is normally sharp but locally characterized by scouring or channeling of underlying lithologies. Initial deposits of the unit vary locally from fine grained volcanic arenites to coarse



breccia-conglomerate. The breccia-conglomerate is normally polyolithic, coarse grained (average 2-20 cm) and very poorly sorted. Only locally can a bimodal size distribution be recognized. Most fragments are angular to subrounded and any given fragment type can vary from angular to rounded. Fragments of Intermediate Footwall, pyritic siltstone and dacite are rarely rounded whereas rhyolite fragments (though normally subangular) are more commonly rounded than the previous lithologies. Only granitoid boulders are everywhere rounded. Sphericity of all fragment types (except granodiorite) is low. Packing varies dramatically but the coarser units are normally matrix-supported, to somewhat fragment-supported. Where elongate tabular fragments are present a chaotic clast fabric is locally visible, though more normally clasts are elongate subparallel to the base of the unit, thus defining a crude bedding. In rare underground bedding plane exposures in MacLean, the long axes of pebbles parallel the long axis of the orebody. A weak imbrication of larger elongate fragments is locally visible and defines current directions from the vicinity of the Buchans Inlier. Although sorting is normally poor, some thick beds display a gradual vertical grading over several meters. Elsewhere, chaotic unsorted breccia beds several meters thick grade rapidly over the uppermost meter from coarse breccia to fine grained arenite or wacke. In some instances some beds are graded inversely at the base and normally graded at the top (e.g., DDH 1440).

The matrix of the breccia-conglomerate consists of a heterogeneous arenaceous mixture of quartz, plagioclase and small lithic fragments similar in composition to the larger breccia fragments. Fine grained detrital pyrite is common and small massive sulphide fragments occur locally. Pumice shreds are a rare local occurrence. Thin beds (1 m and less) of arenaceous material are locally interbedded with breccia-conglomerate and may be massive, or display a crude stratification or be graded. Cross stratification has not been observed.

Further comments on the ore horizon breccias accompany and follow description of the transported ores.

4.3.4. MacLean, Rothermere Two Level Orebodies and Lucky Strike North; transported ores: The MacLean, Rothermere, and Two Level Orebodies occur in a linear arrangement extending northwest from Lucky Strike (Fig. 4-1). Two Level Orebody was discovered shortly after the discovery of Lucky Strike, Rothermere in 1947 and MacLean (formerly Rothermere No. 4) in 1950. MacLean and Rothermere have provided the bulk of production during recent years. Tonnage and grade figures are given in Table 4-1. Because of the gross similarity of the occurrence of these deposits, they are discussed together in this section.

The orebodies occur within dacitic crystal-vitric tuffs of the Lucky Strike ore horizon sequence at a

stratigraphic horizon very near to that of the Lucky Strike deposit (Fig. 3-2). They occur within paleotopographic depressions within a sinuous paleo-channel and are intimately associated with pyritic siltstone, breccia-conglomerate and "granite conglomerate". The ores comprise several breccia beds composed dominantly of sulphide fragments. MacLean Orebody consists of at least four major, distinct breccia beds and Rothermere, of at least three. Lucky Strike North, Two Level, Rothermere No. 2 and a bed along the northern edge of MacLean are characterized by an abundance of granitoid boulders and barite and are probably genetically linked. This linear arrangement of ores lies slightly northward and parallel to the later major Rothermere-MacLean ores. Ore mineralogy consists of a high grade assemblage of low-Fe sphalerite, galena and barite with lesser chalcopryite, pyrite and minor tetrahedrite and bornite (Strong, 1980). Numerous lines of evidence presented in the following paragraphs indicate that these orebodies consist of mechanically transported sulphide fragments deposited from rapidly moving subaqueous density flows derived from the Lucky Strike Orebody.

The orebodies are a series of elongate tabular deposits conformable within the enclosing host rocks. This conformability is evident both in cross section (Fig. 3-2) and on a detailed scale underground where primary features in the ores may be seen to parallel ore-host contacts and



Plate 4-6: Typical low grade transported ore breccia. Two high grade black ore fragments in centre. Note also siltstone fragment with altered rim and "mariposite"-bearing fragment at bottom right. Large fragments at upper left and right are varieties of rhyolite.

bedding in the host rocks. The author has seen no examples where ore transgresses the host rocks, an observation also recorded by Relly (1960, p. 95). The ore rests directly on either dacitic crystal-vitric tuff, pyritic siltstone or "barren" breccia-conglomerate. The footwall contact is normally knife sharp where ore lies on dacite or siltstone and may be sharp or gradational where ore lies on breccia-conglomerate. Any given orebody may rest on all these lithologies throughout its length. The hangingwall contacts of ore against dacitic crystal-vitric tuff or late diabase sills are sharp whereas contacts with overlying breccia-conglomerate or "granite conglomerate" may be sharp or gradational. Where gradational contacts of ore against breccia, in the footwall or hangingwall are observed the "contact" is marked by an increase in discrete high grade sulphide or barite fragments towards the orebody. These fragments texturally occupy a position in the breccias identical to other lithic fragments (Plate 4-6) and are considered to have been deposited at the same time and by the same mode as the lithic fragments. Locally where ore rests on siltstone or dacitic tuff, scour channels within the underlying rocks are visible. In some places at the hangingwall contact, very fine grained clastic ores are load-casted into hangingwall dacitic crystal-vitric tuff. Late diabase dykes locally cut the ore, display chilled margins and have caused varying degrees of contact metamorphism in the ore.




Plate 4-7: High grade transported ore. Yellow ore fragments and darker, smaller, less distinct black ore clasts. Darker clast above coin is barite.





Plate 4-8: Plastically deformed ore fragments. Light grey irregularly shaped galena clast at right center adjacent to darker galena-sphalerite clast (center) with wispy, frayed edge.

The ore itself is a series of breccia beds composed of varying proportions of sulphide and lithic fragments (Plate 4-7). All fragment types which occur in the breccia-conglomerate and "Granite conglomerate" occur in the orebodies and vice versa. In effect then, the orebodies are members of the "granite conglomerate" - breccia-conglomerate horizon. The sulphides occur as discrete fragments from (probably) submicroscopic grains to boulders in excess of 2 m in diameter. Fragments less than 10 cm in diameter are most common. Many sulphide fragments display a distinct streaky aspect and rarely may be termed banded. Streaky banding within fragments is invariably truncated sharply at fragment margins and is commonly oriented randomly with respect to streaks in adjacent fragments, bedding in enclosing matrix and ore-host rock contacts. Some sulphide fragments vary from angular to rounded to tabular and were evidently coherent and competent fragments at the time of deposition. Tabular fragments are commonly aligned parallel to the margins of the orebody though locally a weak imbrication is visible (indicating transport directions from the southeast). Other sulphide fragments are elongate with wavy but sharp contacts with matrix along their long axes, but with wispy, frayed terminations along their short ends. These fragments are commonly moulded plastically around adjacent more competent sulphide and lithic fragments (Plate 4-8). Streaky banding in these






Plate 4-9: Coin rests on fragment of baritic fragmental ore within fragmental black ore matrix.

fragments normally parallels the long axis of the fragment and mimics any contortions of the fragment margin. These are interpreted as sulphide fragments deposited while still in an only partially indurated, plastic condition. The presence of ores composed of sulphide fragments indicates that induration of sulphides of the source orebody was rapid, a conclusion also reached by Spence (1975). They further indicate that the streaky sulphide banding was produced at, or shortly after, deposition and not by a later stage process.

Fragments composed largely or wholly of barite occur throughout the orebodies. In general, these fragments are better rounded and more spherical than the sulphide fragments and, in the case of MacLean and Rothermere, are concentrated towards the top of the orebodies where they form a discrete transported baritic cap. Spectacular fragmental baritic ore is also exposed in the Lucky Strike glory hole and consists of massive to crudely graded beds composed both of fine grained baritic and massive sulphide fragments and less common large (up to 5 cm) broken barite crystals. These clastic crystals indicate that at least portions of the barite were coarsely crystalline at deposition.

Another feature indicative of rapid lithification is the very rare occurrence of fragments composed of fragmental ore (Plate 4-9). This texture has been observed

only in the Lucky Strike glory hole, a testimony to the probable depositional friability of these boulders.

Lithic fragments in the ore are of identical nature and occurrence to those in the associated "granite conglomerate" and breccia-conglomerate, the only difference being that their proportion has been significantly diluted by the greater abundance of sulphide fragments. All lithic fragments (except granodiorite) can be confidently related to nearby rocks in the footwall. Lithic fragments display the same textures and structures in the ore as in the associated breccia-conglomerate and "granite conglomerate". Locally, lithic fragments in the ore display a crude vertical size grading but due to the difficulty in distinguishing ore fragments from matrix while underground, only rare samples of grading of sulphide fragments have been observed.

The matrix of the fragmental ores appears to consist both of more finely divided clastic particles and possibly of some sulphide chemical precipitate. Sulphide and lithic fragments form part of a completely gradational size sequence from less than 1 mm to the largest boulders. Some of this finer grained material is probably derived from attrition and comminution of larger fragments by collisions during density flow. Several polished slabs of fine grained ore exhibit a sand-sized clastic texture which is highlighted by fine grained lithic fragments




Plate 4-10: Fine grained bedded high grade clastic ore  
containing some larger silicate and sulphide  
clasts.






Plate 4-11: Siltstone fragments (S) with altered rims. Larger rhyolite (R) and altered Intermediate Footwall (I) fragments in top photo. Alteration rim on siltstone clast in bottom photo is broken.

(Plate 4-10). However, many samples of lithic-free, fine grained, high grade, streaky black ore show no evidence of clastic textures. All ore sulphides have been slightly annealed during low grade metamorphism and thus, destruction of silt sized (and finer) clastic textures during metamorphism has probably occurred.

If some chemically precipitated matrix sulphide exists it may have been deposited from heavy brines which collected in the depressions which host the ores (c.f. Sato, 1972).

One important feature common to lithic fragments in the ore, breccia-conglomerate and "granite conglomerate" is the presence of alteration rims on some fragments (Plate 4-11). This feature is most commonly developed in pyritic siltstone fragments, with less common development of rims on fragments of rhyolite and Intermediate Footwall and rare rims on dacitic tuff. Very rare rims have been observed on granitoid and sulphide fragments. Siltstone fragments with altered rims are of three types, i.e.,

- 1) fragments with altered rims and fresh cores
- 2) fragments with concentric alteration zones (rare), and
- 3) broken fragments with only partial alteration rims.

In addition to these types there occur siltstone fragments which have been homogeneously altered from rim to core and fragments which have undergone no alteration. In fragments of



Plate 4-12: Pyritic siltstone cut by fractures with adjacent alteration. See text and Fig. 4-3 for schematic and explanation.



Intermediate Footwall with altered rims, the alteration rim is superimposed on previously pervasively altered lithologies, indicating formation of the rim after a period of significant alteration. The similarity of the occurrence of these types of alteration rims to those in the Shitabanhi Orebodies in Japan (Lee et al., p. 61, 1974) is remarkable.

1  
The origin of these fragments has important implications regarding the mode of fragmentation of lithic, sulphide and sulphate fragments in the ore horizon. In several drill cores (e.g., DDH 1910) and underground the author has seen massive, competent, non-brecciated, bedded pyritic siltstone which is transected by hairline fractures in a reticulate pattern (Fig. 4-3, Plate 4-12). Extending in from these fractures is the same light greyish alteration visible in the rimmed fragments in the ore. It is evident that if this rock was broken into fragments along the hairline fractures, a breccia composed of fragments with altered rims would result. The author proposes that hairline fracturing, permeation by hydrothermal fluids and subsequent violent explosion are responsible for the rimmed siltstone, rhyolite and Intermediate Footwall fragments in the ore horizon. These same explosions may have been largely responsible for fragmentation of all other lithologies which occur as clasts in the ore horizon, including massive sulphide. In the case of the Kuroko deposits of Japan, there is evidence of deep-rooted explosions within the white rhyolite domes which have explosively






Figure 4-3: Sketch showing bedded siltstone transected by hairline fractures with alteration extending into the siltstone from the fractures. Fragmentation of this rock would produce breccia fragments with altered rims.

transported a variety of lithic fragments (including accidental basement fragments) to surface with subsequent deposition in the ore horizon (T. Satō, pers. comm. 1975).

The evidence presented in the preceding paragraphs suggests that the Lucky Strike North, MacLean, Rothermere and Two Level Orebodies and their related breccias were emplaced as sulphide fragments from rapidly moving subaqueous density flows. The most compelling evidence in favour of this interpretation is the universal presence of competent to plastic sulphide fragments with chaotically disoriented streaky banding truncated at the fragment margins. The presence of chaotic internal structure, to crude grading to imbrication, the presence of scour channels in underlying rocks and the close association with sulphide-bearing polyolithic breccia-conglomerate of density flow origin is further evidence in support of this hypothesis.

The interpretation of the ores as mechanically transported sulphides deposited from a series of subaqueous debris flows explains a large number of features, i.e.,

- 1) the tabular, elongate shape of the deposits in a south-east-northwest direction,
- 2) the occurrence of the ores in paleotopographic depressions within a paleo-channel,
- 3) the lack of directly associated stockwork mineralization (Fig. 4-2),

- 4) the lack of significant alteration (especially at MacLean) which can be confidently and consistently related to the presence of sulphide mineralization,
- 5) the progressive thinning of MacLean Orebody away from Lucky Strike,
- 6) the presence of coarse breccia beds on finer breccia beds,
- 7) the abundance and variety of lithic fragments in the ore and the presence of fragments with altered rims,
- 8) the occurrence of the sulphides at the same stratigraphic horizon as a major "in situ" deposit,
- 9) the presence of unmineralized siltstone and tuff beds within the ore,
- 10) the absence of well banded (bedded) massive sulphide,
- 11) the relative scarcity of yellow ore fragments,
- 12) the presence of a transported baritic cap and preferential accumulation (in the MacLean case) of barite towards the proximal (southeast) end of the deposit, and
- 13) the occurrence of metal ratios at MacLean and Rothermere which are consistent with their being derived from the upper portions of the zoned Lucky Strike Orebody.

The general mode of transport of the fragmental ores and related breccias can be deduced from previous description. The size of the breccia fragments, thickness of the breccia beds, distances of transport and scour channels at the base are indicative of a powerful and fast-moving transporting agent.

The generally poorly sorted, chaotic, and polyolithic nature of the breccias and ores is further evidence of deposition from rapidly moving flows. The occurrence of long axes of clasts parallel to the direction of flow indicates that clasts did not roll on the bed as in fluvial deposits (e.g., Rust, 1972) but instead were aligned in suspension by clast collisions similar to that proposed by Rees (1968). The author envisages transport by a series of rapidly moving debris flows, not unlike the submarine equivalents of landslides. Calculations by Davies and Walker (1974) on deposits of this type in Quebec indicate flow velocities on the order of tens of meters per second.

The following interpretive sequence of events at Buchans can be constructed from all the preceeding:

- 1) Near-surface explosions occurred in the vicinity of Lucky Strike during black ore formation. Exploded lithic and sulphide fragments collected on the slope and were dislodged by subsequent explosions, by earthquakes or simply by instability of the accumulation on the volcanic slope. The direction of mass flow was controlled by a topographic channel which existed in a northwesterly direction downslope from Lucky Strike. Density flows scoured the underlying beds picking up additional fragments during transport. Debris was preferentially deposited in depressions along the path of flow.

- 2) Black ores continued forming at Lucky Strike and subsequent explosions resulted in further debris flows.
- 3) As baritic ores formed at Lucky Strike, subsequent explosions resulted in transport of increasingly more baritic debris flows which collected on top of underlying transported black ores.
- 4) Explosions in the vicinity of Lucky Strike ceased.
- 5) Red hematite-rich barite formed a cap on Lucky Strike and formation of the orebodies was complete.

4.3.5. Clementine Prospect: The Clementine Prospect is a significant transported sulphide deposit occurring on the Lucky Strike ore horizon, approximately 4 km west of Lucky Strike (Fig. 4-1). The deposit has recently been studied by Calhoun (1979) and Calhoun and Hutchinson (1980) and further description will not be undertaken here.

#### 4.4. Oriental Ore Horizon Sequence

4.4.1. Oriental No. 1 Orebody: The Oriental No. 1 Orebody was discovered in 1926 and produced 2,738,664 tons of ore with an average grade of 15.8% Zn, 8.47% Pb, 1.71% Cu, 3.93 oz/ton Ag and 0.061 oz/ton Au. Apart from Lucky Strike and the small, fault-disrupted Old Buchans deposit it is the only other known major massive sulphide deposit containing "in situ" sulphides associated with stockwork mineralization. However, study of the deposit is hampered by post-ore



Plate 4-13: Banded Oriental No. 1 ore. Interbanded very high grade chalcopyrite and darker galena-sphalerite with minor bornite filling fractures.

imbricate (thrust?) faults of small displacement which have effectively sliced the ore zone into hundreds of sulphide bodies.

The Oriental deposits occur along the north side of the Buchans Inlier and lie above a major (thrust?) fault which separates rocks of the Inlier from the Oriental ore horizon sequence (Fig. 2-2). This same fault may be responsible for repetition of the Lucky Strike ore horizon sequence in the Oriental area, implying that the Oriental deposits lie in approximately the same stratigraphic position as the Lucky Strike deposits.

Virtually all lithologic contacts in the Oriental area are sheared and thus the original stratigraphic relationships are obscured. The deposit is closely associated with altered dacitic crystal-vitric tuffs of the Oriental ore horizon sequence and with highly silicified and mineralized breccias of the Oriental Intermediate Footwall. The ores consist of both "in situ" banded massive sulphide and mechanically transported ores. Ore mineralogy consists of sphalerite, barite and galena with less abundant chalcopyrite, bornite, pyrite and tetrahedrite. Sulphides of the Oriental No. 1 deposit are among the finest in grain size in the Buchans area and banded sulphides are better displayed at Oriental No. 1 than anywhere else in the area (Plate 4-13). Oriental ores display a number of sedimentary structures including slump folds and bedding sag structures.

4.4.2. Oriental No. 2 Orebody: The 0-2 deposit is a complex sulphide body located 150 m north of 0-1 and occurs at, or very nearly in, the same stratigraphic horizon as 0-1. The deposit produced 928,863 tons of massive and fragmental ore at 9.41% Zn, 6.20% Pb, 0.76% Cu, 6.15 oz/ton Ag, and .046 oz/ton Au. The deposit was discovered in 1953, commenced production in 1955 and was mined out in 1969. At present the mine is flooded and information is available only from drill core. The deposit is tabular, varying in thickness up to 10 m and averaging about 5 m. It is approximately 500 m long, 75 m wide and ranges in depth below surface from 40 m to 120 m (Swanson, Perkins and Higgins, 1955).

The ore-bearing sequence conformably overlies dacitic crystal-vitric tuffs which everywhere form the footwall of the deposit. These tuffs vary from green to grey and normally consist of 1-2 mm quartz and plagioclase crystals in a variably sericitized and clay mineralized vitric matrix. This footwall dacite tuff may be overlain conformably in the immediate 0-2 area by any one, or combination of six distinctive rock types, i.e., massive sulphide, poly lithic breccia-conglomerate, monolithic breccia, clastic ore, pyritic siltstone or dacitic vitric-crystal tuff. The relationships between these units are complex.



The footwall polyolithic breccia-conglomerate, clastic ore and siltstone are all underlain and overlain by dacitic tuffs in the immediate footwall of the deposit. The polyolithic breccia-conglomerate consists of angular to rounded fragments of basalt, andesite, dacitic pyroclastic, silicified sugary-textured acid (?) volcanic fragments, fragments similar to portions of the Intermediate Footwall and rare granitoid and jasper pebbles. Pink rhyolite fragments are absent in this horizon. Baritic black ore, yellow ore and less common siliceous or chloritic stockwork fragments occur in varying quantity and sizes up to 25 cm and locally reach ore grade over 60 cm thickness. The footwall breccia-conglomerate is thick between 0-1 and 0-2 but thins rapidly northward. The horizon is very poorly sorted and bedded, although a few graded beds with fragments grading from 50 mm to 1 mm over 2 m are present. The lack of very fine grained sediments at the top of such beds suggests that these debris flows were very dense (i.e., contained little water) and that minimal mixing with water occurred during transport.

Less than one meter of grey pyritic siltstone occupies the same stratigraphic horizon as the breccia-conglomerate in the footwall of the western portion of the orebody.

The polyolithic breccia-conglomerate-clastic ore-siltstone horizon is commonly overlain by a few centimeters

of variably altered dacitic vitric-crystal tuff which is in turn locally overlain by a monomictic breccia containing fragments of strongly altered rocks derived from the Intermediate Footwall. This breccia apparently thickens northward, possibly suggesting a northerly source.

Massive sulphides of the 0-2 orebody everywhere lie in sharp conformable contact with footwall rocks which variably consist of the main footwall dacitic tuff, poly-lithic Breccia-conglomerate, the thin dacitic tuff above the breccia-conglomerate or monomictic breccia. The ore contains both massive (chemically?) deposited sulphide and clastic sulphide facies. Black ore and baritic black ore are dominant in both clastic and massive facies as yellow ore fragments in clastic ore and as wisps in massive ore are volumetrically much less significant. The ore mineralogy consists of barite, sphalerite, galena, pyrite and chalcopryrite with lesser tetrahedrite and bornite (Alcock, 1960).

Massive 0-2 ore, like most other massive sulphides in the area is poorly banded and contains local discontinuous streaks of chalcopryrite-rich ore in black ore. These streaks are commonly randomly oriented in core, suggesting large scale slumping of massive ore. The massive ore commonly contains a variety of strongly altered lithic fragments possibly incorporated into the slumping sulphide during movement. Near the transition from massive to clastic ore,

massive ore commonly takes on an autobrecciated appearance and grades into clastic ore containing both competent angular sulphide fragments and flattened and frayed sulphide fragments which apparently behaved plastically. Local thin fine grained lithic-rich beds are present in clastic ore. In general, clastic sulphide ore is more common at the base and top of the orebody and occurs virtually to the exclusion of massive ore at the northern (distal) periphery of the orebody. The hangingwall contact between high grade clastic ore and overlying volcanic breccia-conglomerate is locally sharp but is commonly characterized by a gradual decrease in abundance and size of sulphide fragments with increasing dilution of lithic fragments. Zoning in the orebody is very poorly developed. Barite-rich zones occur randomly throughout the ore and show little consistent enrichment towards the top of the orebody. No chalcopyrite-rich yellow ore zones are known in the orebody. The ore is not underlain by stockwork mineralization.

All of the above evidence suggests that 0-2 is a transported orebody derived from the nearby 0-1. The distribution of massive and clastic facies and textures in both suggest that 0-2 moved as a large, slowly advancing sulphide mud flow.

The immediate hangingwall of the 0-2 orebody may consist of any of three distinct lithologies which from

stratigraphic bottom to top include 1) monomictic volcanic breccia with fragments derived from the Intermediate Footwall, b) polyolithic volcanic breccia-conglomerate and c) granitoid-bearing volcanic breccia-conglomerate. Strongly altered felsic pyroclastics also locally form the immediate hangingwall but their exact stratigraphic relationship to other hangingwall lithologies is unknown.

The monomictic Intermediate Footwall-bearing breccia is only locally present and contains numerous black ore fragments and boulders. It is conformably overlain by a polyolithic volcanic breccia-conglomerate which bears a lithologic similarity to its counterpart in the footwall. This unit is the most common hangingwall lithology and normally contains a few granitoid pebbles but only rare pink rhyolite pebbles. Angular and plastically deformed sulphide clasts are ubiquitous and yellow ore fragments are normally relatively more abundant than in the orebody itself. The breccia-conglomerate is conformably, and in most places gradationally overlain by a similar breccia-conglomerate but containing abundant subrounded to rounded, pink to white granitoid boulders up to several centimeters in diameter. Other fragment and boulder lithologies in the "granite conglomerate" include green dacitic crystal-vitric tuff, basalt, a variety of fragments derived from the Intermediate Footwall Formation (some bearing the mariposite-like sericite), and rare chert and pumice

fragments. Sulphide fragments less than one centimeter in diameter are common and may consist of baritic black ore, black ore, yellow ore and rare fine grained siliceous pyritic stockwork fragments. Rare isolated sulphide boulders may exceed 10 cm diameter. Sulphide fragments generally decrease in abundance with distance above the ore zone. The unit on the whole is poorly sorted and bedded although a crude sorting into relatively boulder-rich and boulder-poor zones is locally observed. Graded bedding and reverse grading are rare occurrences. Fragments and boulders are most commonly matrix-supported though local fragment-supported zones are present.

North of the 0-2 zone, the "granite conglomerate" rests conformably on footwall dacite and the unit is conformably overlain by hangingwall dacite.

4.4.3. Sandfill Prospect: The Sandfill Prospect is an subeconomic sulphide deposit which occurs 1 km north-northeast of Oriental (Fig. 2-1) and at approximately the same stratigraphic horizon as the Oriental deposits. The deposit, which is exposed only in drill core, occurs at 350 m depth and comprises 75,000 possible tons of 1.5% Cu, 4.7% Pb, 10.0% Zn, 0.01 oz/ton Au and 2.5 oz/ton Ag. It is a mechanically deposited "orebody" consisting of a concentration of sulphide fragments surrounded by a lower grade zone of sulphide fragments diluted by abundant lithic fragments.

The Sandfill and Middle Branch deposits provide excellent examples of the paleotopographic control exerted on the distribution of the transported ores.

The deposit occurs in a sequence of debris flows of the Oriental ore horizon sequence and lies approximately 100 m above the fault-bounded upper contact of Footwall Arkose. In the footwall of the deposit there occurs a small (100 m long and 15 m thick) pervasively silicified and lightly pyritized grey-white rhyolite body with a strongly brecciated top. This rhyolite breccia grades upward into a thin (less than 10 m) sequence of white rhyolitic lapillistone and lesser bedded rhyolitic wacke with lightly disseminated pyrite and rare films of galena and sphalerite in the matrix. Matrix pyrite is generally coarser in coarser wackes suggesting a detrital origin. Within a few meters of the overlying "ore", extraneous lithic fragments become abundant. These include pyritic siltstone (with altered rims), altered intermediates (with "mariposite") and a few small Cu-Pb-Zn sulphide fragments which increase upwards in size and abundance to a maximum of 80% sulphide fragments. (Weak silicification occurs for one meter below "ore" in a single drill core.) This "ore" consists mainly of galena-sphalerite-chalcopyrite fragments up to 10 cm in size, with lesser chalcopyrite and/or pyrite, barite and barite-sphalerite fragments. Many of the fine grained banded sulphide fragments bear a strong

resemblance to Oriental ores. Lithic fragments occur throughout the ore and a few fine grained (1 cm) essentially barren, polyolithic breccia beds are intercalated with the "ore". Within one meter of the top of the "ore" the size of both lithic and sulphide fragments decreases dramatically and the ratio of lithic to sulphide fragments increases rapidly as the sequence grades into a pyritic wacke bed. This "ore" sequence is subsequently overlain by at least nine graded breccia beds ranging in thickness from less than one meter to 20 m thick, each of which grade from polyolithic breccia at the base to greywacke or siltstone at the top. Small sulphide fragments are a component of each breccia bed but their abundance and size decreases with each successive breccia bed above the major sulphide concentration. This sequence of subaqueous debris flows is subsequently overlain by dacitic pyroclastic flows which locally contain a few granitic pebbles.

Considerable evidence exists to suggest that the Sandfill Prospect is a mechanically deposited sulphide accumulation. This evidence includes:

- 1) the presence of angular sulphide fragments
- 2) banding in sulphide fragments is terminated at fragment boundaries
- 3) the abundance of silicate fragments
- 4) the vertical and lateral, increase in size and relative abundance of sulphide fragments towards "ore"

- 5) the lack of metal zoning in the sulphide body
- 6) the absence of stockwork mineralization
- 7) the lack of visible alteration consistently related to the presence of sulphides
- 8) the occurrence of graded sulphide and silicate fragments at the top of the "ore"
- 9) the occurrence of the ore in a sequence of graded sulphide-bearing subaqueous debris flows.

4.4.4. Middle Branch Prospect: The Middle Branch prospect is a small sulphide deposit exposed in a single drill hole at a depth of 555 m. The single intersection over 2 m assayed 0.4% Cu, 3.1% Pb, 4.8% Zn, 0.02 oz/ton Au, 1.4 oz/ton Ag and 8.0% Fe in sulphides.

The deposit occurs at a stratigraphic horizon very close to that of the Sandfill deposit although rapid lateral facies changes and minor faults filled with diabase have hampered detailed correlation. The deposit is bounded on both the footwall and the hangingwall by diabase dykes. Beneath the footwall diabase there occurs a sequence of fractured and altered siliceous siltstones which contain detrital pyrite and traces of epigenetic galena-sphalerite in thin fractures. A distinct whitish alteration of siltstone occurs adjacent to the network of epigenetic fractures and pervades into the siltstone for short distances. These siltstones are interbedded with rhyolitic tuff-breccia containing grey to pink rhyolite fragments with



lesser siltstone fragments and minor pyrite and galena-sphalerite fragments up to one centimeter. Traces of Pb-Zn veining as films in fractures occur locally and a single vein of pyrite-barite occurs 15 m below "ore".

The sulphide body itself is not so obviously fragmental as the Sandfill although a transported origin is suspected. The "ore" contains 20% lithic breccia fragments but no sulphide fragments are macroscopically visible. The gangue minerals are much coarser grained than Sandfill with pyrite commonly of 1-2 mm size and barite up to 1 cm. Galena and sphalerite are fine grained and the ore contains relatively little chalcopryrite. Overlying the hangingwall diabase there occurs a polyolithic breccia which contains a few sulphide-bearing fragments.

The ultimate source of the Sandfill and Middle Branch Prospects is of considerable exploration interest. As with all transported ores, an analysis of paleotopography may be valuable in tracing such ores to their sources. An isopach map of the Oriental ore horizon sequence, a schematic cross section of the Sandfill-Middle Branch area and a paleotopographic plan are presented in Figures 4-4, 4-5 and 4-6, respectively. It can be seen from these diagrams that the Sandfill and Middle Branch Prospects occupy opposite flanks of a ridge of coalescing rhyolite-breccia domes which traverse the length of a deep paleotrough. This trough extends from Old Buchans to Oriental




Figure 4-4. Isopach map of the Oriental ore horizon sequence illustrating the nature of the Oriental-Sandfill Middle Branch channel. In situ ore, solid black; transported ore, stippled. Contour interval = 30 m




Figure 4-5

Schematic stratigraphy in the Sandy Lake - Sandfill - Middle Branch Area




Figure 4-6

Paleotopographic plan  
at the time of ore  
deposition in the Sandfill  
Middle Branch area.

and subsequently northward to the Sandfill-Middle Branch area, and beyond. The presence of this trough linking the in situ Oriental No. 1 orebody to the Sandfill-Middle Branch Prospects provides compelling evidence that Oriental No. 1 constitutes the source of the transported Prospects. Other arguments in favour of an Oriental source are listed below:

- 1) Correlation between Middle Branch-Sandfill and Oriental indicate that all three zones occur at essentially the same stratigraphic horizon.
- 2) The Oriental and Sandfill-Middle Branch transported ores both consist of mainly black ore fragments in polyolithic breccias.
- 3) The broad thick sequence which hosts the Oriental ores funnels rapidly northward into a relatively narrow, northeast trending trough which hosts the Sandfill-Middle Branch mineralization (Fig. 4-4).
- 4) Though alteration and minor sulphide veining exist in the Sandfill area, typical stockwork similar to that at Oriental or Lucky Strike is not known to occur.
- 5) Sorting and grading of host rocks in the Sandfill-Middle Branch zones is better than that at Oriental. This, combined with the relatively lithic-diluted nature of the ores, suggests that they are relatively distal deposits and that Oriental could be the distant source.

Evidence in favour of an undiscovered source is as follows:

- 1) Points 1), 2), and 5) above, could be equally explained by an undiscovered source.
- 2) Altered rhyolite fragments are abundant in the Sandfill-Middle Branch ore horizon and are relatively much less abundant in breccias of the Oriental area. These fragments are demonstrably of local origin, and therefore suggest that the sulphide fragments could also be local
- 3) The size of lithic fragments (some over 120 cm, or approximately 2600 kg) and sulphide fragments (up to 45 cm in diameter or approximately 225 kg) in the ore horizon breccias is suggestive of a local source.
- 4) It can be seen from Fig. 4-2 that zones of sulphide veining and alteration in rhyolite occur near the Sandfill and Middle Branch deposits. If these deposits were derived from Oriental this would have to be ascribed to coincidence.
- 5) Metal ratios of the Sandfill deposit are dissimilar from the Oriental No. 1 or Oriental No. 2 orebodies. In comparison, metal ratios from Rothermere and MacLean are virtually identical to their presumed source at Lucky Strike.
- 6) Barite-pyrite boulders occur in the ore breccias and Barite-pyrite veinlets occur in altered rhyolite in the

Sandfill-Middle Branch area. This barite-pyrite association is relatively poorly developed at Oriental.

- 7) It is entirely possible that the Middle Branch channel (Fig. 4-6) does not connect with the Sandfill-Oriental channel. A separate source might thus be suggested for at least the Middle Branch prospect.
- 8) If a major thrust fault is present on top of the Footwall Arkose a number of further arguments can be invoked. It is beyond the scope of this thesis to fully document the reasoning for these arguments.

4.4.5. Old Buchans Orebodies: The Old Buchans Orebodies are among the most poorly understood deposits in the Buchans area due to the antiquity of mine workings, poor core recovery, strong faulting and lack of study. The Old Buchans Conglomerate Orebody was transported, occurring on the Oriental horizon and grossly similar to O-2. The Old Buchans East and West Orebodies are associated with stockworks and numerous faults. Their stratigraphic position is uncertain and conceivably they could lie in the Lucky Strike thrust block, or even possibly in a lower thrust block.

#### 4.5. Selected Comments

4.5.1. Miscellaneous Occurrences of Sulphide Clasts: A noteworthy feature of the Buchans area is the widespread distribution of sulphide clasts occurring over several




Figure 4-7: Plan showing distribution of high grade sulphide clasts in the Buchans area. Dots indicate vertical drill holes which have intersected sulphide clasts. In situ orebodies are shown in solid black, transported orebodies as outlined.





Plate 4-14: Central greyish black ore pebble and smaller  
clastic sulphide grains in pyroclastic breccia.  
Enlargement approximately 1.5X.

square kilometers in the Lucky Strike and Oriental ore horizon sequences. The distribution of such clasts as intersected in drill core is shown in Fig. 4-7. They range from grains and pebbles to 30 cm boulders of high grade black ore (Plate 4-14). Clasts of yellow ore are much less common. They normally occur in polyolithic breccias or pyroclastic breccias at the same stratigraphic horizon as major in situ and transported orebodies, but significant concentrations of clasts are also found within the ore horizon sequences, at stratigraphic positions in which no in situ ore is known.

Many such clasts display diffuse, dark rims of alteration suggestive of diagenetic chemical attack. Smaller grains locally are almost entirely degraded. The significance of this is discussed in Chapter Seven.

Minor occurrences of sulphide clasts are known within the Intermediate Footwall and Ski Hill Sequence. These are grossly explicable in terms of the minor stratiform mineralization occurring at these horizons. An inexplicable occurrence of small, high grade sulphide grains and pebbles occur within the Wiley's Sequence, 6.7 km distant from Lucky Strike Orebody.

4.5.2. Comments on Granitoid Boulders at Buchans:  
The Feeder Granodiorite is lithologically and chemically similar to some Buchans Group rhyolites and to granitoid

boulders in the ore horizon. It probably represents the source of the granitoid boulders occurring throughout the Buchans Group and in the ore horizon.

Granitoid boulders in volcanic terranes are not uncommon (e.g., Cady, 1975; Helwig and Sarpf, 1969) and in some instances, such boulders are thought to represent the plutonic equivalents of cogenetic volcanics (e.g. Gabrielse and Reesor, 1974). Cobbing and Pitcher (1972) suggest that magma pressure may force up fault blocks of solidified magma chamber thus exposing them to erosion at the same time as intrusion-extrusion. However no such fault blocks and no logical source, exposed during ore deposition, are known at Buchans.

More plausibly, the author suggests that the granitoid boulders were rounded during gas-transport to surface in "pebble dykes", such as those that occur in close association with the Japanese Kuroko deposits and many porphyry Cu-Mo deposits. No such dykes are known at Buchans, but the most likely source area, in the Lucky Strike region, is now eroded. An origin of this nature would explain the unique sphericity and rounding of the granitoid boulders as well as their anomalous abundance in the areally and stratigraphically restricted ore horizon breccias.

4.5.3. Buchans as Compared to Other Occurrences of Mechanically Transported Ores: The occurrence of mechanically transported ores was predicted as early as 1965 (Suffel, 1965, p. 1062) and since have been recognized in association with several massive sulphide deposits, e.g. at Noranda, Sinclair (1971); Spence (1975); at Kidd Creek, Walker et al. (1975); in the Iberian pyrite belt (Schermerhorn 1970, 1971); and in the Kuroko deposits (see next section). Sulphide fragments have been noted in many other deposits, some of which may be transported.

The Buchans transported ores are proximal by comparison to those of the Iberian pyrite belt. Within this class, they represent (to the author's knowledge) the world's best preserved, most extensively developed, farthest travelled and highest grade transported ores.

#### 4.6. Comparison of Buchans and Kuroko Deposits

The Buchans deposits and the famous Kuroko deposits of Japan are so very similar that a few words of comparison are warranted. Both types occur as stratiform polymetallic deposits associated with breccia horizons in felsic volcanic rocks (Sato, 1974; Lambert and Sato, 1974). The ore and gangue mineralogy of both types is similar, although neither gypsum nor anhydrite have been identified at Buchans. Both types have generally sharp boundaries with their wall



Plate 4-15: Fragmental ore from the Japanese Aina i Kuroko deposit. The bulk of the same is composed of black and yellow ore clasts.

rocks and display numerous internal submarine sedimentary structures and textures consistent with a volcanic-exhalative mode of origin. Both appear to be relatively shallow water phenomena though this has been questioned in the Kuroko case by Urabe and Sato (1978). Stringer and stockwork mineralization is common to the footwall of both types. Sediments associated with both types are dominantly volcaniclastic as opposed to chemical. Some Kuroko ore-bodies (e.g., Shakanai Mine, Ohtagaki et al., 1974, p. 135) contain interbedded barren sedimentary layers like Rothermere and MacLean Orebodies. Though not a commonly mentioned feature, some Kuroko deposits occupy depressions in the underlying volcanic surface (e.g., Ohtagaki et al., 1974, p. 137; N. Sato and Kusaka, 1974, p. 146; Tanimura et al., 1974, p. 153; Hirabayashi, 1974, p. 200; Honishi, 1974, p. 218). Pebbles of granitic basement rock occur in close association with ores of the Yoshino Mine (Osada et al., 1974, p. 185) and are apparently related to contemporaneous block faulting with basement uplift. It is interesting to note that these same deposits are not associated with white rhyolite domes and contain no gypsum.

Mechanically transported sulphide ores have been described from several Kuroko mines (Plate 4-15), e.g., Shakanai Mine, (Kajiwara, 1970; Ohtagaki et al., 1974), Kosaka Mine (Sato, 1968; Oshima et al., 1974), Ainal Mine (Ishikawa and Yanagisawa, 1974), Hanaoka Mine (Takahashi






Plate 4-16: Motoyama volcanic breccia, Kosaka Mine. Note altered rims on rhyolite fragments, similar to those occurring at Buchans.


and Suga, 1974; Ito et al., 1974), Kamikita Mine (Lee et al., 1974), Kurosawa Mine (Motegi, 1974) and Fukazawa Mine (Tanimura et al., 1974). Imbrication, bedding sag structures, graded sulphide breccias, scouring at the base of the deposits and ore balls surrounded by clay are features of transported Kuroko ores which have also been identified at Buchans. Both plastic and competent sulphide fragments have been identified in both the Kuroko (e.g., Ishikawa and Yanagisawa, 1974, p. 87) and Buchans deposits. The occurrence at Buchans of lithic fragments with all three types of alteration rims as described by Lee et al. (1974, p. 61) is remarkable (Plate 4-16).

Despite the absence of gypsum and anhydrite at Buchans, and the limited development of yellow ore, zoning is very similar to that of a typical Kuroko deposit as described by T. Sato (1974). The stratigraphically lowest stockwork mineralization at Buchans is pyritic, much like that of the Kuroko deposits. Stratigraphically higher stockwork mineralization at Buchans is pyritic and polymetallic like some Kuroko stockworks. Such mineralization at Buchans is associated with strong chloritization and/or silicification; where silicification is dominant the Buchans stockwork closely resembles Keiko. Copper is relatively more abundant in the stockwork at Buchans than in the overlying massive sulphide, a feature in common with the Kuroko deposits. Yellow ore is poorly developed at Buchans but,



where present, is concentrated towards the base of the massive sulphide (George, 1937), i.e., in a similar position to the Oko. Pb-Zn-rich black ore occupies the upper portion of the "in situ" deposits at Buchans in a similar stratigraphic position to the Kuroko. Gold and silver minerals are concentrated in this zone in both cases. Bedded barite overlies the black ore in both Buchans and Kuroko examples. Though no red hematite-quartz layer is present at Buchans, this horizon is locally occupied by hematitized barite and hematitized tuff. Several Kuroko deposits are subsequently overlain by tuffaceous felsic volcanic rocks.

Several features of the Buchans deposit differ from those of typical Kuroko deposits. Though present at Buchans, white rhyolite bodies do not display the intimate spatial relationship to mineralization which is so characteristic of the Kuroko deposits. Gypsum is absent at Buchans. Yellow ores are poorly developed and the abundance of pyrite throughout is less than that of typical Kuroko deposits. Well banded ores and ores displaying colloform banding are relatively less abundant at Buchans. Though mechanically transported sulphide ores are characteristic of both examples, the Buchans deposits are anomalous in both the high quantity and quality of transported ores as well as transport distances which exceed any of the Kuroko deposits (T. Sato, pers. comm., 1975).



## CHAPTER FIVE

### GENERAL PETROCHEMICAL NATURE OF THE BUCHANS GROUP

#### 5.1. Post-Magmatic Chemical Changes of Buchans Group Rocks

The ongoing spilitic debate (e.g., Amstutz, 1974) and an interest in ocean floor chemical changes has fostered abundant research into post-depositional changes in submarine volcanics (especially basalts). Documented chemical changes as a result of low temperature depositional and diagenetic alteration range from essentially nil (e.g., Clarke, 1975) to substantial additions and removals of certain elements. On the basis of petrographic, geochemical and experimental data (and with a few discrepancies),  $\text{SiO}_2$ ,  $\text{CaO}$ ,  $\text{MnO}$ , total Fe and Cu are commonly leached during low temperature submarine alteration of basaltic rocks whereas  $\text{H}_2\text{O}$ ,  $\text{K}_2\text{O}$ ,  $\text{MgO}$ , Rb, Cs,  $\text{Fe}^{++}/\text{Fe}^{+}$  and K/Rb commonly increase (e.g., S.R. Hart, 1969, 1973; Philpotts *et al.*, 1969; Cann, 1970; R. Hart, 1973; Hajash, 1975; Bischoff and Dickson, 1975). The same authors have generally concluded that Ti and Al have remained relatively inert during alteration but little consensus has been reached on the behavior of more mobile elements such as Na.

Under conditions of prehnite-pumpellyite metamorphism, such as that experienced by the Buchans Group, volcanic sequences commonly undergo varying degrees of

hydration and redistribution of elements, especially Na and Ca (Smith, 1968; Jolly and Smith, 1972; Jolly, 1972). Among the base metals, Jolly (1974) demonstrated considerable redistribution of Cu and removal of Zn in low rank basaltic rocks.

In view of the variety of probable post-magmatic chemical changes, any attempt to relate rock chemistry to mineralization must demonstrate that anomalous conditions are indeed a function of mineralization and not due to other processes. Apart from the effects of mineralization in proximity to ore, rocks of the Buchans Group have undoubtedly been subject to chemical changes as a result of depositional, diagenetic and metamorphic alteration. Perhaps the most obvious effect of post-magmatic alteration is the widespread depositional-metamorphic introduction of water combining to form hydrous minerals (chlorite, sericite, pumpellyite, prehnite, etc.). Apart from increasing loss on ignition, this has the effect of "diluting" the abundance of other elements.

Local diagenetic redistribution of several elements is evident in virtually all lithologies. Concordant hematitization and calcite introduction and abundant vein and amygdale calcite are characteristic of many mafic pyroclastics and breccias. In these rocks crustification, comb and cockade structures, indicative of open space

filling, are visible locally. Such volcanoclastics commonly alternate with finer grained tuffaceous rocks which show no such alteration, indicating that calcite introduction occurred at, or shortly after deposition. The variety of vesicle-fillings further indicate post-depositional mobility of  $H_2O$ , Si, Ca, Mg, Fe and Al. Analyses of rocks containing significant vesicle and fracture filling was not attempted in this study. It is shown in subsequent sections that diagenetic-metamorphic alteration of crystal-vitric tuffs involved hydration and addition of MgO with removal of  $Na_2O$ . Secondary quartz halos around quartz crystals in these rocks indicate mobility of silica probably during diagenetic alteration of glass to clay minerals. Dispersion of very mobile elements such as Hg from the orebodies possibly provides an indication of the extent of secondary migrations. In the case of Hg, depositional, diagenetic and metamorphic processes have caused dispersion only on the scale of a few meters (Thurlow, 1973). If this is the case, then elements much less mobile than Hg have probably migrated proportionally shorter distances.

With respect to the elemental abundances discussed in subsequent sections, it is apparent that an average of a large number of samples will tend to nullify the effects of local migrations as the population will presumably include samples at all phases of migration. In this regard, the averages are probably a reasonable approximation of the

original composition. However, the effects of depositional enrichments of elements and of hydration are superimposed on the averages and quantitative estimates of their contribution are difficult if not impossible.

#### 5.2. General Elemental Abundances and Chemical Variations of Buchans Group Rocks

The average chemical composition of all significant lithologic units of the Buchans Group analysed in this study are given in Table 5-1.

The volcanics of the Buchans Group display little alkaline affinity. Neither olivine nor feldspathoids are present in the mode or the norm and pyroxenes are neither sodic nor titaniferous. Phosphorus and titanium are present in abundances well below normal for any alkaline suite and the Buchans volcanics are generally subalkaline on the alkali: silica diagram of Irvine and Baragar (1971) (Fig. 5-1).

Similarly, the volcanic rocks of the Buchans Group bear little lithologic or chemical resemblance to oceanic tholeiites. The relatively shallow water environment, the relative abundance of felsic volcanics and the total absence of ultramafic rocks and sheeted dykes render an ocean floor origin extremely unlikely. Chemically, the Buchans Group volcanics display abundances of large cations (K, Rb, Ba, Sr, etc.) which are much greater than those of

Table 5-1: Average Chemical Composition of Buchans Area Rocks

|                                | Footwall<br>Basalt<br>Flows | Footwall<br>Basalt<br>Fragmentals | Prominent<br>Quartz<br>Sequence<br>Rhyolites | Prominent<br>Quartz Sequence<br>Dacitic Tuffs<br>(Large Quartz) | Prominent<br>Quartz Sequence<br>Dacitic Tuffs<br>(Small Quartz) | Prominent<br>Quartz Sequence<br>Tuffaceous<br>Siltstone | Ski Hill<br>Sequence<br>Flows | Ski Hill<br>Sequence<br>Fragmentals | Intermediate<br>Footwall | Mineralized<br>Intermediate<br>Footwall | Pyritic<br>Siltstone<br>Rhyolite<br>Nucleon |
|--------------------------------|-----------------------------|-----------------------------------|----------------------------------------------|-----------------------------------------------------------------|-----------------------------------------------------------------|---------------------------------------------------------|-------------------------------|-------------------------------------|--------------------------|-----------------------------------------|---------------------------------------------|
| n=                             | 3                           | 8                                 | 9                                            | 11                                                              | 19                                                              | 8                                                       | 31                            | 29                                  | 14                       | 18                                      |                                             |
| SiO <sub>2</sub>               | 48.8                        | 46.6                              | 69.0                                         | 68.5                                                            | 69.8                                                            | 67.4                                                    | 57.4                          | 55.4                                | 54.4                     | 47.2                                    | 65.4                                        |
| TiO <sub>2</sub>               | .83                         | .87                               | .39                                          | .35                                                             | .33                                                             | .33                                                     | .70                           | .64                                 | .53                      | .21                                     | .34                                         |
| Al <sub>2</sub> O <sub>3</sub> | 15.47                       | 16.81                             | 13.36                                        | 14.02                                                           | 14.44                                                           | 13.93                                                   | 15.23                         | 16.18                               | 16.15                    | 8.61                                    | 13.58                                       |
| Fe <sub>2</sub> O <sub>3</sub> | 8.63                        | 11.54                             | 3.78                                         | 3.95                                                            | 2.73                                                            | 4.25                                                    | 8.31                          | 9.20                                | 9.84                     | 17.16                                   | 3.86                                        |
| MnO                            | .14                         | .13                               | .11                                          | .06                                                             | .05                                                             | .17                                                     | .16                           | .16                                 | .32                      | .35                                     | .11                                         |
| MgO                            | 5.88                        | 6.72                              | 1.25                                         | 1.53                                                            | 1.47                                                            | 2.61                                                    | 3.50                          | 3.65                                | 6.34                     | 7.17                                    | 3.18                                        |
| CaO                            | 9.90                        | 6.27                              | 2.39                                         | 2.48                                                            | 2.27                                                            | 2.06                                                    | 6.28                          | 6.42                                | 1.95                     | 2.09                                    | 2.13                                        |
| Na <sub>2</sub> O              | 3.08                        | 1.82                              | 3.69                                         | 2.66                                                            | 2.32                                                            | 1.96                                                    | 3.54                          | 2.58                                | 1.90                     | .29                                     | 2.87                                        |
| K <sub>2</sub> O               | .97                         | 1.40                              | 2.32                                         | 2.46                                                            | 2.62                                                            | 4.13                                                    | 1.04                          | 1.38                                | 2.00                     | .79                                     | 2.18                                        |
| P <sub>2</sub> O <sub>5</sub>  | -                           | -                                 | -                                            | -                                                               | -                                                               | -                                                       | -                             | -                                   | .10                      | .09                                     | .08                                         |
| Loss, Ig.                      | 3.41                        | 7.12                              | 2.83                                         | 3.46                                                            | 3.27                                                            | 2.99                                                    | 2.93                          | 3.91                                | 5.58                     | 11.78                                   | 4.07                                        |
| Total                          | 98.92                       | 99.13                             | 99.32                                        | 99.47                                                           | 99.30                                                           | 99.44                                                   | 99.16                         | 99.74                               | 99.11                    | 95.74                                   | 97.72                                       |
| As (ppm)                       | 410                         | 721                               | 708                                          | 493                                                             | 688                                                             | 873                                                     | 617                           | 1062                                | 1541                     | 1898                                    | 1848                                        |
| Be (SO <sub>4</sub> -)         | -                           | -                                 | -                                            | -                                                               | -                                                               | -                                                       | -                             | -                                   | 29                       | 3492                                    | 1327                                        |
| Pb                             | 25                          | 30                                | 13                                           | 7                                                               | 8                                                               | 14                                                      | 19                            | 28                                  | 79                       | 5470                                    | 88                                          |
| Pb (B-)                        | 7                           | 6                                 | 16                                           | 18                                                              | 11                                                              | 10                                                      | 13                            | 14                                  | 70                       | 5400                                    | 113                                         |
| Zn                             | 64                          | 90                                | 36                                           | 52                                                              | 33                                                              | 37                                                      | 75                            | 88                                  | 235                      | 11700                                   | 364                                         |
| Zn (B-)                        | 13                          | 16                                | 13                                           | 15                                                              | 12                                                              | 12                                                      | 19                            | 21                                  | 68                       | 12000                                   | 325                                         |
| Cu                             | 52                          | 66                                | 3                                            | 3                                                               | 5                                                               | 14                                                      | 49                            | 58                                  | 64                       | 1670                                    | 21                                          |
| Cu (B-)                        | 45                          | 45                                | 6                                            | 5                                                               | 4                                                               | 7                                                       | 43                            | 51                                  | 63                       | 1514                                    | 16                                          |
| S                              | -                           | -                                 | -                                            | -                                                               | -                                                               | -                                                       | -                             | -                                   | -                        | -                                       | -                                           |
| Ag                             | -                           | -                                 | -                                            | -                                                               | -                                                               | -                                                       | -                             | -                                   | .93                      | 4.1                                     | .43                                         |
| Hg (ppb)                       | -                           | -                                 | -                                            | -                                                               | -                                                               | -                                                       | -                             | -                                   | 44                       | 234                                     | 67                                          |
| Pb                             | 22                          | 34                                | 45                                           | 61                                                              | 69                                                              | 71                                                      | 18                            | 32                                  | 39                       | 28                                      | 34                                          |
| Ba                             | 256                         | 245                               | 153                                          | 144                                                             | 265                                                             | 144                                                     | 267                           | 310                                 | 129                      | 68                                      | 219                                         |
| Sr                             | 72                          | 80                                | 122                                          | 140                                                             | 127                                                             | 110                                                     | 87                            | 82                                  | 81                       | 59                                      | 129                                         |
| Yb                             | 43                          | 43                                | -                                            | -                                                               | 43                                                              | 43                                                      | 43                            | 43                                  | 43                       | 92                                      | 43                                          |
| Ni                             | 48                          | 40                                | 4                                            | 6                                                               | 5                                                               | 3                                                       | 21                            | 12                                  | 13                       | 27                                      | 5                                           |
| Co                             | 25                          | 34                                | 27                                           | 18                                                              | 9                                                               | 5                                                       | 27                            | 22                                  | 20                       | 19                                      | 7                                           |
| Cr                             | 243                         | 134                               | 10                                           | 13                                                              | 16                                                              | 40                                                      | 76                            | 47                                  | 53                       | 122                                     | 28                                          |
| V                              | 177                         | 326                               | 25                                           | 24                                                              | 21                                                              | 27                                                      | 170                           | 249                                 | 245                      | 136                                     | 54                                          |
| Th                             | -                           | -                                 | 11(4)                                        | 10(8)                                                           | 14(3)                                                           | -                                                       | -                             | -                                   | 11                       | -                                       | -                                           |
| U                              | -                           | -                                 | 3(4)                                         | 1(8)                                                            | 2(3)                                                            | -                                                       | -                             | -                                   | 2                        | -                                       | -                                           |
| Ge                             | -                           | -                                 | 13(4)                                        | 15(8)                                                           | 17(3)                                                           | -                                                       | -                             | -                                   | 18                       | -                                       | -                                           |
| n =                            | 3                           | 8                                 | 9                                            | 11                                                              | 19                                                              | 8                                                       | 31                            | 29                                  | 14                       | 18                                      |                                             |

POOR COPY  
COPIE DE QUALITEE INFRIEURE

Table 5-1 (continued)

|                                | Oriental and<br>Lucky Strike<br>Population | Clementine<br>New<br>Population | White<br>Population<br>(Oriental) | Lucky Strike<br>New<br>Population | Oriental<br>New<br>Population | Clementine<br>New<br>Population | "Marion"<br>New<br>Population | Upper Buckhorn<br>Subgroup<br>Population | Upper Buckhorn<br>Subgroup<br>Population |
|--------------------------------|--------------------------------------------|---------------------------------|-----------------------------------|-----------------------------------|-------------------------------|---------------------------------|-------------------------------|------------------------------------------|------------------------------------------|
|                                |                                            |                                 |                                   |                                   |                               |                                 |                               |                                          |                                          |
| Si                             | 7                                          | 2                               | 2                                 | 94                                | 26                            | 32                              | 4                             | 16                                       | 16                                       |
| SiO <sub>2</sub>               | 72.6                                       | 72.4                            | 74.5                              | 71.0                              | 67.0                          | 71.7                            | 61.7                          | 69.2                                     | 61.6                                     |
| TiO <sub>2</sub>               | .37                                        | .16                             | .31                               | .23                               | .32                           | .24                             | .28                           | .29                                      | .43                                      |
| Al <sub>2</sub> O <sub>3</sub> | 13.07                                      | 13.72                           | 11.60                             | 13.64                             | 15.56                         | 13.25                           | 12.86                         | 14.52                                    | 15.43                                    |
| Fe <sub>2</sub> O <sub>3</sub> | 2.09                                       | 1.54                            | .02                               | 2.23                              | 2.66                          | 2.15                            | 4.12                          | 3.27                                     | 6.49                                     |
| MnO                            | .03                                        | .05                             | .01                               | .08                               | .06                           | .06                             | .10                           | .07                                      | .12                                      |
| MgO                            | .28                                        | .44                             | .31                               | 1.23                              | 2.56                          | 1.20                            | 1.88                          | 1.68                                     | 2.35                                     |
| CaO                            | 1.21                                       | .92                             | .60                               | 2.17                              | 2.05                          | 1.97                            | 2.27                          | 1.50                                     | 2.92                                     |
| Na <sub>2</sub> O              | 4.77                                       | 3.77                            | .94                               | 2.69                              | 2.17                          | 1.79                            | 2.36                          | 2.77                                     | 3.87                                     |
| K <sub>2</sub> O               | 2.65                                       | 4.03                            | 7.79                              | 2.82                              | 2.72                          | 3.20                            | 2.30                          | 2.43                                     | 1.09                                     |
| P <sub>2</sub> O <sub>5</sub>  | -                                          | -                               | -                                 | .06                               | -                             | -                               | .10                           | -                                        | -                                        |
| Loss, 1%                       | 1.35                                       | 1.83                            | 1.65                              | 2.97                              | 3.90                          | 3.76                            | 5.74                          | 3.06                                     | 4.43                                     |
| Total                          | 99.23                                      | 98.90                           | 97.73                             | 99.16                             | 99.00                         | 99.32                           | 93.71                         | 98.79                                    | 98.73                                    |
| Si                             | 1133                                       | 1274                            | 3810                              | 1657                              | 1388                          | 1042                            | 31600                         | 866                                      | 352                                      |
| SiO <sub>2</sub>               | -                                          | -                               | -                                 | -                                 | -                             | -                               | -                             | -                                        | -                                        |
| Ti                             | 15                                         | 14                              | 18                                | 29                                | 41                            | 14                              | 559                           | 15                                       | 19                                       |
| TiO <sub>2</sub>               | -                                          | -                               | -                                 | -                                 | -                             | -                               | -                             | -                                        | -                                        |
| Al                             | 15                                         | 18                              | 38                                | 24                                | 77                            | 24                              | -                             | 55                                       | 15                                       |
| Al <sub>2</sub> O <sub>3</sub> | 30                                         | 24                              | 237                               | 109                               | 160                           | 25                              | 1323                          | 65                                       | 78                                       |
| Fe                             | 13                                         | 14                              | 241                               | 34                                | 120                           | 8                               | -                             | 13                                       | 18                                       |
| Fe <sub>2</sub> O <sub>3</sub> | 6                                          | 5                               | 1                                 | 5                                 | 13                            | 1                               | 596                           | 5                                        | 52                                       |
| Ca                             | 7                                          | 4                               | 4                                 | 2                                 | 16                            | 2                               | -                             | 5                                        | 54                                       |
| CaO                            | -                                          | -                               | -                                 | -                                 | -                             | -                               | -                             | -                                        | -                                        |
| Mg                             | -                                          | -                               | -                                 | -                                 | -                             | -                               | -                             | -                                        | -                                        |
| MgO                            | -                                          | -                               | -                                 | -                                 | -                             | -                               | -                             | -                                        | -                                        |
| Mn                             | 43                                         | 72                              | 63                                | 70                                | 83                            | 99                              | 99                            | 59                                       | 28                                       |
| MnO                            | 224                                        | 118                             | 84                                | 222                               | 250                           | 127                             | 625                           | 111                                      | 144                                      |
| Na                             | 164                                        | 101                             | 157                               | 122                               | 194                           | 100                             | 116                           | 124                                      | 93                                       |
| Na <sub>2</sub> O              | -                                          | -                               | -                                 | -3                                | -                             | -                               | -3                            | -                                        | -                                        |
| K                              | 2                                          | 5                               | 2                                 | 3                                 | 4                             | 4                               | 5                             | 5                                        | 9                                        |
| K <sub>2</sub> O               | 30                                         | 17                              | 35                                | 5                                 | 16                            | 19                              | 6                             | 21                                       | 25                                       |
| Cl                             | 10                                         | 6                               | 0                                 | 24                                | 9                             | 2                               | 33                            | 10                                       | 26                                       |
| V                              | 26                                         | 10                              | 13                                | 19                                | 16                            | 15                              | 65                            | 10                                       | 118                                      |
| Tr                             | 16                                         | 19(11)                          | 17                                | 16(12)                            | 19(21)                        | 16                              | -                             | 8(2)                                     | -                                        |
| U                              | 3                                          | 5(1)                            | 3                                 | 4(12)                             | 4(21)                         | 4                               | -                             | 1(2)                                     | -                                        |
| C                              | 13                                         | 13(1)                           | 6                                 | 13(12)                            | 16(23)                        | 13                              | -                             | 15(2)                                    | -                                        |
| W                              | 7                                          | 5                               | 2                                 | 94                                | 26                            | 13                              | 4                             | 16                                       | 6                                        |

POOR COPY  
COPIE DE QUALITEE INFERIEURE

Table 5-1 (continued)

|                                      | Granite Boulders<br>In Ore Horizon | Feeder<br>Granodiorite | Low Titanium<br>Diabase | High<br>Titanium<br>Diabase | Alkali-<br>Feldspar<br>Granite | Paralkaline<br>Granite | Hungry Mountain<br>Complex |
|--------------------------------------|------------------------------------|------------------------|-------------------------|-----------------------------|--------------------------------|------------------------|----------------------------|
| $\text{SiO}_2$                       | 72.8                               | 74.5                   | 48.3                    | 46.8                        | 74.7                           | 76.0                   | 70.6                       |
| $\text{TiO}_2$                       | .30                                | .23                    | .01                     | 2.01                        | .20                            | .18                    | .34                        |
| $\text{Al}_2\text{O}_3$              | 12.45                              | 12.59                  | 15.84                   | 15.48                       | 12.68                          | 11.32                  | 13.96                      |
| $\text{FeO}$                         | 1.63                               | 2.14                   | 10.45                   | 11.75                       | 1.66                           | 2.39                   | 3.06                       |
| $\text{MnO}$                         | .05                                | .04                    | .18                     | .20                         | .02                            | .05                    | .05                        |
| $\text{MgO}$                         | .91                                | .48                    | 7.07                    | 6.25                        | .15                            | .01                    | 1.11                       |
| $\text{CaO}$                         | 1.27                               | 1.38                   | 8.69                    | 8.97                        | .43                            | .23                    | 2.30                       |
| $\text{Na}_2\text{O}$                | 4.02                               | 4.08                   | 2.32                    | 3.05                        | 3.84                           | 3.90                   | 3.45                       |
| $\text{K}_2\text{O}$                 | 1.94                               | 2.58                   | .94                     | .46                         | 4.88                           | 4.51                   | 3.34                       |
| $\text{P}_2\text{O}_5$               | -                                  | -                      | .23                     | .36                         | -                              | -                      | -                          |
| Loss, Ig.                            | 2.18                               | .64                    | 5.48                    | 4.77                        | .73                            | .62                    | 1.19                       |
| Total                                | 97.17                              | 98.68                  | 100.31                  | 100.10                      | 99.29                          | 99.21                  | 99.40                      |
| $\text{H}_2\text{O (ppm)}$           | 1758                               | 983                    | 593                     | 450                         | 416                            | 403                    | 1454                       |
| $\text{H}_2\text{O (H}_2\text{O}^+)$ | 806                                | 0                      | -                       | -                           | 0                              | 0                      | 0                          |
| $\text{Fe (S)}$                      | 571                                | 7                      | 17                      | 24                          | 17                             | 24                     | 16                         |
| $\text{Fe (S)}$                      | 683                                | 8                      | 8                       | 15                          | 10                             | 13                     | 14                         |
| $\text{Fe (S)}$                      | 849                                | 21                     | 70                      | 96                          | 34                             | 65                     | 28                         |
| $\text{Fe (S)}$                      | 587                                | 5                      | 19                      | 19                          | 12                             | 44                     | 9                          |
| $\text{Fe (S)}$                      | 60                                 | 3                      | 57                      | 40                          | 1                              | 3                      | 3                          |
| $\text{Fe (S)}$                      | 58                                 | 2                      | 42                      | 37                          | 1                              | 3                      | 3                          |
| $\text{Fe (S)}$                      | -                                  | -                      | -                       | -                           | -                              | -                      | -                          |
| $\text{Fe (S)}$                      | -                                  | -                      | -                       | -                           | -                              | -                      | -                          |
| $\text{Fe (S)}$                      | -                                  | -                      | -                       | -                           | -                              | -                      | -                          |
| $\text{Fe (S)}$                      | 18                                 | 54                     | 21                      | 19                          | 139                            | 108                    | 85                         |
| $\text{Fe (S)}$                      | 172                                | 114                    | 314                     | 345                         | 42                             | 19                     | 255                        |
| $\text{Fe (S)}$                      | 117                                | 107                    | 81                      | 168                         | 248                            | 386                    | 116                        |
| $\text{Fe (S)}$                      | -                                  | -                      | -3                      | -3                          | -                              | -                      | -                          |
| $\text{Fe (S)}$                      | 4                                  | 5                      | 56                      | 48                          | 2                              | 3                      | 17                         |
| $\text{Fe (S)}$                      | 35                                 | 49                     | 31                      | 31                          | 43                             | 48                     | 56                         |
| $\text{Fe (S)}$                      | 10                                 | 9                      | 143                     | 77                          | 7                              | 8                      | 10                         |
| $\text{Fe (S)}$                      | 14                                 | 45                     | 356                     | 262                         | 14                             | 10                     | 41                         |
| $\text{Fe (S)}$                      | -                                  | 16(1)                  | -                       | -                           | 23(1)                          | 14(1)                  | -                          |
| $\text{Fe (S)}$                      | -                                  | 0(1)                   | -                       | -                           | 3(1)                           | 4(1)                   | -                          |
| $\text{Fe (S)}$                      | -                                  | 12(1)                  | -                       | -                           | 14(1)                          | 24(1)                  | -                          |
| $\text{Fe (S)}$                      | 5                                  | 3                      | 9                       | 18                          | 12                             | 5                      | 3                          |






Figure 5-1: Alkali: silica diagrams for Buchans Group flows (small dots), low-titanium diabase (stars) and Feeder Granodiorite (large dots). Dividing line between alkaline and subalkaline fields after Irvine and Baragar (1971).

oceanic tholeiites and have correspondingly lower Ni, Cr and K/Rb (Table 5-1).

In contrast, the lithologies and depositional environments of the Buchans Group volcanics, combined with the general chemical nature of Buchans Group flows, indicate that these rocks are most similar to modern calc-alkaline suites. A triangular FMA diagram, presented in Fig. 5-2 demonstrates a lack of iron enrichment, a characteristic of calc-alkaline suites.

The general elemental abundances of Buchans Group flows are also similar to those of calc-alkaline suites of modern island arcs.  $K_2O$ , an important element in discriminating volcanic suites, averages uniformly around 1% in Buchans Group mafic flows, similar to that of high alumina basalts (Fig. 5-3). According to the classification scheme of MacKenzie and Chappel (1972) for calc-alkaline rocks, Buchans Group mafic and intermediate flows span a range from high Alumina basalt to dacite. Other elemental abundances of Buchans Group flows are closely comparable to those of calc-alkaline suites of modern island arcs.

Diagrams involving the relatively immobile elements Ti, Zr (Figs. 5-4 and 5-5) show the bulk of Buchans Group mafic flows plotted in the field of calc-alkali basalts.

K:Rb ratios of Buchans Group flows encompass a range from 250 to 900, centering around 450 (Fig. 5-6), a




Figure 5-2: AFM diagram for Buchans Group flows (small dots), low-titanium diabase (stars) and Feeder Granodiorite (large dots). A =  $\text{Na}_2\text{O} + \text{K}_2\text{O}$ , F = total Fe as  $\text{FeO}$ , M =  $\text{MgO}$ . Dividing line between calc-alkaline and tholeiitic suites after Irvine and Baragar (1971).




Figure 5-3: K<sub>2</sub>O: SiO<sub>2</sub> classification of calc-alkaline rocks after Mackenzie and Chappel (1972). Buchans Group flows (dots) and low-titanium diabase (stars).




Figure 5-4: Ti:Zr diagram for Buchans Group basalts (dots), low-titanium diabase (solid stars) and high-titanium diabase (open stars). Fields A to D after Pearce and Cann (1973). A = low potash tholeiites, C = calc-alkali basalts, D = ocean floor basalts, B = overlap of fields A, C and D.




Figure 5-5:  $\text{Ti}/100$ :  $\text{Zr}$ :  $\text{Sr}/2$  diagram for Buchans Group basalts. Fields A, B and C after Pearce and Cann (1973). A = island arc basalts, B = calc-alkali basalts, C = ocean floor basalts.




Figure 5-6: K: Rb diagram for Buchans Group Flows (small dots) low titanium diabase (stars) and Feeder Granodiorite (large dots).

fairly typical average for modern island arc sequences.

A histogram of  $\text{SiO}_2$  abundances in Buchans Group flows is given in Fig. 5-7. The diagram shows a relative paucity of samples of "intermediate" composition yielding a weakly bimodal distribution. Sampling bias obviously has a major effect on this diagram, however from general observation the author considers the dominance of mafic and felsic volcanics over those of intermediate composition to be a reality. Compared to other Newfoundland calc-alkaline mafic to felsic volcanic suites, the Buchans Group has relatively more abundant intermediate volcanics (Strong, 1975).

A correlation matrix for Buchans Group flows is given in Table 5-2. In general, the predictable elemental associations are observed amongst both major and trace elements. Of particular interest to this study is the behavior of the base metals and barium. Pb shows a near perfect correlation with Pb(S) and little correlation with any other element. This would seem to indicate that the bulk of Pb in Buchans Group flows occurs almost exclusively as minute sulphides which show preference for no specific host mineral phase. Cu shows a very strong correlation with Cu(S) and weaker association with Fe, Ca and Mg. This seems to indicate that it occurs as minute sulphide within Fe, Ca and Mg-bearing phases (e.g. chlorite,






Figure 5-7: Histogram of  $\text{SiO}_2$  contents of Buchans Group flows ( $n = 56$ ). <sup>2</sup> The relative lack of samples of intermediate composition is not a function of sampling error.



epidote, calcite, plagioclase) and also shows limited substitution for Fe and Mg in these minerals. The correlation of Zn to Zn(S) is not nearly as strong as the previous associations suggesting a complex occurrence of Zn both as minute sulphide and also widely substituted for Fe in various phases.

The behavior of Ba in flows is important to later discussions. K is the closest associate of Ba, with other elements showing little consistent correlation. This would seem to indicate that the major controlling factor on Ba distribution is the abundance of  $K_2O$  and that Ba most commonly occupies K lattice sites.

Harker diagrams, for Buchans Group flows, plotting anhydrous weight percent and ppm of the various oxides, elements and loss on ignition against silica, are presented in Fig. 5-8. Also plotted on these diagrams are samples of low-titanium diabase and the Feeder Granodiorite. Though Harker diagrams were previously presented by the author (Thurlow, 1973), the diagrams presented herein are considered much more reliable as fragmental rocks and altered flows are omitted from the data.

The Harker diagrams are, for the most part, self-explanatory and general comments are outlined in point form below.




Figure 5-8: Anhydrous Harker diagrams for Buchans Group flows (small dots), low-titanium diabase (stars) and Feeder Granodiorite (large dots).

Figure 5-8 (continued)




Figure 5-8 (continued)




Figure 5-8 (continued)




Figure 5-8 (continued)






Figure 5-8 (continued)




Figure 5-8 (continued)





Figure 5-8 (continued)



- 1) Buchans Group volcanic rocks span a compositional range from basalt to rhyolite but andesitic varieties are somewhat less common than more mafic and felsic end members.
- 2) The variation of all elements is that which is expected from normal petrological considerations. Most major elements decrease with increasing silica, in part because of simple dilution by silica (the "constant sum effect"). Only  $\text{Na}_2\text{O}$  and  $\text{K}_2\text{O}$  increase with increasing silica content. Among the trace elements, Ni, Cr, V, Sr, Cu and Cu(S) decrease and the large cations Rb, Ba and Zr increase with increasing silica. Somewhat surprisingly, Zn, Zn(S), Pb, Pb(S) and Co show little variation across the range of silica concentrations.
- 3) Anomalous scatter in some diagrams is probably largely due to post-depositional migration of elements. The elements most strongly affected (e.g. K, Rb, Ca, Sr) are those which are most easily mobilized in the secondary environment.
- 4) Despite scatter on some diagrams it is evident that all Buchans Group flows form a consanguineous mafic to felsic sequence. As such, there is no evidence to suggest that mafic and felsic members are not magmatically related nor is there any evidence to suggest that any portion of the Buchans Group has evolved from a magma distinct from the major magma source.

- 5) Both the low-titanium diabase and the Feeder Granodiorite plot entirely within the trends of the Buchans Group flows. This suggests a genetic link between these three lithologic groups.
- 6) Two types of variations are visible among elements which decrease in abundances with increasing silica. Some elements (e.g. Fe) show consistent and regular decrease whereas other elements (e.g. Mg, Ni, Cr, V) decrease rapidly at lower silica levels and less rapidly at higher silica levels. In the latter case this must indicate rapid removal of a specific phase(s) with increasing differentiation. Since no olivine has been identified at Buchans, this probably reflects fractionation of augite.

### 5.3. Chemical Variations in Flows as a Function of Stratigraphic Position

The previous sections have dealt with overall compositional variations within the Buchans Group. In this section, variations in the composition of flows are examined in order to determine whether variations in chemistry have any stratigraphic significance. Such variations, if present, presumably would reflect changing magmatic conditions. If such variations are present, it is important to determine whether they are of significance with respect to mineralization or merely a result of normal petrologic processes. Of

particular interest in this study is whether unaltered rhyolite flows in the ore-bearing Lucky Strike and Oriental ore horizon sequences are chemically different from flows elsewhere in the stratigraphic sequence.

Chemical data on mafic and intermediate flows of the Footwall Basalt, Ski Hill Sequence, and Upper Buchans Subgroup were plotted on one set of diagrams and data for felsic flows of the Lucky Strike-Oriental Sequences, Prominent Quartz Sequence and Upper Buchans Subgroup on a second series of diagrams, yielding a total of over 50 diagrams. All elements plotted were ratioed to  $Al_2O_3$  based on the assumption that significantly greater abundances of any element in a given sample may "dilute" the abundances of elements under scrutiny and consequently introduce apparent scatter to the data.

#### 5.3.1. Variations in Compositions of Mafic and Inter-

##### mediate Flows Through the Stratigraphic Sequence:

The distribution and number of samples (34) of these groups of rocks is not wholly adequate to offer firm conclusions as to whether any given stratigraphic unit differs significantly from others. In general, as determined from extensive core logging throughout the Buchans Group, there is no secular change in the composition of flows through the stratigraphy. Most mafic units are wholly basaltic and only the Ski Hill sequence contains significant amounts of andesite. Within

the basaltic units, the chemical diagrams plotted indicate that there is no significant difference for any elements between units at differing stratigraphic levels. For example, the large cations (e.g. K, Rb, etc.) do not increase in abundance with stratigraphic height as has been found in some island arc sequences (e.g. Kuno, 1966; Sugimura, 1968; Jakes and White, 1971; Kean and Strong, 1973). Samples of more intermediate flows from the Ski Hill sequence form a continuum on the variation diagrams with more mafic counterparts from other units indicating that differences in elemental abundances are not due to any geochemical peculiarity but instead are a function of normal magmatic processes.

In essence, the limited amount of geochemical data seems to indicate that there is no geochemical anomaly associated with any particular mafic unit. The Ski Hill sequence, which is the mafic unit most closely associated with mineralization, is "anomalous" only in that it contains a greater abundance of truly intermediate rocks as opposed to the other largely basaltic sequences.

#### 5.3.2. Variations in Compositions of Felsic Flows

Through the Stratigraphic Sequence: Almost all elements show no essential differences in composition between felsic flows of the Lucky Strike-Oriental sequences, Prominent Quartz sequence and Upper Buchans Subgroup. In

general, flows of the ore-bearing Lucky Strike-Oriental sequences are compositionally more homogeneous than other flows though this may only reflect the volumetrically less abundant and spatially more restricted nature of these flows. MgO and MnO tend to be less abundant in the ore horizon flows than in the two other groups (Fig. 5-9). This probably reflects lesser quantities of chlorite after amphibole. CaO and  $\text{Fe}_2\text{O}_3$  tend to be slightly more abundant in the Prominent Quartz Sequence rhyolites than in the other two groups (Fig. 5-9) probably reflecting the greater abundance of epidote after plagioclase visible in these rocks.

Apart from the above-mentioned small differences, felsic flows of the Lucky Strike-Oriental sequences are similar to other felsic flows of the Buchans group. The major ore-forming elements, Zn, Pb and Cu are not present in greater abundances. Ba is marginally higher in ore horizon rhyolites but shows a strong affinity to  $\text{K}_2\text{O}$  (correlation = .87).  $\text{K}_2\text{O}$  is likewise marginally higher in the ore horizon rhyolites.

Tasumi and Clark (1972) and Ishihara (1974) have shown (on the basis of a limited number of analyses) that felsic lavas genetically related to mineralization in the Hokuroku district are somewhat more differentiated than lavas unrelated to mineralization. A plot of silica vs. Thornton and Tuttle's differentiation index for Buchans Group






Figure 5 - 9 : CaO, MgO and MnO variations in flows as a function of stratigraphic position.

felsic flows is presented in Fig. 5-10. On the basis of a limited number of analyses, it appears that for a given silica content, felsic flows of the ore-bearing Lucky Strike-Oriental ore horizon sequences are more "differentiated" than similar flows of other formations. The number of samples involved in this comparison is not sufficient, to firmly establish whether this distinction is actually statistically significant. Many more analyses combined with detailed petrographic work are required to determine the validity of this hypothesis. Such work could prove fruitful for the outcome would have a bearing on the overall source of metals, whether magmatic or leached from country rock.

5.3.3. Some Comments on the Chemistry of "White Rhyolite" at Buchans: The presence of "white rhyolite" associated with many massive sulphide deposits is of theoretical and practical exploration interest. As indicated in Chapter Four, "white rhyolite" at Buchans displays a more tenuous relationship to mineralization than in the Kuroko deposits but, nonetheless, is generally linked to the presence of ore.

Two analyses of Buchans "white rhyolite" are presented in Table 5-3. Compared to associated unaltered rhyolite, and holding alumina constant, the "white rhyolite" has suffered massive introduction of  $K_2O$ ,  $SiO_2$  and Ba and removal of  $Fe_2O_3$ ,  $Na_2O$  and  $CaO$ . Mineralogically this is




Figure 5-10: SiO<sub>2</sub>: differentiation index plot comparing unaltered ore horizon rhyolites to rhyolites at other stratigraphic positions.

Table 5-3: Composition of "white rhyolite" compared to other ore horizon rhyolites

|                                | WHITE RHYOLITE |           | 10 Ore Horizon<br>Rhyolites |
|--------------------------------|----------------|-----------|-----------------------------|
|                                | 1083-586       | 1730-1203 |                             |
| SiO <sub>2</sub>               | 73.3           | 75.6      | 72.7                        |
| TiO <sub>2</sub>               | .39            | .22       | .31                         |
| Al <sub>2</sub> O <sub>3</sub> | 12.91          | 10.29     | 13.68                       |
| Fe <sub>2</sub> O <sub>3</sub> | .03            | .01       | .1.94                       |
| MnO                            | .01            | .01       | .04                         |
| MgO                            | .60            | .01       | .34                         |
| CaO                            | .44            | .76       | 1.12                        |
| Na <sub>2</sub> O              | 1.52           | .36       | 4.47                        |
| K <sub>2</sub> O               | 7.64           | 7.93      | 3.06                        |
| L.O.I.                         | 1.68           | 1.62      | 1.49                        |
| Total                          | 98.52          | 96.81     | 99.15                       |
| Ba                             | 2982           | 4639      | 1175                        |
| Pb                             | 3              | 32        | 15                          |
| Pb(S)                          | 6              | 69        | 16                          |
| Zn                             | 19             | 455       | 28                          |
| Zn(S)                          | 22             | 460       | 13                          |
| Cu                             | 1              | 1         | 6                           |
| Cu(S)                          | 2              | 5         | 6                           |
| Rb                             | 62             | 63        | 52                          |
| Sr                             | 49             | 119       | 192                         |
| Zr                             | 166            | 147       | 146                         |
| Ni                             | 1              | 2         | 3                           |
| Co                             | 27             | 42        | 26                          |
| Cr                             | 0              | 0         | 9                           |
| V                              | 10             | 15        | 22                          |

reflected in conversion of normal rhyolite to an entirely secondary assemblage composed essentially of quartz and K-feldspar. This mineralogy contrasts with the quartz-sericite-rich assemblage of the Japanese "white rhyolite" (Date and Taniwura, 1974).

5.3.4. Summary: Based upon a limited number of analyses of mafic intermediate and felsic flows (total 54 analyses) there appear to be no distinct changes in composition in Buchans Group volcanics throughout the stratigraphic sequence. The mafic and felsic units most closely associated with mineralization are chemically very similar to other such units throughout the Buchans Group and do not appear to contain abnormal amounts of base metal or barium. As such, there appear to be no magmatic changes associated with ore horizon rocks that have a bearing on the presence of mineralization. Similar results to these were obtained at Noranda by Larson and Webber (1977).

#### 5.4. Geochemical Comparison of Flows and Cogenetic Fragmental Rocks

In any rock geochemical study it is important to distinguish and separate those processes which cause chemical change as a result of mineralization from other processes unrelated to mineralization. In an effort to understand the gross post-magmatic chemical changes involved in

pyroclastic rocks of the Buchans Group, a study was undertaken to compare the chemistry of massive flows with associated flanking pyroclastic rocks. This study was based on two reasonably sound assumptions, i.e., 1) that flows and stratigraphically related pyroclastic rocks were originally derived from chemically similar magmas (i.e., are cogenetic), and 2) that the composition of flows more closely approximates that of the original magma. In view of the overall chemical coherence of all Buchans Group volcanic rocks and in view of similarities of phenocryst composition between flows and associated pyroclastics, assumption No. 1 appears valid. Petrographic comparison of flows and related pyroclastics indicates that the more permeable fragmental rocks have invariably undergone greater degrees of alteration than related flows and thus, assumption No. 2 also appears valid.

Chemical data on flows and related pyroclastics were plotted for five groups of lithologies, i.e., mafic and intermediate rocks of the Footwall Basalt and Ski Hill horizons and felsic rocks of the Lucky Strike-Oriental sequences, Prominent Quartz sequence and Upper Buchans Subgroup, yielding a total of over 100 diagrams. Flows and related fragmentals were plotted on the same diagrams to assist visual comparison. All elements were ratioed to  $Al_2O_3$  based on the assumption that  $Al_2O_3$  remained relatively inert during chemical change. The effect of this ratio is

7

to eliminate effects of variable addition of extraneous elements (and consequent variable dilution of elements under scrutiny) which would cause apparent increased scatter of data points.

In virtually all cases, flows form more coherent groups than related pyroclastics, supporting the contention that they have undergone less post-magmatic chemical change. Apart from differences in scatter, differences in general abundances between the two groups were normally not significant. Only Co showed consistent behavior in all groups, being higher in flows relative to cogenetic pyroclastic rocks.

Among the mafic and intermediate rocks of the Footwall Basalt and Ski Hill sequence, Ca is generally higher in flows (unavoidable calcite amygdules) as is Mn and Co. There is a tendency for Zn and K to have been added to the related fragmentals in each case.

Among felsic sequences compared, all had higher Na and Co in felsic flows as compared to cogenetic crystal-vitric tuffaceous rocks. Apart from these general tendencies, the felsic rocks of the Prominent Quartz sequence and Upper Buchans Subgroup showed no essential differences for all other elements.

In contrast, the Lucky Strike-Oriental felsic flows and crystal-vitric tuffs show a number of additional




Figure 5-11: Geochemical comparison of ore horizon dacitic tuffs (dots) and rhyolite flows (stars).



differences. Dacitic crystal-vitric tuffs tend to be higher than flows in Mn, Mg, Ca, loss on ignition, Rb and Cu and lower in Co, Si and Na (the more significant of these are shown in Fig. 5-11). In view of the similar phenocryst mineralogy and proportions, and similar alteration of phenocrysts in these two groups of rocks, it can be reasonably assumed that all of these elemental differences reflect post-magmatic changes in chemistry of matrix glass in the crystal-vitric tuffaceous rocks. From Fig. 5-11 it can be seen that hydration and subsequent diagenetic-metamorphic alteration of groundmass glass to sericite and chlorite has resulted in addition of volatiles (principally  $H_2O$ ) and MgO and removal of  $Na_2O$  from the glass. In Section 6.4. it is demonstrated using factor analysis that variations in the degree of these chemical changes are responsible for a large proportion of the total chemical variation within the dacite.

These results bear a strong similarity to those of Sopuck (1977), who compared matrix and clasts in the same rocks, from a sequence of unmineralized volcanics. His data show very marked differences in silica content between matrix and clasts either because they were not similar originally and/or because the matrix has undergone extreme alteration. Clast-matrix study at Buchans is not possible as coarse fragmental rocks are normally polyolithic. It is emphasized that flows and related pyroclastics in this

study have essentially similar silica contents allowing  
meaningful comparison.

## CHAPTER SIX

### GEOCHEMISTRY OF THE ORE HORIZON DACITIC TUFFS

#### 6.1. Introduction

Dacitic crystal-vitric tuffs of the Lucky Strike-Oriental ore horizon sequences are the most widespread lithologic unit which display intimate relationships to massive sulphide mineralization. These units therefore represent the lithologic units most likely to exhibit regional chemical variations which may be a function of mineralization and thus of interest to exploration. The chemistry of the ore horizon tuffs is examined somewhat more closely in this chapter in an effort to understand the processes responsible for chemical variation and to derive geochemical criteria of use in exploration.

#### 6.2. General Chemical Abundances in the Ore Horizon Dacitic Tuffs

The average chemical abundances of the ore horizon dacitic tuffs from the Lucky Strike, Clementine and Oriental areas is presented in Table 6-1 and histograms of abundances for each element in Fig. 6-1. Most elements show normal population distributions with the exceptions of more lognormal Cu, Cu(S), Pb, Pb(S), Zn, Zn(S), Mg, Ni, Co and V.

Table 6-1: Average chemical composition of ore horizon dacitic tuffs.

|                                | Lucky Strike<br>Horizon<br>Dacitic Tuffs | Oriental<br>Horizon<br>Dacitic Tuffs | Clementine<br>Area<br>Dacitic Tuffs |
|--------------------------------|------------------------------------------|--------------------------------------|-------------------------------------|
| SiO <sub>2</sub>               | 71.0                                     | 67.0                                 | 71.7                                |
| TiO <sub>2</sub>               | .23                                      | .32                                  | .24                                 |
| Al <sub>2</sub> O <sub>3</sub> | 13.68                                    | 15.56                                | 13.75                               |
| Fe <sub>2</sub> O <sub>3</sub> | 2.23                                     | 2.66                                 | 2.15                                |
| MnO                            | .08                                      | .06                                  | .06                                 |
| MgO                            | 1.23                                     | 2.56                                 | 1.20                                |
| CaO                            | 2.17                                     | 2.05                                 | 1.97                                |
| Na <sub>2</sub> O              | 2.69                                     | 2.17                                 | 1.79                                |
| K <sub>2</sub> O               | 2.82                                     | 2.72                                 | 3.20                                |
| P <sub>2</sub> O <sub>5</sub>  | .06                                      | -                                    | -                                   |
| Loss. Ig.                      | 2.97                                     | 3.90                                 | 3.76                                |
| Total                          | 99.16                                    | 99.00                                | 99.32                               |
| Ba                             | 1657                                     | 1380                                 | 1042                                |
| Ba (SO <sub>4</sub> =)         | -                                        | -                                    | -                                   |
| Pb                             | 29                                       | 41                                   | 14                                  |
| Pb(S)                          | 24                                       | 77                                   | 24                                  |
| Zn                             | 109                                      | 160                                  | 25                                  |
| Zn(S)                          | 34                                       | 120                                  | 8                                   |
| Cu                             | 5                                        | 13                                   | 1                                   |
| Cu(S)                          | 2                                        | 16                                   | 2                                   |
| S                              | -                                        | -                                    | -                                   |
| Ag                             | -                                        | -                                    | -                                   |
| Hg                             | -                                        | -                                    | -                                   |
| Rb                             | 70                                       | 83                                   | 99                                  |
| Sr                             | 222                                      | 250                                  | 127                                 |
| Zr                             | 122                                      | 156                                  | 140                                 |
| Mo                             | 43                                       | -                                    | -                                   |
| Ni                             | 3                                        | 4                                    | 4                                   |
| Co                             | 5                                        | 16                                   | 19                                  |
| Cr                             | 24                                       | 9                                    | 2                                   |
| V                              | 19                                       | 16                                   | 15                                  |
| Th                             | 16(12)                                   | 19(23)                               | 16                                  |
| U                              | 4(12)                                    | 4(23)                                | 4                                   |
| Ga                             | 13(12)                                   | 16(23)                               | 13                                  |

Fig. 6-1: Histogram for all elements determined in ore horizon dacitic tuffs.




Fig. 6-1 (con't)




Fig. 6-1 (con't)





Fig. 6-1 (con't)





### 6.3. Correlation Matrix for Ore Horizon Dacitic Tuffs

A correlation matrix for ore horizon dacitic tuffs is given in Table 6-2. In general, most elemental associations are those normally predictable on petrochemical grounds. Nonetheless, there exist a few unusual features including a complete lack of Ba correlation with all other elements including  $K_2O$ . This is anomalous in view of the strong Ba:  $K_2O$  coherence in associated rhyolites and indicates that the process(es) controlling Ba distribution are quite different in these two suites of closely related rocks. Among the base metals, there is little correlation between Cu and the strongly correlative Pb-Zn pair.

Lengthy study of the correlation matrix reveals many element interactions which can be related to geochemical processes which acted upon the dacitic tuffs. Due to the complexity of these interactions, factor analyses was undertaken to better understand inter-element associations.

### 6.4. R-Mode Factor Analysis of the Ore Horizon Dacitic Tuffs - Major Elements

A large number of geochemical studies have proven the merit of factor analysis as a valuable tool in understanding processes responsible for geochemical variations. In order to identify geologic processes associated with various factors, it is necessary to understand the mode of

- 255 -

[illegible]

occurrence of the various elements. Certainly, in most geochemical studies, the mode of occurrence of the major elements is much better understood than that of the trace elements. For this reason, R-mode factor models were computed first for the major elements in order to facilitate identification of geochemical processes responsible for chemical variation in the dacites. Apart from this reason, major elements only were included in the first factor models in order to reduce the number of factors required to reasonably explain the variance in the dacite (factor analysis treats all variables as statistical equals, though in geochemical studies, the major elements comprise, by definition, the bulk of the elements of which the rock is composed). Factor matrices presented have undergone a varimax rotation similar to that described by Koch and Link (1971).

A variety of major element factor models were computed for the ore horizon dacite. The six factor model accounting for 94% of the variance in the dacite was selected as (by far) the most geologically explicable model. The rotated factor matrix is presented in Table 6-3.

Factor 1 (accounting for 16% of the total variance) accounts for almost all the MgO variation and a significant proportion of the Na<sub>2</sub>O and loss on ignition in the ore horizon dacite. Microprobe examination of the dacite indicates that the bulk of the MgO occurs in chlorite after matrix glass

Table 6-3: Rotated factor matrix for dacitic tuffs of the Lucky Strike ore horizon sequence. The six factor matrix accounts for 94% of the data variability.

| Variable                       | 1<br>Chlorite<br>After<br>Glass<br>(15.73%) | 2<br>Hematite,<br>Leucoxene,<br>etc.<br>(16.95%) | 3<br>Secondary<br>Calcite<br>(17.17%) | 4<br>Silica-<br>Alumina<br>(17.49%) | 5<br>Sericite<br>After<br>Glass<br>(14.78%) | 6<br>MnO<br>(11.96%) |
|--------------------------------|---------------------------------------------|--------------------------------------------------|---------------------------------------|-------------------------------------|---------------------------------------------|----------------------|
| Fe <sub>2</sub> O <sub>3</sub> | .15                                         | .83                                              | -.14                                  | -.19                                | .04                                         | .35                  |
| TiO <sub>2</sub>               | -.02                                        | .93                                              | .06                                   | -.13                                | -.09                                        | -.20                 |
| SiO <sub>2</sub>               | -.15                                        | -.30                                             | .41                                   | .80                                 | .15                                         | -.05                 |
| CaO                            | -.09                                        | .07                                              | -.95                                  | -.09                                | .01                                         | .20                  |
| K <sub>2</sub> O               | -.01                                        | .08                                              | .04                                   | -.38                                | -.90                                        | -.05                 |
| MgO                            | .96                                         | .10                                              | .03                                   | -.15                                | -.06                                        | -.05                 |
| Al <sub>2</sub> O <sub>3</sub> | .13                                         | .12                                              | .01                                   | -.89                                | -.23                                        | -.18                 |
| Na <sub>2</sub> O              | -.54                                        | .03                                              | .34                                   | -.05                                | .71                                         | .16                  |
| L.O.I.                         | .53                                         | -.11                                             | -.71                                  | -.26                                | -.25                                        | -.10                 |
| MnO                            | -.09                                        | .01                                              | -.12                                  | .11                                 | .10                                         | .96                  |

(minor amounts also occur as chlorite after hornblende). In Section 5.4. it was demonstrated that devitrification of matrix glass involved hydration, removal of soda and addition of MgO. This factor is thus easily identified as representing alteration of original matrix glass to chlorite with release of  $\text{Na}_2\text{O}$ .

Factor 2 (accounting for 17% of the total variance) contains large loadings for  $\text{Fe}_2\text{O}_3$  and  $\text{TiO}_2$ . These elements occur almost exclusively in trace quantities of rutile, ilmenite, leucoxene, hematite and pyrite, all of which, in the author's experience, are randomly distributed in the unaltered tuffs. Factor 2 indicates that the occurrence of these minerals is not related to any geochemical process which has affected the dacitic tuffs nor is their occurrence linked to the presence of any other minerals.

Factor 3 (accounting for 17% of the total variance) accounts for almost all the CaO variation in the dacite and half of the loss on ignition variation. Calcite is the only mineral which can satisfy the loadings of this factor with only a minor contribution from prehnite. Calcite occurs principally as alteration of plagioclase and as minute calcite veinlets transecting all rocks. Factor 3 is thus a "secondary calcite factor".

Factor 4 (accounting for 17% of the total variance) has high antipathetic loadings for  $\text{SiO}_2$  and  $\text{Al}_2\text{O}_3$  and accounts

for most of the variation of these two elements. As no other elements are significantly affected, this factor represents substitution of  $\text{Al}_2\text{O}_3$  for  $\text{SiO}_2$  (and vice versa) as a result of the varying presence of some mineral or as a result of some secondary geochemical process. Such substitutions are possible as a result of variation in both plagioclase and chlorite compositions and this factor is tentatively correlated to such compositional variations. Variations in the proportions of quartz and plagioclase may also contribute to this factor.

Factor 5 (accounting for 15% of the total variance) accounts for almost all the  $\text{K}_2\text{O}$  variation in the dacite. Potash occurs mainly in sericite (after matrix glass) with only minor amounts in the much less abundant K-feldspar. Identification of this factor as a "sericite after glass" factor satisfactorily explains the sympathetic relationship between  $\text{K}_2\text{O}$ , loss on ignition and  $\text{Al}_2\text{O}_3$ . Their antipathetic relationship with  $\text{Na}_2\text{O}$  indicates that soda has been expelled from glass during sericitic alteration, a fact confirmed by electron microprobe scans.

Factor 6 (accounting for 12% of the total variance) accounts for virtually all MnO variation with no other elements significantly affected. As such, it must be interpreted as post depositional solution and redistribution of MnO.

The release of  $\text{Na}_2\text{O}$  during alteration of matrix glass to chlorite and sericite (factors 1 and 5 above) is of

interest because none of the other factors involve addition of soda. It must be concluded that soda originally present in glass was largely removed from the dacitic tuffs during alteration. In this regard, it is interesting to refer back to Figure 5-12 (Section 5-4) in which it was demonstrated that soda has been significantly depleted from the dacitic tuffs as compared to the rather sodic cogenetic rhyolite flows. Since none of the lithologies overlying or underlying the dacite have undergone spilitic alteration it appears that loss of soda from the dacite must have occurred at deposition, with soda in glass being released to sea water. Thus it appears that alteration of glass (probably originally to clay minerals) occurred shortly after deposition and prior to deposition of overlying basalt flows. Regional depositional soda depletion of originally glassy pyroclastics has been suggested as a process possibly operative in the vicinity of the New Brunswick massive sulphide deposits (Govett *et al.*, 1974, p. 83). In Section 7.2. it is shown that this process, though operative at Buchans, bears no relationship to mineralization.

A consequence of this interpretation is that the inverse correlation between  $\text{Na}_2\text{O}$  and  $\text{K}_2\text{O}$  in the dacitic tuffs is not a function of local redistribution in glass of  $\text{Na}_2\text{O}$  and  $\text{K}_2\text{O}$  as previously suggested by the author (Thurlow, 1973; Thurlow *et al.*, 1975). Instead, glass-rich samples contain high  $\text{K}_2\text{O}$  and low  $\text{Na}_2\text{O}$  whereas glass-poor samples (i.e., crystal-

rich samples) contain relatively lower  $K_2O$  and higher  $Na_2O$  ( $Na_2O$  in plagioclase). Consequently the inverse soda:potash correlation is a function of the glass:crystal ratio in the dacite. It follows that anomalously low soda:potash ratios in the dacite are a function both of the degree of depositional alteration of glass and of the glass:crystal ratio and that low values of this ratio may not bear any relationship to the presence of mineralization. It is demonstrated in Section 7.2. that none of the above six factors display any spatial distribution related to the presence of ore.

It is evident from the above that factor analysis is a powerful aid in identifying and understanding processes causing chemical variation.

#### 6.5. R-mode Factor Analysis of the Ore Horizon Dacitic Tuffs - Major and Trace Elements

In this section, the behavior of trace elements is considered with regard to the processes responsible for chemical variation in the dacite as determined in the previous section. Factor models were computed first for all major elements and barium and subsequently for all major and trace elements combined.

When barium is included in the major element factor models it forms a factor independent of all other elements. When the number of factors is reduced so as to reveal barium



associations with other elements, it does not show correlation with any of the previously determined factors. It is concluded that the process(es) responsible for variation in barium concentration is not among those processes mentioned in the previous section.

The remainder of the trace elements behave fairly predictably in the combined major and trace elements factor models (a representative nine factor model is shown in Table 6-4). Rubidium variation is virtually entirely incorporated into the "sericite after glass" factor and zirconium enters the "rutile-leucoxene" factor. Strontium forms a weak independent factor which, in some factor models, also accounts for up to one third of the barium variation. The meaning of this factor is obscure, as Buchans barite is Sr-poor.

Factor analysis reveals that the variation in the base metals Pb and Zn is quite independent from that of Cu. Pb, Zn and Zn(S) show moderate to strong coherence with the "chlorite after glass" factor. Pb(S) variation is somewhat more complex than that of Zn(S) as it is associated with both the "chlorite after glass" and "calcite" factors. It is concluded that silicate zinc and lead are largely incorporated into the chlorite lattice, that sulphide zinc occurs primarily in association with chlorite and that sulphide lead occurs primarily in association with chlorite and calcite.

In contrast, Cu variation is strongly linked to that of Ni and V as these elements form a strong factor which is mildly linked to a Cu(S)-Cu factor (Table 6-4). This may indicate that these factors represent variations in proportion of a submicroscopic sulphide phase.

Table 6-4: Rotated factor matrix for dacitic tuffs of the Lucky Strike ore horizon sequence. The nine factor matrix accounts for 83% of the data variability.

|                                | FACTOR |     |     |     |     |     |     |     |     |
|--------------------------------|--------|-----|-----|-----|-----|-----|-----|-----|-----|
|                                | 1      | 2   | 3   | 4   | 5   | 6   | 7   | 8   | 9   |
| Fe <sub>2</sub> O <sub>3</sub> | 00     | -02 | 00  | 23  | 90  | 03  | 12  | 04  | -15 |
| TiO <sub>2</sub>               | -18    | -02 | 01  | -16 | 86  | 04  | -13 | -09 | 10  |
| SiO <sub>2</sub>               | 53     | -03 | -08 | -50 | -42 | -14 | -04 | 27  | 02  |
| CaO                            | 00     | -17 | -03 | 89  | 07  | 14  | 11  | 04  | 13  |
| K <sub>2</sub> O               | -85    | -04 | 08  | 00  | 08  | -17 | -03 | 07  | 07  |
| MgO                            | -21    | 78  | 20  | 17  | 18  | -02 | -04 | 00  | -07 |
| Al <sub>2</sub> O <sub>3</sub> | -74    | 07  | 12  | 06  | 21  | 27  | 00  | -22 | -03 |
| Na <sub>2</sub> O              | 60     | -37 | -03 | -37 | 06  | 24  | 12  | -27 | -12 |
| L.I.                           | -40    | 35  | -01 | 75  | -02 | -20 | -03 | -03 | 03  |
| MnO                            | 36     | -14 | 08  | 25  | 13  | 05  | 67  | 26  | 05  |
| Zr                             | -44    | -08 | -01 | -18 | 40  | 62  | -12 | -03 | -02 |
| Sr                             | 10     | -08 | 11  | 10  | -01 | 82  | 02  | 15  | 18  |
| Rb                             | -87    | -03 | 06  | 13  | 03  | 10  | -03 | 15  | -06 |
| Zn                             | 02     | 91  | 05  | -13 | -08 | -15 | 03  | 02  | -01 |
| Cu                             | 01     | 11  | 89  | 07  | -03 | 00  | 05  | -37 | 00  |
| Ba                             | 00     | 07  | 01  | 11  | -04 | 09  | 02  | 05  | 96  |
| Cu(s)                          | 11     | -05 | 16  | 02  | 06  | -11 | -03 | -86 | -07 |
| Pb(s)                          | 01     | 56  | -03 | 43  | 04  | 06  | -12 | -23 | 11  |
| Zn(s)                          | 24     | 62  | -02 | -17 | 00  | -37 | -30 | 16  | 08  |
| Pb                             | -02    | 93  | 02  | 06  | -08 | 10  | 04  | 00  | 10  |
| Ni                             | -10    | 02  | 96  | 00  | 02  | 06  | -06 | 08  | 00  |
| Co                             | 06     | -04 | 44  | -02 | -06 | -11 | -70 | 27  | -04 |
| Cr                             | -01    | 00  | 08  | -09 | -04 | 16  | 73  | 08  | -02 |
| V                              | -09    | 04  | 97  | 03  | 02  | 02  | 04  | 01  | 00  |

## CHAPTER SEVEN

### APPLIED ASPECTS OF THE GEOCHEMISTRY OF ORE HORIZON DACITIC TUFFS

#### 7.1. Geochemical Comparison of Ore Horizon Dacitic Tuffs to Other Dacitic Tuffs

The average chemical composition of ore horizon dacitic tuffs as compared to dacitic tuffs at other non ore-bearing horizons is given in Table 5-1. A quick scan of this table reveals little visible difference in composition for most elements except possibly for Zn and Pb and a strong apparent enrichment of Ba in ore horizon dacitic tuffs. Histograms were prepared in order to more accurately assess the validity of these comparisons.

A histogram of Ba abundances in ore horizon dacitic tuffs as compared to other petrographically similar dacitic tuffs is given in Fig. 7-1. For reasons given in subsequent sections, all samples of ore horizon tuff from within 30 m of ore were excluded from this comparison. The histogram shows two distinct normally distributed Ba populations, highest in the ore horizon dacitic tuffs.

Discriminant functions analysis was undertaken in order to enhance the distinction between ore horizon and non ore horizon dacitic tuffs, and to identify other elements which might significantly differ between these groups.

First, using dacitic tuffs from the Oriental and Lucky Strike




Figure 7-1: Histogram of Ba abundances in ore horizon dacitic tuffs (n=111) as compared to non-ore horizon dacitic tuffs (n=55). Ore horizon samples from within 30 m of ore are omitted. Nine samples of ore horizon tuff exceed 2500 ppm and are not plotted. Area of overlap not coloured.

areas, a discriminant function was calculated which indicated that ore horizon tuffs could be distinguished with a high degree of confidence from non-ore horizon tuffs. In order to test this practical validity of this formula, discriminant scores of 13 samples of Clementine area tuff (from areas 3 or more kilometers from Lucky Strike) were calculated. The results show that all of these samples were successfully classified as ore horizon tuffs. A new discriminant function was thus calculated including the samples of Clementine area dacitic tuff.

A histogram of discriminant scores, given in Fig. 7-2 shows the separation of ore horizon and non-ore horizon dacitic tuffs. The data from this plot (Table 7-1) indicates that lower  $\text{Fe}_2\text{O}_3$  and higher Mn in the ore horizon dacitic tuffs are significant as well as Ba in contributing to the discrimination.

From these data, it is evident that ore horizon dacitic tuffs can be distinguished from other lithologically similar dacitic tuffs with a fair degree of confidence either by simply comparing Ba abundances or by using a 24 element discriminant function. The latter can be easily computed for any new samples using any calculator. The degree of confidence ascribed to classification of unknowns can be empirically determined from the existing test cases. Using Ba abundances only, it can be stated that any sample in excess of 1000 ppm Ba belongs to the ore horizon, with 95% confidence. Using




Figure 7-2: Histogram of discriminant scores.  
Ore horizon dacitic tuffs, stippled (n = 134)  
Non-ore horizon dacitic tuffs, ruled (n = 46)

Table 7-1: Discriminant function data. Sign convention in first column; elements with positive values are more abundant in the ore horizon dacitic tuffs.

| Variable                | Difference Of Means | Discriminant Function Coefficients | Percentage Contribution To Discrimination |
|-------------------------|---------------------|------------------------------------|-------------------------------------------|
| $\text{Fe}_2\text{O}_3$ | -000.863%           | -1.45400                           | 26.387                                    |
| $\text{TiO}_2$          | -000.066%           | -2.12364                           | 2.945                                     |
| $\text{SiO}_2$          | 000.674%            | -0.13161                           | 0.445                                     |
| CaO                     | 000.080%            | 0.71228                            | 1.189                                     |
| $\text{K}_2\text{O}$    | 000.441%            | 0.28694                            | 2.639                                     |
| MgO                     | -000.053%           | 0.95599                            | 1.050                                     |
| $\text{Al}_2\text{O}_3$ | -000.192%           | -0.37748                           | 1.512                                     |
| $\text{Na}_2\text{O}$   | -000.100%           | 0.58322                            | 1.212                                     |
| L.O.I.                  | 000.004%            | -0.23810                           | 0.022                                     |
| MnO                     | 000.017%            | 26.66167                           | 9.540                                     |
| Zr                      | 005.300 ppm         | 0.02850                            | 3.154                                     |
| Sr                      | 024.718 ppm         | -0.00462                           | 2.387                                     |
| Rb                      | 015.453 ppm         | 0.02783                            | 8.979                                     |
| Zn                      | 060.314 ppm         | 0.00023                            | 0.291                                     |
| Cu                      | -003.152 ppm        | 0.01215                            | 0.800                                     |
| Ba                      | 988.062 ppm         | 0.00056                            | 11.648                                    |
| Cu(s)                   | -006.523 ppm        | -0.02070                           | 2.819                                     |
| Pb(s)                   | 020.354 ppm         | -0.01330                           | 5.653                                     |
| Zn(s)                   | 033.407 ppm         | 0.00258                            | 1.802                                     |
| Pb                      | 019.005 ppm         | 0.01415                            | 5.615                                     |
| Ni                      | -002.117 ppm        | -0.10467                           | 4.626                                     |
| Co                      | -004.909 ppm        | 0.03095                            | 3.173                                     |
| Cr                      | 007.093 ppm         | 0.01159                            | 1.716                                     |
| V                       | -002.971 ppm        | 0.00639                            | 0.396                                     |



the discriminant functions, the probability of an ore horizon sample being classified as non-ore horizon is 7.5% whereas the probability of a non-ore horizon sample being classified as ore horizon is 10.9%.

The exploration applications of this are fairly obvious. Already, the stratigraphy has been re-examined and revised to account for two anomalous "non-ore horizon" samples such that the member from which these come is now known to form part of the ore horizon.

#### 7.2. Chemical Variations in the Ore Horizon Dacitic Tuffs as a Function of Distance to Ore


7.2.1. Introduction: In an effort to further understand the nature and distribution of anomalous chemical characteristics of the ore horizon dacite, studies were undertaken to determine whether consistent and significant chemical variations exist with proximity to ore. If such relationships could be established, then the distance to ore of unknown samples (e.g. Clementine area samples) could be approximated from observed data.

Plots of chemical variations versus lateral stratigraphic distance to ore were constructed for the ore horizon dacite in two steps: 1) Plots of distance to nearest orebody, whether transported or in situ, and 2) plots of distance to nearest in situ orebody (a measure of the distance to centers of ore-forming stockwork activity). Plots for all

elements determined were constructed as well as a wide variety of elemental ratios which could conceivably bear a relationship to proximity to ore. Also plotted versus distance to ore were factor scores for each sample as derived in previous sections. In total, over 100 plots of chemical variations as a function of distance to ore were constructed.

7.2.2. Chemical Variations versus Distance to Nearest Ore (whether transported or in situ): Plots of this nature were constructed largely to ascertain whether widespread post-depositional dispersion of elements has occurred from the orebodies. Data plotted consisted of 120 samples of ore horizon dacitic tuff ranging from 1 m to 2.9 km from ore. Plots constructed included single element plots as well as numerous combinations and ratios of variables (e.g.  $\text{Cu(S)}/\text{Cu}$ ,  $\text{Fe}/\text{Mn}$ ,  $\text{Mg}/\text{Ca}$ ,  $\text{Na}/\text{K}$ , etc.) as well as the factor scores as determined in a previous section.

The resultant plots show that virtually no consistent variations exist with proximity to ore (some typical examples are shown in Fig. 7-3). Interestingly, none of the plots of the scores for the six factors accounting for most of the chemical variations in the dacitic tuffs show any apparent spatial relationship to ore, e.g., chlorite after glass and sericite after glass are not related to the ore-forming process. However, in plots involving Ba, Pb and Zn, samples within 50 m of ore display erratic behavior with a



Distance to Nearest Ore (log meters)

Figure 7-3: Lateral variations of elements in ore horizon dacitic tuffs as a function of distance to nearest ore.




Figure 7-3 (con't)

large proportion of anomalously high values (Fig. 7-3). Zn displays the "best" trend of all elements with virtually all values in excess of 100 ppm occurring within 50 m of ore. In this regard, it is interesting that all samples obtained from the thick sequence of dacitic tuffs south of the Clementine prospect contain less than 50 ppm Zn. Unlike the other major ore-forming elements, Cu gives no indication of nearby ore (Fig. 7-3). It is interesting to note that Ba also shows no relationship to proximity to ore and thus the anomalous Ba content of the ore horizon is a regional characteristic of the unit.

The "nearest ore" for most samples within 50 m, and all samples with 10 m of ore, is mechanically transported ore. Since some of these deposits occur beyond the limits of significant alteration associated with the main stockworks, and since these samples are derived from both footwall and hangingwall rocks, it is logical to conclude that small amounts of Ba, Pb and Zn have migrated limited distances from mechanically transported ores. It is also logical to conclude (though it cannot be proven) that such migrations occurred during compaction - diagenesis when the host dacitic crystalline tuffs were water-saturated and more permeable than they were after lithification.

7.2.3. Chemical Variations versus Distance to in situ ore: The occurrence of mechanically transported ore at Buchans provides an excellent opportunity to study

dispersion of ore-forming elements from massive sulphide as all evidence indicates that wall rock anomalies can conceivably be caused only by post-ore dispersion directly from the orebodies. It is demonstrated in the previous section that such dispersion is erratic and limited only to areas within 50 m of ore. Nevertheless, it is possible that other dispersion processes not related to proximity to transported ores may have acted upon the ore horizon dacite. Dispersion patterns related to such processes would be superposed upon local dispersion from transported ore.

A number of alteration studies (e.g. papers in Ishihara, 1974) have demonstrated that hydrothermal activity and alteration of host rocks continues after massive sulphide deposition, affecting both footwall and hangingwall rocks alike. Such activity raises the possibility of lateral dispersion of elements in unconsolidated crystal-vitric tuffs as a function of the distance to the center of the geothermal activity. If post-ore hydrothermal activity reached the sea water - sediment interface, local sea water may be slightly enriched in ore-forming constituents which may subsequently be adsorbed on to clay minerals or deposited syngenetically. Such processes could yield widespread anomalies whose intensity would presumably be some function of distance to the hydrothermal center.

In order to test such possibilities, chemical variations of dacitic tuff samples were plotted as a function

of distance to in situ ore. In order to eliminate local dispersion effects from mechanically transported ores, all samples within 50 m of transported ore and displaying evidence of enrichment related to their proximity to such ore were eliminated from consideration. Samples in this study range from less than 100 m to more than 3.6 km from in situ ore.

The results of these plots are similar to the results of the previous section. No meaningful consistent trends are visible for any element, element ratios or for the six factors causing the bulk of the geochemical variation in the dacitic tuffs. Typical results are shown in Fig. 7-4.

Previous work by the author (Thurlow, 1973) in the immediate vicinity of the orebodies showed that there were no visible trends across stratigraphy within dacitic tuffs except within 30 m of ore. The results of this work appear to show that there are correspondingly no consistent lateral trends relating to the presence of ore.

#### 7.2.4. Discriminant Scores as a Function of Distance

to Ore: The discriminant function which effectively separates ore horizon dacitic tuffs from non-ore horizon dacitic tuffs did not yield a perfect 100% separation. A number of samples of ore horizon were effectively "misclassified" as non-ore horizon dacitic tuff (see Section 7-1). In checking which samples were misclassified, it was noticed that some of these were samples relatively remote




Figure 7-4: Lateral variations of elements in ore horizon dacitic tuffs as a function of distance to nearest in situ ore.






Figure 7-4 (con't)

from ore; i.e., several hundred meters or more. Plots were constructed (Fig. 7-5) to determine whether in fact the ore horizon dacitic tuffs actually become chemically more like non-ore horizon dacitic tuffs with increasing distance from ore. The diagrams show no obvious trend but, with some imaginative license, one could say that ore horizon dacitic tuffs become slightly more like non-ore horizon tuffs with increasing distance from ore. If one could prove this more convincingly, it would be possible to state that the anomalous chemistry of the ore horizon dacitic tuffs is indeed related to the presence of ore.

### 7.3. The Cause of the Ba Anomaly in Ore Horizon Dacitic Tuffs

The anomalous Ba content of ore horizon dacitic tuffs is the most important geochemical distinction between the ore horizon and other similar horizons throughout the Buchans Group. Some pertinent facts regarding the distribution of Ba, as outlined in previous sections are:

- 1) Ba in cogenetic rhyolite is controlled by  $K_2O$  distribution (correlation coefficient = .87)
- 2) Ba in other Buchans Group dacitic tuffs is also largely controlled by  $K_2O$  distribution (correlations coefficients for three such felsic units are .62, .46 and .58)
- 3) Ba in ore horizon dacitic tuffs shows no correlation with  $K_2O$  (correlation coefficient = .02)




Figure 7-5: Lateral variations of discriminant scores in ore horizon dacitic tuffs as a function of distance to ore.

- 4) Barium variation is apparently not related to any of the major post-depositional chemical changes that have affected the dacitic tuffs.

These criteria seem to suggest that Ba in the ore horizon tuffs has been enriched by a unique process. Though it cannot be proven, it is reasonable to suggest that this process might have some connection with the formation of the barite-rich orebodies.

In an attempt to identify the cause of the Ba anomaly a number of ore horizon and non-ore horizon dacitic tuffs were examined under the scanning electron microprobe. The following observations were recorded:

- 1) Ba is evenly and randomly distributed in non-ore horizon tuffs both in plagioclase and altered sericitic matrix glass
- 2) Ba has two modes of occurrence in samples of ore horizon tuffs: a) evenly and randomly distributed, as above and b) as tiny isolated concentrations
- 3) The isolated concentrations are composed of a Ba and sulphur-bearing mineral; probably barite
- 4) The isolated concentrations occur: a) as specks in matrix glass, b) as inclusions? in plagioclase, c) in small fractures in quartz, plagioclase and amphibole crystals (these relationships are schematically shown in Fig. 7-6).




Figure 7-6: Schematic diagram of thin section showing modes of occurrence of barite in ore horizon dacitic tuffs as determined from electron microprobe scans. Plagioclase (P), quartz (Q), and hornblende, (A) sit in matrix of glass shards altered to sericite and chlorite.

It is evident from the above that the barium anomaly in the ore horizon tuffs is likely caused by trace amounts of very fine grained disseminated barite.

The origin of this barite is enigmatic.

Any model must explain the widespread distribution of barite throughout the ore horizon with no apparent enrichment except very close to ore, take into account the extreme insolubility of barite in sea water, and account for the lack of a concomitant base metal anomaly.

Many of the ore horizon tuffs were deposited from subaqueous pyroclastic flows which locally contain minor concentrations of sulphide and barite clasts and grains (see Section 4.5.1.). Such clasts occur in the ore horizon at distances exceeding 4 km from the nearest known in situ deposit. It is very probable that, in addition to the visible clasts, microscopic barite and sulphide were also transported and homogenized by these density flows. Once emplaced, the sulphide and barite grains and clasts were distributed throughout porous tuff incontact with matrix sea water with a temperature, Eh and pH probably similar to that of ambient sea water. Compaction of the unit would result in a slow upward movement of sea water through the unit.

Sea water is a powerful oxidizing electrolyte capable of attacking and breaking down the common sulphides pyrite, sphalerite, galena and chalcopyrite. Constantinou (1976) has described several of the Cyprus orebodies as being



Plate 7-1: Drill core sample showing yellow ore clast in pyroclastic breccia of Lucky Strike ore horizon sequence. Note dark rim around margin of the clast, dark fracture cutting clast and dark spots within the clast. Enlargement approximately 2X.

in an advanced stage of leaching due to oxidation by sea water. Complete destruction of the ores was prevented by deposition of the ochre and overlying pillow lavas. It is likely that the Buchans ores were subject to the same attack, though almost immediate burial by thick pyroclastics would have halted this process at a very early stage. It is likely that the base metal anomalies recorded within 30 m of transported ore are related to this process combined with simple expulsion of matrix water and possible electro-chemical dispersion as described by Sato and Mooney (1960).

Tiny sulphide grains and larger clasts throughout the ore horizon pyroclastics would also have been subject to this relatively brief period of oxidation. Conversely, barite would remain stable. It is proposed that the barium anomaly and lack of associated base metal enrichment in the ore horizon pyroclastics is related to this process.

This proposal finds support in field evidence. Many of the larger sulphide clasts display a dark sooty margin which merges into the host pyroclastics (Plate 7-1). Smaller Pb-Zn grains in tuff commonly have a thin, anomalous, blue-grey halo. Instances of anomalous blue-grey specks with no visible sulphide are common in several areas. The material sampled in this study contained none of the above features and thus may represent the complete-removal end member of this process.



## CHAPTER EIGHT

### SUMMARY AND EXPLORATION APPLICATION

#### 8.1. Summary of Thesis

The purposes for undertaking this study, as outlined in Chapter 1, have been largely achieved though many old and new questions remain unresolved.

The relationships of the Buchans Group to adjacent rock units has been newly established in some cases, and clarified in others. The superposition of the Hungry Mountain Complex upon the Buchans Group by southeastward-directed thrusting has been established and the significance of thrusting within the Buchans Group has been documented. The recognition of thrusting has resulted in the establishment of a new stratigraphic succession based upon indisputably conformable contacts. A plausible plutonic equivalent of the Buchans Group volcanics has been identified and suggested as a source for the enigmatic granitoid boulders within the Buchans Group. The peralkaline nature of a large part of the Topsails Granite has been recognized and the intrusive relationship with the Buchans Group of all phases of the granite and related gabbro has been clarified.

Study of the Buchans ores has resulted in a genetic model comprising stockwork, in situ and transported

ores. The latter are particularly well developed and preserved at Buchans. Their nature has been documented and their origin interpreted as a result of brecciation at source, transport downslope in paleotopographic channels by gravity-driven debris flows and preferential deposition in depressions.

Whole-rock chemical analyses of unaltered Buchans Group volcanics was undertaken largely to determine whether this geochemical approach has exploration applications. It has been shown that Buchans Group volcanic rocks are similar in chemical abundances and variations to other calc-alkaline volcanic suites. There was apparently little chemical evolution in Buchans Group flows through the stratigraphic column and flows within the ore horizon sequences are essentially similar to their counterparts elsewhere in the Buchans Group. Study of flows and cogenetic pyroclastics at various stratigraphic horizons has indicated some chemical differences between these groups, probably due to greater permeability of pyroclastics during diagenesis and metamorphism. Ore horizon pyroclastics are chemically similar to those at other stratigraphic horizons and show no anomaly related to mineralization with the exception of anomalously high Ba concentrations in the ore horizon.

The major causes of geochemical variations in the ore horizon pyroclastic rocks have been quantified by factor analysis and shown to bear no relationship to mineralization. A variety of other elemental abundances and ratios within unaltered ore horizon pyroclastics similarly show no relationship to ore, whether in situ or transported. Erratic base metal anomalies are present in ore horizon pyroclastics within 50 m of ore and appear to be the result of clastic concentration combined with chemical and electrochemical dispersion.

The anomalous abundances of Ba in ore horizon pyroclastics is a regional feature which provides a means of distinguishing ore horizon pyroclastics from similar lithologies at other stratigraphic horizons. The Ba anomaly is caused by microscopic barite which was widely distributed by subaqueous pyroclastic flows which transported most ore horizon pyroclastics.

## 8.2. Exploration Applications

Assuming that no ore remains undiscovered within the depth capability of current geophysical technology, then geological observation, interpretation and application is the most powerful exploration tool at Buchans. (This statement is substantiated by the recent discovery of the MacLean Extension deposit.) An understanding of the

stratigraphy and of the myraid of new exploration possibilities posed by thrust faults is essential. A detailed knowledge of the nature and distribution of significant lithologies, alteration and paleotopography in the vicinity of the known ores is a prerequisite to effective exploration. Recognition of analogous lithologies and alteration in new exploration areas constitutes an important exploration lead.

Rock geochemistry, as studied in this thesis, has only limited exploration application. The relatively fresh rocks of the Buchans Group provide an excellent opportunity to determine whether geochemically anomalous conditions existed in rocks which are not anomalous in hand specimen. The anomalous Ba content of ore horizon pyroclastics is a regional feature related to the presence of ore and serves to distinguish pyroclastics which host mineralization from those which do not. Accurate rock analysis for Ba in the Buchans and Roberts Arm Groups, and their correlatives, should be seriously considered as a viable reconnaissance method of identifying potentially mineralized felsic volcanic horizons.

Previous work by the author (Thurlow, 1973; Thurlow, et al., 1975) showed that altered rocks of the Intermediate Footwall were depleted in  $\text{Na}_2\text{O}$ ,  $\text{CaO}$  and  $\text{K}_2\text{O}$ , and enriched in  $\text{SiO}_2$ ,  $\text{MgO}$  and many trace elements. These

results are typical of many massive sulphide terranes (see Table 1-1). The application of these parameters in exploration has become widespread, especially in areas where subsequent deformation and metamorphism have masked the appearance of altered rocks. However, at Buchans, the relatively pristine condition of the rocks makes even incipient stages of alteration evident in hand specimen.

A problem more common than identifying alteration at Buchans is the significance of the recognized alteration. The experienced eye can identify some alteration types as having no economic importance. However, some instances of interesting alteration occur at horizons known to be barren and there is at least one instance of alteration of uncertain significance in ore horizon pyroclastics. A fruitful line of future research would be an integrated mineralogical and geochemical study of clay mineral alteration related to ore as compared to alteration at the known barren horizons. This type of study may provide an important means of assessing instances of alteration of uncertain significance.

# REFERENCES

- ABBEY, S., 1968. Analysis of rocks and minerals by atomic absorption spectroscopy; Part 2; Determination of total iron, magnesium, calcium, sodium and potassium. G.S.C. Paper 68-20.
- ABBEY, S., 1975. Studies in "standard samples". Geol. Surv. Can. Pap. 74-41.
- ALCOCK, J.B., 1961. Oriental No. Two Orebody. Unpub. B.Sc. Thesis, Royal School of Mines, London, 56p.
- AMSTUTZ, G.C. (Ed.), 1974. Spilites and spilitic rocks. International Union of Geological Sciences, Series A, Number 4; Springer-Verlag, Berlin, 471p.
- ANDERSON, F.D., 1972a. Unpublished Geological Survey of Canada maps (Buchans 12A/15, Badger 12A/16).
- ANDERSON, F.D., 1972b. Buchans and Badger areas, Newfoundland. Geol. Surv. Can. Pap. 72-1, Part A, p. 2.
- ANGER, C., 1963. The lead-zinc-copper deposits of Buchans, middle Newfoundland. Neues Jahrbuch fur Minerologie, Montashefte, pp. 126-136.
- BAIRD, D.M., 1960. Massive sulphide deposits in Newfoundland. Can. Inst. Min. Metall. Bull., 53, pp. 77-80.
- BELL, K. and BLENKINSOP, J., 1981. A geochronological study of the Buchans Area, Newfoundland. In: "The Buchans Orebodies - Fifty Years of Mining and Exploration", E.A. Swanson, D.P. Strong and J.G. Thurlow (eds.). Geol. Assoc. Can. Special Paper No. 22, (in press).
- BELT, E.S., 1969. Newfoundland Carboniferous stratigraphy and its relation to the Maritimes and Ireland. In: "North Atlantic - Geology and Continental Drift", M. Kay (ed.). Am. Assoc. Petro. Geol. Mem. 12, pp. 734-753.
- BENNETT, R.A. and ROSE, W.I., 1973. Some compositional changes in Archean felsic volcanic rocks related to massive sulphide mineralization. Econ. Geol., 68, pp. 886-891.
- BISCHOFF, J.L. and DICKSON, F.W., 1975. Seawater-basalt interaction at 200°C and 500 bars: implication for origin of sea-floor heavy-metal deposits and regulation of seawater chemistry. Earth and Planet Sci. Lett., 25, pp. 385-397.

- BOSTOCK, H.H., 1978. The Roberts Arm Group, Newfoundland: Geological notes on a Middle or Upper Ordovician island arc environment. Geol. Surv. Can. Pap. 78-15, 21p.
- BOSTOCK, H.H., CURRIE, K.L., and WANLESS, R.K., 1979. The age of the Roberts Arm Group, north-central Newfoundland. Can. Jour. Earth Sci., 16, pp. 599-606.
- BOYLE, R.W. and GARRETT, R.G., 1970. Geochemical Prospecting - A review of its status and future. Earth Sci. Rev., 6, pp. 51-75.
- BRADSHAW, P.M.D., CLEWS, D.R., and WALKER, J.L., 1970. "Exploration Geochemistry - Part 4: Primary dispersion". Mining in Canada 1970.
- BUCHANS STAFF, 1955. Buchans operation, Newfoundland. Can. Inst. Min. Metall. Bull., 48, pp. 349-353.
- CADY, W.M., 1975. Tectonic setting of the Tertiary volcanic rocks of the Olympic Peninsula, Washington. Jour. Research U.S. Geol. Surv., 3, (5), pp. 573-582.
- CALHOUN, T., 1979. A study of a fragmental ore occurrence, Clementine prospect, Buchans, Newfoundland. Unpub. M.Sc. Thesis, University of Western Ontario, 95 p.
- CALHOUN, T.A. and HUTCHINSON, R.W., 1981. Determination of flow direction and source of fragmental sulphides, Clementine deposit, Buchans, Newfoundland. In: "The Buchans Orebodies - Fifty Years of Mining and Exploration", E.A. Swanson, D.F. Strong and J.G. Thurlow (eds.). Geol. Assoc. Can. Special Paper No. 22, (in press).
- CANN, J.R., 1970. "Rb, Sr, Y, Zr and Nb in some ocean floor basaltic rocks". Earth and Planet. Sci. Let., 10, pp. 7-11.
- CARLISLE, D., 1963. Pillow breccias and their aquagene tuffs, Quadra Island, British Columbia. Jour. Geol., 71, pp. 48-71.
- CATHERALL, D.J., 1960. Engine House Orebody. Unpub. B.Sc. Thesis, Royal School of Mines, London, 92p.

- CLARKE, D.B., 1975. Tertiary basalts dredged from Baffin Bay. Can. J. Earth Sci., 12, pp. 1396-1405.
- COBBING, E.J. and PITCHER, W.S., 1972. The coastal batholith of central Peru. Jour. Geol. Soc. Lond., 128, pp. 421-460.
- CONSTANTINOU, G., 1976. Genesis of the conglomerate structure, porosity and collomorphic textures of the massive sulphide ores of Cyprus. In: "Metallogeny and plate tectonics", D.F. Strong (ed.). Geol. Assoc. Can. Spec. Paper 14, pp. 187-210.
- COOMBS, D.S., 1960. Lower grade mineral facies in New Zealand. Int. Geol. Congress, 21st. Copenhagen, Pt. 13, pp. 339-351.
- DATE, J. and TANIMURA, S., 1974. Dacite and rhyolite associated with the Kuroko mineralization. In: "Geology of Kuroko Deposits", S. Ishihara (ed.). Soc. Min. Geol. Japan Spec. Issue 6, pp. 261-266.
- DAVENPORT, P.H. and I. NICHOL, 1972. Bedrock geochemistry as a guide to areas of base metal potential in volcano-sedimentary belts of the Canadian Shield. In: "Geochemical Exploration 1972, Proceedings of the fourth International Geochemical Exploration Symposium, London", M.J. Jones (ed.). pp. 45-57.
- DAVIES, I.C. and WALKER, R.G., 1974. Transport and deposition of resedimented conglomerates: the Cap Enrage Formation, Cambro-Ordovician, Gaspé, Quebec. J. Sed. Petro., 44, pp. 1200-1216.
- DEAN, P.L., 1977. A report on the geology and metallogeny of the Notre Dame Bay area. Nfld. Dept. of Mines and Energy, 77-10, 17p.
- DESCARREUX, J., 1973. A petrochemical study of the Abitibi volcanic belt and its bearing on the occurrence of massive sulphide ores. Can. Inst. Min. Metall. Bull., 66, pp. 61-69.
- DUNNING, G.R. and HERD, R.K., 1980. The Annieopsquotch ophiolite complex, southwest Newfoundland and its regional relationships. In: "Current Research, Part A". Geol. Surv. Can. Paper 80-1A, pp. 227-234.



- ELLISTON, J., 1963. Gravitational sediment movements in the Warranunga Geosyncline. In: "Syntaphral Tectonics and Diagenesis, A Symposium (S.W. Carey: Convener)", University of Tasmania, pp. C1-C18.
- ENTWISTLE, L.P. and BARNES, M.P., 1971. Report on exploration possibilities at the Buchans unit, Newfoundland. ASARCO unpub. company paper.
- FISHER, R.V., 1960. Classification of volcanic breccias. G.S.A. Bull., 71, pp. 973-982.
- FISKE, R.S., 1963. Subaqueous pyroclastic flows in the Ohanapecosh formation, Washington. G.S.A. Bull., 74, pp. 391-406.
- FISKE, R.S., 1969. Recognition and significance of pumice in massive pyroclastic rocks. Bull. Geol. Soc. Amer., 80, pp. 1-8.
- FISKE, R.S. and MATSUDA, T., 1964. Submarine equivalents of ash flows in the Tokiva Formation, Japan. Amer. Jour. Sci., 262, pp. 76-106.
- FISKE, R.S., HOPSON, C.A., and WATERS, A.C., 1963. Geology of Mt. Ranier National Park, Washington. U.S.G.S. Prof. Pap. 444, 93p.
- FRANKLIN, J.M., KASARDA, J., and POULSEN, K.H., 1975. Petrology and chemistry of the alteration zone of the Mattabi massive sulphide deposit. Econ. Geol., 70, pp. 63-79.
- GABRIELSE, H. and REESON, J.E., 1974. Nature and setting of granitic plutons in the central and eastern parts of the Canadian Cordillera. In: "Pacific Geology", 8, Tokai Univ. Press., pp. 109-138.
- GALE, G.H., 1969. The primary dispersion of Cu, Zn, Ni, Co, Mn and Na adjacent to sulphide deposits, Springale Peninsula, Newfoundland. Unpub. M. Sc. Thesis, Memorial University.
- GEORGE, P.W., 1937. Geology of lead-zinc-copper deposits at Buchans, Newfoundland. Amer. Inst. Min. Met. Eng., T.P. 816 (Class I, Mining Geology, No. 70).
- GJELSVIK, T., 1968. Distribution of major elements in wall rocks and the silicate fraction of the Skorovass pyrite deposit, Grong area, Norway. Econ. Geol., 63, pp. 217-231.

- GOODFELLOW, W.D., 1975. Major and minor element haloes in volcanic rocks at Brunswick No. 12 sulphide deposit, N.B., Canada. In: "Geochemical Exploration", 1974, Amsterdam, Elsevier, pp. 279-295.
- GOODWIN, A.M., 1968. Archean protocontinental growth and early crustal history of the Canadian Shield. Int. Geol. Congr., 23rd, 1, pp. 69-89.
- GOVETT, J.G.S., 1972. Interpretation of a rock geochemical exploration survey in Cyprus - statistical and graphical techniques. J. Geochem. Explor., 1, pp. 77-102.
- GOVETT, G.J.S. and GOODFELLOW, W.D., 1975. Use of rock geochemistry in detecting blind sulphide deposits: a discussion. Transactions Institution of Mining and Metallurgy, Section B, 85, pp. 134-140.
- GOVETT, G.J.S. and NICHOL, I., 1979. Lithogeochemistry in mineral exploration. In: "Geophysics and geochemistry in the search for metallic ores", P.J. Hood (ed.). Geol. Surv. Can., Econ. Geol. Report 31, pp. 339-362.
- GOVETT, G.J.S. and PANTAZIS, Th. M., 1971. Distribution of Cu, Zn, Ni and Co in the Troodos Pillow Lava Series, Cyprus. IMM Trans., 80, pp. B27-B46.
- GOVETT, G.J.S., WHITEHEAD, R.E.S., CROSBY, R.M., and AUSTRIA, V.B., 1974. Exploration geochemistry in New Brunswick. Can. Inst. Min. Metall., 67, pp. 76-84.
- GRAF, J.L., 1977. Rare earth elements as hydrothermal tracers during the formation of massive sulphide deposits in volcanic rocks. Econ. Geol., 72, 527-548.
- HAJASH, A., 1975. Hydrothermal processes along mid-ocean ridges: an experimental investigation. Contrib. Mineral. Petrol., 53, pp. 205-226.
- HART, R., 1970. Chemical exchange between sea water and deep ocean basalts. Earth Planet Sci. Lett., 9, pp. 269-279.

- HART, R.A., 1973. A model for chemical exchange in the basalt-seawater system of oceanic layer II. *Can. Jour. Earth Sci.*, 10, pp. 799-816.
- HART, S.R., 1969. K, Rb, Cs contents and K/Rb, K/Cs ratios of fresh and altered submarine basalts. *Earth Planet Sci. Lett.*, 6, pp. 295-303.
- HART, S.R., 1973. Submarine basalts from Kilauea rift, Hawaii: nondependence of trace element composition on extrusion depth. *Earth Planet Sci. Lett.*, 20, pp. 201-203.
- HAWKES, H.E. and WEBB, J.S., 1962. Geochemistry in mineral exploration. Harper and Row, N.Y., 415p.
- HELWIG, J. and SARPI, E., 1969. Plutonic-pebble conglomerates, New World Island, Newfoundland, and history of eugeosynclines. In: "North Atlantic - Geology and Continental Drift", M. Kay (ed.). Amer. Assoc. Petrol. Geol., Mem. 12, pp. 443-466.
- HENLEY, R.W. and THORNLEY, P., 1981. Low grade metamorphism and the geothermal environment of massive sulphide ore formation, Buchans, Newfoundland. In: "The Buchans Orebodies - Fifty Years of Mining and Exploration", E.A. Swanson, D.F. Strong and J.G. Thurlow (eds.). Geol. Assoc. Can. Special Paper No. 22, (in press).
- HIRABAYASHI, T., 1974. On the Kuroko-type orebodies in the Yokota mine area, Mishi-Aizu district, Fukushima Prefecture. In: "Geology of Kuroko Deposits", S. Ishihara (ed.). Soc. Min. Geol. Japan Spec. Issue 6, pp. 195-202.
- HONISHI, O., 1974. Geology of the Wanibuchi gypsum deposits, Shimane Prefecture. In: "Geology of Kuroko Deposits", S. Ishihara (ed.). Society of Mining Geologists of Japan, Spec. Issue 6, pp. 213-220.
- IRVINE, T.N. and BARAGAR, W.R.A., 1971. A guide to the chemical classification of the common volcanic rocks. *Can. Jour. Earth Sci.*, 8, pp. 523-548.
- ISHIHARA, S. (Ed.), 1974. Geology of the Kuroko deposits. Society of Mining Geologists of Japan, Spec. Issue No. 6, 435p.

- ISHIKAWA, Y. and YANAGISAWA, Y., 1974. Geology of the Aina mine, with special reference to syngenetic origin of the Daikoku deposits. In: "Geology of Kuroko deposits", S. Ishihara (ed.). Soc. Min. Geol. Japan, Spec. Issue 6, pp. 79-88.
- ITO, T., TAKAHASHI, T., and OMORI, Y., 1974. Submarine volcanic-sedimentary features in the Matsumine Kuroko deposits, Hanaoka mine, Japan. In: "Geology of the Kuroko deposits", S. Ishihara (ed.). Soc. Min. Geol. Japan, Spec. Issue 6, pp. 115-130.
- JAKES, P. and WHITE, A.J.R., 1971. Composition of island arcs and continental growth. Earth and Planetary Sci. Lett., 12, pp. 224-230.
- JOLLY, W.T., 1972. Degradation (hydration) - aggradation (dehydration) and low-rank metamorphism of mafic volcanic sequences. 24th Int. Geol. Cong., Section 2, pp. 11-18.
- JOLLY, W.T., 1974. Behavior of Cu, Zn and Ni during prehnite-pumpellyite rank metamorphism of the Keweenaw basalts, Northern Michigan. Econ. Geol., 69, pp. 1118-1125.
- JOLLY, W.T. and SMITH, R.E., 1972. Degradation and metamorphic differentiation of the Keweenaw Tholeiitic Lavas of Northern Michigan, USA. Jour. Pet., 13, (2), pp. 273-309.
- JONES, J.G., 1969. Pillow lavas as depth indicators. Amer. Jour. Sci., 267, pp. 181-195.
- KAJIWARA, Y., 1970. Syngenetic features of the Kuroko ore from the Shakanai. In: "Volcanism and Ore Genesis", Tatsumi, T. (ed.). Univ. of Tokyo Press, pp. 197-206.
- KALLIOKOWSKI, J., 1955. Gull Pond, Newfoundland. Geol. Surv. Can. Paper 54-4.
- KEAN, B.F., 1977. Geology of the Lake Ambrose sheet, west half. In: "Report of Activities, 1976", R.V. Gibbons (ed.). Newfoundland Dept. of Mines and Energy, Report 77-1, pp. 21-26.
- KEAN, B.F., 1979. Buchans map area (12A/15), Newfoundland. In: "Report of Activities for 1978", R.V. Gibbons (ed.). Newfoundland Dept. of Mines and Energy, Report 79-1, pp. 18-24.

- KEAN, B.F., 1980. Buchans map sheet (12A/15), Newfoundland. Nfld. Dept. of Mines and Energy, Map 79-125.
- KEAN, B.F. and JAYASINGHE, N.R., 1980. Badger map area (12A/16) Newfoundland. In: "Current research", O'Driscoll, C. and Gibbons, R. (eds.). Newfoundland Dept. of Mines and Energy, Report 80-1, pp. 37-43.
- KEAN, B.F., DEAN, P.L. and STRONG, D.R., 1981. Regional geology of the central volcanic belt in Newfoundland. In: "The Buchans Orebodies - Fifty Years of Mining and Exploration", E.A. Swanson, D.F. Strong and J.G. Thurlow (eds.). Geol. Assoc. Can. Special Paper No. 22, (in press).
- KOCH, G.S. and LINK, 1971. "Statistical Analysis of Geological Data",
- KOWALIK, J., RYE, R.O., and SAWKINS, F.J., 1981. Stable isotope study of the Buchans polymetallic sulphide deposits. In: "The Buchans Orebodies - Fifty Years of Mining and Exploration", E.A. Swanson, D.F. Strong and J.G. Thurlow (eds.). Geol. Assoc. Can. Special Paper No. 22, (in press).
- KUNO, H., 1966. Lateral variation of basalt magma type across continental margins and island arcs. Bull. Volcan., 29, pp. 195-222.
- LAMBERT, I.B. and SATO, T., 1974. The Kuroko and associated ore deposits of Japan: a review of their features and metallogenesis. Econ. Geol., 69, pp. 1215-1236.
- LANGMHYR, F.J. and PAUS, P.E., 1968. The analysis of inorganic siliceous materials by atomic absorption spectrophotometry and the hydrofluoric acid decomposition technique, Part I: The analysis of silicate rocks. Anal. Chim. Acta., 43, pp. 397-408.
- LARGE, R.R., 1977. Chemical evolution and zonation of massive sulphide deposits in volcanic terrains. Econ. Geol., 72, pp. 549-572.
- LARSEN, K.B., 1973. The formation and penecontemporaneous deformation of effaceous sediments in Buchans, Newfoundland. Unpub. ASARCO report.
- LARSON, L. and WEBBER, G.R., 1977. Chemical and petrographic variations in rhyolitic zones in the Noranda area, Quebec. Can. Inst. Min. Metall. Bull., 70, pp. 80-93.

- LAVIN, O.P., 1976. Lithogeochemical discrimination between mineralized and unmineralized cycles of volcanism in the Sturgeon Lake and Ben Nevis areas of the Canadian Shield. Unpub. M.Sc. thesis, Queens University, 249p.
- LEE, M.S., MIYASIMA, T., and MIZUMOTO, H., 1974. Geology of the Kamaikita mine, Aomori Prefecture, with special reference to genesis of fragmental ores. In: "Geology of Kuroko Deposits", S. Ishihara (ed.). Soc. Min. Geol. Japan, Spec. Issue 6, pp. 53-66.
- LEVINSON, A.A., 1974. Introduction to exploration geochemistry, Second Edition, Applied Publishing, Illinois, 924p.
- LUNDBERG, H., 1957. The discovery of large lead-zinc deposits at Buchans, Newfoundland. Mining Geophysics, 6th Commonwealth Min. Met. Congress., 1957, pp. 141-154.
- MACKENZIE, D.E. and CHAPPELL, B.W., 1972. Shoshonite and calc-alkaline lavas from the highlands of Papua, New Guinea. Contrib. Min. Petrol., 35, pp. 50-62.
- MACLEAN, H.J., 1941. Memorandum on prospecting in the Little Sandy Brook area. Unpub. ASARCO report.
- MACLEAN, H.J., 1943. Memorandum on prospecting in the Woodman's Brook - No. 2 Siding area. Unpub. ASARCO report.
- MARCOTTE, D. and DAVID, M., 1978. Targets definition of Kuroko type deposits in Abitibi by statistical analysis of geochemical data. Paper presented to Can. Inst. Min. Metall.
- MCCONNEL, J.W., 1976. Geochemical dispersion in wallrocks of Archean massive sulphide deposits. Unpub. M.Sc. thesis, Queens University, 230p.
- MOTEGI, M., 1974. Kuroko deposits of the Kurosawa mine, Fukushima Prefecture. In: "Geology of Kuroko deposits", S. Ishihara (ed.). Soc. Min. Geol. Japan, Spec. Issue 6, pp. 203-208.
- MURRAY, A., 1871. Annual report. Newfoundland Geological Survey.

- MUTTI, E., 1965. Submarine flood tuffs associated with turbidites in Oligocene deposits of Rhodes Island, Greece. *Sedimentology*, 5, pp. 265-288.
- NAIRIS, B., 1971. Endogene dispersion aureoles around the Rudtjebacken sulphide ore in the Adak area, Northern Sweden. In: "Geochemical Exploration", Can. Inst. Min. Metall., Spec. Vol. 11, pp. 357-374.
- NEALE, E.R.W. and NASH, W.A., 1963. Sandy Lake (East Half), Newfoundland. *Geol. Surv. Can. Paper* 62-28.
- NEARY, G.N., 1981. Mining History of the Buchans Area. In: "The Buchans Orebodies - Fifty Years of Mining and Exploration", E.A. Swanson, D.F. Strong and J.G. Thurlow (eds.). *Geol. Assoc. Can. Special Paper No. 22*, (in press).
- NEWHOUSE, W.H., 1931. Geology and ore deposits of Buchans, Newfoundland. *Econ. Geol.*, 26, pp. 399-414.
- NILSSON, C.A., 1968. Wall rock alteration at Boliden deposit, Sweden. *Econ. Geol.*, 63, pp. 472-494.
- OHTAGAKI, T., ONO, T., and KIMURA, A., 1974. Geology of the Tashiro mine, Fukushima Prefecture. In: "Geology of Kuroko deposits", S. Ishihara (ed.). *Soc. Min. Geol. Japan, Spec. Issue* 6, pp. 189-194.
- OSADA, T., ABE, M., and DAIMARU, K., 1974. Geology of the Yoshino mine, Yamagata Prefecture. In: "Geology of Kuroko deposits", S. Ishihara (ed.). *Soc. Min. Geol. Japan, Spec. Issue* 6, pp. 183-188.
- OSHIMA, T., HASHIMOTO, T., KAMOND, K., KAWABE, S., SUCA, K., TANIMURA, S., and ISHIKAWA, Y., 1974. Geology of the Kosada mine, Akita Prefecture. In: "Geology of Kuroko deposits", S. Ishihara (ed.). *Soc. Min. Geol. Japan, Spec. Issue* 6, pp. 89-100.
- PANTAZIS, T.L.M. and GOVETT, G.J.S., 1973. Interpretation of a detailed rock geochemical survey around Mathiati mine, Cyprus. *Jour. Geochem. Exploration*, 2, pp. 25-36.
- PEARCE, J.A. and CANN, J.R., 1973. Tectonic setting of basic volcanic rocks determined using trace element analysis. *Earth. Planet. Sci. Lett.*, 19, pp. 290-300.
- PEARCE, T.H., 1974. Quench plagioclase from some Archean basalts. *Can. J. Earth Sci.*, 11, pp. 715-719.

- PHILPOTTS, J.A., SCHNETZLER, C.C., and HART, S.R., 1969. Submarine basalts: some K, Rb, Sr, Ba, rare earth, H<sub>2</sub>O and CO<sub>2</sub> data bearing on their alteration, modification by plagioclase and possible source materials. *Earth. Planet. Sci. Lett.*, 7, pp. 293-299.
- RELLEY, B.H., 1960. The geology of the Buchans mine. Unpub. Ph.D. Thesis, McGill University, 281p.
- RIDLER, R.H., 1973. Exhalite concept, a new tool for exploration. *Northern Miner*, Nov. 29, pp. 59-61.
- RILEY, G.C., 1957. Red Indian Lake (west half) map area, Newfoundland. *Geol. Surv. Can. Map* 8-1957.
- SAKRISON, H.C., 1967. Chemical studies of the host rocks of the Lake Dufault mine, Quebec. Unpub. Ph.D. thesis, McGill University.
- SAKRISON, H.C., 1971. Rock geochemistry - its current usefulness on the Canadian Shield. *CIM Bull.*, Nov., pp. 28-31.
- SANGSTER, D.F., 1972. Precambrian volcanogenic massive sulphide deposits in Canada: A review. *Geol. Surv. Can. Paper* 72-22, 44p.
- SANGSTER, D.F. and SCOTT, S.D., 1976. Precambrian, stratbound, massive Cu-Zn-Pb sulfide ores of North America. In: "Handbook of stratabound and stratiform ore deposits", K. Wolf (ed.), 6, pp. 129-222.
- SATO, M. and MOONEY, H.M., 1960. The electrochemical mechanism of sulphide self potential. *Geophysics*, 25, pp. 226-249.
- SATO, N. and KUSAKA, H., 1974. Geology of the Matsuki mine, Ohdate city, Akita prefecture. In: "Geology of Kuroko deposits", S. Ishihara (ed.). *Soc. Min. Geol. Japan, Spec. Issue* 6, pp. 141-146.
- SATO, T., 1968. Ore showing graded bedding from Uchinotai western ore body, Kosaka mine, Akita prefecture. *Bull. Geol. Surv. Japan*, 19, pp. 7-20.
- SATO, T., 1972. Behaviors of ore-forming solutions in sea water. *Mining Geology*, 22, pp. 31-42.
- SATO, T., 1974. Distribution and geological setting of the Kuroko deposits. In: "Geology of Kuroko deposits", S. Ishihara (ed.). *Soc. Min. Geol. Japan, Spec. Issue* 6, pp. 1-10.



- SCHERMERHORN, L.J.G., 1970. The deposition of volcanics and pyritite in the Iberian pyrite belt. Mineral. Deposita, 5, pp. 273-279.
- SCHERMERHORN, L.J.G., 1971. Pyrite emplacement by gravity flow. Bull. Geol. Min., 72, pp. 304-308.
- SHAPIRO, L. and BRANNOCK, W.M., 1962. Rapid analysis of silicate, carbonate and phosphate rocks. U.S.G.S. Bulletin, 1144A, A31.
- SIMMONS, B.D. and the GEOLOGICAL STAFF, Lake Dufault Divin, Falconbridge Copper Ltd., 1973. Geology of the Millenbach massive sulphide deposit, Noranda, Quebec. Can. Inst. Min. Metall. Bull., 66, pp. 67-78.
- SINCLAIR, W.D., 1971. A volcanic origin for the No. 5 zone of the Horne mine, Noranda, Quebec. Econ. Geol., 66, pp. 1225-1231.
- SMITH, R.E., 1968. Redistribution of major elements in the alteration of some basic lavas during burial metamorphism. Jour. Petrol., 9, pp. 191-219.
- SNELGROVE, A.K., 1928. The geology of the central mineral belt of Newfoundland. Can. Min. and Met. Bull., 21, pp. 1057-1127.
- SOPUCK, V.J., 1977. A lithogeochemical approach in the search for areas of felsic volcanic rocks associated with mineralization in the Canadian Shield. Unpub. Ph.D. thesis, Queens Univ.
- SPENCE, C.D., 1975. Volcanogenic features of the Vauze sulfide deposit, Noranda, Quebec. Econ. Geol., 70, pp. 102-114.
- SPITZ, G. and DARLING, R., 1978. Major and minor element lithogeochemical anomalies surrounding the Louvem copper deposit, Val d'Or, Quebec. Can. Jour. Earth Sci., 15, pp. 1161-1169.
- STAUFFER, P.H., 1967. Grain-flow deposits and their implications, Santa Ynez Mountains, California. Jour. Sed. Petrology, 37, pp. 487-508.
- STRECKEISEN, A., 1976. To each plutonic rock its proper name. Earth Sci. Rev., 12, pp. 1-33.
- STRONG, D.F., 1973. Lushs Bight and Roberts Arm Groups of central Newfoundland: possible juxtaposed oceanic and island-arc volcanic suites. Bull. Geol. Soc. Amer., 84, pp. 3917-3928.

- STRONG, D.F., 1977. Volcanic regimes of the Newfoundland Appalachians. Geol. Assoc. Can. Spec. Pap. 16, pp. 61-90.
- STRONG, D.F., 1981. Mineralographic notes on ore samples of the Buchans district. In: "The Buchans Orebodies - Fifty Years of Mining and Exploration", E.A. Swanson, D.F. Strong, and J.G. Thurlow (eds.). Geol. Assoc. Can. Special Paper No. 22, (in press).
- SUFFEL, G.G., 1965. Remarks on some sulphide deposits in volcanic extrusives. Can. Inst. Min. Metall. Bull., 58, pp. 1057-1063.
- SUGIMURA, A., 1968. Spatial relations of basaltic magmas. In: "Basalts", H.H. Hess and A. Poldervaart (eds.). New York, Interscience, pp. 573-577.
- SWANSON, E.A. and BROWN, R.L., 1962. Geology of the Buchans orebodies. Can. Inst. Min. Metall. Bull., 55, pp. 618-626.
- SWANSON, E.A., PERKINS, E.W., and HIGGINS, W.H., 1956. Diamond drill electrical surveys, Oriental #2 orebody, Buchans mine. Paper presented at A.I.M.E. meeting, Feb. 1956.
- TAKAHASHI, T. and SUGA, K., 1974. Geology and ore deposits of the Hanaoka Kuroko belt, Akita Prefecture. In: "Geology of Kuroko deposits", S. Ishihara (ed.). Soc. Min. Geol. Japan, Spec. Issue 6, pp. 101-114.
- TANIMURA, S., SHIMODA, T., and SAWAGUCHI, T., 1974. On the Fukazawa orebodies, Akita Prefecture. In: "Geology of the Kuroko deposits", S. Ishihara (ed.). Soc. Min. Geol. Japan, Spec. Issue 6, pp. 147-156.
- TATSUMI, T. and CLARKE, L.A., 1972. Chemical composition of acid volcanic rocks genetically related to formation of the Kuroko deposit. Jour. Geol. Soc. Japan, 78, pp. 191-201.
- TAYLOR, R.P., STRONG, D.F., and KEAN, B.F., 1980. The Topsails igneous complex: Silurian-Devonian peralkaline magmatism in western Newfoundland. Can. Jour. Earth Sci., 17, pp. 425-439.
- THURLOW, J.G., 1973. Lithogeochemical studies in the vicinity of the Buchans massive sulphide deposits, central Newfoundland. Unpub. M.Sc. thesis, Memorial Univ., 172p.

- THURLOW, J.G., 1975. Buchans area report. In: Geology, mineral deposits and mineral potential of the Buchans volcanic belt", by B.P. Kean and J.G. Thurlow. Report submitted to the Buchans Task Force, 56p.
- THURLOW, J.G., 1977. The occurrence, origin and significance of mechanically transported ores at Buchans, Newfoundland. In: "Volcanic processes in ore genesis". Special Pub., 7, Geol. Soc. London., p. 127.
- THURLOW, J.G., SWANSON, E.A., and STRONG, D.F., 1975. Geology and lithogeochemistry of the Buchans polymetallic sulphide deposits, Newfoundland. Econ. Geol., 70, pp. 130-144.
- TONO, N., 1974. Minor elements distribution around Kuroko deposits in northern Akita, Japan. In: "Geology of Kuroko deposits", S. Ishihara (ed.). Soc. Min. Geol. Japan, Spec. Issue 6, pp. 399-420.
- TUTTLE, O.F. and BOWEN, N.L., 1958. Origin of granite in the light of experimental studies in the system  $\text{NaAlSi}_3\text{O}_8$  -  $\text{KAlSi}_3\text{O}_8$  -  $\text{SiO}_2$  -  $\text{H}_2\text{O}$ . Geol. Soc. Amer. Memoir, 74, 38p.
- URABE, T. and SATO, T., 1978. Kuroko deposits of the Kosaka Mine, northeast Honshu, Japan - products of submarine hot springs on Miocene sea floor. Econ. Geol., 73, pp. 161-179.
- WAHL, J.L., GOVETT, G.J.S., and GOODFELLOW, W.D., 1975. Anomalous element distribution in volcanic rocks around Key Anacon, Heath Steele B-zone and Brunswick No. 12 sulphide deposits (abstract). Can. Inst. Min. Metall. Bull., 68, p. 49.
- WALKER, P.N. and KING, J., 1975. Preliminary report, Buchans Project. Unpub. ASARCO report, 4p.
- WALKER, R.R., MATULICH, A., AMOS, A.C., WATKINS, J.J., and MANNARD, G.W., 1975. The geology of the Kidd Creek mine. Econ. Geol., 70, pp. 80-89.
- WHITEHEAD, R.E., 1973. Environment of stratiform sulphide deposition; variation in Mn:Fe ratio in host rocks at Heath Steele mine, New Brunswick, Canada. Mineralium Deposita, 8, pp. 148-160.

- WHITEHEAD, R.E.S. and GOVETT, G.J.S., 1974. Exploration rock geochemistry - detection of trace element haloes at Heath Steele mines (N.B., Canada) by discriminant analysis. *J. Geochem. Explor.*, 3, pp. 371-386.
- WILLIAMS, H., 1967. Silurian rocks of Newfoundland. G.A.C. Special Paper 4, pp. 93-137.
- WILLIAMS, H., 1970. Red Indian Lake (east half), Newfoundland. Geological Survey of Canada, Map 1196A.
- WILLIAMS, H., 1979. Appalachian orogen in Canada. *Can. Jour. Earth Sci.*, 16, pp. 792-807.
- WOAKES, M., 1954. A microscopic study of the ore sulphides from Buchans mine, Buchans, Newfoundland. Unpub. B.Sc. thesis, Royal School of Mines, London, 59p.
- YAMADA, E., 1973. Subaqueous pumice flow deposits in the Orikobe caldera, Miyaji Prefecture, Japan. *J. Geol. Soc. Japan*, 79, pp. 585-597.

# APPENDIX I

## Classification of Plutonic Rocks (after Steckelsan, 1976)




Fig. 1a. General classification and nomenclature of plutonic rocks according to mineral content (in vol. %).

$Q + A + P = 100$ , or  $A + P + P = 100$ .

1a, quartzolite (silexite); 1b, quartz-rich granitoids; 2, alkali-feldspar granite; 3, granite; 4, granodiorite; 5, tonalite; 6°, quartz alkali-feldspar syenite; 7°, quartz syenite; 8°, quartz monzonite; 9°, quartz monzodiorite/quartz monzogabbro; 10°, quartz diorite/quartz gabbro/quartz anorthosite; 6, alkali-feldspar syenite; 7, syenite; 8, monzonite; 9, monzodiorite/monzogabbro; 10, diorite/gabbro/anorthosite; 6', foid-bearing alkali-feldspar syenite; 7', foid-bearing syenite; 8', foid-bearing monzonite; 9', foid-bearing monzodiorite/monzogabbro; 10', foid-bearing diorite/gabbro; 11, foid syenite; 12, foid monzosyenite (syn. foid plagiisyenite); 13, foid monzodiorite/foid monzogabbro (essexite = nepheline monzodiorite/monzogabbro); 14, foid diorite/foid gabbro (theralite = nepheline gabbro, teschenite = analcime gabbro); 15, foidolites; 16, ultramafic plutonic rocks (ultramafitolites).

## APPENDIX II

### FIELD AND LABORATORY METHODS

- (i) Sample Collection
- (ii) Sample Preparation
- (iii) Analytical Procedures - Major elements
- (iv) Analytical Procedures - Trace elements

#### (1) Sample Collection

A small (less than 1 kg) reference sample was collected from almost all outcrops visited. The decision to take larger samples for geochemical analysis was based on a number of factors, i.e., the stratigraphic and aerial coverage desired, suitability of the lithology for geochemical analysis and accessibility of the outcrop. Samples taken for geochemical analysis ranged from one to five kilograms (varying with sample grain size) and were collected with a 3 kg sledge hammer. Visibly altered samples were not collected unless the alteration was characteristic of the area or the specific unit sampled. Similarly, samples were not collected from outcrops displaying lithologic heterogeneity, deep weathering, secondary veining, abundant amygdules, etc. Where pillow lava was sampled, care was taken to ensure that no selvedge or interpillow material was included. Samples were trimmed of weathering as best as possible in the field and remaining weathering was removed with

the diamond saw at Buchans. All field samples, regardless of weathering, were slabbed with the diamond saw in order to macroscopically assess their suitability for analysis.

The philosophy governing sample collection from core was similar to that in the field except that weathering was not a factor. The visual assessment of the suitability of core samples for geochemical analysis was based on a number of factors. Sections of core containing veins, alteration or sulphides were not sampled. Some sections of core contain a visible sheen of steel (or rust) derived from abrasion with the core barrel during drilling. Samples were not taken from such sections. Core samples taken were deemed to be representative of the section of core under study. Sampling of felsic pyroclastic rocks was especially critical as these rocks were known to best distinguish between mineralized and barren sequences (Thurlo, 1973). Pyroclastic rocks sampled were fine grained (1-5 mm) and homogeneous. Crystal-vitric tuffs (the most widespread pyroclastic type) were sampled in preference to other types such that texturally and mineralogically similar lithologies were being geochemically compared. Pyroclastic rocks containing numerous lithic fragments were not sampled.

In general, diamond drill core allows a better visual estimation of the suitability for analysis than does outcrop and also provides more exact stratigraphic positioning.

For these reasons, core samples were analysed in preference to surface samples where both were available from the same stratigraphic unit in the same area.

#### (11) Sample Preparation

Samples selected for analysis were washed, logged, dried and put into clean bags before crushing. One half to two thirds of each sample was subjected first to a coarse crush, then a fine crush on a Denver steel jaw crusher. The crushed samples were rolled, quartered and placed in glass jars to await pulverizing.

Pulverizing was effected by two different methods. Those samples prepared during the previous study (Thurlow, 1973) were pulverized in alumina ball mills by a method described by Thurlow (op. cit.).

Additional samples prepared during the present study were pulverized in a Tema swing mill with a tungsten carbide liner. Approximately 50 grams of sample was pulverized for two minutes which yielded a powder of minus 200 mesh.

Contamination during sample reduction is inevitable but all possible efforts were taken to ensure that it was minimized. Samples were ordered according to lithology before crushing and pulverizing to ensure that the effects of inter-sample contamination were minimized. Samples



containing sulphides were crushed and pulverized after the sulphide-free samples. After each sample was crushed, the removable jaw of the crusher was taken off and the entire apparatus was brushed and blown clean with compressed air. The internal parts of the swing mill were scrubbed with water after each sample was pulverized. Pulverized samples were stored in new glass jars. Care at all intervening stages of the crushing and grinding procedure ensured that inter-sample contamination was negligible.

Contamination from the crushing and pulverizing apparatus is inevitable. The removable steel jaw of the jaw crusher at Buchans was weighed prior to, and after crushing. The jaw weighed 12 grams less after 625 samples were crushed, an average contamination of approximately 20 ppm steel per sample from that jaw. Alumina contamination from the alumina ball mills used for pulverizing was determined by a similar method and found to be approximately 0.1%  $\text{Al}_2\text{O}_3$  per sample. As tungsten was not determined in the present study tungsten carbide contamination from the Tema swing mill was not estimated.

#### (iii) Analytical Procedures - Major Elements

All major elements were determined by the author using a Perkin Elmer 303 atomic absorption spectrophotometer equipped with a recorder readout. The digestion technique used was similar to that described by Langhyr and Paus (1968).

Exactly 0.1000 g of sample was weighed and placed in a poly carbonate digestion bottle. Digestion was effected by addition of 5 ml of concentrated HF and heating on a water bath for 20 minutes. Samples were subsequently cooled and 50 ml boric acid added to complex undissolved fluorides. Following a second heating, samples were diluted with 145 ml distilled and de-ionized water. Standards were prepared in a manner similar to that described by Abbey (1968).

Calcium and magnesium were determined using a air-acetelyne flame by adding a lanthanum oxide - HCl solution to supress interferences by aluminum. Phosphorus was determined by a colormetric method similar to that described by Shapiro and Brannock (1962). Loss on ignition was determined by heating a known weight of sample at  $1050^{\circ}\text{C}$  for 90 minutes and measuring the weight loss due to escape of volatiles (principally  $\text{H}_2\text{O}$ ,  $\text{CO}_2$  and  $\text{SO}_2$ ).

Precision of major element analyses was determined by choosing a sample of intermediate composition and preparing solutions of this sample at intervals throughout the analytical period. The results of thirteen such runs are as follows:

PRECISION OF ATOMIC ABSORPTION MAJOR ELEMENT DETERMINATIONS

| <u>Element</u>                 | <u>Number Of<br/>Determinations</u> | <u>Mean<br/>(Weight %)</u> | <u>Standard<br/>Deviation</u> | <u>Coefficient of<br/>Variation (%)</u> |
|--------------------------------|-------------------------------------|----------------------------|-------------------------------|-----------------------------------------|
| SiO <sub>2</sub>               | 13                                  | 65.4                       | .82                           | 1.25                                    |
| TiO <sub>2</sub>               | 13                                  | .40                        | .01                           | 2.50                                    |
| Al <sub>2</sub> O <sub>3</sub> | 13                                  | 14.29                      | .21                           | 1.47                                    |
| Fe <sub>2</sub> O <sub>3</sub> | 13                                  | 4.59                       | .10                           | 2.18                                    |
| MnO                            | 13                                  | .11                        | .01                           | 9.09                                    |
| MgO                            | 13                                  | 1.71                       | .03                           | 1.76                                    |
| CaO                            | 13                                  | 3.47                       | .05                           | 1.44                                    |
| Na <sub>2</sub> O              | 13                                  | 1.88                       | .03                           | 1.60                                    |
| K <sub>2</sub> O               | 13                                  | 3.40                       | .05                           | 1.47                                    |
| P <sub>2</sub> O <sub>5</sub>  | 13                                  | .05                        | .03                           | 60.0                                    |
| Loss Ig.                       | 13                                  | 3.89                       | .14                           | 3.60                                    |

Accuracy was determined by comparison to the U.S.

G.S. standard BCR-1. The results of five analyses are as follows:

ACCURACY OF ATOMIC ABSORPTION MAJOR ELEMENT DETERMINATIONS

| <u>Wt. %</u>                   | <u>A*</u> | <u>Mean<br/>(Weight %)</u> | <u>Standard<br/>Deviation</u> |
|--------------------------------|-----------|----------------------------|-------------------------------|
| SiO <sub>2</sub>               | 54.85     | 55.38                      | 0.37                          |
| TiO <sub>2</sub>               | 2.22      | 2.35                       | 0.18                          |
| Al <sub>2</sub> O <sub>3</sub> | 13.68     | 13.50                      | 0.27                          |
| Fe <sub>2</sub> O <sub>3</sub> | 13.52     | 13.00                      | 0.28                          |
| CaO                            | 6.98      | 6.63                       | 0.07                          |
| MgO                            | 3.49      | 3.57                       | 0.06                          |
| Na <sub>2</sub> O              | 3.29      | 3.23                       | 0.05                          |
| K <sub>2</sub> O               | 1.68      | 1.73                       | 0.05                          |
| MnO                            | 0.19      | 0.18                       | 0.01                          |

\*Proposed values after Abbey (1975).

(iv) Analytical Procedures - Trace Elements

Rock chips sent to the ASARCO laboratory at Salt Lake City for trace element analysis were ground at  $-50^{\circ}\text{C}$  to prevent loss of Hg vapour. Hg was subsequently determined on a modified Perkin Elmer 303 using a flameless method. Zn, Mo, Cu, Pb, Ni, Co, V and Ag were determined by conventional atomic absorption methods using a background correction for non-specific absorption (L.D. James, personal communication).

Precision of trace element analyses were determined by replicate analysis of every tenth sample. The results of the precision test are as follows:

PRECISION OF ATOMIC ABSORPTION TRACE ELEMENT DETERMINATIONS

| <u>Element</u> | <u>Number of Determinations</u> | <u>Range of Analysed Samples (ppm)</u> | <u>Average Coefficient of Variance (5)</u> |
|----------------|---------------------------------|----------------------------------------|--------------------------------------------|
| Cu             | 25                              | 2 - 9700                               | 2.39                                       |
| Pb             | 23                              | 1 - 23000                              | 5.30                                       |
| Zn             | 26                              | 10 - 115000                            | 4.74                                       |
| Mo             | 9                               | 3 - 124                                | 14.57                                      |
| Ag             | 12                              | 0 - 140                                | 4.37                                       |
| Hg             | 24                              | 4 - 3466 (ppb)                         | 11.21                                      |
| Co             | 24                              | 2 - 48                                 | 3.35                                       |
| Ni             | 26                              | 0 - 80                                 | 1.74                                       |
| V              | 23                              | 0 - 520                                | 6.31                                       |

Rb, Ba, Sr, Zr and Cr were determined at Memorial University by X-Ray fluorescence. Ten g of sample powder was mixed with 1.3 g of phenolformaldehyde binding agent and pressed into a 40 mm pellet at 25 tons pressure.

Precision and accuracy data are as follows:

PRECISION AND ACCURACY OF XRF TRACE ELEMENT ANALYSES (13 DETERMINATIONS, SINGLE PELLET, W-1)

| <u>Element</u> | <u>Accepted*<br/>Value</u> | <u>Mean</u> | <u>Standard<br/>Deviation</u> | <u>Difference</u> |
|----------------|----------------------------|-------------|-------------------------------|-------------------|
| Zr             | 105                        | 98          | 2                             | -6.7%             |
| Sr             | 190                        | 189         | 6                             | -0.5              |
| Rb             | 21                         | 22          | 2                             | +4.8              |
| Ba             | 160                        | 171         | 12                            | +6.9              |
| Cr             | 120                        | 92          | 6                             | -23.3             |

\*Abbey (1975)

## FOOTWALL BASALT FLOWS

## MAJOR ELEMENTS (WT%)

| SAMPLE                         | JGT75139 | JGT75219 | 9781479 |
|--------------------------------|----------|----------|---------|
| SiO <sub>2</sub>               | 47.98    | 48.58    | 50.20   |
| TiO <sub>2</sub>               | .84      | .84      | .82     |
| Al <sub>2</sub> O <sub>3</sub> | 17.18    | 16.79    | 15.77   |
| Fe <sub>2</sub> O <sub>3</sub> | 8.78     | 9.32     | 7.65    |
| MnO                            | .15      | .13      | .10     |
| MgO                            | 5.12     | 7.48     | 5.36    |
| CaO                            | 10.98    | 8.84     | 10.70   |
| Na <sub>2</sub> O              | 2.39     | 3.29     | 2.33    |
| K <sub>2</sub> O               | 1.22     | 1.59     | 1.11    |
| LOI                            | 5.98     | 3.18     | 4.58    |
| TOTAL                          | 100.30   | 99.08    | 98.54   |

## TRACE ELEMENTS (PPM)

|                   |     |     |     |
|-------------------|-----|-----|-----|
| Zr                | 92  | 77  | 79  |
| Sr                | 343 | 223 | 334 |
| Rb                | 31  | 33  | 21  |
| Zn                | 52  | 35  | 65  |
| Cu                | 63  | 15  | 74  |
| Ba                | 435 | 461 | 678 |
| CuS               | 54  | 16  | 72  |
| PbS               | 6   | 7   | 5   |
| ZnS               | 12  | 7   | 19  |
| BaSO <sub>4</sub> | 0   | 0   | 0   |
| Pb                | 35  | 15  | 21  |
| Ni                | 35  | 48  | 64  |
| Co                | 27  | 17  | 26  |
| Cr                | 144 | 231 | 478 |
| V                 | 195 | 95  | 170 |
| Hg                |     |     |     |
| Ag                |     |     |     |
| Nb                | 16  | 6   | 2   |

## FOOTWALL BASALT - ALTERED MAFIC FLOWS

## MAJOR ELEMENTS (WT%)

| SAMPLE | J0T75134 | SK-5 350 | 542 550 | 542 625 | 14062440 |
|--------|----------|----------|---------|---------|----------|
| S102   | 46.70    | 46.10    | 44.80   | 54.10   | 43.30    |
| T102   | 1.18     | 1.18     | .90     | .63     | 1.00     |
| Al2O3  | 15.83    | 15.26    | 16.28   | 11.70   | 15.89    |
| Fe2O3  | 12.19    | 9.89     | 9.84    | 6.06    | 11.44    |
| MnO    | .20      | .08      | .20     | .14     | .21      |
| MgO    | 5.42     | 14.32    | 8.78    | 4.18    | 6.85     |
| CaO    | 5.16     | 1.86     | 9.84    | 10.03   | 9.85     |
| Na2O   | 3.51     | 2.65     | 2.10    | 3.91    | 3.00     |
| K2O    | 1.07     | .10      | .97     | .11     | .83      |
| LOI    | 7.70     | 7.10     | 5.10    | 7.30    | 6.10     |
| TOTAL  | 98.96    | 98.54    | 98.81   | 98.16   | 98.55    |

## TRACE ELEMENTS (PPM)

|       |     |     |     |     |     |
|-------|-----|-----|-----|-----|-----|
| Zr    | 60  | 50  | 96  | 46  | 66  |
| Sr    | 92  | 41  | 352 | 144 | 239 |
| Rb    | 25  | 6   | 20  | 9   | 10  |
| Zn    | 96  | 47  | 105 | 94  | 72  |
| Cu    | 28  | 72  | 53  | 44  | 64  |
| Ba    | 270 | 71  | 614 | 162 | 310 |
| CuS   | 28  | 85  | 42  | 34  | 51  |
| PbS   | 9   | 5   | 20  | 11  | 5   |
| ZnS   | 20  | 2   | 12  | 0   | 17  |
| BaSO4 | 0   | 0   | 0   |     | 0   |
| Pb    | 27  | 33  | 33  | 28  | 24  |
| Ni    | 4   | 00  | 00  | 35  | 50  |
| Co    | 25  | 45  | 37  | 27  | 28  |
| Cr    | 27  | 622 | 208 | 103 | 262 |
| V     | 235 | 230 | 275 | 260 | 165 |
| Hg    |     |     | 12  | 12  |     |
| Ag    |     |     | 16  | 13  |     |
| Nb    | 0   | 6   |     |     | 6   |

## FOOTWALL BASALT - MAFIC FRAGMENTALS

## MAJOR ELEMENTS (WT%)

| SAMPLE | 542 700 | 5421000 | 05421150 | 5421225 | 05421350 |
|--------|---------|---------|----------|---------|----------|
| SiO2   | 37.90   | 48.00   | 48.10    | 46.50   | 44.70    |
| TiO2   | .81     | .86     | .85      | .80     | .93      |
| Al2O3  | 14.15   | 16.48   | 16.00    | 15.22   | 17.97    |
| Fe2O3  | 10.32   | 11.64   | 11.89    | 10.15   | 12.27    |
| MnO    | .21     | .12     | .13      | .15     | .12      |
| MgO    | 7.64    | 8.22    | 7.99     | 8.72    | 8.48     |
| CaO    | 13.34   | 4.53    | 5.00     | 7.25    | 4.45     |
| Na2O   | 1.98    | 1.98    | 2.29     | 1.94    | 1.63     |
| K2O    | .40     | 1.05    | .63      | .24     | 1.18     |
| LOI    | 11.70   | 6.40    | 6.30     | 8.30    | 7.50     |
| TOTAL  | 98.45   | 99.28   | 99.18    | 99.27   | 99.23    |

## TRACE ELEMENTS (PPM)

|       |     |     |     |     |     |
|-------|-----|-----|-----|-----|-----|
| Zr    | 82  | 87  | 75  | 75  | 83  |
| Sr    | 244 | 283 | 223 | 254 | 224 |
| Rb    | 18  | 26  | 17  | 16  | 33  |
| Zn    | 91  | 88  | 80  | 73  | 91  |
| Cu    | 50  | 66  | 59  | 58  | 55  |
| Ba    | 257 | 373 | 340 | 276 | 429 |
| CuS   | 32  | 23  | 30  | 32  | 31  |
| PbS   | 12  | 2   | 7   | 3   | 2   |
| ZnS   | 11  | 8   | 14  | 13  | 10  |
| BaSO4 | 0   |     |     |     |     |
| Pb    | 30  | 34  | 36  | 42  | 34  |
| Ni    | 42  | 59  | 54  | 40  | 64  |
| Co    | 32  | 41  | 40  | 30  | 43  |
| Cr    | 157 | 180 | 161 | 151 | 188 |
| V     | 278 | 320 | 260 | 270 | 290 |
| Hg    | 10  | 8   | 16  | 11  | 17  |
| Ag    | 15  | 12  | 13  | 13  | 11  |
| Nb    |     |     |     |     |     |



## FOOTWALL BASALT - MAFIC FRAGMENTALS

## MAJOR ELEMENTS (WTX)

| SAMPLE | 05421475 | 12141450 | 12141475 |
|--------|----------|----------|----------|
| SiO2   | 50.40    | 52.40    | 44.50    |
| TiO2   | .94      | .78      | .99      |
| Al2O3  | 17.65    | 16.28    | 20.69    |
| Fe2O3  | 10.96    | 10.70    | 14.41    |
| MnO    | .07      | .12      | .16      |
| HgO    | 2.66     | 4.12     | 5.96     |
| CaO    | 4.22     | 6.24     | 5.10     |
| Na2O   | .75      | 1.89     | 2.14     |
| K2O    | 3.44     | 1.92     | 2.35     |
| LOI    | 7.50     | 4.40     | 4.50     |
| TOTAL  | 98.59    | 98.85    | 100.80   |

## TRACE ELEMENTS (PPM)

|       |     |      |      |
|-------|-----|------|------|
| Zr    | 83  | 68   | 90   |
| Sr    | 128 | 292  | 318  |
| Rb    | 73  | 40   | 49   |
| Zn    | 75  | 95   | 127  |
| Cu    | 18  | 94   | 128  |
| Ba    | 495 | 1619 | 1981 |
| CuS   | 8   | 81   | 112  |
| PbS   | 6   | 12   | 6    |
| ZnS   | 2   | 31   | 38   |
| BaSO4 | 0   | -    | 0    |
| Pb    | 28  | 16   | 20   |
| Ni    | 39  | 13   | 16   |
| Co    | 20  | 28   | 32   |
| Cr    | 155 | 37   | 43   |
| V     | 420 | 381  | 390  |
| Hg    | 8   | 40   | 50   |
| Ag    | 9   | 2    | 1    |
| Nb    |     |      |      |

## FOOTWALL BASALT - ALTERED MAFIC FRAGMENTALS

## MAJOR ELEMENTS (WT%)

| SAMPLE | 02021225 | 542 775 | 5421075 | 05421575 | 05421625 |
|--------|----------|---------|---------|----------|----------|
| S102   | 31.90    | 59.90   | 42.10   | 31.80    | 42.50    |
| TiO2   | .37      | .36     | .83     | .77      | .65      |
| Al2O3  | 8.25     | 10.89   | 14.63   | 13.27    | 11.30    |
| Fe2O3  | 5.47     | 10.67   | 9.52    | 6.34     | 15.24    |
| MnO    | .20      | .24     | .17     | .15      | .20      |
| MgO    | 3.62     | 4.22    | 6.81    | 5.04     | 3.16     |
| CaO    | 27.40    | 1.22    | 13.30   | 20.70    | 12.50    |
| Na2O   | 1.22     | 1.82    | 2.29    | 1.13     | .85      |
| K2O    | .43      | .96     | .65     | .94      | 1.29     |
| LOI    | 21.60    | 6.60    | 9.30    | 20.20    | 12.10    |
| TOTAL  | 100.46   | 96.88   | 99.60   | 100.34   | 99.79    |

## TRACE ELEMENTS (PPM)

|       |     |      |     |     |     |
|-------|-----|------|-----|-----|-----|
| Zr    | 36  | 52   | 88  | 90  | 57  |
| Sr    | 121 | 61   | 285 | 194 | 101 |
| Rb    | 19  | 29   | 18  | 29  | 34  |
| Zn    | 78  | 125  | 87  | 75  | 77  |
| Cu    | 44  | 33   | 66  | 39  | 31  |
| Ba    | 129 | 2010 | 285 | 205 | 436 |
| CuS   | 29  | 27   | 28  | 21  | 29  |
| PbS   | 24  | 10   | 5   | 15  | 15  |
| ZnS   | 17  | 19   | 8   | 14  | 6   |
| BaSO4 | 0   |      | 0   |     |     |
| Pb    | 38  | 15   | 44  | 38  | 40  |
| Ni    | 36  | 50   | 51  | 28  | 47  |
| Co    | 20  | 42   | 33  | 33  | 27  |
| Cr    | 76  | 50   | 171 | 105 | 110 |
| V     | 163 | 161  | 270 | 290 | 168 |
| Hg    | 43  | 15   | 10  | 21  | 19  |
| Ag    | 15  | 0    | 15  | 13  | 14  |
| Nb    |     |      |     |     |     |

## FOOTWALL BASALT - ALTERED MAFIC FRAGMENTALS

## MAJOR ELEMENTS (WT%)

| SAMPLE | 05421700 | 05421875 | 05421950 | 05422025 | 05422100 |
|--------|----------|----------|----------|----------|----------|
| SiO2   | 35.50    | 34.50    | 44.20    | 43.90    | 35.50    |
| TiO2   | .80      | .66      | .97      | .73      | .71      |
| Al2O3  | 14.62    | 11.82    | 17.65    | 14.60    | 11.65    |
| Fe2O3  | 8.39     | 8.67     | 12.47    | 9.41     | 9.93     |
| MnO    | .17      | .17      | .13      | .19      | .18      |
| MgO    | 7.02     | 4.76     | 2.98     | 5.04     | 6.65     |
| CaO    | 15.33    | 19.36    | 6.00     | 13.30    | 16.85    |
| Na2O   | 1.37     | 1.82     | 1.19     | 2.66     | 1.50     |
| K2O    | 1.05     | .72      | 6.25     | .99      | 1.17     |
| LOI    | 15.70    | 17.10    | 7.30     | 7.70     | 15.80    |
| TOTAL  | 99.95    | 99.58    | 99.14    | 98.52    | 99.94    |

## TRACE ELEMENTS (PPM)

|       |     |     |     |     |     |
|-------|-----|-----|-----|-----|-----|
| Zr    | 76  | 103 | 87  | 94  | 66  |
| Sr    | 162 | 619 | 196 | 472 | 161 |
| Rb    | 31  | 26  | 110 | 27  | 37  |
| Zn    | 73  | 59  | 62  | 85  | 68  |
| Cu    | 41  | 24  | 55  | 47  | 52  |
| Ba    | 203 | 164 | 969 | 486 | 150 |
| CuS   | 37  | 19  | 41  | 30  | 51  |
| PbS   | 11  | 10  | 2   | 8   | 11  |
| ZnS   | 4   | 1   | 2   | 11  | 5   |
| BaSO4 |     |     |     |     | 0   |
| Pb    | 33  | 36  | 31  | 37  | 30  |
| Ni    | 21  | 94  | 82  | 77  | 56  |
| Co    | 26  | 37  | 29  | 36  | 32  |
| Cr    | 94  | 224 | 231 | 205 | 170 |
| V     | 310 | 290 | 310 | 255 | 300 |
| Hg    | 40  | 32  | 23  | 28  | 27  |
| Ag    | 12  | 13  | 11  | 13  | 11  |
| Nb    |     |     |     |     |     |

## FOOTWALL BASALT - ALTERED MAFIC FRAGMENTALS

## MAJOR ELEMENTS (WT%)

| SAMPLE | 05422150 | 05422200 | 08692020 | 14930000 | 14930100 |
|--------|----------|----------|----------|----------|----------|
| SiO2   | 37.80    | 39.60    | 33.50    | 36.00    | 42.10    |
| TiO2   | .59      | .79      | .63      | .63      | .94      |
| Al2O3  | 11.96    | 14.34    | 11.50    | 14.10    | 16.90    |
| Fe2O3  | 7.56     | 10.36    | 5.30     | 8.84     | 15.26    |
| MnO    | .14      | .18      | .15      | .11      | .06      |
| MgO    | 6.67     | 8.48     | 2.60     | 1.76     | 1.70     |
| CaO    | 15.74    | 13.63    | 23.24    | 19.86    | 9.00     |
| Na2O   | .64      | 2.45     | 1.95     | 2.59     | 1.90     |
| K2O    | 1.90     | .50      | .99      | 1.13     | 3.35     |
| LOI    | 16.30    | 9.27     | 19.60    | 15.00    | 8.70     |
| TOTAL  | 99.32    | 99.60    | 99.46    | 100.02   | 99.91    |

## TRACE ELEMENTS (PPM)

|       |     |     |     |     |     |
|-------|-----|-----|-----|-----|-----|
| Zr    | 38  | 93  | 69  | 81  | 95  |
| Sr    | 152 | 411 | 253 | 342 | 260 |
| Rb    | 41  | 19  | 27  | 34  | 66  |
| Zn    | 64  | 87  | 59  | 63  | 30  |
| Cu    | 21  | 54  | 91  | 43  | 4   |
| Ba    | 669 | 230 | 125 | 140 | 320 |
| CuS   | 11  | 43  | 77  | 34  | 1   |
| PbS   | 11  | 7   | 17  | 12  | 10  |
| ZnS   | 3   | 9   | 15  | 9   | 11  |
| BaSO4 |     |     | 0   | 0   |     |
| Pb    | 29  | 35  | 23  | 19  | 17  |
| Ni    | 56  | 97  | 45  | 55  | 53  |
| Co    | 28  | 44  | 26  | 20  | 17  |
| Cr    | 164 | 257 | 127 | 170 | 240 |
| V     | 270 | 260 | 290 | 170 | 129 |
| Hg    | 23  | 23  | 33  | 16  | 9   |
| Ag    | 11  | 10  | 0   | 0   | 0   |
| Nb    |     |     |     |     |     |

## FOOTWALL BASALT - ALTERED MAFIC FRAGMENTALS

## MAJOR ELEMENTS (WT%)

SAMPLE 14930250

|                                |       |
|--------------------------------|-------|
| SiO <sub>2</sub>               | 36.70 |
| TiO <sub>2</sub>               | .62   |
| Al <sub>2</sub> O <sub>3</sub> | 11.67 |
| Fe <sub>2</sub> O <sub>3</sub> | 7.25  |
| MnO                            | .11   |
| H <sub>2</sub> O               | 1.65  |
| CaO                            | 21.39 |
| Na <sub>2</sub> O              | 2.34  |
| K <sub>2</sub> O               | 2.01  |
| LOI                            | 16.80 |

TOTAL 100.54

## TRACE ELEMENTS (PPM)

|                   |     |
|-------------------|-----|
| Zr                | 82  |
| Sr                | 28  |
| Rb                | 50  |
| Zn                | 49  |
| Cu                | 22  |
| Ba                | 327 |
| CuS               | 15  |
| PbS               | 9   |
| ZnS               | 8   |
| BaSO <sub>4</sub> |     |
| Pb                | 19  |
| Ni                | 37  |
| Co                | 13  |
| Cr                | 123 |
| V                 | 16  |
| Hg                | 14  |
| Ag                | 2   |
| Nb                |     |

## FOOTBALL BASALT SEQUENCE - RHYOLITE FLOWS

## MAJOR ELEMENTS (WT%)

SAMPLE JGT 160B

|                                |       |
|--------------------------------|-------|
| SiO <sub>2</sub>               | 77.30 |
| TiO <sub>2</sub>               | .10   |
| Al <sub>2</sub> O <sub>3</sub> | 11.40 |
| Fe <sub>2</sub> O <sub>3</sub> | 2.20  |
| MnO                            | .03   |
| H <sub>2</sub> O               | 1.82  |
| CaO                            | .16   |
| Na <sub>2</sub> O              | 4.49  |
| K <sub>2</sub> O               | .50   |
| LOI                            | 1.50  |

|       |       |
|-------|-------|
| TOTAL | 99.58 |
|-------|-------|

## TRACE ELEMENTS (PPM)

|                   |     |
|-------------------|-----|
| Zr                | 125 |
| Sr                | 72  |
| Rb                | 11  |
| Zn                | 39  |
| Cu                | 1   |
| Ba                | 60  |
| CuS               | 1   |
| PbS               | 6   |
| ZnS               | 1   |
| BaSO <sub>4</sub> | 0   |
| Pb                | 6   |
| Ni                | 3   |
| Co                | 13  |
| Cr                | 11  |
| V                 | 10  |
| Hg                |     |
| Ag                |     |
| Nb                | 6   |

## BASALT FLOWS BELOW PROMINENT QUARTZ SEQUENCE

## MAJOR ELEMENTS (WT%)

| SAMPLE | 21051526 | 2820 153 |
|--------|----------|----------|
| SiO2   | 45.30    | 46.70    |
| TiO2   | 1.04     | 1.20     |
| Al2O3  | 16.61    | 16.75    |
| Fe2O3  | 9.38     | 9.90     |
| MnO    | .14      | .20      |
| MgO    | 9.36     | 8.59     |
| CaO    | 11.55    | 10.35    |
| Na2O   | 1.85     | 2.00     |
| K2O    | .12      | 1.15     |
| LOI    | 4.00     | 3.10     |
| TOTAL  | 99.35    | 99.94    |

## TRACE ELEMENTS (PPM)

|       |     |     |
|-------|-----|-----|
| Zr    | 73  | 57  |
| Sr    | 148 | 150 |
| Rb    | 2   | 22  |
| Zn    | 50  | 52  |
| Cu    | 70  | 54  |
| Ba    | 57  | 276 |
| CuS   | 135 | 53  |
| PbS   | 20  | 62  |
| ZnS   | 20  | 18  |
| BaSO4 |     | 0   |
| Pb    | 20  | 92  |
| Ni    | 133 | 50  |
| Co    | 36  | 35  |
| Cr    | 157 | 258 |
| V     | 125 | 130 |
| Hg    |     |     |
| Ag    |     |     |
| Nb    |     | 1   |

## ALTERED BASALTIC FLOWS BELOW PROMINENT QUARTZ SEQUENCE

## MAJOR ELEMENTS (WT%)

SAMPLE 2867 143

|                                |       |
|--------------------------------|-------|
| SiO <sub>2</sub>               | 49.88 |
| TiO <sub>2</sub>               | .91   |
| Al <sub>2</sub> O <sub>3</sub> | 15.24 |
| Fe <sub>2</sub> O <sub>3</sub> | 18.95 |
| MnO                            | .16   |
| H <sub>2</sub> O               | 8.93  |
| CaO                            | 4.69  |
| Na <sub>2</sub> O              | 4.85  |
| K <sub>2</sub> O               | .89   |
| LOI                            | 5.48  |

|       |       |
|-------|-------|
| TOTAL | 98.42 |
|-------|-------|

## TRACE ELEMENTS (PPM)

|                   |     |
|-------------------|-----|
| Zr                | 144 |
| Sr                | 166 |
| Rb                | 5   |
| Zn                | 136 |
| Cu                | 16  |
| Ba                | 97  |
| CuS               | 19  |
| PbS               | 14  |
| ZnS               | 26  |
| BaSO <sub>4</sub> | 8   |
| Pb                | 35  |
| Ni                | 27  |
| Co                | 31  |
| Cr                | 156 |
| V                 | 285 |
| Hg                |     |
| Ag                |     |
| Nb                | 5   |



3-12

## FOOTWALL ARKOSE SEQUENCE - DACITIC TUFFS

## MAJOR ELEMENTS (WT%)

SAMPLE JBT75216 R1-5 153

|                                |       |       |
|--------------------------------|-------|-------|
| SiO <sub>2</sub>               | 72.88 | 64.28 |
| TiO <sub>2</sub>               | .24   | .24   |
| Al <sub>2</sub> O <sub>3</sub> | 12.55 | 17.97 |
| Fe <sub>2</sub> O <sub>3</sub> | 2.52  | 2.57  |
| MnO                            | .87   | .88   |
| HgO                            | 1.18  | 3.49  |
| CaO                            | 1.24  | 1.18  |
| Na <sub>2</sub> O              | 3.59  | 1.27  |
| K <sub>2</sub> O               | 2.57  | 3.78  |
| LOI                            | 1.78  | 4.88  |
| TOTAL                          | 98.46 | 99.58 |

## TRACE ELEMENTS (PPM)

|                   |     |     |
|-------------------|-----|-----|
| Zr                | 71  | 148 |
| Sr                | 138 | 248 |
| Rb                | 49  | 182 |
| Zn                | 66  | 41  |
| Cu                | 2   | 2   |
| Ba                | 615 | 989 |
| CuS               | 1   | 3   |
| PbS               | 4   | 6   |
| ZnS               | 5   | 8   |
| BaSO <sub>4</sub> | 8   | 8   |
| Pb                | 17  | 17  |
| Ni                | 1   | 8   |
| Co                | 8   | 6   |
| Cr                | 12  | 8   |
| V                 | 25  | 15  |
| Hg                |     |     |
| Ag                |     |     |
| Nb                | 7   | 12  |

## WILBY'S SEQUENCE - FLOWS

## MAJOR ELEMENTS (WT%)

| SAMPLE | JGT75243 | JGT75307 | 724 303 | 2622 60 | 2622 335 |
|--------|----------|----------|---------|---------|----------|
| S102   | 72.60    | 66.50    | 66.80   | 59.20   | 70.40    |
| TiO2   | .11      | .40      | .54     | .86     | .12      |
| Al2O3  | 10.96    | 13.92    | 14.13   | 15.30   | 15.49    |
| Fe2O3  | 1.57     | 5.00     | 5.23    | 6.14    | 2.09     |
| MnO    | .10      | .19      | .23     | .16     | .02      |
| HgO    | .11      | 1.95     | 1.50    | 2.99    | .01      |
| CaO    | 4.00     | 1.38     | 1.83    | 3.76    | 1.10     |
| Na2O   | 4.22     | 4.99     | 4.49    | 4.39    | 2.10     |
| K2O    | 1.29     | 2.13     | 2.91    | 2.61    | 3.59     |
| LOI    | 4.20     | 2.10     | 1.80    | 3.60    | 3.50     |
| TOTAL  | 99.16    | 98.64    | 99.54   | 99.09   | 99.30    |

## TRACE ELEMENTS (PPM)

|       |     |     |      |     |     |
|-------|-----|-----|------|-----|-----|
| Zr    | 96  | 91  | 92   | 218 | 78  |
| Sr    | 247 | 103 | 119  | 418 | 95  |
| Rb    | 26  | 37  | 47   | 49  | 82  |
| Zn    | 21  | 44  | 61   | 73  | 7   |
| Cu    | 1   | 1   | 1    | 33  | 1   |
| Ba    | 506 | 966 | 1127 | 819 | 432 |
| CuS   | 1   | 1   | 2    | 3   | 2   |
| PbS   | 17  | 3   | 23   | 4   | 7   |
| ZnS   | 10  | 1   | 14   | 15  | 8   |
| BaSO4 | 4   | 0   | 0    | 0   | 0   |
| Pb    | 31  | 14  | 23   | 17  | 5   |
| Ni    | 1   | 1   | 5    | 50  | 8   |
| Co    | 21  | 21  | 18   | 22  | 11  |
| Cr    | 10  | 8   | 14   | 115 | 7   |
| V     | 10  | 15  | 10   | 120 | 30  |
| Hg    |     |     |      |     |     |
| Ag    |     |     |      |     |     |
| Nb    | 9   | 9   | 5    | 17  | 12  |

## WILBY'S BEQUENCE - FLOWS

## MAJOR ELEMENTS (WT%)

SAMPLE 2622 365

|       |       |
|-------|-------|
| SiO2  | 55.88 |
| TiO2  | .62   |
| Al2O3 | 19.28 |
| Fe2O3 | 8.87  |
| MnO   | .17   |
| HgO   | 2.74  |
| CaO   | 2.54  |
| Na2O  | 4.55  |
| K2O   | 2.22  |
| LOI   | 4.58  |

TOTAL 99.61

## TRACE ELEMENTS (PPM)

|       |     |
|-------|-----|
| Zr    | 45  |
| Sr    | 255 |
| Rb    | 61  |
| Zn    | 161 |
| Cu    | 4   |
| Ba    | 657 |
| CuS   | 2   |
| PbS   | 8   |
| ZnS   | 35  |
| BaSO4 | 8   |
| Pb    | 21  |
| Ni    | 58  |
| Co    | 45  |
| Cr    | 77  |
| V     | 128 |
| Hg    |     |
| Ag    |     |
| Nb    | 8   |

PROMINENT QUARTZ SEQUENCE - RHYOLITE FLOWS

MAJOR ELEMENTS (WT%)

| SAMPLE | 19402111 | 2090 242 | 2105 177 | 2753 505 | 2762 101 |
|--------|----------|----------|----------|----------|----------|
| S102   | 76.80    | 69.30    | 67.80    | 77.30    | 70.70    |
| T102   | .26      | .35      | .45      | .40      | .29      |
| Al2O3  | 11.54    | 13.55    | 12.43    | 11.06    | 13.28    |
| Fe2O3  | 1.73     | 4.08     | 4.70     | 1.97     | 3.20     |
| MnO    | .04      | .06      | .10      | .05      | .05      |
| KgO    | .64      | 1.49     | 1.51     | .11      | .71      |
| CaO    | 1.00     | 2.57     | 2.96     | 1.72     | 2.21     |
| Na2O   | 2.42     | 2.99     | 2.68     | 3.82     | 3.43     |
| K2O    | 4.22     | 3.57     | 2.52     | 1.34     | 2.74     |
| LOI    | 1.80     | 2.10     | 4.20     | .80      | 2.90     |
| TOTAL  | 100.45   | 100.06   | 99.35    | 98.57    | 99.51    |

TRACE ELEMENTS (PPM)

|       |      |     |     |     |     |
|-------|------|-----|-----|-----|-----|
| Zr    | 154  | 149 | 148 | 135 | 146 |
| Sr    | 90   | 143 | 93  | 229 | 137 |
| Rb    | 50   | 63  | 57  | 22  | 63  |
| Zn    | 28   | 37  | 52  | 20  | 17  |
| Cu    | 1    | 1   | 15  | 4   | 1   |
| Ba    | 1216 | 662 | 419 | 782 | 491 |
| CuS   | 1    | 3   | 29  | 4   | 1   |
| PbS   | 23   | 25  | 20  | 11  | 12  |
| ZnS   | 16   | 12  | 11  | 20  | 22  |
| BaSO4 |      |     |     |     |     |
| Pb    | 12   | 6   | 13  | 1   | 3   |
| Ni    | 3    | 5   | 8   | 3   | 5   |
| Co    | 22   | 19  | 22  | 58  | 29  |
| Cr    | 0    | 11  | 16  | 3   | 9   |
| V     | 10   | 50  | 55  | 10  | 25  |
| Hg    |      |     |     |     |     |
| Ag    |      |     |     |     |     |
| Nb    |      |     |     |     |     |

## WILBY'S SEQUENCE - DACITIC TUFFS

## MAJOR ELEMENTS (WT%)

| SAMPLE | JGT75190 | 718   | 37 | 2622  | 242 | 2660  | 767 | 2754  | 728 |
|--------|----------|-------|----|-------|-----|-------|-----|-------|-----|
| SiO2   | 66.00    | 69.50 |    | 64.90 |     | 67.90 |     | 64.70 |     |
| TiO2   | .40      | .56   |    | .50   |     | .33   |     | .32   |     |
| Al2O3  | 13.82    | 13.51 |    | 16.54 |     | 13.13 |     | 16.97 |     |
| Fe2O3  | 5.14     | 3.06  |    | 3.75  |     | 3.88  |     | 3.26  |     |
| MnO    | .08      | .05   |    | .05   |     | .08   |     | .11   |     |
| MgO    | 1.72     | .95   |    | .79   |     | 1.25  |     | 1.15  |     |
| CaO    | 3.71     | 4.10  |    | 2.41  |     | 3.09  |     | 4.14  |     |
| Na2O   | 1.24     | 3.01  |    | 1.77  |     | 2.66  |     | 3.93  |     |
| K2O    | 3.04     | 1.04  |    | 3.52  |     | 2.60  |     | 2.72  |     |
| LOI    | 4.30     | 3.00  |    | 5.40  |     | 4.20  |     | 2.10  |     |
| TOTAL  | 99.45    | 98.78 |    | 99.71 |     | 99.12 |     | 99.40 |     |

## TRACE ELEMENTS (PPM)

|       |     |     |     |     |     |
|-------|-----|-----|-----|-----|-----|
| Zr    | 118 | 128 | 161 | 150 | 97  |
| Sr    | 528 | 220 | 173 | 146 | 348 |
| Rb    | 74  | 31  | 95  | 80  | 60  |
| Zn    | 56  | 55  | 35  | 41  | 51  |
| Cu    | 2   | 9   | 2   | 9   | 1   |
| Ba    | 767 | 314 | 601 | 310 | 759 |
| CuS   | 2   | 8   | 4   | 12  | 1   |
| PbB   | 5   | 8   | 23  | 19  | 3   |
| ZnS   | 6   | 11  | 18  | 17  | 12  |
| BaSO4 | 6   | 8   | 8   |     | 8   |
| Pb    | 17  | 14  | 7   | 4   | 13  |
| Ni    | 5   | 12  | 24  | 7   | 3   |
| Co    | 9   | 25  | 13  | 18  | 17  |
| Cr    | 10  | 11  | 10  | 15  | 11  |
| V     | 20  | 25  | 30  | 25  | 45  |
| Hg    |     |     |     |     |     |
| Ag    |     |     |     |     |     |
| Nb    | 7   | 11  | 0   |     | 6   |

## WILBY'S SEQUENCE - DACITIC TUFFS

## MAJOR ELEMENTS (WT%)

| SAMPLE                         | 2769 219 | 2777 545 | 2888 311 | BB 229 | BB 232 |
|--------------------------------|----------|----------|----------|--------|--------|
| SiO <sub>2</sub>               | 67.78    | 71.98    | 78.48    | 76.88  | 62.88  |
| TiO <sub>2</sub>               | .38      | .51      | .26      | .82    | .26    |
| Al <sub>2</sub> O <sub>3</sub> | 18.28    | 12.99    | 13.43    | 13.17  | 18.56  |
| Fe <sub>2</sub> O <sub>3</sub> | 1.95     | 2.38     | 2.31     | .89    | 5.84   |
| MnO                            | .81      | .83      | .86      | .81    | .18    |
| HgO                            | 1.81     | .99      | 3.34     | .61    | 1.54   |
| CaO                            | .32      | 1.54     | 1.11     | 1.88   | .54    |
| Na <sub>2</sub> O              | .43      | 1.93     | .74      | 3.51   | 8.51   |
| K <sub>2</sub> O               | 4.82     | 3.39     | 3.28     | 2.51   | .69    |
| LOI                            | 4.28     | 3.88     | 4.68     | 2.58   | 1.58   |
| TOTAL                          | 99.82    | 99.46    | 99.45    | 101.82 | 99.54  |

## TRACE ELEMENTS (PPM)

|                   |     |     |     |    |     |
|-------------------|-----|-----|-----|----|-----|
| Zr                | 169 | 122 | 187 |    | 133 |
| Sr                | 78  | 112 | 89  | 97 | 59  |
| Rb                | 187 | 88  | 85  | 88 | 15  |
| Zn                | 14  | 18  | 98  |    |     |
| Cu                | 1   | 2   | 11  |    |     |
| Ba                | 774 | 438 | 668 | 19 | 76  |
| CuS               | 1   | 4   | 14  |    |     |
| PbS               | 17  | 28  | 38  |    |     |
| ZnS               | 17  | 17  | 22  |    |     |
| BaSO <sub>4</sub> |     |     |     |    |     |
| Pb                | 1   | 11  | 28  |    |     |
| Ni                | 3   | 2   | 6   |    |     |
| Co                | 11  | 22  | 19  |    |     |
| Cr                | 8   | 5   | 6   | 8  | 18  |
| V                 | 15  | 18  | 28  |    |     |
| Hg                |     |     |     |    |     |
| Ag                |     |     |     |    |     |
| Nb                |     |     |     |    |     |

## WILBY'S SEQUENCE - DACITIC TUFFS

## MAJOR ELEMENTS (WT%)

| SAMPLE                         | BB    | 246    | BB     | 247 | BB | 257 |
|--------------------------------|-------|--------|--------|-----|----|-----|
| SiO <sub>2</sub>               | 69.48 | 69.18  | 76.98  |     |    |     |
| TiO <sub>2</sub>               | .28   | .32    | .14    |     |    |     |
| Al <sub>2</sub> O <sub>3</sub> | 13.71 | 14.65  | 11.45  |     |    |     |
| Fe <sub>2</sub> O <sub>3</sub> | 4.11  | 4.48   | 1.39   |     |    |     |
| MnO                            | .85   | .87    | .83    |     |    |     |
| MgO                            | 1.29  | 1.37   | 2.63   |     |    |     |
| CaO                            | 2.83  | 2.84   | .58    |     |    |     |
| Na <sub>2</sub> O              | 2.94  | 3.55   | .31    |     |    |     |
| K <sub>2</sub> O               | 2.75  | 2.73   | 3.29   |     |    |     |
| LOI                            | 2.78  | 2.88   | 3.48   |     |    |     |
| TOTAL                          | 99.26 | 101.83 | 108.84 |     |    |     |

## TRACE ELEMENTS (PPM)

|                   |     |     |     |
|-------------------|-----|-----|-----|
| Zr                | 154 | 147 | 111 |
| Sr                | 178 | 217 | 69  |
| Rb                | 67  | 54  | 91  |
| Zn                |     |     |     |
| Cu                |     |     |     |
| Ba                | 558 | 648 | 482 |
| CuS               |     |     |     |
| PbS               |     |     |     |
| ZnS               |     |     |     |
| BaSO <sub>4</sub> |     |     |     |
| Pb                |     |     |     |
| Ni                |     |     |     |
| Co                |     |     |     |
| Cr                | 13  | 13  | 8   |
| V                 |     |     |     |
| Hg                |     |     |     |
| Ag                |     |     |     |
| Nb                |     |     |     |

# PROMINENT QUARTZ SEQUENCE - DACITIC TUFFS

## MAJOR ELEMENTS (WT%)

| SAMPLE                         | 1908  | 85 | 1940  | 217 | 1960  | 233 | 2090  | 35 | 2490  | 527 |
|--------------------------------|-------|----|-------|-----|-------|-----|-------|----|-------|-----|
| SiO <sub>2</sub>               | 68.80 |    | 64.70 |     | 68.60 |     | 69.70 |    | 63.90 |     |
| TiO <sub>2</sub>               | .20   |    | .44   |     | .41   |     | .30   |    | .45   |     |
| Al <sub>2</sub> O <sub>3</sub> | 14.09 |    | 15.50 |     | 14.45 |     | 13.76 |    | 15.40 |     |
| Fe <sub>2</sub> O <sub>3</sub> | 3.69  |    | 4.46  |     | 3.66  |     | 3.64  |    | 5.46  |     |
| MnO                            | .13   |    | .05   |     | .05   |     | .07   |    | .04   |     |
| HgO                            | 1.55  |    | 2.02  |     | 1.11  |     | 1.00  |    | 3.37  |     |
| CaO                            | 2.07  |    | 2.95  |     | 3.82  |     | 2.61  |    | 1.68  |     |
| Na <sub>2</sub> O              | 2.59  |    | 2.24  |     | 2.82  |     | 3.42  |    | 1.63  |     |
| K <sub>2</sub> O               | 2.26  |    | 2.78  |     | 1.67  |     | 1.94  |    | 3.96  |     |
| LOI                            | 4.00  |    | 4.40  |     | 2.90  |     | 3.10  |    | 3.40  |     |
| TOTAL                          | 99.38 |    | 99.62 |     | 99.49 |     | 99.54 |    | 99.29 |     |

## TRACE ELEMENTS (PPM)

|                   |     |     |     |     |     |
|-------------------|-----|-----|-----|-----|-----|
| Zr                | 99  | 170 | 158 | 153 | 140 |
| Sr                | 91  | 144 | 199 | 191 | 177 |
| Rb                | 53  | 66  | 46  | 46  | 143 |
| Zn                | 227 | 65  | 29  | 35  | 23  |
| Cu                | 1   | 2   | 1   | 1   | 4   |
| Ba                | 663 | 515 | 469 | 482 | 923 |
| CuS               | 1   | 9   | 4   | 1   | 5   |
| PbS               | 10  | 19  | 12  | 22  | 11  |
| ZnS               | 31  | 23  | 7   | 17  | 15  |
| BaSO <sub>4</sub> | 0   |     |     |     |     |
| Pb                | 1   | 11  | 4   | 7   | 1   |
| Ni                | 6   | 9   | 4   | 6   | 7   |
| Co                | 16  | 19  | 19  | 17  | 15  |
| Cr                | 11  | 16  | 12  | 12  | 16  |
| V                 | 10  | 15  | 50  | 30  | 10  |
| Hg                |     |     |     |     |     |
| Ag                |     |     |     |     |     |
| Nb                | 12  |     |     |     |     |



# PROHINENT QUARTZ SEQUENCE - DACITIC TUFFS

## MAJOR ELEMENTS (WT%)

| SAMPLE                         | 2501 234 | 2501 469 | 2528 113 | 2530 110 | 2571 77 |
|--------------------------------|----------|----------|----------|----------|---------|
| SiO <sub>2</sub>               | 74.20    | 72.30    | 68.40    | 62.90    | 69.20   |
| TiO <sub>2</sub>               | .34      | .28      | .33      | .31      | .24     |
| Al <sub>2</sub> O <sub>3</sub> | 12.21    | 14.43    | 13.79    | 16.97    | 13.31   |
| Fe <sub>2</sub> O <sub>3</sub> | 2.57     | 1.25     | 3.41     | 4.10     | 3.31    |
| MnO                            | .03      | .02      | .07      | .05      | .05     |
| HgO                            | 4.43     | 1.24     | 1.87     | 1.84     | .88     |
| CaO                            | 1.50     | 1.52     | 2.46     | 2.11     | 2.95    |
| Na <sub>2</sub> O              | 1.68     | 2.12     | 2.85     | 1.91     | 2.05    |
| K <sub>2</sub> O               | 3.76     | 3.80     | 2.02     | 2.97     | 2.67    |
| LOI                            | 2.00     | 1.90     | 3.80     | 5.30     | 4.70    |
| TOTAL                          | 99.72    | 98.86    | 99.00    | 98.54    | 99.36   |

## TRACE ELEMENTS (PPM)

|                   |     |     |     |     |     |
|-------------------|-----|-----|-----|-----|-----|
| Zr                | 91  | 107 | 117 | 125 | 151 |
| Sr                | 154 | 151 | 187 | 100 | 9   |
| Rb                | 124 | 86  | 64  | 66  | 53  |
| Zn                | 45  | 36  | 44  | 30  | 34  |
| Cu                | 4   | 1   | 1   | 5   | 2   |
| Ba                | 894 | 853 | 829 | 793 | 374 |
| CuS               | 2   | 2   | 2   | 4   | 2   |
| PbS               | 8   | 9   | 8   | 9   | 29  |
| ZnS               | 17  | 13  | 14  | 12  | 10  |
| BaSO <sub>4</sub> | 0   | 0   | 6   | 0   |     |
| Pb                | 12  | 5   | 8   | 7   | 17  |
| Ni                | 4   | 4   | 7   | 5   | 5   |
| Co                | 8   | 8   | 11  | 10  | 17  |
| Cr                | 10  | 11  | 14  | 13  | 11  |
| V                 | 10  | 10  | 10  | 50  | 15  |
| Hg                |     |     |     |     |     |
| Ag                |     |     |     |     |     |
| Nb                | 11  | 7   | 10  | 15  |     |

# PROMINENT QUARTZ SEQUENCE - DACITIC TUFFS

## MAJOR ELEMENTS (WTZ)

| SAMPLE | 26660048 | 26660200 | 26660300 | 26660350 | 26660475 |
|--------|----------|----------|----------|----------|----------|
| SiO2   | 66.50    | 70.60    | 74.40    | 73.60    | 71.90    |
| TiO2   | .37      | .32      | .20      | .10      | .36      |
| Al2O3  | 15.17    | 14.80    | 12.60    | 13.60    | 13.35    |
| Fe2O3  | 4.47     | 2.04     | 1.41     | 1.89     | 3.64     |
| MnO    | .10      | .02      | .02      | .03      | .04      |
| HgO    | 1.72     | .93      | .64      | 1.37     | 1.48     |
| CaO    | 2.27     | 3.55     | 3.15     | 1.22     | 1.98     |
| Na2O   | 3.10     | 3.14     | 2.65     | 2.20     | 3.14     |
| K2O    | 2.61     | 1.32     | 1.15     | 2.03     | 1.86     |
| LOI    | 2.20     | 3.40     | 2.90     | 2.60     | 2.00     |
| TOTAL  | 98.59    | 100.20   | 99.20    | 98.72    | 99.75    |

## TRACE ELEMENTS (PPH)

|       |     |     |     |     |     |
|-------|-----|-----|-----|-----|-----|
| Zr    | 181 | 171 | 120 | 126 | 114 |
| Sr    | 211 | 401 | 514 | 597 | 342 |
| Rb    | 60  | 43  | 38  | 74  | 69  |
| Zn    | 20  | 22  | 20  | 22  | 34  |
| Cu    | 27  | 4   | 5   | 5   | 4   |
| Ba    | 696 | 564 | 519 | 832 | 619 |
| CuS   | 25  | 1   | 1   | 2   | 1   |
| PbS   | 6   | 7   | 3   | 8   | 8   |
| ZnS   | 17  | 13  | 0   | 9   | 10  |
| BaSO4 |     |     |     |     | 0   |
| Pb    | 6   | 10  | 7   | 1   | 5   |
| Ni    | 4   | 2   | 1   | 1   | 4   |
| Co    | 4   | 2   | 2   | 1   | 4   |
| Cr    | 30  | 22  | 23  | 17  | 31  |
| V     | 54  | 24  | 10  | 0   | 39  |
| Hg    | 20  | 14  | 20  | 27  | 0   |
| Ag    | 0   | 1   | 0   | 0   | 0   |
| Nb    |     |     |     |     |     |

# PROMINENT QUARTZ SEQUENCE - DACITIC TUFFS

## MAJOR ELEMENTS (WT%)

| SAMPLE | 26660600 | 26660725 | 26660850 | 2680 267 | 28511475 |
|--------|----------|----------|----------|----------|----------|
| S102   | 68.20    | 72.10    | 75.40    | 71.50    | 70.40    |
| T102   | .28      | .24      | .20      | .25      | .36      |
| Al2O3  | 14.05    | 14.97    | 12.71    | 12.84    | 13.10    |
| Fe2O3  | 2.23     | 1.88     | 1.82     | 3.17     | 3.40     |
| MnO    | .03      | .03      | .05      | .03      | .06      |
| MgO    | 1.58     | 1.63     | 2.31     | .63      | 2.39     |
| CaO    | 2.10     | 2.71     | 1.17     | 1.84     | 1.11     |
| Na2O   | 3.03     | 3.20     | 1.87     | 3.54     | 3.46     |
| K2O    | 1.67     | 1.54     | 2.48     | 2.01     | 2.64     |
| LOI    | 3.00     | 2.40     | 2.50     | 2.70     | 2.00     |
| TOTAL  | 96.17    | 100.70   | 100.51   | 98.51    | 98.92    |

## TRACE ELEMENTS (PPM)

|       |     |     |     |     |     |
|-------|-----|-----|-----|-----|-----|
| Zr    | 117 | 120 | 126 | 146 | 111 |
| Sr    | 248 | 365 | 233 | 134 | 98  |
| Rb    | 52  | 47  | 58  | 40  | 48  |
| Zn    | 12  | 8   | 13  | 17  | 28  |
| Cu    | 6   | 1   | 3   | 3   | 5   |
| Ba    | 740 | 744 | 670 | 226 | 545 |
| CuS   | 3   | 0   | 1   | 4   | 8   |
| PbS   | 7   | 8   | 15  | 17  | 13  |
| ZnS   | 4   | 9   | 7   | 14  | 6   |
| BaSO4 | 0   | 0   | 0   | 0   | 0   |
| Pb    | 3   | 3   | 2   | 7   | 7   |
| Ni    | 2   | 2   | 2   | 4   | 3   |
| Co    | 2   | 2   | 1   | 20  | 21  |
| Cr    | 23  | 21  | 20  | 8   | 13  |
| V     | 15  | 9   | 2   | 35  | 30  |
| Hg    | 0   | 6   | 7   |     |     |
| Ag    | 0   | 0   | 0   |     |     |
| Nb    |     |     |     |     | 13  |

# PROMINENT QUARTZ SEQUENCE - DACITIC TUFFS

## MAJOR ELEMENTS (WT%)

SAMPLE 28511955 2858 65

|       |       |       |
|-------|-------|-------|
| S102  | 69.38 | 68.98 |
| T102  | .48   | .52   |
| Al2O3 | 14.38 | 14.27 |
| Fe2O3 | 4.76  | 3.98  |
| MnO   | .85   | .85   |
| HgO   | .98   | 1.59  |
| CaO   | 1.68  | 3.49  |
| Na2O  | 2.11  | 2.75  |
| K2O   | 3.12  | 1.43  |
| LOI   | 3.58  | 2.78  |

TOTAL 188.12 99.68

## TRACE ELEMENTS (PPM)

|       |     |     |
|-------|-----|-----|
| Zr    | 186 | 168 |
| Sr    | 68  | 242 |
| Rb    | 65  | 33  |
| Zn    | 43  | 29  |
| Cu    | 5   | 1   |
| Ba    | 511 | 484 |
| CuS   | 8   | 1   |
| PbS   | 22  | 28  |
| ZnS   | 18  | 4   |
| BaSO4 | 8   |     |
| Pb    | 8   | 5   |
| Ni    | 6   | 6   |
| Co    | 17  | 21  |
| Cr    | 12  | 14  |
| V     | 38  | 18  |
| Hg    |     |     |
| Ag    |     |     |
| Nb    | 15  |     |

# PROMINENT QUARTZ SEQUENCE - TUFFACEOUS SILTSTONE

## MAJOR ELEMENTS (WT%)

| SAMPLE | 26661850 | 08691625 | 08691650 | 08691740 | 08691850 |
|--------|----------|----------|----------|----------|----------|
| SiO2   | 76.50    | 67.20    | 67.10    | 75.20    | 70.30    |
| TiO2   | .10      | .31      | .47      | .18      | .21      |
| Al2O3  | 12.67    | 14.20    | 13.60    | 11.10    | 13.60    |
| Fe2O3  | 1.31     | 3.22     | 6.23     | 3.25     | 2.59     |
| MnO    | .03      | .34      | .14      | .10      | .14      |
| MgO    | 2.29     | 1.34     | 1.82     | 1.34     | .77      |
| CaO    | .55      | 1.71     | 1.94     | .73      | 1.29     |
| Na2O   | .39      | 1.90     | 3.13     | 2.73     | 2.98     |
| K2O    | 3.20     | 6.34     | 2.97     | 3.38     | 6.10     |
| LOI    | 3.30     | 3.20     | 1.80     | 1.40     | 1.20     |
| TOTAL  | 100.42   | 99.76    | 99.20    | 99.41    | 99.26    |

## TRACE ELEMENTS (PPM)

|       |     |      |     |     |     |
|-------|-----|------|-----|-----|-----|
| Zr    | 54  | 136  | 135 | 116 | 120 |
| Sr    | 11  | 77   | 224 | 87  | 110 |
| Rb    | 62  | 91   | 38  | 66  | 79  |
| Zn    | 23  | 43   | 51  | 23  | 29  |
| Cu    | 52  | 17   | 6   | 10  | 4   |
| Ba    | 811 | 1126 | 502 | 751 | 861 |
| CuS   | 25  | 14   | 2   | 6   | 1   |
| PbS   | 4   | 25   | 5   | 14  | 12  |
| ZnS   | 1   | 12   | 17  | 8   | 8   |
| BaSO4 | 6   |      | 0   |     | 4   |
| Pb    | 1   | 29   | 11  | 19  | 19  |
| Ni    | 1   | 4    | 4   | 4   | 2   |
| Co    | 1   | 6    | 10  | 5   | 3   |
| Cr    | 22  | 24   | 33  | 27  | 25  |
| V     | 6   | 44   | 78  | 26  | 22  |
| Hg    | .6  | 81   | 23  | 33  | 19  |
| Ag    | 2   | .1   | 2   | 1   | 0   |
| Nb    |     |      |     |     |     |

# PROMINENT QUARTZ SEQUENCE - TUFFACEOUS SILTSTONE

## MAJOR ELEMENTS (WT%)

| SAMPLE | 26660150 | 26660900 | 08691477 |
|--------|----------|----------|----------|
| SiO2   | 70.40    | 47.70    | 64.60    |
| TiO2   | .10      | .90      | .40      |
| Al2O3  | 14.16    | 17.44    | 14.70    |
| Fe2O3  | 1.70     | 10.80    | 4.70     |
| MnO    | .00      | .20      | .31      |
| MgO    | 3.43     | 7.39     | 2.49     |
| CaO    | .92      | 7.63     | 1.71     |
| Na2O   | .06      | 3.31     | 1.20     |
| K2O    | 4.61     | 1.02     | 5.26     |
| LOI    | 4.90     | 4.10     | 3.70     |
| TOTAL  | 100.44   | 100.49   | 99.15    |

## TRACE ELEMENTS (PPM)

|       |      |     |      |
|-------|------|-----|------|
| Zr    | 71   | 103 | 134  |
| Sr    | 0    | 530 | 114  |
| Rb    | 114  | 25  | 95   |
| Zn    | 41   | 14  | 72   |
| Cu    | 3    | 2   | 19   |
| Ba    | 1023 | 774 | 1135 |
| CuS   | 1    | 0   | 11   |
| PbS   | 4    | 7   | 10   |
| ZnS   | 19   | 0   | 22   |
| BaSO4 | 0    |     |      |
| Pb    | 5    | 4   | 21   |
| Ni    | 2    | 1   | 0    |
| Co    | 2    | 1   | 10   |
| Cr    | 10   | 154 | 21   |
| V     | 0    | 0   | 42   |
| Hg    | 32   | 0   | 21   |
| Ag    | 0    | 0   | 2    |
| Nb    |      |     |      |

# PROMINENT QUARTZ SEQUENCE - MAFIC FRAGMENTALS

## MAJOR ELEMENTS (WT%)

| SAMPLE | 26661325 | 26661725 | 26661850 | 26662000 | 26662125 |
|--------|----------|----------|----------|----------|----------|
| SiO2   | 56.70    | 49.60    | 54.30    | 65.80    | 61.00    |
| TiO2   | .46      | .40      | .63      | .33      | .43      |
| Al2O3  | 17.84    | 16.72    | 17.69    | 14.00    | 17.70    |
| Fe2O3  | 9.15     | 10.48    | 3.66     | 5.70     | 5.34     |
| MnO    | .05      | .00      | .09      | .09      | .07      |
| HgO    | 3.65     | 8.80     | .92      | 5.20     | 5.62     |
| CaO    | 2.71     | 4.02     | 7.45     | .30      | .30      |
| Na2O   | 2.03     | 1.93     | 7.25     | 3.70     | 2.87     |
| K2O    | 4.95     | 2.38     | 1.67     | 1.15     | 2.84     |
| LOI    | 3.80     | 5.80     | 6.70     | 4.10     | 4.30     |
| TOTAL  | 101.34   | 100.21   | 100.36   | 100.37   | 100.55   |

## TRACE ELEMENTS (PPM)

|       |     |     |     |    |    |
|-------|-----|-----|-----|----|----|
| Zr    | 54  | 43  | 50  | 23 | 30 |
| Sr    | 180 | 272 | 181 | 3  | 0  |
| Rb    | 113 | 53  | 24  | 15 | 27 |
| Zn    | 24  | 38  | 43  | 52 | 58 |
| Cu    | 9   | 54  | 16  | 10 | 9  |
| Ba    | 463 | 246 | 115 | 74 | 96 |
| CuS   | 4   | 48  | 17  | 4  | 8  |
| PbS   | 5   | 7   | 7   | 30 | 7  |
| ZnS   | 3   | 6   | 12  | 23 | 17 |
| BaSO4 | 0   |     |     |    |    |
| Pb    | 4   | 0   | 0   | 13 | 1  |
| Ni    | 50  | 178 | 1   | 3  | 3  |
| Co    | 10  | 39  | 6   | 7  | 5  |
| Cr    | 490 | 532 | 19  | 30 | 28 |
| V     | 96  | 156 | 68  | 57 | 84 |
| Hg    | 6   | 6   | 10  | 39 | 20 |
| Ag    | 0   | 0   | 0   | 3  | 0  |
| Nb    |     |     |     |    |    |

PROMINENT QUARTZ SEQUENCE - MAFIC FRAGMENTALS

MAJOR ELEMENTS (WT%)

| SAMPLE                         | 26662450 | 26662575 | 26662700 | 26662825 | 26662950 |
|--------------------------------|----------|----------|----------|----------|----------|
| SiO <sub>2</sub>               | 60.80    | 63.80    | 59.10    | 51.50    | 51.90    |
| TiO <sub>2</sub>               | .32      | .32      | .30      | .40      | .43      |
| Al <sub>2</sub> O <sub>3</sub> | 11.14    | 15.76    | 14.21    | 17.71    | 17.19    |
| Fe <sub>2</sub> O <sub>3</sub> | 12.03    | 4.54     | 7.70     | 9.37     | 9.34     |
| MnO                            | .10      | .17      | .15      | .12      | .13      |
| MgO                            | 1.07     | 3.54     | 3.51     | 5.10     | 5.04     |
| CaO                            | .82      | 1.41     | 2.70     | 2.75     | 2.31     |
| Na <sub>2</sub> O              | 4.21     | 2.60     | 2.78     | 3.39     | 2.74     |
| K <sub>2</sub> O               | 1.31     | 3.02     | 3.16     | 3.36     | 4.54     |
| LOI                            | 7.40     | 4.30     | 4.70     | 5.90     | 5.20     |
| TOTAL                          | 99.20    | 99.46    | 98.39    | 99.60    | 98.82    |

TRACE ELEMENTS (PPH)

|                   |     |     |     |     |     |
|-------------------|-----|-----|-----|-----|-----|
| Zr                | 50  | 42  | 31  | 28  | 27  |
| Sr                | 98  | 62  | 79  | 97  | 97  |
| Rb                | 23  | 30  | 25  | 31  | 33  |
| Zn                | 24  | 24  | 35  | 29  | 35  |
| Cu                | 43  | 13  | 36  | 56  | 55  |
| Ba                | 71  | 111 | 201 | 201 | 245 |
| CuS               | 23  | 11  | 44  | 68  | 70  |
| PbS               | 40  | 13  | 4   | 9   | 8   |
| ZnS               | 20  | 12  | 17  | 13  | 24  |
| BaSO <sub>4</sub> |     | 0   |     | 0   |     |
| Pb                | 46  | 3   | 1   | 2   | 3   |
| Ni                | 4   | 2   | 4   | 5   | 10  |
| Co                | 26  | 6   | 8   | 10  | 13  |
| Cr                | 36  | 32  | 27  | 32  | 43  |
| V                 | 29  | 58  | 161 | 260 | 215 |
| Hg                | 191 | 12  | 32  | 36  | 22  |
| Ag                | 25  | 0   | 1   | 0   | 0   |
| Nb                |     |     |     |     |     |



PROMINENT QUARTZ SEQUENCE - ALTERED MAFIC FRAGMENTALS

MAJOR ELEMENTS (WT%)

| SAMPLE                         | 26661475 | 26661625 | 26662250 | 26662350 |
|--------------------------------|----------|----------|----------|----------|
| SiO <sub>2</sub>               | 53.30    | 41.10    | 40.30    | 19.60    |
| TiO <sub>2</sub>               | .32      | .36      | .28      | .24      |
| Al <sub>2</sub> O <sub>3</sub> | 12.55    | 13.82    | 10.00    | 7.60     |
| Fe <sub>2</sub> O <sub>3</sub> | 7.32     | 6.96     | 23.10    | 37.30    |
| MnO                            | .09      | .20      | .08      | .08      |
| HgO                            | 6.15     | 5.11     | 8.51     | 9.21     |
| CaO                            | 10.18    | 15.62    | .56      | .65      |
| Na <sub>2</sub> O              | 1.64     | 4.26     | 1.78     | .40      |
| K <sub>2</sub> O               | .91      | .51      | 1.34     | .45      |
| LOI                            | 7.30     | 11.20    | 14.20    | 21.80    |
| TOTAL                          | 99.76    | 99.22    | 100.15   | 97.41    |

TRACE ELEMENTS (PPM)

|                   |     |     |     |      |
|-------------------|-----|-----|-----|------|
| Zr                | 21  | 30  | 49  | 55   |
| Sr                | 229 | 222 | 60  | 70   |
| Rb                | 23  | 16  | 19  | 23   |
| Zn                | 43  | 34  | 870 | 5600 |
| Cu                | 62  | 6   | 59  | 101  |
| Ba                | 112 | 78  | 94  | 83   |
| CuS               | 45  | 3   | 31  | 61   |
| PbS               | 7   | 8   | 167 | 1350 |
| ZnS               | 7   | 12  | 680 | 6300 |
| BaSO <sub>4</sub> |     | 0   | 0   |      |
| Pb                | 2   | 2   | 185 | 1390 |
| Ni                | 116 | 95  | 2   | 2    |
| Co                | 26  | 23  | 12  | 10   |
| Cr                | 277 | 317 | 36  | 72   |
| V                 | 166 | 151 | 57  | 32   |
| Hg                | 10  | 8   | 426 | 2516 |
| Ag                | 0   | 0   | 21  | 122  |
| Nb                |     |     |     |      |

## INTERMEDIATE FOOTWALL - LUCKY STRIKE AREA

## MAJOR ELEMENTS (WT%)

| SAMPLE | 02020650 | 02020700 | 02020750 | 02020800 | 02020875 |
|--------|----------|----------|----------|----------|----------|
| SiO2   | 56.70    | 52.60    | 56.70    | 52.50    | 56.80    |
| TiO2   | .55      | .29      | .40      | .49      | .35      |
| Al2O3  | 14.99    | 15.11    | 14.45    | 16.87    | 13.75    |
| Fe2O3  | 7.81     | 11.05    | 8.01     | 8.58     | 9.90     |
| MnO    | .45      | .44      | .46      | .64      | .56      |
| MgO    | 8.69     | 9.36     | 8.57     | 9.48     | 8.45     |
| CaO    | .35      | .29      | .65      | .27      | .69      |
| Na2O   | .09      | 3.31     | 2.36     | 2.07     | 1.86     |
| K2O    | 1.64     | .03      | .59      | 1.27     | .41      |
| LOI    | 6.40     | 6.30     | 5.50     | 5.90     | 5.50     |
| TOTAL  | 97.67    | 98.78    | 97.69    | 98.07    | 98.27    |

## TRACE ELEMENTS (PPM)

|       |      |     |     |     |     |
|-------|------|-----|-----|-----|-----|
| Zr    | 46   | 33  | 28  | 33  | 22  |
| Sr    | 0    | 9   | 9   | 17  | 2   |
| Rb    | 42   | 11  | 18  | 30  | 15  |
| Zn    | 310  | 284 | 260 | 450 | 270 |
| Cu    | 25   | 5   | 117 | 260 | 55  |
| Ba    | 1037 | 135 | 282 | 594 | 331 |
| CuS   | 26   | 1   | 156 | 330 | 64  |
| PbS   | 21   | 17  | 27  | 15  | 30  |
| ZnS   | 49   | 35  | 44  | 74  | 42  |
| BaSO4 |      |     | 0   |     |     |
| Pb    | 34   | 36  | 41  | 32  | 51  |
| Ni    | 12   | 30  | 30  | 34  | 14  |
| Co    | 16   | 23  | 22  | 27  | 20  |
| Cr    | 70   | 112 | 91  | 100 | 69  |
| V     | 180  | 234 | 264 | 258 | 242 |
| Hg    |      | 25  | 27  | 29  | 19  |
| Ag    | 7    | 10  | 6   | 7   | 11  |
| Nb    |      |     |     |     |     |

## INTERMEDIATE FOOTWALL - LUCKY STRIKE AREA

## MAJOR ELEMENTS (WT%)

| SAMPLE | 02020950 | 02021050 | 02021150 | 07U2 100 | 542 150 |
|--------|----------|----------|----------|----------|---------|
| SiO2   | 56.70    | 45.20    | 55.00    | 61.40    | 54.10   |
| TiO2   | .31      | .40      | .40      | .11      | .42     |
| Al2O3  | 12.00    | 12.00    | 14.99    | 14.25    | 17.60   |
| Fe2O3  | 10.67    | 10.20    | 8.99     | 7.80     | 9.85    |
| MnO    | .53      | .72      | .58      | .45      | .67     |
| MgO    | 8.53     | 11.19    | 8.63     | 5.12     | 5.91    |
| CaO    | .39      | .68      | .72      | .23      | .39     |
| Na2O   | 1.89     | .25      | 2.50     | 2.71     | 3.18    |
| K2O    | .12      | .27      | .43      | 1.26     | 1.57    |
| LOI    | 6.00     | 10.20    | 5.60     | 4.40     | 5.10    |
| TOTAL  | 97.14    | 98.87    | 97.84    | 97.73    | 98.79   |

## TRACE ELEMENTS (PPM)

|       |     |     |     |     |     |
|-------|-----|-----|-----|-----|-----|
| Zr    | 28  | 40  | 36  | 93  | 85  |
| Sr    | 6   | 19  | 10  | 45  | 65  |
| Rb    | 12  | 18  | 19  | 28  | 32  |
| Zn    | 294 | 740 | 295 | 280 | 280 |
| Cu    | 7   | 100 | 14  | 13  | 5   |
| Ba    | 143 | 212 | 196 | 652 | 856 |
| CuS   | 2   | 119 | 12  | 1   | 1   |
| PbS   | 17  | 120 | 16  | 100 | 22  |
| ZnS   | 32  | 355 | 28  | 48  | 26  |
| BaSO4 | 0   | 0   | 0   | 0   | 4   |
| Pb    | 33  | 120 | 36  | 122 | 30  |
| Ni    | 13  | 11  | 15  | 12  | 12  |
| Co    | 23  | 29  | 24  | 17  | 17  |
| Cr    | 64  | 54  | 63  | 77  | 60  |
| V     | 268 | 274 | 300 | 132 | 230 |
| Hg    | 27  | 70  | 33  | 82  | 17  |
| Ag    | 10  | 15  | 10  | 10  | 10  |
| Nb    |     |     |     |     |     |

## INTERMEDIATE FOOTWALL - LUCKY STRIKE AREA

## MAJOR ELEMENTS (WT%)

| SAMPLE | 542 225 | 542 300 | 542 375 | 542 450 | 542 500 |
|--------|---------|---------|---------|---------|---------|
| S102   | 55.00   | 37.80   | 74.10   | 53.80   | 51.00   |
| T102   | .59     | .40     | .15     | .62     | .77     |
| Al2O3  | 17.42   | 20.54   | 12.36   | 18.75   | 17.16   |
| Fe2O3  | 9.47    | 13.85   | 1.89    | 8.26    | 8.87    |
| MnO    | .20     | .60     | .04     | .31     | .16     |
| MgO    | 5.13    | 12.57   | .60     | 6.26    | 7.02    |
| CaO    | .25     | 1.09    | .27     | .76     | 1.48    |
| Na2O   | 1.05    | 2.73    | 1.02    | 2.05    | 1.28    |
| K2O    | 3.26    | .90     | 3.15    | 2.42    | 5.08    |
| LOI    | 6.00    | 8.10    | 2.80    | 5.50    | 5.20    |
| TOTAL  | 98.37   | 98.66   | 96.38   | 98.73   | 98.02   |

## TRACE ELEMENTS (PPM)

|       |      |     |      |      |      |
|-------|------|-----|------|------|------|
| Zr    | 40   | 55  | 119  | 40   | 50   |
| Sr    | 17   | 52  | 0    | 59   | 93   |
| Rb    | 62   | 23  | 58   | 39   | 53   |
| Zn    | 162  | 340 | 930  | 270  | 360  |
| Cu    | 13   | 5   | 48   | 40   | 127  |
| Ba    | 1101 | 470 | 1427 | 2943 | 3491 |
| CuS   | 5    | 1   | 44   | 35   | 102  |
| PbS   | 15   | 21  | 1085 | 210  | 86   |
| ZnS   | 12   | 28  | 695  | 123  | 59   |
| BaSO4 |      | 0   |      | 106  |      |
| Pb    | 24   | 34  | 1060 | 190  | 111  |
| Ni    | 23   | 19  | 1    | 17   | 14   |
| Co    | 35   | 26  | 3    | 25   | 30   |
| Cr    | 60   | 74  | 19   | 61   | 37   |
| V     | 290  | 420 | 16   | 260  | 330  |
| Hg    | 21   | 23  | 70   | 70   | 25   |
| Ag    | 8    | 11  | 10   | 16   | 15   |
| Nb    |      |     |      |      |      |

## INTERMEDIATE FOOTWALL - LUCKY STRIKE AREA

## MAJOR ELEMENTS (WT%)

| SAMPLE | 542   | 525    | 77921428 | 07921575 | 07921649 | 08681655 |
|--------|-------|--------|----------|----------|----------|----------|
| B102   | 37.30 | 48.40  | 59.50    | 54.10    | 49.50    |          |
| T102   | .06   | .76    | .56      | .50      | .90      |          |
| Al2O3  | 6.52  | 19.33  | 20.33    | 14.20    | 19.20    |          |
| Fe2O3  | 25.40 | 10.86  | 3.16     | 9.30     | 12.30    |          |
| MnO    | .24   | .20    | .05      | .47      | .17      |          |
| HgO    | 2.06  | 10.38  | 2.41     | 9.21     | 5.04     |          |
| CaO    | .61   | .97    | 1.60     | .43      | 3.76     |          |
| Na2O   | 0.00  | .64    | 1.77     | .37      | 1.47     |          |
| K2O    | .24   | 2.94   | 5.19     | 1.32     | 4.72     |          |
| LOI    | 15.80 | 6.70   | 5.00     | 5.90     | 4.30     |          |
| TOTAL  | 93.23 | 101.18 | 99.57    | 95.88    | 101.36   |          |

## TRACE ELEMENTS (PPM)

|       |      |      |      |     |      |
|-------|------|------|------|-----|------|
| Zr    | 45   | 48   | 197  | 40  | 62   |
| Sr    | 92   | 103  | 45   | 0   | 239  |
| Rb    | 23   | 38   | 71   | 30  | 72   |
| Zn    | 162  | 82   | 24   | 61  | 105  |
| Cu    | 101  | 29   | 1    | 6   | 94   |
| Ba    | 7550 | 1381 | 1615 | 509 | 3917 |
| CuS   | 83   | 29   | 0    | 2   | 91   |
| PbS   | 27   | 13   | 10   | 9   | 7    |
| ZnS   | 42   | 24   | 9    | 22  | 23   |
| BaSO4 | 51   |      | 7    |     |      |
| Pb    | 44   | 14   | 11   | 12  | 19   |
| Ni    | 8    | 15   | 1    | 4   | 16   |
| Co    | 38   | 18   | 2    | 5   | 28   |
| Cr    | 40   | 48   | 19   | 38  | 40   |
| V     | 310  | 490  | 17   | 175 | 470  |
| Hg    | 46   | 18   | 12   | 13  | 25   |
| Ag    | 23   | 0    | 6    | 0   | 1    |
| Nb    |      |      |      |     |      |

## INTERMEDIATE FOOTWALL - LUCKY STRIKE AREA

## MAJOR ELEMENTS (WT%)

| SAMPLE | 08681725 | 08681825 | 08681900 | 08681975 | 972 825 |
|--------|----------|----------|----------|----------|---------|
| SiO2   | 55.80    | 49.10    | 51.20    | 48.70    | 64.80   |
| TiO2   | .82      | 1.02     | .80      | .83      | .10     |
| Al2O3  | 15.80    | 18.80    | 17.30    | 17.40    | 15.13   |
| Fe2O3  | 10.80    | 13.49    | 11.64    | 13.87    | 5.08    |
| MnO    | .17      | .19      | .16      | .20      | .05     |
| MgO    | 4.58     | 5.21     | 4.97     | 6.34     | 3.95    |
| CaO    | 6.00     | 7.26     | 6.15     | 6.67     | .58     |
| Na2O   | 1.66     | 2.92     | 1.80     | 2.72     | 1.08    |
| K2O    | 2.33     | 1.28     | 2.03     | .73      | 4.08    |
| LOI    | 3.20     | 4.20     | 3.90     | 4.90     | 3.90    |
| TOTAL  | 101.24   | 103.47   | 99.95    | 102.36   | 98.75   |

## TRACE ELEMENTS (PPM)

|       |      |     |      |      |      |
|-------|------|-----|------|------|------|
| Zr    | 46   | 82  | 49   | 68   | 107  |
| Sr    | 274  | 285 | 346  | 266  | 60   |
| Rb    | 28   | 26  | 29   | 20   | 80   |
| Zn    | 103  | 115 | 104  | 106  | 105  |
| Cu    | 105  | 171 | 103  | 91   | 4    |
| Ba    | 1407 | 985 | 1951 | 5632 | 1109 |
| CuS   | 93   | 158 | 81   | 94   | 3    |
| PbS   | 16   | 9   | 7    | 5    | 15   |
| ZnS   | 28   | 27  | 27   | 24   | 24   |
| BaSO4 | 0    | 0   | 0    | 0    | 0    |
| Pb    | 27   | 26  | 24   | 23   | 34   |
| Ni    | 13   | 15  | 14   | 14   | 8    |
| Co    | 30   | 31  | 30   | 33   | 18   |
| Cr    | 38   | 45  | 42   | 52   | 14   |
| V     | 380  | 350 | 340  | 360  | 15   |
| Hg    | 25   | 19  | 27   | 29   |      |
| Ag    | 0    | 1   | 0    | 1    |      |
| Nb    |      |     |      |      |      |

## INTERMEDIATE FOOTWALL - LUCKY STRIKE AREA

## MAJOR ELEMENTS (WT%)

| SAMPLE | 12141320 | 12141340 | 12141360 | 12141360 | 12141425 |
|--------|----------|----------|----------|----------|----------|
| SiO2   | 56.20    | 67.00    | 61.40    | 66.00    | 54.20    |
| TiO2   | .67      | .49      | .56      | .44      | .82      |
| Al2O3  | 21.10    | 16.44    | 17.74    | 15.43    | 15.60    |
| Fe2O3  | 6.74     | 4.29     | 4.61     | 4.02     | 11.52    |
| MnO    | .10      | .09      | .12      | .13      | .14      |
| HgO    | 3.00     | 1.79     | 2.09     | 1.50     | 4.83     |
| CaO    | 2.17     | 2.19     | 2.78     | 2.04     | 5.43     |
| Na2O   | 2.06     | 2.69     | 2.71     | 4.13     | 1.69     |
| K2O    | 4.82     | 3.04     | 3.12     | 1.92     | 1.57     |
| LOI    | 4.20     | 2.60     | 3.90     | 3.40     | 4.20     |
| TOTAL  | 101.06   | 100.62   | 99.03    | 99.01    | 100.00   |

## TRACE ELEMENTS (PPM)

|       |      |      |      |     |      |
|-------|------|------|------|-----|------|
| Zr    | 249  | 218  | 202  | 224 | 65   |
| Sr    | 213  | 367  | 433  | 426 | 243  |
| Rb    | 103  | 71   | 74   | 55  | 31   |
| Zn    | 86   | 72   | 79   | 79  | 110  |
| Cu    | 5    | 8    | 4    | 218 | 92   |
| Ba    | 1557 | 1138 | 1100 | 963 | 1116 |
| CuS   | 1    | 3    | 1    | 210 | 72   |
| PbS   | 9    | 12   | 7    | 67  | 16   |
| ZnS   | 18   | 18   | 20   | 17  | 30   |
| BaSO4 | 0    |      | 245  |     | 0    |
| Pb    | 15   | 18   | 15   | 55  | 24   |
| Ni    | 4    | 2    | 4    | 3   | 13   |
| Co    | 3    | 3    | 3    | 5   | 25   |
| Cr    | 21   | 24   | 34   | 26  | 30   |
| V     | 12   | 25   | 32   | 21  | 420  |
| Hg    | 28   | 50   | 29   | 308 | 52   |
| Ag    | 0    | 0    | 0    | 28  | 2    |
| Nb    |      |      |      |     |      |

## INTERMEDIATE FOOTWALL - LUCKY STRIKE AREA

## MAJOR ELEMENTS (WT%)

| SAMPLE | 19111158 | 20431126 | 26663625 | 26663725 | 26663775 |
|--------|----------|----------|----------|----------|----------|
| S102   | 64.80    | 66.00    | 50.80    | 49.70    | 51.20    |
| Ti02   | .58      | .58      | .82      | .80      | .77      |
| Al2O3  | 13.97    | 13.91    | 17.76    | 18.13    | 17.95    |
| Fe2O3  | 5.15     | 5.26     | 12.05    | 11.87    | 10.88    |
| MnO    | .09      | .13      | .08      | .12      | .16      |
| HgO    | .98      | 1.10     | 2.55     | 4.95     | 4.40     |
| CaO    | 4.42     | 4.85     | 3.47     | 2.97     | 6.97     |
| Na2O   | 4.21     | 3.43     | 2.00     | 1.55     | 2.37     |
| K2O    | 1.20     | .92      | 4.92     | 2.40     | 1.00     |
| LOI    | 4.80     | 2.40     | 5.10     | 6.50     | 3.00     |
| TOTAL  | 100.20   | 98.50    | 99.55    | 99.07    | 98.70    |

## TRACE ELEMENTS (PPM)

|       |     |     |     |     |     |
|-------|-----|-----|-----|-----|-----|
| Zr    | 112 | 107 | 70. | 71  | 64  |
| Sr    | 198 | 402 | 200 | 204 | 266 |
| Rb    | 21  | 19  | 90  | 55  | 21  |
| Zn    | 100 | 74  | 48  | 58  | 113 |
| Cu    | 14  | 2   | 15  | 103 | 127 |
| Ba    | 856 | 731 | 938 | 864 | 797 |
| CuS   | 13  | 5   | 6   | 89  | 85  |
| PbS   | 31  | 7   | 13  | 12  | 8   |
| ZnS   | 31  | 25  | 9   | 22  | 24  |
| BaSO4 |     | 0   |     | 0   |     |
| Pb    | 18  | 29  | 9   | 8   | 19  |
| Ni    | 6   | 3   | 7   | 13  | 15  |
| Co    | 15  | 17  | 11  | 23  | 33  |
| Cr    | 0   | 10  | 43  | 41  | 42  |
| V     | 50  | 45  | 820 | 574 | 512 |
| Hg    |     |     | 8   | 10  | 15  |
| Ag    |     |     | 0   | 0   | 0   |
| Nb    |     | 5   |     |     |     |



## INTERMEDIATE FOOTWALL - LUCKY STRIKE AREA

## MAJOR ELEMENTS (WT%)

SAMPLE 27040975

|       |       |
|-------|-------|
| SiO2  | 58.70 |
| TiO2  | .60   |
| Al2O3 | 15.90 |
| Fe2O3 | 8.95  |
| MnO   | .10   |
| MgO   | 2.41  |
| CaO   | 4.23  |
| Na2O  | 2.50  |
| K2O   | 2.42  |
| LOI   | 2.70  |

|       |       |
|-------|-------|
| TOTAL | 98.51 |
|-------|-------|

## TRACE ELEMENTS (PPM)

|       |      |
|-------|------|
| Zr    | 70   |
| Sr    | 256  |
| Rb    | 63   |
| Zn    | 46   |
| Cu    | 42   |
| Ba    | 1075 |
| CuS   | 37   |
| PbS   | 4    |
| ZnS   | 18   |
| BaSO4 | 24   |
| Pb    | 15   |
| Ni    | 10   |
| Co    | 10   |
| Cr    | 37   |
| V     | 301  |
| Hg    | 22   |
| Ag    | 0    |
| Nb    |      |

## INTERMEDIATE FOOTWALL - VARIABLY MINERALIZED AND ALTERED

## MAJOR ELEMENTS (WT%)

| SAMPLE | 01750275 | 01750350 | 01750417 | 02020200 | 02020250 |
|--------|----------|----------|----------|----------|----------|
| SiO2   | 71.90    | 59.60    | 73.10    | 29.10    | 16.70    |
| TiO2   | .20      | .77      | .17      | .37      | .01      |
| Al2O3  | 7.91     | 17.21    | 11.86    | 15.05    | 1.15     |
| Fe2O3  | 6.85     | 9.29     | 1.19     | 12.00    | 42.00    |
| MnO    | .07      | .44      | .04      | .52      | .30      |
| MgO    | 1.81     | 1.07     | 1.51     | 23.30    | 1.02     |
| CaO    | .20      | 5.60     | 1.04     | .39      | 8.78     |
| Na2O   | .05      | .57      | 3.39     | .01      | 0.00     |
| K2O    | 1.99     | 3.50     | 2.50     | 0.00     | .01      |
| LOI    | 6.00     | 8.30     | 1.90     | 12.20    | 25.90    |
| TOTAL  | 97.06    | 98.15    | 96.70    | 92.94    | 95.95    |

## TRACE ELEMENTS (PPM)

|       |      |      |      |      |      |
|-------|------|------|------|------|------|
| Zr    | 25   | 63   | 152  | 60   | 53   |
| Sr    | 20   | 60   | 373  | 39   | 124  |
| Rb    | 48   | 57   | 13   | 25   | 20   |
| Zn    | 9999 | 9999 | 505  | 9999 | 890  |
| Cu    | 2820 | 1790 | 76   | 2225 | 109  |
| Ba    | 3104 | 2319 | 311  | 191  | 9999 |
| CuS   | 1540 | 1350 | 84   | 1540 | 84   |
| PbS   | 9999 | 3650 | 4500 | 999  | 1340 |
| ZnS   | 9999 | 9999 | 200  | 9999 | 650  |
| BaSO4 |      | 6750 |      |      | 999  |
| Pb    | 999  | 2500 | 2300 | 9999 | 795  |
| Ni    | 9    | 6    | 9    | 60   | 3    |
| Co    | 14   | 13   | 17   | 43   | 14   |
| Cr    | 69   | 35   | 61   | 351  | 54   |
| V     | 61   | 98   | 173  | 270  | 16   |
| Hg    | 278  | 144  | 36   | 1078 | 278  |
| Ag    | 40   | 28   | 46   | 24   | 125  |
| Nb    |      |      |      |      |      |

## INTERMEDIATE FOOTWALL - VARIABLY MINERALIZED AND ALTERED

## MAJOR ELEMENTS (WT%)

| SAMPLE | 02020300 | 02020350 | 02020400 | 02020450 | 02020500 |
|--------|----------|----------|----------|----------|----------|
| SiO2   | 16.00    | 26.30    | 48.70    | 53.40    | 53.30    |
| TiO2   | .16      | .14      | 0.00     | .26      | .20      |
| Al2O3  | 8.55     | 6.91     | 3.00     | 7.91     | 10.90    |
| Fe2O3  | 42.30    | 31.60    | 23.70    | 17.19    | 12.31    |
| MnO    | .37      | .20      | .22      | .39      | .01      |
| HgO    | 10.86    | 10.95    | 3.57     | 7.47     | 14.14    |
| CaO    | .10      | .81      | 1.98     | 1.27     | .35      |
| Na2O   | 0.00     | 0.00     | 0.00     | 0.00     | .02      |
| K2O    | 0.00     | 0.00     | .03      | .30      | .01      |
| LOI    | 21.40    | 19.30    | 14.40    | 10.90    | 8.15     |
| TOTAL  | 99.74    | 96.29    | 95.60    | 99.17    | 100.19   |

## TRACE ELEMENTS (PPH)

|       |      |      |      |      |     |
|-------|------|------|------|------|-----|
| Zr    | 51   | 66   | 51   | 52   | 5   |
| Sr    | 61   | 61   | 48   | 20   | 0   |
| Rb    | 20   | 28   | 23   | 19   | 12  |
| Zn    | 280  | 4800 | 9999 | 320  | 590 |
| Cu    | 6650 | 2200 | 4000 | 1360 | 20  |
| Ba    | 189  | 341  | 130  | 320  | 123 |
| CuS   | 7600 | 1900 | 3100 | 1570 | 15  |
| PbS   | 57   | 1100 | 8900 | 390  | 67  |
| ZnS   | 55   | 2500 | 999  | 71   | 72  |
| BaSO4 |      |      | 47   |      |     |
| Pb    | 99   | 8450 | 9500 | 305  | 80  |
| Ni    | 14   | 3    | 6    | 3    | 136 |
| Co    | 17   | 17   | 22   | 20   | 31  |
| Cr    | 99   | 62   | 43   | 30   | 464 |
| V     | 156  | 118  | 60   | 116  | 266 |
| Hg    | 134  | 154  | 412  | 96   | 25  |
| Ag    | 39   | 88   | 67   | 22   | 11  |
| Nb    |      |      |      |      |     |

## INTERMEDIATE FOOTWALL - VARIABLY MINERALIZED AND ALTERED

## MAJOR ELEMENTS (WT%)

| SAMPLE | 02020550 | 02020600 | 0542  | 0542 50 | 27853355 |
|--------|----------|----------|-------|---------|----------|
| SiO2   | 48.00    | 56.00    | 56.20 | 61.10   | 42.30    |
| TiO2   | .06      | .38      | .02   | .08     | .90      |
| Al2O3  | 7.65     | 11.37    | 5.05  | 5.90    | 15.20    |
| Fe2O3  | 10.18    | 10.29    | 12.59 | 8.76    | 9.58     |
| MnO    | .71      | .47      | .13   | .07     | .17      |
| KgO    | 9.31     | 9.44     | 3.64  | 1.50    | 7.24     |
| CaO    | 6.45     | 1.79     | .50   | .02     | 12.37    |
| Na2O   | 0.00     | .01      | .01   | .03     | 2.69     |
| K2O    | .01      | .67      | .62   | 1.30    | .84      |
| LOI    | 11.60    | 7.80     | 9.30  | 6.90    | 7.90     |
| TOTAL  | 93.97    | 98.22    | 88.06 | 85.66   | 99.19    |

## TRACE ELEMENTS (PPM)

|       |      |     |      |      |     |
|-------|------|-----|------|------|-----|
| Zr    | 35   | 40  | 67   | 98   | 74  |
| Sr    | 28   | 0   | 40   | 63   | 231 |
| Rb    | 19   | 22  | 36   | 51   | 15  |
| Zn    | 999  | 710 | 2900 | 9999 | 65  |
| Cu    | 485  | 19  | 250  | 1360 | 56  |
| Ba    | 87   | 599 | 366  | 4405 | 188 |
| CuS   | 950  | 17  | 350  | 1100 | 42  |
| PbS   | 7150 | 400 | 1090 | 9999 | 14  |
| ZnS   | 360  | 270 | 3500 | 999  | 18  |
| BaSO4 | 20   |     | 2600 |      | 0   |
| Pb    | 2450 | 243 | 748  | 9999 | 27  |
| Ni    | 72   | 24  | 15   | 3    | 85  |
| Co    | 21   | 19  | 14   | 4    | 32  |
| Cr    | 266  | 117 | 34   | 27   | 339 |
| V     | 184  | 194 | 148  | 44   | 150 |
| Hg    | 322  | 25  | 77   | 211  |     |
| Ag    | 23   | 14  | 16   | 32   |     |
| Nb    |      |     |      |      | 8   |

## SKI HILL SEQUENCE FLOWS

## MAJOR ELEMENTS (WTZ)

| SAMPLE | JGT75 41 | JGT75 43 | JGT75170 | 08680250 | 08680525 |
|--------|----------|----------|----------|----------|----------|
| S102   | 58.90    | 61.20    | 53.60    | 61.40    | 51.30    |
| T102   | .73      | .77      | .80      | .91      | .58      |
| Al2O3  | 15.81    | 13.90    | 16.46    | 13.50    | 16.70    |
| Fe2O3  | 7.80     | 8.02     | 11.44    | 11.00    | 10.00    |
| MnO    | .17      | .20      | .16      | .21      | .18      |
| HgO    | 2.49     | 2.40     | 3.61     | 3.05     | 6.32     |
| CaO    | 6.50     | 6.82     | 7.18     | 4.68     | 9.79     |
| Na2O   | 4.03     | 3.12     | 3.13     | 3.27     | 2.42     |
| K2O    | .47      | .57      | 1.61     | 1.22     | .65      |
| LOI    | 1.00     | .90      | 1.30     | 2.40     | 2.60     |
| TOTAL  | 97.90    | 97.90    | 99.29    | 101.64   | 100.62   |

## TRACE ELEMENTS (PPH)

|       |     |     |      |      |     |
|-------|-----|-----|------|------|-----|
| Zr    | 81  | 91  | 73   | 47   | 31  |
| Sr    | 305 | 334 | 228  | 140  | 255 |
| Rb    | 20  | 16  | 24   | 18   | 17  |
| Zn    | 57  | 73  | 65   | 124  | 87  |
| Cu    | 91  | 23  | 136  | 88   | 62  |
| Ba    | 363 | 351 | 1546 | 1465 | 398 |
| CuS   | 75  | 21  | 125  | 78   | 47  |
| PbS   | 12  | 16  | 19   | 15   | 10  |
| ZnS   | 12  | 21  | 19   | 25   | 9   |
| BaSO4 | 0   | 0   | 0    |      |     |
| Pb    | 22  | 19  | 28   | 25   | 24  |
| Ni    | 4   | 3   | 8    | 8    | 33  |
| Co    | 35  | 34  | 19   | 36   | 30  |
| Cr    | 23  | 16  | 19   | 36   | 124 |
| V     | 145 | 145 | 215  | 340  | 310 |
| Hg    |     |     |      | 27   | 19  |
| Ag    |     |     |      | 0    | 0   |
| Nb    | 4   | 0   | 10   |      |     |

## SKI HILL SEQUENCE FLOWS

## MAJOR ELEMENTS (WT%)

| SAMPLE | 88692188 | 14930150 | 27851517 | 28371950 | 28372575 |
|--------|----------|----------|----------|----------|----------|
| SiO2   | 46.80    | 48.80    | 51.60    | 47.10    | 53.50    |
| TiO2   | .80      | .83      | .48      | .88      | .65      |
| Al2O3  | 15.10    | 15.95    | 16.59    | 15.39    | 17.48    |
| Fe2O3  | 10.42    | 10.19    | 9.81     | 9.31     | 8.63     |
| MnO    | .15      | .17      | .16      | .22      | .18      |
| HgO    | 9.37     | 6.20     | 4.97     | 4.81     | 2.10     |
| CaO    | 9.32     | 11.58    | 9.56     | 9.83     | 10.10    |
| Na2O   | 1.57     | 2.44     | 2.69     | 2.30     | 2.09     |
| K2O    | .25      | .49      | 1.07     | .62      | 1.07     |
| LOI    | 4.90     | 5.10     | 3.60     | 4.60     | 2.90     |
| TOTAL  | 98.68    | 100.95   | 99.73    | 95.06    | 98.70    |

## TRACE ELEMENTS (PPM)

|       |     |     |     |     |     |
|-------|-----|-----|-----|-----|-----|
| Zr    | 109 | 98  | 71  | 86  | 89  |
| Sr    | 345 | 432 | 39  | 275 | 401 |
| Rb    | 17  | 17  | 15  | 15  | 22  |
| Zn    | 83  | 64  | 62  | 45  | 42  |
| Cu    | 58  | 56  | 73  | 63  | 60  |
| Ba    | 213 | 271 | 538 | 384 | 710 |
| CuS   |     | 36  | 67  | 52  | 43  |
| PbS   |     | 5   | 11  | 14  | 14  |
| ZnS   |     | 6   | 19  | 9   | 16  |
| BaSO4 |     | 0   | 0   | 0   | 0   |
| Pb    | 23  | 15  | 30  | 16  | 25  |
| Ni    | 79  | 69  | 34  | 92  | 12  |
| Co    | 39  | 35  | 27  | 31  | 23  |
| Cr    | 294 | 168 | 181 | 355 | 27  |
| V     | 275 | 368 | 230 | 150 | 150 |
| Hg    | 50  | 14  |     |     |     |
| Ag    | 0   | 2   |     |     |     |
| Nb    |     |     | 11  | 4   | 9   |

## SKI HILL SEQUENCE FLOWS

## MAJOR ELEMENTS (WT%)

| SAMPLE                         | 2841 50 | 2841 150 | 2851 283 | 2851 876 | 1925 100 |
|--------------------------------|---------|----------|----------|----------|----------|
| SiO <sub>2</sub>               | 70.10   | 65.50    | 65.90    | 58.00    | 61.30    |
| TiO <sub>2</sub>               | .66     | .66      | .54      | .72      | .80      |
| Al <sub>2</sub> O <sub>3</sub> | 13.12   | 14.03    | 14.40    | 12.52    | 16.25    |
| Fe <sub>2</sub> O <sub>3</sub> | 5.07    | 6.30     | 4.89     | 9.53     | 6.19     |
| MnO                            | .09     | .11      | .07      | .18      | .23      |
| HgO                            | .96     | 2.27     | 1.71     | 2.62     | 1.85     |
| CaO                            | 3.75    | 2.70     | 2.17     | 7.86     | 2.23     |
| Na <sub>2</sub> O              | 3.74    | 3.91     | 5.60     | 4.90     | 5.93     |
| K <sub>2</sub> O               | .58     | 1.63     | 1.56     | .09      | .71      |
| LOI                            | 1.00    | 2.20     | 2.50     | 1.80     | 2.80     |
| TOTAL                          | 99.07   | 99.39    | 99.50    | 98.22    | 98.29    |

## TRACE ELEMENTS (PPM)

|                   |     |     |     |     |     |
|-------------------|-----|-----|-----|-----|-----|
| Zr                | 110 | 111 | 86  | 44  | 155 |
| Sr                | 316 | 27  | 214 | 97  | 203 |
| Rb                | 10  | 22  | 16  | 5   | 16  |
| Zn                | 36  | 71  | 63  | 72  | 121 |
| Cu                | 1   | 4   | 7   | 92  | 2   |
| Ba                | 396 | 763 | 976 | 741 | 600 |
| CuS               | 1   | 6   | 9   | 87  | 2   |
| PbS               | 5   | 25  | 19  | 16  | 8   |
| ZnS               | 19  | 37  | 18  | 33  | 19  |
| BaSO <sub>4</sub> |     |     | 0   | 0   | 0   |
| Pb                | 14  | 18  | 18  | 23  | 10  |
| Ni                | 5   | 10  | 7   | 5   | 3   |
| Co                | 31  | 24  | 21  | 27  | 26  |
| Cr                | 18  |     | 9   | 18  | 17  |
| V                 | 90  | 65  | 60  | 165 | 40  |
| Hg                |     |     |     |     |     |
| Ag                |     |     |     |     |     |
| Nb                |     |     | 6   | 4   | 8   |

## SKI HILL SEQUENCE FLOWS

## MAJOR ELEMENTS (WT%)

SAMPLE 1925 140

|                                |       |
|--------------------------------|-------|
| SiO <sub>2</sub>               | 61.80 |
| TiO <sub>2</sub>               | .80   |
| Al <sub>2</sub> O <sub>3</sub> | 15.75 |
| Fe <sub>2</sub> O <sub>3</sub> | 7.07  |
| MnO                            | .17   |
| HgO                            | 2.20  |
| CaO                            | 2.18  |
| Na <sub>2</sub> O              | 5.37  |
| K <sub>2</sub> O               | .30   |
| LOI                            | 2.60  |

|       |       |
|-------|-------|
| TOTAL | 98.24 |
|-------|-------|

## TRACE ELEMENTS (PPM)

|                   |     |
|-------------------|-----|
| Zr                | 154 |
| Sr                | 291 |
| Rb                | 24  |
| Zn                | 134 |
| Cu                | 2   |
| Ba                | 288 |
| CuS               | 33  |
| PbS               | 6   |
| ZnS               | 25  |
| BaSO <sub>4</sub> | 0   |
| Pb                | 14  |
| Ni                | 5   |
| Co                | 22  |
| Cr                | 16  |
| V                 | 55  |
| Hg                |     |
| Ag                |     |
| Nb                | 5   |



## SKI HILL SEQUENCE FRAGMENTALS

## MAJOR ELEMENTS (WTX)

| SAMPLE | JGT75 40 | JGT75 42 | 07920050 | 07920150 | 08680150 |
|--------|----------|----------|----------|----------|----------|
| SiO2   | 59.50    | 60.20    | 50.40    | 45.00    | 62.50    |
| TiO2   | .78      | .77      | .79      | .75      | .56      |
| Al2O3  | 15.41    | 15.01    | 17.54    | 16.92    | 13.80    |
| Fe2O3  | 7.92     | 8.01     | 11.48    | 11.13    | 7.25     |
| MnO    | .15      | .15      | .19      | .21      | .21      |
| MgO    | 2.57     | 2.44     | 3.72     | 4.02     | 2.81     |
| CaO    | 4.99     | 4.69     | 6.12     | 7.24     | 5.22     |
| Na2O   | 2.73     | 2.78     | 2.87     | 3.81     | 2.47     |
| K2O    | 1.88     | 1.90     | 2.08     | .83      | 2.12     |
| LOI    | 2.40     | 2.30     | 4.50     | 6.60     | 2.60     |
| TOTAL  | 98.33    | 98.25    | 99.69    | 96.51    | 99.54    |

## TRACE ELEMENTS (PPM)

|       |      |      |      |      |      |
|-------|------|------|------|------|------|
| Zr    | 110  | 105  | 76   | 81   | 83   |
| Sr    | 320  | 321  | 284  | 288  | 340  |
| Rb    | 47   | 36   | 35   | 24   | 39   |
| Zn    | 69   | 74   | 111  | 119  | 85   |
| Cu    | 41   | 39   | 90   | 94   | 48   |
| Ba    | 1722 | 1697 | 1860 | 2238 | 9999 |
| CuS   | 34   | 28   | 101  | 105  | 31   |
| PbS   | 17   | 14   | 26   | 17   | 17   |
| ZnS   | 17   | 8    | 42   | 27   | 21   |
| BaSO4 | 0    | 0    | 3    |      | 0    |
| Pb    | 31   | 25   | 31   | 33   | 29   |
| Ni    | 3    | 8    | 12   | 12   | 14   |
| Co    | 40   | 32   | 28   | 30   | 19   |
| Cr    | 17   | 16   | 46   | 44   | 63   |
| V     | 110  | 115  | 370  | 380  | 185  |
| Hg    |      |      | 31   | 15   | 54   |
| Ag    |      |      | 0    | 1    | 1    |
| Nb    | 9    | 7    |      |      |      |

## SKI HILL SEQUENCE FRAGMENTALS

## MAJOR ELEMENTS (WTX)

| SAMPLE | 08680325 | 08680425 | 08680625 | 08680725 | 08680825 |
|--------|----------|----------|----------|----------|----------|
| SiO2   | 52.30    | 50.40    | 52.70    | 50.60    | 63.40    |
| TiO2   | .74      | .74      | .60      | .63      | .63      |
| Al2O3  | 16.90    | 17.00    | 15.40    | 16.00    | 13.50    |
| Fe2O3  | 11.00    | 11.09    | 10.08    | 9.00     | 9.63     |
| MnO    | .16      | .17      | .20      | .16      | .18      |
| HgO    | 4.00     | 4.58     | 4.74     | 2.00     | 2.75     |
| CaO    | 7.29     | 7.84     | 8.40     | 10.04    | 5.22     |
| Na2O   | 2.15     | 2.15     | 2.08     | 4.09     | 2.78     |
| K2O    | 1.43     | 1.12     | .52      | .50      | .88      |
| LOI    | 4.40     | 4.10     | 5.50     | 7.10     | 2.10     |
| TOTAL  | 100.37   | 99.19    | 100.22   | 100.92   | 101.07   |

## TRACE ELEMENTS (PPM)

|       |     |     |     |     |     |
|-------|-----|-----|-----|-----|-----|
| Zr    | 70  | 64  | 83  | 70  | 80  |
| Sr    | 305 | 334 | 357 | 298 | 323 |
| Rb    | 18  | 23  | 31  | 18  | 22  |
| Zn    | 111 | 106 | 112 | 85  | 85  |
| Cu    | 82  | 59  | 52  | 71  | 65  |
| Ba    | 372 | 837 | 932 | 312 | 991 |
| CuS   | 71  | 45  | 43  | 75  | 57  |
| PbS   | 10  | 10  | 24  | 16  | 14  |
| ZnS   | 23  | 16  | 21  | 23  | 18  |
| BaSO4 |     | 0   | 0   |     | 0   |
| Pb    | 26  | 26  | 40  | 25  | 23  |
| Ni    | 14  | 13  | 10  | 8   | 5   |
| Co    | 27  | 26  | 22  | 22  | 20  |
| Cr    | 38  | 50  | 43  | 37  | 34  |
| V     | 335 | 300 | 275 | 300 | 265 |
| Hg    | 23  | 25  | 36  | 34  | 29  |
| Ag    | 0   | 0   | 1   | 1   | 0   |
| Nb    |     |     |     |     |     |

## SKI HILL SEQUENCE FRAGMENTALS

## MAJOR ELEMENTS (WT%)

| SAMPLE                         | 08680925 | 08681125 | 08681225 | 08681325 | 08681425 |
|--------------------------------|----------|----------|----------|----------|----------|
| SiO <sub>2</sub>               | 43.90    | 47.70    | 61.50    | 49.20    | 52.50    |
| TiO <sub>2</sub>               | .75      | .72      | .65      | .75      | .60      |
| Al <sub>2</sub> O <sub>3</sub> | 19.80    | 19.60    | 15.40    | 18.00    | 16.00    |
| Fe <sub>2</sub> O <sub>3</sub> | 13.35    | 12.15    | 7.61     | 11.90    | 8.92     |
| MnO                            | .21      | .18      | .15      | .15      | .12      |
| HgO                            | 6.61     | 6.27     | 2.21     | 7.21     | 4.74     |
| CaO                            | 9.00     | 9.24     | 4.68     | 6.37     | 7.53     |
| Na <sub>2</sub> O              | 1.70     | 1.73     | 2.39     | 1.76     | 2.43     |
| K <sub>2</sub> O               | 1.51     | 1.24     | 2.03     | 1.22     | .93      |
| LOI                            | 4.90     | 3.40     | 5.10     | 4.30     | 6.50     |
| TOTAL                          | 101.73   | 102.23   | 101.72   | 100.86   | 100.27   |

## TRACE ELEMENTS (PPM)

|                   |      |     |     |      |     |
|-------------------|------|-----|-----|------|-----|
| Zr                | 87   | 60  | 88  | 77   | 75  |
| Sr                | 393  | 341 | 252 | 314  | 393 |
| Rb                | 31   | 20  | 46  | 28   | 27  |
| Zn                | 150  | 101 | 113 | 104  | 87  |
| Cu                | 73   | 52  | 18  | 70   | 81  |
| Ba                | 1116 | 863 | 965 | 1056 | 580 |
| CuS               | 71   | 36  | 18  | 61   | 84  |
| PbS               | 13   | 4   | 10  | 19   | 20  |
| ZnS               | 32   | 18  | 20  | 57   | 27  |
| BaSO <sub>4</sub> |      | 0   |     | 0    |     |
| Pb                | 30   | 21  | 21  | 25   | 31  |
| Ni                | 18   | 22  | 3   | 22   | 32  |
| Co                | 31   | 33  | 12  | 20   | 27  |
| Cr                | 56   | 67  | 26  | 84   | 75  |
| V                 | 300  | 300 | 59  | 350  | 200 |
| Hg                | 20   | 17  | 13  | 31   | 27  |
| Ag                | 0    | 0   | 1   | 1    | 0   |
| Nb                |      |     |     |      |     |

## SKI HILL SEQUENCE FRAGMENTALS

## MAJOR ELEMENTS (WT%)

| SAMPLE                         | 08681450 | 08681475 | 08681525 | 08690075 | 08690260 |
|--------------------------------|----------|----------|----------|----------|----------|
| SiO <sub>2</sub>               | 74.40    | 65.70    | 65.00    | 54.50    | 51.30    |
| TiO <sub>2</sub>               | .23      | .46      | .50      | .77      | .81      |
| Al <sub>2</sub> O <sub>3</sub> | 10.70    | 16.60    | 12.90    | 27.21    | 17.90    |
| Fe <sub>2</sub> O <sub>3</sub> | 4.17     | 4.50     | 5.34     | 10.15    | 11.07    |
| MnO                            | .12      | .13      | .17      | .16      | .17      |
| MgO                            | .64      | 1.50     | .79      | 4.20     | 4.57     |
| CaO                            | 3.90     | 4.10     | 8.37     | 6.75     | 6.64     |
| Na <sub>2</sub> O              | 2.42     | 2.29     | 2.82     | 2.64     | 2.06     |
| K <sub>2</sub> O               | 1.44     | 3.16     | .42      | 1.36     | 1.30     |
| LOI                            | 3.00     | 2.70     | 2.50     | 1.00     | 4.10     |
| TOTAL                          | 101.02   | 101.22   | 98.87    | 98.82    | 100.72   |

## TRACE ELEMENTS (PPM)

|                   |     |     |     |      |     |
|-------------------|-----|-----|-----|------|-----|
| Zr                | 128 | 233 | 136 | 93   | 76  |
| Sr                | 244 | 422 | 443 | 430  | 39  |
| Rb                | 42  | 80  | 16  | 30   | 26  |
| Zn                | 10  | 50  | 55  | 96   | 105 |
| Cu                | 3   | 3   | 11  | 65   | 111 |
| Ba                | 375 | 854 | 224 | 1017 | 893 |
| CuS               | 0   | 0   | 6   | 52   | 97  |
| PbS               | 0   | 4   | 21  | 0    | 31  |
| ZnS               | 2   | 0   | 13  | 28   | 22  |
| BaSO <sub>4</sub> | 0   | 0   | 0   | 0    | 0   |
| Pb                | 19  | 27  | 34  | 19   | 36  |
| Ni                | 1   | 1   | 2   | 13   | 19  |
| Co                | 1   | 2   | 7   | 24   | 31  |
| Cr                | 21  | 21  | 37  | 46   | 43  |
| V                 | 5   | 31  | 68  | 275  | 305 |
| Hg                | 10  | 10  | 15  | 41   | 52  |
| Ag                | 0   | 0   | 0   | 2    | 1   |
| Nb                |     |     |     |      |     |

## SKI HILL SEQUENCE FRAGMENTALS

## MAJOR ELEMENTS (WT%)

| SAMPLE                         | 08690348 | 08690450 | 08690554 | 08690750 | 08690850 |
|--------------------------------|----------|----------|----------|----------|----------|
| SiO <sub>2</sub>               | 57.40    | 55.80    | 49.40    | 52.70    | 48.80    |
| TiO <sub>2</sub>               | .58      | .58      | .54      | .73      | .73      |
| Al <sub>2</sub> O <sub>3</sub> | 16.00    | 15.70    | 16.80    | 17.00    | 18.90    |
| Fe <sub>2</sub> O <sub>3</sub> | 6.85     | 9.74     | 10.30    | 10.34    | 11.60    |
| MnO                            | .18      | .20      | .23      | .15      | .15      |
| K <sub>2</sub> O               | 3.41     | 4.28     | 6.10     | 4.37     | 5.85     |
| CaO                            | 8.21     | 8.03     | 9.56     | 8.32     | 8.94     |
| Na <sub>2</sub> O              | 3.88     | 2.24     | 2.70     | 2.77     | 1.72     |
| K <sub>2</sub> O               | 1.17     | 1.10     | .66      | .64      | .59      |
| LOI                            | 2.90     | 2.40     | 3.10     | 4.60     | 4.20     |
| TOTAL                          | 100.58   | 100.07   | 99.47    | 101.62   | 101.48   |

## TRACE ELEMENTS (PPM)

|                   |     |     |     |     |     |
|-------------------|-----|-----|-----|-----|-----|
| Zr                | 72  | 79  | 78  | 81  | 72  |
| Sr                | 384 | 306 | 421 | 355 | 387 |
| Rb                | 31  | 28  | 25  | 22  | 21  |
| Zn                | 83  | 92  | 96  | 114 | 89  |
| Cu                | 72  | 62  | 114 | 59  | 73  |
| Ba                | 695 | 741 | 533 | 526 | 451 |
| CuS               | 60  | 51  | 120 | 54  | 57  |
| PbS               | 4   | 16  | 5   | 21  | 8   |
| ZnS               | 34  | 15  | 37  | 32  | 25  |
| BaSO <sub>4</sub> | 0   |     |     | 0   |     |
| Pb                | 16  | 20  | 20  | 26  | 20  |
| Ni                | 28  | 16  | 35  | 21  | 12  |
| Co                | 22  | 24  | 29  | 26  | 28  |
| Cr                | 115 | 74  | 122 | 58  | 39  |
| V                 | 230 | 326 | 315 | 315 | 405 |
| Hg                | 43  | 40  | 40  | 30  | 30  |
| Ag                | 6   | 4   | 5   | 0   | 0   |
| Nb                |     |     |     |     |     |

## SKI HILL SEQUENCE FRAGMENTALS

## MAJOR ELEMENTS (WT%)

| SAMPLE                         | 08690950 | 08691050 | 08691150 | 08691250 | 08691325 |
|--------------------------------|----------|----------|----------|----------|----------|
| SiO <sub>2</sub>               | 57.90    | 60.10    | 49.40    | 46.60    | 53.00    |
| TiO <sub>2</sub>               | .60      | .58      | .65      | .72      | .72      |
| Al <sub>2</sub> O <sub>3</sub> | 16.80    | 17.20    | 16.60    | 17.30    | 16.20    |
| Fe <sub>2</sub> O <sub>3</sub> | 8.59     | 8.13     | 10.68    | 12.57    | 8.70     |
| MnO                            | .15      | .15      | .12      | .11      | .22      |
| HgO                            | 3.18     | 3.44     | 7.17     | 7.50     | 2.39     |
| CaO                            | 3.33     | 5.21     | 5.59     | 9.25     | 8.60     |
| Na <sub>2</sub> O              | 7.24     | 2.92     | 1.59     | 2.34     | 5.00     |
| K <sub>2</sub> O               | .05      | 1.25     | .73      | .35      | .05      |
| LOI                            | 1.90     | 2.70     | 4.50     | 3.60     | 4.50     |
| TOTAL                          | 99.74    | 101.60   | 97.03    | 100.34   | 99.38    |

## TRACE ELEMENTS (PPM)

|                   |     |      |     |     |     |
|-------------------|-----|------|-----|-----|-----|
| Zr                | 70  | 125  | 91  | 65  | 53  |
| Sr                | 151 | 533  | 409 | 279 | 215 |
| Rb                | 12  | 27   | 26  | 17  | 2   |
| Zn                | 86  | 97   | 107 | 113 | 89  |
| Cu                | 24  | 42   | 53  | 55  | 72  |
| Ba                | 141 | 1065 | 511 | 340 | 717 |
| CuS               | 25  | 41   | 54  | 71  | 35  |
| PbS               | 18  | 64   | 12  | 10  | 0   |
| ZnS               | 24  | 23   | 30  | 20  | 02  |
| BaSO <sub>4</sub> |     |      |     | 0   | 7   |
| Pb                | 21  | 61   | 23  | 23  | 1   |
| Ni                | 6   | 5    | 20  | 28  | 9   |
| Co                | 19  | 17   | 26  | 33  | 02  |
| Cr                | 37  | 29   | 100 | 150 | 13  |
| V                 | 100 | 150  | 250 | 410 | 41  |
| Hg                | 37  | 41   | 41  | 49  | 3   |
| Ag                | 1   | 0    | 2   | 1   | 6   |
| Nb                |     |      |     |     | 0   |

## SKI HILL SEQUENCE FRAGMENTALS

## MAJOR ELEMENTS (WT%)

| SAMPLE                         | 08691350 | 12140105 | 1214 195 | 12140300 | 12140500 |
|--------------------------------|----------|----------|----------|----------|----------|
| SiO <sub>2</sub>               | 76.50    | 64.10    | 61.80    | 53.70    | 46.60    |
| TiO <sub>2</sub>               | .19      | .35      | .73      | .74      | .70      |
| Al <sub>2</sub> O <sub>3</sub> | 5.65     | 11.84    | 15.38    | 16.00    | 17.23    |
| Fe <sub>2</sub> O <sub>3</sub> | 5.74     | 4.83     | 5.21     | 9.97     | 11.46    |
| MnO                            | .21      | .12      | .00      | .21      | .18      |
| MgO                            | 1.55     | 1.51     | 1.38     | 3.18     | 4.37     |
| CaO                            | 2.56     | 3.86     | 4.35     | 6.58     | 7.78     |
| Na <sub>2</sub> O              | .29      | .52      | .23      | 3.08     | .78      |
| K <sub>2</sub> O               | 1.67     | 1.97     | 3.31     | 1.20     | 3.10     |
| LOI                            | 3.00     | 6.00     | 6.00     | 4.40     | 9.90     |
| TOTAL                          | 97.36    | 95.10    | 99.27    | 99.06    | 102.10   |

## TRACE ELEMENTS (PPM)

|                   |   |     |     |      |     |
|-------------------|---|-----|-----|------|-----|
| Zr                | 3 | 66  | 57  | 87   | 34  |
| Sr                |   | 39  | 5   | 365  | 200 |
| Rb                |   | 50  | 63  | 28   | 17  |
| Zn                |   | 101 | 74  | 124  | 121 |
| Cu                |   | 44  | 22  | 47   | 60  |
| Ba                |   | 823 | 958 | 1121 | 241 |
| CuS               |   | 32  | 17  | 42   | 58  |
| PbS               |   | 19  | 2   | 17   | 9   |
| ZnS               |   | 7   | 6   | 11   | 13  |
| BaSO <sub>4</sub> |   |     | 10  |      |     |
| Pb                |   | 45  | 25  | 51   | 35  |
| Ni                |   | 6   | 1   | 9    | 31  |
| Co                |   | 12  | 10  | 22   | 26  |
| Cr                |   | 35  | 22  | 47   | 36  |
| V                 |   | 106 | 138 | 254  | 226 |
| Hg                |   | 38  | 31  | 32   | 34  |
| Ag                |   | 7   | 5   | 12   | 10  |
| Nb                |   |     |     |      |     |

## SKI HILL SEQUENCE FRAGMENTALS

## MAJOR ELEMENTS (WT%)

| SAMPLE                         | 12140600 | 12140700 | 12140900 | 12140975 | 28371275 |
|--------------------------------|----------|----------|----------|----------|----------|
| SiO <sub>2</sub>               | 49.40    | 56.10    | 49.40    | 64.30    | 55.70    |
| TiO <sub>2</sub>               | .45      | .66      | .85      | .40      | .89      |
| Al <sub>2</sub> O <sub>3</sub> | 16.67    | 17.28    | 20.00    | 13.07    | 16.45    |
| Fe <sub>2</sub> O <sub>3</sub> | 10.59    | 9.41     | 11.86    | 3.13     | 9.83     |
| MnO                            | .16      | .21      | .22      | .10      | .13      |
| MgO                            | 6.00     | 3.35     | 3.90     | .52      | 3.20     |
| CaO                            | 6.85     | 7.43     | 5.63     | 7.10     | 3.12     |
| Na <sub>2</sub> O              | 1.89     | 2.01     | 3.35     | 5.10     | 1.19     |
| K <sub>2</sub> O               | .90      | 1.00     | 1.12     | .57      | 2.72     |
| LOI                            | 3.70     | 2.20     | 1.40     | 4.60     | 3.80     |
| TOTAL                          | 96.61    | 100.45   | 97.73    | 98.82    | 97.03    |

## TRACE ELEMENTS (PPM)

|                   |     |     |     |      |      |
|-------------------|-----|-----|-----|------|------|
| Zr                | 58  | 53  | 114 | 79   | 96   |
| Sr                | 304 | 289 | 505 | 448  | 243  |
| Rb                | 24  | 19  | 29  | 17   | 79   |
| Zn                | 112 | 95  | 151 | 19   | 73   |
| Cu                | 69  | 47  | 89  | 26   | 73   |
| Ba                | 978 | 679 | 801 | 3763 | 1869 |
| CuS               | 53  | 30  | 75  | 24   | 57   |
| PbS               | 7   | 7   | 7   | 8    | 12   |
| ZnS               | 20  | 12  | 21  | 8    | 20   |
| BaSO <sub>4</sub> | 0   |     |     | 0    | 0    |
| Pb                | 39  | 40  | 44  | 32   | 27   |
| Ni                | 27  | 7   | 5   | 1    | 7    |
| Co                | 29  | 22  | 29  | 6    | 35   |
| Cr                | 85  | 36  | 43  | 27   | 38   |
| V                 | 241 | 194 | 202 | 130  | 215  |
| Hg                | 32  | 29  | 95  | 103  |      |
| Ag                | 11  | 11  | 11  | 7    |      |
| Nb                |     |     |     |      | 9    |



## SKI HILL SEQUENCE FRAGMENTALS

## MAJOR ELEMENTS (WT%)

| SAMPLE | 28371398 | 28371588 | 28373200 | 28373550 | 2851 820 |
|--------|----------|----------|----------|----------|----------|
| SiO2   | 58.10    | 57.00    | 52.90    | 48.00    | 58.20    |
| TiO2   | .60      | .61      | .68      | .69      | .78      |
| Al2O3  | 17.20    | 16.90    | 17.07    | 19.09    | 15.07    |
| Fe2O3  | 7.43     | 7.14     | 9.29     | 10.09    | 8.00     |
| MnO    | .12      | .11      | .17      | .15      | .19      |
| HgO    | 2.81     | 2.89     | 4.21     | 4.43     | 3.24     |
| CaO    | 6.70     | 6.69     | 8.49     | 8.51     | 4.68     |
| Na2O   | 1.90     | 2.88     | 2.13     | 3.55     | 2.77     |
| K2O    | .71      | .70      | .74      | .47      | 1.80     |
| LOI    | 1.90     | 2.60     | 3.50     | 3.50     | 3.70     |
| TOTAL  | 97.47    | 97.52    | 99.18    | 98.48    | 98.43    |

## TRACE ELEMENTS (PPM)

|       |     |     |     |     |     |
|-------|-----|-----|-----|-----|-----|
| Zr    | 93  | 75  | 80  | 71  | 77  |
| Sr    | 424 | 373 | 318 | 276 | 263 |
| Rb    | 17  | 31  | 19  | 20  | 23  |
| Zn    | 52  | 70  | 57  | 82  | 84  |
| Cu    | 41  | 62  | 78  | 83  | 32  |
| Ba    | 65  | 718 | 688 | 424 | 874 |
| CuS   | 38  | 39  | 50  | 69  | 34  |
| PbS   | 15  | 14  | 20  | 20  | 25  |
| ZnS   | 16  | 14  | 14  | 17  | 31  |
| BaSO4 | 0   | 0   | 0   | 0   | 0   |
| Pb    | 29  | 25  | 30  | 37  | 25  |
| Ni    | 7   | 6   | 46  | 9   | 6   |
| Co    | 24  | 23  | 32  | 26  | 23  |
| Cr    | 27  | 39  | 78  | 44  | 13  |
| V     | 140 | 165 | 170 | 280 | 85  |
| Hg    |     |     |     |     |     |
| Ag    |     |     |     |     |     |
| Nb    | 5   | 4   | 3   | 2   | 2   |

## ALTERED SKI HILL SEQUENCE FRAGMENTALS

## MAJOR ELEMENTS (WT%)

| SAMPLE | 2851 882 | 28371500 | 28373200 | 28373550 | 2851 820 |
|--------|----------|----------|----------|----------|----------|
| SiO2   | 51.40    | 57.00    | 52.90    | 48.00    | 50.20    |
| TiO2   | .96      | .61      | .60      | .69      | .70      |
| Al2O3  | 15.33    | 16.90    | 17.07    | 19.09    | 15.07    |
| Fe2O3  | 12.82    | 7.14     | 9.29     | 10.09    | 8.00     |
| MnO    | .22      | .11      | .17      | .15      | .19      |
| MgO    | 3.39     | 2.89     | 4.21     | 4.43     | 3.24     |
| CaO    | 6.12     | 6.69     | 8.49     | 8.51     | 4.68     |
| Na2O   | 3.49     | 2.80     | 2.13     | 3.55     | 2.77     |
| K2O    | 2.39     | .70      | .74      | .47      | 1.80     |
| LOI    | 2.80     | 2.60     | 3.50     | 3.50     | 3.70     |
| TOTAL  | 98.92    | 97.52    | 99.18    | 98.48    | 98.43    |

## TRACE ELEMENTS (PPM)

|       |     |     |     |     |     |
|-------|-----|-----|-----|-----|-----|
| Zr    | 56  | 75  | 80  | 71  | 77  |
| Sr    | 167 | 373 | 318 | 276 | 263 |
| Rb    | 21  | 31  | 19  | 20  | 23  |
| Zn    | 78  | 70  | 57  | 82  | 84  |
| Cu    | 117 | 62  | 70  | 83  | 32  |
| Ba    | 691 | 718 | 688 | 424 | 874 |
| CuS   | 108 | 39  | 50  | 69  | 34  |
| PbS   | 20  | 14  | 20  | 20  | 25  |
| ZnS   | 41  | 14  | 14  | 17  | 31  |
| BaSO4 | 0   | 0   | 0   | 0   | 0   |
| Pb    | 24  | 25  | 30  | 37  | 25  |
| Ni    | 11  | 6   | 46  | 9   | 6   |
| Co    | 29  | 23  | 32  | 26  | 23  |
| Cr    | 21  | 39  | 78  | 44  | 13  |
| V     | 240 | 165 | 170 | 280 | 85  |
| Hg    |     |     |     |     |     |
| Ag    |     |     |     |     |     |
| Nb    | 0   | 4   | 3   | 2   | 2   |

## SKI HILL SEQUENCE FRAGMENTALS

## MAJOR ELEMENTS (WT%)

SAMPLE 2851 882

|                                |       |
|--------------------------------|-------|
| SiO <sub>2</sub>               | 51.48 |
| TiO <sub>2</sub>               | .96   |
| Al <sub>2</sub> O <sub>3</sub> | 15.33 |
| Fe <sub>2</sub> O <sub>3</sub> | 12.82 |
| MnO                            | .22   |
| MgO                            | 3.39  |
| CaO                            | 6.12  |
| Na <sub>2</sub> O              | 3.49  |
| K <sub>2</sub> O               | 2.39  |
| LOI                            | 2.88  |

|       |       |
|-------|-------|
| TOTAL | 98.92 |
|-------|-------|

## TRACE ELEMENTS (PPM)

|                   |     |
|-------------------|-----|
| Zr                | 36  |
| Sr                | 167 |
| Rb                | 21  |
| Zn                | 78  |
| Cu                | 117 |
| Ba                | 691 |
| CuS               | 188 |
| PbS               | 28  |
| ZnS               | 41  |
| BaSO <sub>4</sub> | 8   |
| Pb                | 24  |
| Ni                | 11  |
| Co                | 29  |
| Cr                | 21  |
| V                 | 248 |
| Hg                |     |
| Ag                |     |
| Nb                | 8   |

## LUCKY STRIKE ORE HORIZON SEQUENCE - HANGING WALL DACITIC TUFFS

## MAJOR ELEMENTS (WT%)

| SAMPLE                         | JGT75328 | 643 290 | 07920225 | 07920400 | 07920475 |
|--------------------------------|----------|---------|----------|----------|----------|
| SiO <sub>2</sub>               | 79.90    | 71.90   | 69.20    | 70.80    | 70.20    |
| TiO <sub>2</sub>               | .23      | .17     | .35      | .35      | .23      |
| Al <sub>2</sub> O <sub>3</sub> | 9.92     | 12.55   | 13.27    | 13.26    | 15.44    |
| Fe <sub>2</sub> O <sub>3</sub> | .81      | 2.14    | 2.85     | 2.90     | 2.14     |
| MnO                            | .03      | .04     | .09      | .11      | .05      |
| MgO                            | .11      | .91     | .93      | .92      | 1.07     |
| CaO                            | .73      | 2.85    | 3.78     | 2.87     | 1.65     |
| Na <sub>2</sub> O              | 3.14     | 2.66    | 3.13     | 2.20     | 1.80     |
| K <sub>2</sub> O               | 2.60     | 2.45    | 2.08     | 2.94     | 3.57     |
| LOI                            | 1.20     | 3.90    | 3.30     | 2.50     | 3.20     |
| TOTAL                          | 98.67    | 99.57   | 98.98    | 98.85    | 100.15   |

## TRACE ELEMENTS (PPM)

|                   |      |      |      |      |      |
|-------------------|------|------|------|------|------|
| Zr                | 70   | 93   | 162  | 187  | 107  |
| Sr                | 95   | 133  | 280  | 354  | 334  |
| Rb                | 38   | 57   | 53   | 77   | 87   |
| Zn                | 115  | 20   | 56   | 60   | 29   |
| Cu                | 2    | 1    | 3    | 3    | 4    |
| Ba                | 1624 | 1437 | 4081 | 1070 | 1232 |
| CuS               | 1    | 1    | 0    | 0    | 0    |
| PbS               | 1    | 10   | 32   | 13   | 9    |
| ZnS               | 125  | 11   | 19   | 9    | 3    |
| BaSO <sub>4</sub> | 4    | 6    | 3    | 0    |      |
| Pb                | 12   | 11   | 49   | 30   | 13   |
| Ni                | 1    | 2    | 2    | 2    | 1    |
| Co                | 56   | 15   | 2    | 3    | 2    |
| Cr                | 10   | 11   | 31   | 21   | 22   |
| V                 | 10   | 10   | 2    | 0    | 0    |
| Hg                |      |      | 10   | 13   | 11   |
| Ag                |      |      | 1    | 2    | 1    |
| Nb                | 7    | 15   |      |      |      |

## LUCKY STRIKE ORE HORIZON SEQUENCE - HANGING WALL DACITIC TUFFS

## MAJOR ELEMENTS (WT%)

| SAMPLE                         | 07920575 | 07920675 | 07920775 | 07920925 | 07920950 |
|--------------------------------|----------|----------|----------|----------|----------|
| SiO <sub>2</sub>               | 74.00    | 72.20    | 70.50    | 74.90    | 69.50    |
| TiO <sub>2</sub>               | .20      | .20      | .18      | .20      | .22      |
| Al <sub>2</sub> O <sub>3</sub> | 12.40    | 13.40    | 13.20    | 12.04    | 15.90    |
| Fe <sub>2</sub> O <sub>3</sub> | 1.85     | 2.19     | 2.25     | 1.67     | 1.86     |
| MnO                            | .10      | .09      | .10      | .09      | .09      |
| MgO                            | .72      | 1.19     | 1.03     | .62      | 2.26     |
| CaO                            | 1.06     | 1.32     | 2.77     | 2.68     | 1.20     |
| Na <sub>2</sub> O              | 4.93     | 2.53     | 2.13     | 3.34     | 1.62     |
| K <sub>2</sub> O               | 1.57     | 4.00     | 3.34     | 1.28     | 3.00     |
| LOI                            | 1.40     | 2.20     | 4.10     | 1.70     | 3.00     |
| TOTAL                          | 98.23    | 99.32    | 99.60    | 98.52    | 98.65    |

## TRACE ELEMENTS (PPM)

|                   |      |      |     |      |      |
|-------------------|------|------|-----|------|------|
| Zr                | 96   | 121  | 115 | 122  | 134  |
| Sr                | 108  | 216  | 151 | 597  | 358  |
| Rb                | 40   | 116  | 84  | 40   | 105  |
| Zn                | 97   | 23   | 20  | 35   | 141  |
| Cu                | 11   | 2    | 2   | 6    | 57   |
| Ba                | 1260 | 1551 | 783 | 1152 | 2194 |
| CuS               | 0    | 0    | 1   | 2    | 1    |
| PbS               | 12   | 12   | 17  | 21   | 12   |
| ZnS               | 4    | 2    | 4   | 8    | 2    |
| BaSO <sub>4</sub> | 0    |      | 0   |      | 5    |
| Pb                | 16   | 18   | 18  | 56   | 33   |
| Ni                | 1    | 1    | 1   | 1    | 57   |
| Co                | 1    | 1    | 1   | 2    | 40   |
| Cr                | 28   | 24   | 79  | 22   | 29   |
| V                 | 0    | 6    | 0   | 0    | 345  |
| Hg                | 13   | 10   | 11  | 18   | 5    |
| Ag                | 2    | 2    | 0   | 0    | 1    |
| Nb                |      |      |     |      |      |

## LUCKY STRIKE ORE HORIZON SEQUENCE - HANGING WALL DACITIC TUFFS

## MAJOR ELEMENTS (WT%)

| SAMPLE                         | 07920975 | 07921000 | 07921025 | 08681550 | 08681590 |
|--------------------------------|----------|----------|----------|----------|----------|
| SiO <sub>2</sub>               | 69.90    | 73.30    | 70.10    | 74.40    | 72.10    |
| TiO <sub>2</sub>               | .22      | .18      | .25      | .18      | .22      |
| Al <sub>2</sub> O <sub>3</sub> | 15.23    | 13.00    | 14.87    | 13.20    | 13.50    |
| Fe <sub>2</sub> O <sub>3</sub> | 1.62     | 1.51     | 2.56     | 1.90     | 1.83     |
| MnO                            | .07      | .10      | .08      | .08      | .08      |
| MgO                            | 1.20     | .75      | 1.82     | .56      | .62      |
| CaO                            | 2.32     | 2.62     | 2.24     | 2.11     | 2.17     |
| Na <sub>2</sub> O              | 2.90     | 3.66     | 2.12     | 2.43     | 2.63     |
| K <sub>2</sub> O               | 2.70     | 1.47     | 3.09     | 3.12     | 3.09     |
| LOI                            | 2.50     | 2.20     | 2.80     | 1.90     | 2.30     |
| TOTAL                          | 98.66    | 98.87    | 99.93    | 99.80    | 98.54    |

## TRACE ELEMENTS (PPM)

|                   |      |      |      |     |      |
|-------------------|------|------|------|-----|------|
| Zr                | 175  | 110  | 108  | 120 | 112  |
| Sr                | 698  | 551  | 357  | 278 | 216  |
| Rb                | 86   | 37   | 72   | 92  | 92   |
| Zn                | 26   | 32   | 127  | 18  | 25   |
| Cu                | 2    | 3    | 7    | 2   | 2    |
| Ba                | 1042 | 7814 | 3316 | 975 | 2439 |
| CuS               | 0    | 0    | 2    | 0   | 0    |
| PbS               | 21   | 9    | 18   | 14  | 21   |
| ZnS               | 1    | 7    | 27   | 5   | 4    |
| BaSO <sub>4</sub> |      |      |      | 0   | 0    |
| Pb                | 40   | 28   | 30   | 18  | 19   |
| Ni                | 1    | 1    | 2    | 0   | 0    |
| Co                | 2    | 2    | 2    | 1   | 1    |
| Cr                | 22   | 22   | 28   | 27  | 19   |
| V                 | 0    | 0    | 27   | 9   | 10   |
| Hg                | 8    | 6    | 12   | 11  | 13   |
| Ag                | 0    | 0    | 0    | 0   | 0    |
| Nb                |      |      |      |     |      |

## LUCKY STRIKE ORE HORIZON SEQUENCE - HANGING WALL DACITIC TUFFS

## MAJOR ELEMENTS (WT%)

| SAMPLE | 08681630 | 08692190 | 08692215 | 08692240 | 08692265 |
|--------|----------|----------|----------|----------|----------|
| S102   | 65.40    | 66.60    | 67.10    | 67.70    | 71.70    |
| TiO2   | .40      | .21      | .25      | .27      | .21      |
| Al2O3  | 14.29    | 15.91    | 14.60    | 15.12    | 13.24    |
| Fe2O3  | 4.50     | 2.09     | 2.17     | 2.44     | 1.63     |
| MnO    | .11      | .05      | .07      | .08      | .06      |
| MgO    | 1.71     | 1.51     | 1.25     | .98      | .69      |
| CaO    | 3.47     | .64      | 2.09     | 1.65     | 1.81     |
| Na2O   | 1.80     | 1.77     | 1.74     | 4.00     | 2.29     |
| K2O    | 3.40     | 4.58     | 4.32     | 3.50     | 3.79     |
| LOI    | 3.80     | 3.30     | 4.10     | 1.90     | 2.80     |
| TOTAL  | 98.88    | 96.66    | 97.69    | 97.64    | 98.22    |

## TRACE ELEMENTS (PPM)

|       |     |      |      |      |      |
|-------|-----|------|------|------|------|
| Zr    | 179 | 164  | 174  | 166  | 145  |
| Sr    | 263 | 32   | 147  | 221  | 139  |
| Rb    | 95  | 123  | 106  | 76   | 95   |
| Zn    | 67  | 25   | 24   | 42   | 37   |
| Cu    | 5   | 3    | 2    | 5    | 3    |
| Ba    | 905 | 1215 | 5058 | 1510 | 1018 |
| CuS   | 2   | 1    | 1    | 2    | 1    |
| PbS   | 12  | 14   | 37   | 18   | 23   |
| ZnS   | 11  | 2    | 2    | 8    | 3    |
| BaSO4 |     |      |      |      |      |
| Pb    | 21  | 19   | 22   | 23   | 26   |
| Ni    | 2   | 3    | 4    | 4    | 3    |
| Co    | 4   | 2    | 2    | 2    | 2    |
| Cr    | 22  | 7    | 8    | 8    | 5    |
| V     | 26  | 25   | 20   | 37   | 18   |
| Hg    | 12  | 0    | 0    | 5    | 4    |
| Ag    | 0   | 0    | 0    | 0    | 0    |
| Nb    |     |      |      |      |      |

## LUCKY STRIKE ORE HORIZON SEQUENCE - HANGING WALL DACITIC TUFFS

## MAJOR ELEMENTS (WT%)

| SAMPLE                         | 08692315 | 08692340 | 08692390 | 08692405 | 08692430 |
|--------------------------------|----------|----------|----------|----------|----------|
| SiO <sub>2</sub>               | 68.00    | 76.30    | 75.90    | 67.50    | 68.80    |
| TiO <sub>2</sub>               | .24      | .19      | .23      | .21      | .24      |
| Al <sub>2</sub> O <sub>3</sub> | 16.38    | 12.40    | 13.48    | 14.94    | 13.62    |
| Fe <sub>2</sub> O <sub>3</sub> | 1.94     | 1.93     | 1.49     | 2.12     | 2.78     |
| MnO                            | .06      | .06      | .04      | .08      | .08      |
| HgO                            | .65      | .67      | .56      | 1.39     | 1.50     |
| CaO                            | 2.04     | 1.00     | 1.00     | 2.31     | 2.52     |
| Na <sub>2</sub> O              | 4.25     | 3.90     | 4.89     | 3.58     | 1.92     |
| K <sub>2</sub> O               | 3.12     | 2.40     | 1.90     | 2.76     | 3.50     |
| LOI                            | 1.90     | 1.60     | 1.70     | 3.60     | 3.80     |
| TOTAL                          | 98.58    | 100.45   | 101.19   | 98.49    | 98.76    |

## TRACE ELEMENTS (PPM)

|                   |      |     |     |     |     |
|-------------------|------|-----|-----|-----|-----|
| Zr                | 197  | 143 | 160 | 167 | 142 |
| Sr                | 233  | 105 | 167 | 81  | 138 |
| Rb                | 86   | 62  | 50  | 66  | 100 |
| Zn                | 33   | 37  | 32  | 39  | 51  |
| Cu                | 8    | 3   | 3   | 5   | 8   |
| Ba                | 1217 | 892 | 428 | 535 | 650 |
| CuS               | 4    | 1   | 1   | 2   | 5   |
| PbS               | 21   | 14  | 15  | 15  | 24  |
| ZnS               | 3    | 8   | 9   | 10  | 12  |
| BaSO <sub>4</sub> |      |     |     |     |     |
| Pb                | 22   | 18  | 21  | 22  | 27  |
| Ni                | 3    | 3   | 3   | 4   | 7   |
| Co                | 2    | 2   | 2   | 2   | 2   |
| Cr                | 6    | 9   | 8   | 11  | 23  |
| V                 | 21   | 14  | 24  | 19  | 47  |
| Hg                | 5    | 5   | 6   | 11  | 5   |
| Ag                | 0    | 0   | 0   | 0   | 0   |
| Nb                |      |     |     |     |     |



## LUCKY STRIKE ORE HORIZON SEQUENCE - HANGING WALL DACITIC TUFFS

## MAJOR ELEMENTS (WT%)

| SAMPLE | 9781268 | 12141888 | 12141825 | 12141850 | 12141875 |
|--------|---------|----------|----------|----------|----------|
| SiO2   | 78.48   | 73.28    | 66.98    | 75.88    | 67.48    |
| TiO2   | .33     | .22      | .23      | .13      | .23      |
| Al2O3  | 13.73   | 13.48    | 15.74    | 18.11    | 15.44    |
| Fe2O3  | 2.98    | 2.84     | 2.25     | 2.88     | 1.83     |
| MnO    | .88     | .18      | .11      | .15      | .88      |
| MgO    | .99     | .79      | .92      | .57      | .78      |
| CaO    | 2.23    | 2.79     | 3.12     | 2.96     | 3.28     |
| Na2O   | 2.77    | 2.31     | 2.68     | 3.21     | 1.98     |
| K2O    | 2.45    | 2.63     | 3.14     | 1.36     | 4.15     |
| LOI    | 3.58    | 2.18     | 3.78     | 2.88     | 4.58     |
| TOTAL  | 99.46   | 99.68    | 98.79    | 99.89    | 99.43    |

## TRACE ELEMENTS (PPM)

|       |     |      |     |      |     |
|-------|-----|------|-----|------|-----|
| Zr    | 81  | 119  | 151 | 81   | 188 |
| Sr    | 68  | 377  | 442 | 311  | 83  |
| Rb    | 68  | 82   | 88  | 41   | 99  |
| Zn    | 83  | 31   | 36  | 27   | 28  |
| Cu    | 1   | 2    | 2   | 2    | 2   |
| Ba    | 688 | 1868 | 177 | 1987 | 977 |
| CuS   | 1   | 8    | 8   | 8    | 8   |
| PbS   | 4   | 3    | 9   | 5    | 15  |
| ZnS   | 13  | 2    | 7   | 2    | 2   |
| BaSO4 | 4   |      |     |      | 3   |
| Pb    | 7   | 28   | 29  | 22   | 26  |
| Ni    | 2   | 1    | 1   | 8    | 8   |
| Co    | 15  | 3    | 3   | 2    | 2   |
| Cr    | 9   | 41   | 38  | 21   | 42  |
| V     | 18  | 8    | 8   | 8    | 6   |
| Hg    |     | 16   | 13  | 22   | 22  |
| Ag    |     | 5    | 5   | 4    | 8   |
| Nb    | 11  |      |     |      |     |

## LUCKY STRIKE ORE HORIZON SEQUENCE - HANGING WALL DACITIC TUFFS

## MAJOR ELEMENTS (WT%)

| SAMPLE                         | 12141100 | 12141125 | 12141150 | 12141175 | 12141200 |
|--------------------------------|----------|----------|----------|----------|----------|
| SiO <sub>2</sub>               | 70.20    | 69.00    | 72.70    | 73.60    | 73.50    |
| TiO <sub>2</sub>               | .23      | .10      | .10      | .03      | .15      |
| Al <sub>2</sub> O <sub>3</sub> | 14.53    | 13.23    | 12.72    | 11.96    | 11.78    |
| Fe <sub>2</sub> O <sub>3</sub> | 2.05     | 1.84     | 1.70     | 2.12     | 1.95     |
| MnO                            | .07      | .09      | .11      | .17      | .13      |
| MgO                            | 1.04     | .71      | .64      | .34      | .39      |
| CaO                            | 2.42     | 2.15     | 3.62     | 2.17     | 3.58     |
| Na <sub>2</sub> O              | 2.24     | 3.45     | 1.85     | 5.82     | 2.91     |
| K <sub>2</sub> O               | 3.69     | 2.50     | 3.26     | .52      | 1.98     |
| LOI                            | 3.60     | 3.30     | 3.70     | 2.10     | 2.60     |
| TOTAL                          | 100.07   | 96.45    | 100.48   | 98.83    | 98.97    |

## TRACE ELEMENTS (PPM)

|                   |      |     |      |      |     |
|-------------------|------|-----|------|------|-----|
| Zr                | 116  | 92  | 117  | 99   | 59  |
| Sr                | 105  | 135 | 355  | 207  | 106 |
| Rb                | 100  | 72  | 95   | 19   | 60  |
| Zn                | 38   | 35  | 35   | 43   | 29  |
| Cu                | 1    | 2   | 1    | 2    | 2   |
| Ba                | 1124 | 967 | 2341 | 1712 | 649 |
| CuS               | 0    | 0   | 0    | 0    | 0   |
| PbS               | 6    | 11  | 13   | 6    | 13  |
| ZnS               | 4    | 2   | 1    | 7    | 3   |
| BaSO <sub>4</sub> |      |     |      |      |     |
| Pb                | 21   | 22  | 23   | 18   | 10  |
| Ni                | 0    | 0   | 1    | 1    | 1   |
| Co                | 3    | 3   | 2    | 2    | 2   |
| Cr                | 26   | 19  | 22   | 25   | 51  |
| V                 | 4    | 5   | 10   | 0    | 10  |
| Hg                | 22   | 16  | 31   | 27   | 32  |
| Ag                | 7    | 6   | 4    | 5    | 6   |
| Nb                |      |     |      |      |     |

## LUCKY STRIKE ORE HORIZON SEQUENCE - HANGING WALL DACITIC TUFFS

## MAJOR ELEMENTS (WT%)

| SAMPLE                         | 12141225 | 12141245 | 12141265 | 12141285 | 12141295 |
|--------------------------------|----------|----------|----------|----------|----------|
| SiO <sub>2</sub>               | 72.00    | 70.30    | 77.20    | 72.70    | 71.10    |
| TiO <sub>2</sub>               | .18      | .17      | .23      | .22      | .17      |
| Al <sub>2</sub> O <sub>3</sub> | 13.63    | 13.69    | 13.90    | 12.60    | 13.96    |
| Fe <sub>2</sub> O <sub>3</sub> | 1.87     | 2.16     | 1.12     | 1.79     | 1.88     |
| MnO                            | .12      | .13      | .03      | .08      | .07      |
| MgO                            | .63      | .79      | .36      | .57      | .60      |
| CaO                            | 2.65     | 2.69     | .29      | 2.85     | 2.28     |
| Na <sub>2</sub> O              | 2.05     | .86      | .56      | 2.15     | 4.86     |
| K <sub>2</sub> O               | 3.33     | 4.14     | 3.32     | 2.51     | 1.47     |
| LOI                            | 3.00     | 4.40     | 1.80     | 3.00     | 2.00     |
| TOTAL                          | 99.46    | 99.33    | 98.81    | 98.55    | 98.39    |

## TRACE ELEMENTS (PPM)

|                   |      |      |      |      |      |
|-------------------|------|------|------|------|------|
| Zr                | 110  | 90   | 116  | 130  | 113  |
| Sr                | 268  | 50   | 27   | 393  | 35   |
| Rb                | 91   | 104  | 75   | 74   | 45   |
| Zn                | 42   | 47   | 17   | 38   | 44   |
| Cu                | 1    | 1    | 2    | 2    | 2    |
| Ba                | 1223 | 1472 | 1930 | 4077 | 1111 |
| CuS               | 0    | 0    | 0    | 0    | 0    |
| PbS               | 24   | 17   | 11   | 0    | 11   |
| ZnS               | 3    | 4    | 1    | 4    | 5    |
| BaSO <sub>4</sub> |      |      | 9    |      |      |
| Pb                | 15   | 11   | 4    | 6    | 7    |
| Ni                | 1    | 1    | 1    | 0    | 0    |
| Co                | 3    | 2    | 1    | 1    | 2    |
| Cr                | 21   | 18   | 24   | 27   | 31   |
| V                 | 4    | 6    | 0    | 0    | 0    |
| Hg                | 32   | 31   | 137  | 31   | 22   |
| Ag                | 5    | 6    | 4    | 5    | 6    |
| Nb                |      |      |      |      |      |

## LUCKY STRIKE ORE HORIZON SEQUENCE - HANGING WALL DACITIC TUFFS

## MAJOR ELEMENTS (WT%)

| SAMPLE                         | 14930400 | 14930450 | 14930500 | 14930550 | 14930600 |
|--------------------------------|----------|----------|----------|----------|----------|
| SiO <sub>2</sub>               | 73.10    | 69.10    | 69.60    | 73.90    | 71.80    |
| TiO <sub>2</sub>               | .23      | .21      | .20      | .18      | .26      |
| Al <sub>2</sub> O <sub>3</sub> | 15.00    | 14.57    | 14.61    | 13.66    | 14.61    |
| Fe <sub>2</sub> O <sub>3</sub> | 1.70     | 1.99     | 2.06     | 1.75     | 2.05     |
| MnO                            | .06      | .08      | .10      | .07      | .08      |
| HgO                            | 2.33     | .88      | .63      | .45      | .68      |
| CaO                            | 1.15     | 3.32     | 3.20     | 2.12     | 1.32     |
| Na <sub>2</sub> O              | 2.42     | 1.95     | 2.00     | 2.38     | 2.93     |
| K <sub>2</sub> O               | 2.33     | 3.16     | 2.68     | 3.64     | 3.39     |
| LOI                            | 3.00     | 4.70     | 3.70     | 3.10     | 2.90     |
| TOTAL                          | 101.32   | 99.96    | 98.78    | 101.25   | 100.02   |

## TRACE ELEMENTS (PPM)

|                   |      |      |      |     |     |
|-------------------|------|------|------|-----|-----|
| Zr                | 133  | 62   |      | 154 | 130 |
| Sr                | 212  | 0    |      | 192 | 41  |
| Rb                | 60   | 51   |      | 94  | 81  |
| Zn                | 14   | 18   | 25   | 26  | 18  |
| Cu                | 1    | 2    | 5    | 4   | 3   |
| Ba                | 2170 | 1654 | 2320 | 970 | 716 |
| CuS               | 0    | 1    | 2    | 2   | 1   |
| PbS               | 11   | 8    | 8    | 18  | 15  |
| ZnS               | 3    | 1    | 2    | 4   | 6   |
| BaSO <sub>4</sub> |      | 9    |      |     | 0   |
| Pb                | 11   | 9    | 11   | 17  | 11  |
| Ni                | 1    | 1    | 1    | 1   | 1   |
| Co                | 1    | 1    | 2    | 2   | 2   |
| Cr                | 20   | 24   |      | 28  | 97  |
| V                 | 0    | 13   | 0    | 0   | 10  |
| Hg                | 12   | 12   | 21   | 12  | 10  |
| Ag                | 3    | 0    | 2    | 0   | 0   |
| Nb                |      |      |      |     |     |

## LUCKY STRIKE ORE HORIZON SEQUENCE - HANGING WALL DACITIC TUFFS

## MAJOR ELEMENTS (WTX)

| SAMPLE                         | 14930725 | 14930750 | 14930775 | 14930800 | 26103011 |
|--------------------------------|----------|----------|----------|----------|----------|
| SiO <sub>2</sub>               | 72.30    | 71.20    | 72.50    | 73.70    | 69.30    |
| TiO <sub>2</sub>               | .29      | .23      | .27      | .17      | .29      |
| Al <sub>2</sub> O <sub>3</sub> | 12.54    | 14.00    | 13.37    | 13.46    | 14.03    |
| Fe <sub>2</sub> O <sub>3</sub> | 2.03     | 2.37     | 2.08     | 2.16     | 2.50     |
| MnO                            | .11      | .12      | .08      | .09      | .04      |
| MgO                            | .64      | .68      | .63      | .64      | 1.86     |
| CaO                            | 3.50     | 1.42     | 2.19     | 2.02     | 2.00     |
| Na <sub>2</sub> O              | 1.03     | 3.21     | 3.20     | 2.30     | 1.74     |
| K <sub>2</sub> O               | 4.13     | 3.84     | 3.02     | 3.51     | 4.05     |
| LOI                            | 4.20     | 1.60     | 1.70     | 2.20     | 3.50     |
| TOTAL                          | 100.77   | 98.67    | 99.04    | 100.25   | 99.31    |

## TRACE ELEMENTS (PPM)

|                   |      |      |      |      |     |
|-------------------|------|------|------|------|-----|
| Zr                | 146  | 106  | 158  | 148  | 133 |
| Sr                | 128  | 163  | 275  | 309  | 143 |
| Rb                | 110  | 81   | 75   | 93   | 137 |
| Zn                | 25   | 31   | 22   | 21   | 28  |
| Cu                | 1    | 6    | 2    | 1    | 2   |
| Ba                | 4178 | 1649 | 1344 | 1140 | 567 |
| CuS               | 0    | 2    | 0    | 0    | 1   |
| PbS               | 15   | 15   | 18   | 12   | 27  |
| ZnS               | 7    | 7    | 5    | 13   | 2   |
| BaSO <sub>4</sub> |      | 4    |      |      |     |
| Pb                | 14   | 12   | 12   | 12   | 13  |
| Ni                | 2    | 2    | 1    | 1    | 2   |
| Co                | 2    | 3    | 1    | 2    | 11  |
| Cr                | 24   | 26   | 20   | 23   | 8   |
| V                 | 18   | 12   | 0    | 15   | 10  |
| Hg                | 10   | 22   | 12   | 10   |     |
| Ag                | 0    | 2    | 3    | 6    |     |
| Nb                |      |      |      |      |     |

## LUCKY STRIKE ORE HORIZON SEQUENCE - HANGING WALL DACITIC TUFFS

## MAJOR ELEMENTS (WT%)

| SAMPLE                         | 26663025 | 26663075 | 26663150 | 26 3250 | 26663325 |
|--------------------------------|----------|----------|----------|---------|----------|
| SiO <sub>2</sub>               | 76.80    | 66.60    | 71.90    | 73.70   | 70.40    |
| TiO <sub>2</sub>               | .17      | .31      | .24      | .24     | .20      |
| Al <sub>2</sub> O <sub>3</sub> | 9.73     | 14.21    | 13.66    | 11.14   | 12.98    |
| Fe <sub>2</sub> O <sub>3</sub> | 2.02     | 2.37     | 1.87     | 1.72    | 2.40     |
| MnO                            | .13      | .07      | .06      | .06     | .09      |
| MgO                            | .33      | .80      | .76      | .52     | .63      |
| CaO                            | 1.86     | 3.53     | 1.30     | .99     | 3.00     |
| Na <sub>2</sub> O              | 4.86     | 3.68     | 5.00     | 2.47    | 2.27     |
| K <sub>2</sub> O               | .39      | 2.94     | 1.50     | 4.86    | 3.36     |
| LOI                            | 1.30     | 3.40     | 1.70     | .80     | 2.20     |
| TOTAL                          | 97.59    | 97.91    | 97.99    | 96.50   | 97.53    |

## TRACE ELEMENTS (PPM)

|                   |      |     |     |      |     |
|-------------------|------|-----|-----|------|-----|
| Zr                | 63   | 101 | 115 | 82   | 108 |
| Sr                | 192  | 198 | 168 | 138  | 282 |
| Rb                | 11   | 75  | 39  | 58   | 89  |
| Zn                | 13   | 18  | 14  | 24   | 21  |
| Cu                | 5    | 16  | 30  | 11   | 3   |
| Ba                | 1062 | 619 | 260 | 1904 | 61  |
| CuS               | 1    | 15  | 35  | 7    | 1   |
| PbS               | 4    | 18  | 10  | 11   | 7   |
| ZnS               | 7    | 7   | 2   | 3    | 1   |
| BaSO <sub>4</sub> |      |     | 0   |      |     |
| Pb                | 1    | 2   | 2   | 10   | 12  |
| Ni                | 1    | 1   | 1   | 3    | 1   |
| Co                | 2    | 2   | 3   | 4    | 2   |
| Cr                | 25   | 25  | 23  | 22   | 23  |
| V                 | 20   | 33  | 18  | 37   | 10  |
| Hg                | 8    | 10  | 10  | 10   | 8   |
| Ag                | 2    | 0   | 1   | 0    | 0   |
| Nb                |      |     |     |      |     |

3-66

## LUCKY STRIKE ORE HORIZON SEQUENCE - HANGING WALL DACITIC TUFFS

## MAJOR ELEMENTS (WT%)

| SAMPLE                         | 26    | 3400  | 26663475 | 26    | 3550  | 26802856 | 27101053 |
|--------------------------------|-------|-------|----------|-------|-------|----------|----------|
| SiO <sub>2</sub>               | 70.10 | 68.60 | 72.00    | 68.00 | 69.50 |          |          |
| TiO <sub>2</sub>               | .23   | .15   | .19      | .11   | .32   |          |          |
| Al <sub>2</sub> O <sub>3</sub> | 13.46 | 13.07 | 12.44    | 14.71 | 13.91 |          |          |
| Fe <sub>2</sub> O <sub>3</sub> | 2.67  | 1.58  | 1.56     | 2.15  | 2.68  |          |          |
| MnO                            | .08   | .06   | .08      | .04   | .07   |          |          |
| MgO                            | .67   | .40   | .97      | .71   | 1.15  |          |          |
| CaO                            | 2.50  | 2.89  | 2.48     | 3.33  | 1.69  |          |          |
| Na <sub>2</sub> O              | 2.82  | 4.35  | 2.72     | 1.40  | 2.31  |          |          |
| K <sub>2</sub> O               | 3.30  | 2.06  | 2.14     | 3.72  | 4.03  |          |          |
| LOI                            | 1.80  | 3.40  | 2.60     | 5.00  | 3.30  |          |          |
| TOTAL                          | 97.63 | 96.56 | 97.18    | 99.17 | 98.96 |          |          |

## TRACE ELEMENTS (PPM)

|                   |      |      |      |      |      |
|-------------------|------|------|------|------|------|
| Zr                | 125  | 104  | 96   | 129  | 155  |
| Sr                | 295  | 172  | 254  | 113  | 17   |
| Rb                | 76   | 51   | 48   | 90   | 88   |
| Zn                | 24   | 15   | 16   | 18   | 17   |
| Cu                | 3    | 3    | 1    | 1    | 1    |
| Ba                | 1177 | 1734 | 1179 | 1090 | 1106 |
| CuS               | 1    | 0    | 0    | 1    | 1    |
| PbS               | 7    | 12   | 10   | 28   | 14   |
| ZnS               | 6    | 5    | 2    | 17   | 12   |
| BaSO <sub>4</sub> | 0    |      | 3    |      |      |
| Pb                | 6    | 5    | 0    | 15   | 13   |
| Ni                | 2    | 2    | 1    | 3    | 2    |
| Co                | 3    | 2    | 1    | 15   | 20   |
| Cr                | 22   | 25   | 20   | 4    | 9    |
| V                 | 22   | 15   | 10   | 20   | 10   |
| Hg                | 10   | 6    | 10   |      |      |
| Ag                | 0    | 0    | 0    |      |      |
| Nb                |      |      |      |      |      |

## LUCKY STRIKE ORE HORIZON SEQUENCE - HANGING WALL DACITIC TUFFS

## MAJOR ELEMENTS (WT%)

| SAMPLE                         | 27101150 | 27711312 | 27711350 | 2800 182 | 2800 387 |
|--------------------------------|----------|----------|----------|----------|----------|
| SiO <sub>2</sub>               | 68.80    | 71.70    | 72.90    | 71.00    | 66.70    |
| TiO <sub>2</sub>               | .26      | .20      | .26      | .20      | .44      |
| Al <sub>2</sub> O <sub>3</sub> | 14.24    | 13.28    | 13.00    | 14.63    | 15.88    |
| Fe <sub>2</sub> O <sub>3</sub> | 3.06     | 2.13     | 2.17     | 1.72     | 4.09     |
| MnO                            | .08      | .04      | .06      | .09      | .07      |
| MgO                            | 1.42     | .95      | .78      | 1.12     | 2.41     |
| CaO                            | 2.17     | 2.10     | 2.58     | 2.20     | 2.62     |
| Na <sub>2</sub> O              | 1.59     | 2.75     | 2.93     | 2.03     | 2.03     |
| K <sub>2</sub> O               | 4.06     | 2.63     | 2.01     | 3.67     | 2.35     |
| LOI                            | 4.20     | 2.70     | 2.50     | 2.40     | 3.30     |
| TOTAL                          | 99.88    | 98.48    | 99.19    | 99.06    | 99.89    |

## TRACE ELEMENTS (PPM)

|                   |     |      |     |      |      |
|-------------------|-----|------|-----|------|------|
| Zr                | 149 | 134  | 140 | 140  | 189  |
| Sr                | 117 | 311  | 360 | 461  | 358  |
| Rb                | 96  | 78   | 58  | 113  | 65   |
| Zn                | 24  | 19   | 17  | 16   | 66   |
| Cu                | 3   | 1    | 1   | 1    | 1    |
| Ba                | 584 | 1359 | 955 | 1254 | 1094 |
| CuS               | 4   | 1    | 1   | 1    | 1    |
| PbS               | 25  | 32   | 27  | 28   | 18   |
| ZnS               | 14  | 2    | 8   | 2    | 17   |
| BaSO <sub>4</sub> |     |      |     |      |      |
| Pb                | 11  | 21   | 12  | 15   | 22   |
| Ni                | 4   | 2    | 2   | 1    | 5    |
| Co                | 11  | 45   | 18  | 16   | 15   |
| Cr                | 11  | 5    | 6   | 6    | 9    |
| V                 | 10  | 15   | 10  | 10   | 10   |
| Hg                |     |      |     |      |      |
| Ag                |     |      |     |      |      |
| Nb                |     |      |     |      |      |



## LUCKY STRIKE ORE HORIZON SEQUENCE - HANGING WALL DACITIC TUFFS

## MAJOR ELEMENTS (WT%)

SAMPLE 14936650

|                                |       |
|--------------------------------|-------|
| SiO <sub>2</sub>               | 72.60 |
| TiO <sub>2</sub>               | .29   |
| Al <sub>2</sub> O <sub>3</sub> | 13.51 |
| Fe <sub>2</sub> O <sub>3</sub> | 2.50  |
| MnO                            | .13   |
| HgO                            | .70   |
| CaO                            | 1.40  |
| Na <sub>2</sub> O              | 3.92  |
| K <sub>2</sub> O               | 2.75  |
| LOI                            | 1.50  |

|       |       |
|-------|-------|
| TOTAL | 99.30 |
|-------|-------|

## TRACE ELEMENTS (PPM)

|                   |      |
|-------------------|------|
| Zr                | 126  |
| Sr                | 203  |
| Rb                | 67   |
| Zn                | 24   |
| Cu                | 5    |
| Ba                | 1437 |
| CuS               | 2    |
| PbS               | 17   |
| ZnS               | 10   |
| BaSO <sub>4</sub> |      |
| Pb                | 15   |
| Ni                | 2    |
| Co                | 2    |
| Cr                | 39   |
| V                 | 14   |
| Hg                | 14   |
| Ag                | 0    |
| Nb                |      |

## ORIENTAL AND LUCKY STRIKE AREA RHYOLITE FLOWS

## MAJOR ELEMENTS (WTZ)

| SAMPLE | JGT75327 | 158 584 | 158 617 | 885 195 | 1589 778 |
|--------|----------|---------|---------|---------|----------|
| SiO2   | 69.88    | 71.50   | 74.90   | 72.80   | 75.00    |
| TiO2   | .31      | .33     | .38     | .28     | .36      |
| Al2O3  | 14.21    | 14.55   | 12.90   | 13.49   | 13.68    |
| Fe2O3  | 2.65     | 2.60    | 2.54    | 1.97    | .95      |
| MnO    | .05      | .04     | .02     | .02     | .02      |
| HgO    | .53      | .43     | .33     | .03     | .14      |
| CaO    | 1.78     | 1.50    | 1.19    | 1.19    | .68      |
| Na2O   | 5.34     | 5.26    | 3.67    | 4.85    | 5.02     |
| K2O    | 1.50     | 2.65    | 3.35    | 2.42    | 2.27     |
| LOI    | 2.60     | .80     | .80     | 1.50    | 1.10     |
| TOTAL  | 98.77    | 99.66   | 100.08  | 98.55   | 99.22    |

## TRACE ELEMENTS (PPM)

|       |     |      |      |     |      |
|-------|-----|------|------|-----|------|
| Zr    | 129 | 165  | 159  | 181 | 187  |
| Sr    | 122 | 27   | 317  | 222 | 237  |
| Rb    | 23  | 45   | 66   | 49  | 39   |
| Zn    | 35  | 33   | 31   | 5   | 30   |
| Cu    | 37  | 2    | 1    | 1   | 1    |
| Ba    | 565 | 1242 | 1535 | 962 | 1126 |
| CuS   | 42  | 1    | 1    | 1   | 1    |
| PbS   | 21  | 8    | 11   | 3   | 32   |
| ZnS   | 7   | 5    | 12   | 4   | 16   |
| BaSO4 | 0   |      |      |     |      |
| Pb    | 24  | 10   | 3    | 1   | 43   |
| Ni    | 1   | 2    | 2    | 2   | 2    |
| Co    | 35  | 29   | 23   | 27  | 24   |
| Cr    | 12  | 17   | 9    | 2   | 0    |
| V     | 10  | 55   | 35   | 15  | 10   |
| Hg    |     |      |      |     |      |
| Ag    |     |      |      |     |      |
| Nb    | 9   |      |      |     |      |

3-70

## ORIENTAL AND LUCKY STRIKE AREA RHYOLITE FLOWS

## MAJOR ELEMENTS (WT%)

SAMPLE 1589 811 17301260

|                                |       |       |
|--------------------------------|-------|-------|
| SiO <sub>2</sub>               | 73.88 | 71.00 |
| TiO <sub>2</sub>               | .41   | .58   |
| Al <sub>2</sub> O <sub>3</sub> | 13.89 | 13.76 |
| Fe <sub>2</sub> O <sub>3</sub> | 1.38  | 2.57  |
| MnO                            | .83   | .84   |
| HgO                            | .01   | .59   |
| CaO                            | .96   | 1.19  |
| Na <sub>2</sub> O              | 4.84  | 4.48  |
| K <sub>2</sub> O               | 2.61  | 3.75  |
| LOI                            | 1.30  | 1.10  |

|       |       |       |
|-------|-------|-------|
| TOTAL | 98.43 | 98.90 |
|-------|-------|-------|

## TRACE ELEMENTS (PPM)

|                   |      |      |
|-------------------|------|------|
| Zr                | 174  | 165  |
| Sr                | 201  | 191  |
| Rb                | 31   | 50   |
| Zn                | 24   | 52   |
| Cu                | 1    | 1    |
| Ba                | 1259 | 1247 |
| CuS               | 1    | 1    |
| PbS               | 21   | 12   |
| ZnS               | 11   | 34   |
| BaSO <sub>4</sub> |      |      |
| Pb                | 21   | 2    |
| Ni                | 3    | 3    |
| Co                | 42   | 28   |
| Cr                | 8    | 32   |
| V                 | 20   | 48   |
| Hg                |      |      |
| Ag                |      |      |
| Nb                |      |      |

## CLEMENTINE AREA RHYOLITE FLOWS

## MAJOR ELEMENTS (WT%)

| SAMPLE | 1903 609 | 23491065 | 2866 516 |
|--------|----------|----------|----------|
| SiO2   | 69.90    | 73.50    | 73.90    |
| TiO2   | .16      | .18      | .14      |
| Al2O3  | 14.48    | 13.26    | 13.42    |
| Fe2O3  | 2.19     | 1.50     | .97      |
| MnO    | .08      | .04      | .02      |
| MgO    | .61      | .49      | .21      |
| CaO    | .96      | 1.59     | .20      |
| Na2O   | 3.74     | 4.49     | 3.09     |
| K2O    | 4.78     | 1.70     | 5.53     |
| LOI    | 1.60     | 2.70     | 1.00     |
| TOTAL  | 98.50    | 99.53    | 98.40    |

## TRACE ELEMENTS (PPM)

|       |      |     |      |
|-------|------|-----|------|
| Zr    | 83   | 89  | 132  |
| Sr    | 125  | 150 | 79   |
| Rb    | 85   | 32  | 100  |
| Zn    | 42   | 22  | 7    |
| Cu    | 13   | 1   | 1    |
| Ba    | 1553 | 568 | 1702 |
| CuS   | 8    | 3   | 1    |
| PbS   | 19   | 15  | 20   |
| ZnS   | 16   | 20  | 5    |
| BaSO4 | 0    | 0   | 0    |
| Pb    | 27   | 13  | 3    |
| Ni    | 7    | 3   | 4    |
| Co    | 17   | 9   | 26   |
| Cr    | 9    | 10  | 0    |
| V     | 10   | 10  | 10   |
| Hg    |      |     |      |
| Ag    |      |     |      |
| Nb    | 6    | 0   |      |

## ORIENTAL ORE HORIZON SEQUENCE 'WHITE RHYOLITE'

## MAJOR ELEMENTS (WT%)

SAMPLE 1083 586 17301203

|       |       |       |
|-------|-------|-------|
| SiO2  | 73.30 | 75.60 |
| TiO2  | .39   | .22   |
| Al2O3 | 12.91 | 10.29 |
| Fe2O3 | .03   | .01   |
| MnO   | .01   | .01   |
| MgO   | .60   | .01   |
| CaO   | .44   | .76   |
| Na2O  | 1.52  | .36   |
| K2O   | 7.64  | 7.93  |
| LOI   | 1.60  | 1.60  |

|       |       |       |
|-------|-------|-------|
| TOTAL | 98.44 | 96.79 |
|-------|-------|-------|

## TRACE ELEMENTS (PPM)

|       |      |      |
|-------|------|------|
| Zr    | 166  | 147  |
| Sr    | 49   | 119  |
| Rb    | 62   | 63   |
| Zn    | 19   | 455  |
| Cu    | 1    | 1    |
| Ba    | 2982 | 4639 |
| CuS   | 2    | 5    |
| PbS   | 6    | 69   |
| ZnS   | 22   | 460  |
| BaSO4 |      |      |
| Pb    | 3    | 32   |
| Ni    | 1    | 2    |
| Co    | 27   | 42   |
| Cr    | 0    | 0    |
| V     | 10   | 15   |
| Hg    |      |      |
| Ag    |      |      |
| Nb    |      |      |

## LUCK STRIKE ORE HORIZON SEQUENCE - PYRITIC SILTSTONE

## MAJOR ELEMENTS (WT%)

| SAMPLE                         | 02020150 | 02020175 | 07921080 | 07921093 | 07921100 |
|--------------------------------|----------|----------|----------|----------|----------|
| SiO <sub>2</sub>               | 61.60    | 53.90    | 68.40    | 64.60    | 62.00    |
| TiO <sub>2</sub>               | .30      | .28      | .41      | .49      | .44      |
| Al <sub>2</sub> O <sub>3</sub> | 11.45    | 11.05    | 13.60    | 16.44    | 15.69    |
| Fe <sub>2</sub> O <sub>3</sub> | 2.46     | 2.37     | 4.22     | 3.94     | 2.90     |
| MnO                            | .28      | .33      | .09      | .08      | .07      |
| MgO                            | 15.11    | 20.51    | 1.42     | 1.19     | 1.61     |
| CaO                            | .28      | 1.86     | 1.23     | 2.60     | 2.73     |
| Na <sub>2</sub> O              | .06      | .06      | 2.65     | 3.30     | 2.65     |
| K <sub>2</sub> O               | .74      | .05      | 3.32     | 3.16     | 2.38     |
| LOI                            | 7.40     | 10.10    | 3.00     | 3.50     | 3.50     |
| TOTAL                          | 99.68    | 100.51   | 98.42    | 99.30    | 93.97    |

## TRACE ELEMENTS (PPM)

|                   |     |     |      |      |      |
|-------------------|-----|-----|------|------|------|
| Zr                | 131 | 109 | 128  | 126  | 148  |
| Sr                | 0   | 0   | 176  | 280  | 302  |
| Rb                | 21  | 2   | 44   | 40   | 43   |
| Zn                | 320 | 480 | 2900 | 345  | 1060 |
| Cu                | 6   | 7   | 46   | 36   | 32   |
| Ba                | 511 | 146 | 2013 | 2670 | 2209 |
| CuS               | 2   | 1   | 40   | 32   | 35   |
| PbS               | 26  | 35  | 530  | 56   | 530  |
| ZnS               | 33  | 102 | 1070 | 410  | 2900 |
| BaSO <sub>4</sub> | 0   |     |      |      |      |
| Pb                | 48  | 132 | 580  | 46   | 171  |
| Ni                | 2   | 2   | 5    | 6    | 4    |
| Co                | 3   | 3   | 8    | 9    | 6    |
| Cr                | 25  | 24  | 27   | 28   | 29   |
| V                 | 33  | 22  | 38   | 54   | 73   |
| Hg                | 20  | 30  | 73   | 59   | 106  |
| Ag                | 8   | 9   | 5    | 0    | 5    |
| Nb                |     |     |      |      |      |

## LUCK STRIKE ORE HORIZON SEQUENCE - PYRITIC SILTSTONE

## MAJOR ELEMENTS (WT%)

| SAMPLE | 07921117 | 07921140 | 07921190 | 07921240 | 07921300 |
|--------|----------|----------|----------|----------|----------|
| S102   | 68.40    | 70.50    | 69.00    | 63.50    | 62.70    |
| T102   | .48      | .18      | .20      | .20      | .20      |
| Al2O3  | 14.67    | 12.81    | 12.15    | 12.78    | 12.36    |
| Fe2O3  | 2.92     | 2.10     | 4.13     | 4.00     | 5.67     |
| MnO    | .07      | .08      | .10      | .22      | .09      |
| HgO    | .97      | .56      | .67      | .79      | 1.49     |
| CaO    | 1.95     | 2.72     | 2.43     | 2.44     | 2.33     |
| Na2O   | 4.24     | 2.46     | 3.74     | 2.67     | 2.51     |
| K2O    | 1.64     | 3.31     | 1.32     | 3.11     | 2.79     |
| LOI    | 2.00     | 3.10     | 2.50     | 2.80     | 4.20     |
| TOTAL  | 97.34    | 96.82    | 96.24    | 92.51    | 94.34    |

## TRACE ELEMENTS (PPM)

|       |      |      |     |      |      |
|-------|------|------|-----|------|------|
| Zr    | 109  | 93   | 115 | 97   | 102  |
| Sr    | 308  | 243  | 280 | 209  | 158  |
| Rb    | 29   | 42   | 22  | 38   | 36   |
| Zn    | 53   | 270  | 123 | 57   | 111  |
| Cu    | 8    | 15   | 11  | 17   | 29   |
| Ba    | 2111 | 6904 | 833 | 1586 | 1298 |
| CuS   | 6    | 11   | 5   | 11   | 18   |
| PbS   | 47   | 207  | 68  | 42   | 45   |
| ZnS   | 20   | 510  | 107 | 69   | 78   |
| BaSO4 |      | 3900 |     |      |      |
| Pb    | 29   | 128  | 44  | 25   | 33   |
| Ni    | 3    | 4    | 4   | 4    | 8    |
| Co    | 4    | 6    | 7   | 7    | 9    |
| Cr    | 39   | 22   | 28  | 28   | 37   |
| V     | 82   | 47   | 41  | 30   | 62   |
| Hg    | 16   | 432  | 41  | 31   | 55   |
| Ag    | 8    | 13   | 11  | 5    | 5    |
| Nb    |      |      |     |      |      |

POOR COPY  
COPIE DE QUALITEE INFERIEURE

3-75

## LUCK STRIKE ORE HORIZON SEQUENCE - PYRITIC SILTSTONE

## MAJOR ELEMENTS (WT%)

| SAMPLE | 07921360 | 07921400 | 08692550 | 08692590 | 14930950 |
|--------|----------|----------|----------|----------|----------|
| SiO2   | 62.50    | 64.30    | 70.50    | 67.50    | 65.40    |
| TiO2   | .34      | .32      | .36      | .41      | .44      |
| Al2O3  | 13.45    | 13.32    | 15.33    | 13.56    | 15.30    |
| Fe2O3  | 4.68     | 4.29     | 3.47     | 4.78     | 4.30     |
| MnO    | .10      | .12      | .02      | .04      | .07      |
| MgO    | 1.90     | 2.30     | .83      | 1.86     | 1.21     |
| CaO    | 2.37     | 3.46     | .50      | 2.43     | 3.26     |
| Na2O   | 2.55     | 2.38     | 1.86     | 4.00     | 4.60     |
| K2O    | 2.30     | 2.25     | 3.93     | 1.24     | 1.64     |
| LOI    | 4.30     | 5.40     | 3.70     | 3.80     | 3.80     |
| TOTAL  | 94.49    | 98.14    | 100.50   | 99.62    | 100.02   |

## TRACE ELEMENTS (PPM)

|       |      |      |     |     |      |
|-------|------|------|-----|-----|------|
| Zr    | 98   | 97   | 140 | 176 | 175  |
| Sr    | 231  | 257  | 0   | 293 | 380  |
| Rb    | 35   | 33   | 63  | 21  | 31   |
| Zn    | 78   | 98   | 122 | 135 | 49   |
| Cu    | 31   | 36   | 20  | 27  | 7    |
| Ba    | 1943 | 4467 | 936 | 622 | 1375 |
| CuS   | 11   | 35   | 34  | 23  | 3    |
| PbS   | 41   | 6    | 56  | 124 | 27   |
| ZnS   | 41   | 124  | 55  | 58  | 38   |
| BaSO4 |      |      |     |     | 62   |
| Pb    | 26   | 35   | 69  | 110 | 10   |
| Ni    | 4    | 5    | 7   | 9   | 4    |
| Co    | 7    | 7    | 2   | 2   | 5    |
| Cr    | 28   | 34   | 17  | 17  | 28   |
| V     | 68   | 65   | 81  | 60  | 49   |
| Hg    | 45   | 51   | 28  | 27  | 33   |
| Ag    | 3    | 3    | 5   | 3   | 0    |
| Nb    |      |      |     |     |      |

POOR COPY  
COPIE DE QUALITEE INFERIEURE



## LUCK STRIKE ORE HORIZON SEQUENCE - PYRITIC SILTSTONE

## MAJOR ELEMENTS (WT%)

| SAMPLE | 14930962 | 27040006 | 27040950 |
|--------|----------|----------|----------|
| SiO2   | 68.00    | 66.60    | 67.80    |
| TiO2   | .40      | .46      | .38      |
| Al2O3  | 13.46    | 13.66    | 13.36    |
| Fe2O3  | 4.33     | 4.00     | 5.08     |
| MnO    | .06      | .06      | .13      |
| MgO    | .88      | .84      | .90      |
| CaO    | 2.54     | 1.98     | 2.00     |
| Na2O   | 4.10     | 3.35     | 4.92     |
| K2O    | 1.41     | 2.95     | 1.07     |
| LOI    | 3.00     | 2.80     | 2.30     |
| TOTAL  | 98.18    | 96.70    | 97.84    |

## TRACE ELEMENTS (PPM)

|       |     |      |      |
|-------|-----|------|------|
| Zr    | 177 | 155  | 152  |
| Sr    | 304 | 238  | 281  |
| Rb    | 32  | 55   | 23   |
| Zn    | 109 | 115  | 125  |
| Cu    | 12  | 12   | 12   |
| Ba    | 71  | 1766 | 1097 |
| CuS   | 8   | 3    | 2    |
| PbS   | 84  | 31   | 33   |
| ZnS   | 105 | 96   | 64   |
| BaSO4 |     |      |      |
| Pb    | 42  | 19   | 42   |
| Ni    | 5   | 5    | 4    |
| Co    | 6   | 21   | 7    |
| Cr    | 36  | 28   | 30   |
| V     | 71  | 52   | 39   |
| Hg    | 56  | 46   | 58   |
| Ag    | 2   | 0    | 0    |
| Nb    |     |      |      |

POOR COPY  
COPIE DE QUALITEE INFERIEURE

## LUCK STRIKE ORE HORIZON SEQUENCE - FOOTWALL DACITIC TUFFS

## MAJOR ELEMENTS (WT%)

| SAMPLE                         | 02020100 | 02020110 | 02020122 | 19170000 | 19170025 |
|--------------------------------|----------|----------|----------|----------|----------|
| SiO <sub>2</sub>               | 77.90    | 68.70    | 74.40    | 74.30    | 72.30    |
| TiO <sub>2</sub>               | .13      | .23      | .15      | .29      | .31      |
| Al <sub>2</sub> O <sub>3</sub> | 9.60     | 16.44    | 11.93    | 11.32    | 13.28    |
| Fe <sub>2</sub> O <sub>3</sub> | 1.76     | 1.70     | 1.60     | 2.45     | 3.26     |
| MnO                            | .00      | .05      | .07      | .10      | .12      |
| MgO                            | 4.10     | 3.51     | 2.31     | .42      | .53      |
| CaO                            | .97      | .68      | 1.59     | 1.56     | 1.05     |
| Na <sub>2</sub> O              | .64      | .91      | 1.75     | 4.35     | 5.56     |
| K <sub>2</sub> O               | 1.14     | 3.67     | 2.74     | 1.53     | 1.99     |
| LOI                            | 4.10     | 4.80     | 4.20     | 1.10     | .60      |
| TOTAL                          | 100.42   | 100.69   | 99.74    | 97.42    | 99.00    |

## TRACE ELEMENTS (PPM)

|                   |     |      |      |      |      |
|-------------------|-----|------|------|------|------|
| Zr                | 44  | 107  | 51   | 113  | 132  |
| Sr                | 43  | 66   | 0    | 16   | 93   |
| Rb                | 28  | 77   | 53   | 24   | 32   |
| Zn                | 650 | 1060 | 460  | 37   | 37   |
| Cu                | 5   | 12   | 4    | 5    | 1    |
| Ba                | 984 | 2009 | 1831 | 1397 | 1154 |
| CuS               | 1   | 1    | 1    | 2    | 0    |
| PbS               | 36  | 55   | 50   | 60   | 15   |
| ZnS               | 62  | 67   | 137  | 14   | 22   |
| BaSO <sub>4</sub> |     |      |      |      | 6    |
| Pb                | 76  | 163  | 100  | 29   | 7    |
| Ni                | 1   | 1    | 1    | 2    | 3    |
| Co                | 1   | 4    | 2    | 2    | 3    |
| Cr                | 22  | 28   | 20   | 24   | 26   |
| V                 | 8   | 20   | 6    | 17   | 29   |
| Hg                | 20  | 31   | 34   | 16   | 18   |
| Ag                | 5   | 10   | 7    | 0    | 0    |
| Nb                |     |      |      |      |      |

POOR COPY  
COPIE DE QUALITE INFERIEURE

## LUCK STRIKE ORE HORIZON SEQUENCE - FOOTWALL DACITIC TUFFS

## MAJOR ELEMENTS (WT%)

| SAMPLE                         | 19170050 | 27040050 | 27040225 | 27040275 | 27040525 |
|--------------------------------|----------|----------|----------|----------|----------|
| SiO <sub>2</sub>               | 73.10    | 75.70    | 67.80    | 64.90    | 72.30    |
| TiO <sub>2</sub>               | .26      | .26      | .27      | .20      | .23      |
| Al <sub>2</sub> O <sub>3</sub> | 11.60    | 11.63    | 14.05    | 13.36    | 13.36    |
| Fe <sub>2</sub> O <sub>3</sub> | 2.72     | 2.21     | 2.42     | 2.69     | 2.06     |
| MnO                            | .08      | .05      | .06      | .09      | .08      |
| MgO                            | .42      | .19      | 1.82     | 2.41     | 1.07     |
| CaO                            | 1.24     | .63      | 2.52     | 4.95     | 3.04     |
| Na <sub>2</sub> O              | 4.84     | 5.02     | 2.06     | .86      | 1.93     |
| K <sub>2</sub> O               | 2.00     | 1.56     | 2.83     | 2.76     | .68      |
| LOI                            | 1.00     | .50      | 4.10     | 7.00     | 3.90     |
| TOTAL                          | 97.23    | 97.75    | 97.93    | 99.22    | 98.65    |

## TRACE ELEMENTS (PPM)

|                   |      |     |      |      |     |
|-------------------|------|-----|------|------|-----|
| Zr                | 113  | 144 | 120  | 113  | 114 |
| Sr                | 116  | 84  | 183  | 249  | 101 |
| Rb                | 33   | 29  | 72   | 70   | 95  |
| Zn                | 66   | 34  | 21   | 40   | 30  |
| Cu                | 3    | 1   | 2    | 8    | 3   |
| Ba                | 1862 | 885 | 1820 | 3454 | 733 |
| CuS               | 1    | 0   | 1    | 8    | 2   |
| PbS               | 19   | 12  | 17   | 131  | 38  |
| ZnS               | 27   | 8   | 10   | 13   | 7   |
| BaSO <sub>4</sub> |      | 6   |      | 500  |     |
| Pb                | 10   | 7   | 7    | 68   | 17  |
| Ni                | 1    | 2   | 3    | 3    | 3   |
| Co                | 3    | 8   | 23   | 2    | 4   |
| Cr                | 25   | 25  | 25   | 20   | 21  |
| V                 | 20   | 0   | 31   | 44   | 37  |
| Hg                | 20   | 8   | 8    | 12   | 18  |
| Ag                | 0    | 0   | 0    | 1    | 0   |
| Nb                |      |     |      |      |     |

POOR COPY  
COPIE DE QUALITE INFERIEURE

## LUCK STRIKE ORE HORIZON SEQUENCE - FOOTWALL DACITIC TUFFS

## MAJOR ELEMENTS (WT%)

| SAMPLE                         | 27040575 | 27040650 | 27040725 | 27040800 | 27040875 |
|--------------------------------|----------|----------|----------|----------|----------|
| SiO <sub>2</sub>               | 71.60    | 72.90    | 70.80    | 69.70    | 73.50    |
| TiO <sub>2</sub>               | .20      | .20      | .24      | .23      | .24      |
| Al <sub>2</sub> O <sub>3</sub> | 11.75    | 11.26    | 13.74    | 13.62    | 13.28    |
| Fe <sub>2</sub> O <sub>3</sub> | 2.26     | 2.40     | 2.82     | 2.90     | 2.53     |
| MnO                            | .13      | .11      | .13      | .12      | .10      |
| MgO                            | .86      | 1.12     | 1.04     | 1.84     | 1.38     |
| CaO                            | 2.36     | 2.45     | 2.09     | 1.42     | 1.50     |
| Na <sub>2</sub> O              | 4.84     | 2.53     | 2.74     | 2.47     | 3.35     |
| K <sub>2</sub> O               | 3.50     | 2.43     | 3.07     | 3.54     | 1.89     |
| LOI                            | 2.30     | 2.80     | 1.60     | 2.20     | 2.00     |
| TOTAL                          | 99.80    | 98.20    | 98.27    | 98.04    | 99.77    |

## TRACE ELEMENTS (PPM)

|                   |     |     |      |      |     |
|-------------------|-----|-----|------|------|-----|
| Zr                | 118 | 86  | 129  | 115  | 93  |
| Sr                | 407 | 224 | 290  | 193  | 222 |
| Rb                | 19  | 57  | 77   | 87   | 48  |
| Zn                | 29  | 30  | 21   | 28   | 22  |
| Cu                | 9   | 5   | 1    | 8    | 6   |
| Ba                | 600 | 096 | 1568 | 1938 | 091 |
| CuS               | 5   | 2   | 1    | 3    | 4   |
| PbS               | 21  | 31  | 9    | 8    | 8   |
| ZnS               | 10  | 12  | 3    | 19   | 12  |
| BaSO <sub>4</sub> |     | 10  |      |      | 2   |
| Pb                | 14  | 19  | 11   | 4    | 11  |
| Ni                | 5   | 4   | 2    | 3    | 3   |
| Co                | 4   | 4   | 3    | 3    | 2   |
| Cr                | 32  | 25  | 23   | 24   | 22  |
| V                 | 32  | 32  | 8    | 33   | 34  |
| Hg                | 16  | 14  | 12   | 12   | 12  |
| Ag                | 0   | 0   | 0    | 0    | 0   |
| Nb                |     |     |      |      |     |

POOR COPY  
COPIE DE QUALITE INFERIEURE

## LUCK STRIKE ORE HORIZON SEQUENCE - DACITIC TUFFS IN PROXIMITY TO ORE

## MAJOR ELEMENTS (WT%)

| SAMPLE | JGT75 44 | 01750025 | 01750125 | 02020070 | 02020075 |
|--------|----------|----------|----------|----------|----------|
| SiO2   | 70.30    | 58.00    | 63.50    | 65.60    | 76.80    |
| TiO2   | .44      | .34      | .25      | .24      | .10      |
| Al2O3  | 14.66    | 20.28    | 17.15    | 18.38    | 10.50    |
| Fe2O3  | 2.76     | 3.18     | 2.92     | 1.00     | 1.66     |
| MnO    | .09      | .07      | .06      | .03      | .06      |
| MgO    | 1.92     | 3.14     | 2.73     | 2.50     | 3.92     |
| CaO    | 2.55     | 1.42     | 1.79     | 1.05     | .68      |
| Na2O   | 1.67     | 1.93     | 1.18     | 1.31     | .73      |
| K2O    | 2.05     | 5.51     | 5.11     | 3.84     | 1.47     |
| LOI    | 3.00     | 5.30     | 4.00     | 5.50     | 4.10     |
| TOTAL  | 99.44    | 99.17    | 98.69    | 99.45    | 100.10   |

## TRACE ELEMENTS (PPM)

|       |      |      |      |      |      |
|-------|------|------|------|------|------|
| Zr    | 236  | 164  | 130  | 71   | 41   |
| Sr    | 429  | 136  | 279  | 297  | 34   |
| Rb    | 71   | 130  | 131  | 21   | 33   |
| Zn    | 161  | 53   | 111  | 620  | 730  |
| Cu    | 3    | 5    | 11   | 38   | 6    |
| Ba    | 2476 | 1820 | 1241 | 1049 | 1132 |
| CuS   | 2    | 0    | 1    | 9    | 1    |
| PbS   | 17   | 31   | 24   | 86   | 19   |
| ZnS   | 78   | 7    | 31   | 131  | 48   |
| BaSO4 | 6    | 2    | 3    | 18   |      |
| Pb    | 18   | 25   | 27   | 410  | 56   |
| Ni    | 3    | 2    | 2    | 1    | 2    |
| Co    | 18   | 2    | 3    | 3    | 2    |
| Cr    | 10   | 19   | 19   | 211  | 20   |
| V     | 35   | 17   | 4    | 22   | 16   |
| Hg    |      | 15   | 23   | 27   | 19   |
| Ag    |      | 0    | 2    | 11   | 6    |
| Nb    | 6    |      |      |      |      |

POOR COPY  
COPIE DE QUALITEE INFERIEURE

## LUCK STRIKE ORE HORIZON SEQUENCE - DACITIC TUFFS IN PROXIMITY TO ORE

## MAJOR ELEMENTS (WT%)

| SAMPLE                         | 02020092 | 08692465 | 14930825 | 14930843 | 14930855 |
|--------------------------------|----------|----------|----------|----------|----------|
| SiO <sub>2</sub>               | 67.80    | 78.00    | 73.30    | 70.60    | 62.10    |
| TiO <sub>2</sub>               | .16      | .16      | .17      | .20      | .44      |
| Al <sub>2</sub> O <sub>3</sub> | 14.46    | 10.13    | 13.51    | 14.61    | 17.35    |
| Fe <sub>2</sub> O <sub>3</sub> | 2.69     | .84      | 1.45     | 1.86     | 5.37     |
| MnO                            | .11      | .03      | .06      | .08      | .08      |
| HgO                            | 5.18     | .68      | .92      | 1.72     | 2.65     |
| CaO                            | 1.51     | 1.65     | 2.37     | 2.31     | 2.52     |
| Na <sub>2</sub> O              | .95      | 4.89     | 2.92     | 1.81     | 2.52     |
| K <sub>2</sub> O               | 2.08     | .31      | 2.13     | 2.50     | 2.70     |
| LOI                            | 5.60     | 1.80     | 2.20     | 2.50     | 3.00     |
| TOTAL                          | 100.54   | 98.49    | 99.03    | 98.19    | 98.73    |

## TRACE ELEMENTS (PPM)

|                   |      |      |      |      |      |
|-------------------|------|------|------|------|------|
| Zr                | 7    | 138  | 132  | 151  | 153  |
| Sr                | 150  | 286  | 460  | 442  | 293  |
| Rb                | 43   | 1    | 70   | 71   | 64   |
| Zn                | 540  | 38   | 99   | 43   | 2600 |
| Cu                | 6    | 4    | 10   | 4    | 16   |
| Ba                | 2434 | 3040 | 3602 | 6614 | 2830 |
| CuS               | 2    | 1    | 3    | 1    | 15   |
| PbS               | 44   | 25   | 39   | 98   | 155  |
| ZnS               | 60   | 11   | 5    | 10   | 1690 |
| BaSO <sub>4</sub> | 7    |      | 9    | 12   |      |
| Pb                | 100  | 16   | 50   | 61   | 91   |
| Ni                | 2    | 2    | 2    | 2    | 4    |
| Co                | 3    | 2    | 2    | 3    | 6    |
| Cr                | 21   | 9    | 21   | 25   | 43   |
| V                 | 6    | 18   | 6    | 0    | 103  |
| Hg                | 17   | 4    | 12   | 16   | 24   |
| Ag                | 14   | 2    | 3    | 3    | 2    |
| Nb                |      |      |      |      |      |

POOR COPY  
COPIE DE QUALITEE INFERIEURE

## LUCK STRIKE ORE HORIZON SEQUENCE - DACITIC TUFFS IN PROXIMITY TO ORE

## MAJOR ELEMENTS (WT%)

SAMPLE 14930929 19170075 1917.118

|       |       |       |       |
|-------|-------|-------|-------|
| SiO2  | 66.30 | 70.70 | 68.00 |
| TiO2  | .40   | .36   | .24   |
| Al2O3 | 14.78 | 13.52 | 15.76 |
| Fe2O3 | 3.54  | 3.56  | 2.09  |
| MnO   | .07   | .11   | .06   |
| MgO   | 3.00  | 1.54  | 1.84  |
| CaO   | 3.41  | 2.75  | 2.27  |
| Na2O  | 1.61  | 4.39  | 1.67  |
| K2O   | 2.07  | 1.34  | 3.70  |
| LOI   | 5.20  | 1.30  | 3.50  |

|       |        |       |       |
|-------|--------|-------|-------|
| TOTAL | 101.18 | 99.57 | 99.13 |
|-------|--------|-------|-------|

## TRACE ELEMENTS (PPM)

|       |      |      |      |
|-------|------|------|------|
| Zr    | 163  | 162  | 116  |
| Sr    | 156  | 456  | 255  |
| Rb    | 62   | 23   | 104  |
| Zn    | 92   | 205  | 27   |
| Cu    | 12   | 11   | 2    |
| Ba    | 1434 | 1829 | 1580 |
| CuS   | 7    | 4    | 1    |
| PbS   | 132  | 87   | 8    |
| ZnS   | 52   | 122  | 14   |
| BaSO4 |      |      |      |
| Pb    | 78   | 59   | 6    |
| Ni    | 5    | 10   | 2    |
| Co    | 5    | 6    | 1    |
| Cr    | 28   | 37   | 21   |
| V     | 41   | 64   | 7    |
| Hg    | 33   | 18   | 12   |
| Ag    | 2    | 0    | 1    |
| Nb    |      |      |      |

POOR COPY  
COPIE DE QUALITEE INFERIEURE

## ORIENTAL ORE HORIZON SEQUENCE - GRANITE BOULDERS

## MAJOR ELEMENTS (WT%)

| SAMPLE                         | 1147 118 | 1202 104 | 1217 192 | 1261 230 | 1273 232 |
|--------------------------------|----------|----------|----------|----------|----------|
| SiO <sub>2</sub>               | 72.30    | 71.00    | 70.50    | 73.70    | 76.30    |
| TiO <sub>2</sub>               | .24      | .24      | .30      | .52      | .20      |
| Al <sub>2</sub> O <sub>3</sub> | 13.04    | 13.22    | 12.62    | 12.45    | 10.92    |
| Fe <sub>2</sub> O <sub>3</sub> | 2.39     | 2.11     | 1.64     | 1.98     | .13      |
| MnO                            | .08      | .05      | .05      | .08      | 0.00     |
| MgO                            | 1.25     | .84      | 1.04     | 1.24     | .18      |
| CaO                            | 1.23     | 1.86     | 1.90     | .96      | .42      |
| Na <sub>2</sub> O              | 4.16     | 4.15     | 3.33     | 4.90     | 3.57     |
| K <sub>2</sub> O               | 1.20     | 2.45     | 2.15     | .53      | 1.36     |
| LOI                            | 2.20     | 1.60     | 2.70     | 1.80     | 2.40     |
| TOTAL                          | 98.09    | 97.52    | 96.23    | 98.16    | 95.48    |

## TRACE ELEMENTS (PPM)

|                   |     |      |      |     |      |
|-------------------|-----|------|------|-----|------|
| Zr                | 116 | 85   | 121  | 148 | 116  |
| Sr                | 173 | 239  | 213  | 140 | 96   |
| Rb                | 15  | 27   | 30   | 7   | 10   |
| Zn                | 73  | 38   | 47   | 890 | 3200 |
| Cu                | 3   | 12   | 4    | 83  | 200  |
| Ba                | 929 | 1090 | 1551 | 958 | 3304 |
| CuS               | 4   | 15   | 4    | 79  | 187  |
| PbS               | 13  | 16   | 18   | 570 | 2800 |
| ZnS               | 15  | 21   | 11   | 590 | 2300 |
| BaSO <sub>4</sub> | 104 | 0    | 86   | 440 | 3400 |
| Pb                | 5   | 4    | 15   | 680 | 2150 |
| Ni                | 7   | 4    | 4    | 3   | 3    |
| Co                | 40  | 38   | 37   | 32  | 28   |
| Cr                | 10  | 9    | 8    | 11  | 10   |
| V                 | 10  | 10   | 10   | 30  | 10   |
| Hg                |     |      |      |     |      |
| Ag                |     |      |      |     |      |
| Nb                | 7   | 6    | 4    | 8   | 0    |

POOR COPY  
COPIE DE QUALITE INFERIEURE



## LUCK STRIKE ORE HORIZON SEQUENCE - ORE HORIZON BARREN BRECCIA

## MAJOR ELEMENTS (WT%)

| SAMPLE                         | 07921050 | 07921065 | 19170092 | 19170100 |
|--------------------------------|----------|----------|----------|----------|
| SiO <sub>2</sub>               | 58.40    | 61.20    | 63.20    | 63.80    |
| TiO <sub>2</sub>               | .20      | .50      | .20      | .21      |
| Al <sub>2</sub> O <sub>3</sub> | 13.51    | 14.63    | 11.53    | 11.78    |
| Fe <sub>2</sub> O <sub>3</sub> | 6.18     | 5.07     | 2.54     | 2.70     |
| MnO                            | .14      | .11      | .07      | .08      |
| MgO                            | 2.01     | 2.07     | 1.71     | 1.75     |
| CaO                            | 2.96     | 1.33     | 2.40     | 2.40     |
| Na <sub>2</sub> O              | 2.63     | 3.15     | 1.80     | 1.87     |
| K <sub>2</sub> O               | 2.36     | 2.51     | 2.12     | 2.21     |
| LOI                            | 6.10     | 4.40     | 6.10     | 6.10     |
| TOTAL                          | 94.49    | 94.97    | 91.67    | 92.90    |

## TRACE ELEMENTS (PPM)

|                   |      |      |     |      |
|-------------------|------|------|-----|------|
| Zr                | 82   | 108  | 139 | 135  |
| Sr                | 368  | 632  | 731 | 772  |
| Rb                | 36   | 33   | 47  | 45   |
| Zn                | 2700 | 1420 | 540 | 630  |
| Cu                | 1490 | 480  | 280 | 134  |
| Ba                | 9999 | 9999 | 999 | 9999 |
| CuS               | 1570 | 495  | 340 | 110  |
| PbS               | 1550 | 490  | 450 | 530  |
| ZnS               | 2800 | 1055 | 830 | 1040 |
| BaSO <sub>4</sub> | 999  |      | 999 |      |
| Pb                | 1010 | 270  | 670 | 285  |
| Ni                | 5    | 5    | 6   | 5    |
| Co                | 10   | 7    | 3   | 4    |
| Cr                | 37   | 32   | 31  | 30   |
| V                 | 84   | 94   | 53  | 28   |
| Hg                | 149  | 45   | 105 | 98   |
| Ag                | 108  | 25   | 20  | 57   |
| Nb                |      |      |     |      |

POOR COPY  
COPIE DE QUALITE INFERIEURE

## LUCKY STRIKE AREA TRANSPORTED ORE

## MAJOR ELEMENTS (WT%)

| SAMPLE                         | 02020025 | 2020050 | 07921075 | 00692480 | 00692520 |
|--------------------------------|----------|---------|----------|----------|----------|
| SiO <sub>2</sub>               | 1.20     | 1.20    | 1.90     | 20.30    | 25.50    |
| TiO <sub>2</sub>               | 0.00     | 0.00    | .01      | .19      | .48      |
| Al <sub>2</sub> O <sub>3</sub> | .36      | .34     | .31      | 5.98     | 8.21     |
| Fe <sub>2</sub> O <sub>3</sub> | 4.35     | 2.14    | 11.16    | 3.94     | 7.41     |
| MnO                            | .03      | .02     | .03      | .08      | .18      |
| HgO                            | .20      | .33     | .04      | 2.50     | 4.09     |
| CaO                            | 2.60     | 1.49    | .03      | 1.42     | 4.71     |
| Na <sub>2</sub> O              | 0.00     | 0.00    | 0.00     | .39      | 1.46     |
| K <sub>2</sub> O               | .07      | 0.00    | .05      | 1.04     | .40      |
| LOI                            | 5.80     | 4.20    | 14.90    | 5.90     | 8.10     |
| TOTAL                          | 14.61    | 9.72    | 28.43    | 41.74    | 60.54    |

## TRACE ELEMENTS (PPM)

|                   |      |      |      |      |      |
|-------------------|------|------|------|------|------|
| Zr                | 154  | 203  | 75   | 218  | 94   |
| Sr                | 1190 | 1527 | 74   | 1000 | 185  |
| Rb                | 27   | 23   | 40   | 15   | 35   |
| Zn                |      |      |      |      |      |
| Cu                | 132  | 61   | 295  | 57   | 155  |
| Ba                | 0    | 0    | 3    | 500  | 1    |
| CuS               | 6000 | 3050 | 2000 | 4800 | 1600 |
| PbS               | 41   | 37   | 78   | 24   | 64   |
| ZnS               | 180  | 245  | 12   | 46   | 9    |
| BaSO <sub>4</sub> | 1    | 1    | 2501 | 120  | 952  |
| Pb                | 9000 | 2300 | 8000 | 9200 | 6000 |
| Ni                | 3    | 5    | 9    | 12   | 53   |
| Co                | 2    | 2    | 3    | 2    | 2    |
| Cr                | 29   | 28   | 61   | 0    | 109  |
| V                 | 16   | 20   | 15   | 98   | 129  |
| Hg                | 2455 | 2820 | 1545 | 3222 | 1967 |
| Ag                | 970  | 820  | 1100 | 820  | 410  |
| Nb                |      |      |      |      |      |

POOR COPY  
COPIE DE QUALITEE INFERIEURE

## LUCKY STRIKE AREA TRANSPORTED ORE

## MAJOR ELEMENTS (Wt%)

SAMPLE, 14930848

|                                |       |
|--------------------------------|-------|
| SiO <sub>2</sub>               | 18.90 |
| TiO <sub>2</sub>               | .12   |
| Al <sub>2</sub> O <sub>3</sub> | 3.94  |
| Fe <sub>2</sub> O <sub>3</sub> | 5.47  |
| MnO                            | .05   |
| MgO                            | .44   |
| CaO                            | .53   |
| Na <sub>2</sub> O              | .88   |
| K <sub>2</sub> O               | .46   |
| LOI                            | 11.20 |

|       |       |
|-------|-------|
| TOTAL | 41.99 |
|-------|-------|

## TRACE ELEMENTS (PPM)

|                   |      |
|-------------------|------|
| Zr                | 131  |
| Sr                | 509  |
| Rb                | 40   |
| Zn                |      |
| Cu                | 107  |
| Ba                | 0    |
| CuS               | 9500 |
| PbS               | 84   |
| ZnS               | 96   |
| BaSO <sub>4</sub> | 4572 |
| Pb                | 2500 |
| Ni                | 5    |
| Co                | 3    |
| Cr                | 33   |
| V                 | 33   |
| Hg                | 3466 |
| Ag                | 1360 |
| Nb                |      |

POOR COPY  
COPIE DE QUALITEE INFERIEURE

## LUCKY STRIKE ORE HORIZON SEQUENCE - CLEMENTINE DACITIC TUFFS

## MAJOR ELEMENTS (WT%)

| SAMPLE                         | 1258 443 | 2866 325 | 2869 938 | 2879 437 | 2882 298 |
|--------------------------------|----------|----------|----------|----------|----------|
| SiO <sub>2</sub>               | 71.20    | 68.40    | 77.60    | 68.00    | 73.10    |
| TiO <sub>2</sub>               | .23      | .25      | .16      | .18      | .19      |
| Al <sub>2</sub> O <sub>3</sub> | 12.41    | 15.00    | 11.80    | 14.88    | 12.87    |
| Fe <sub>2</sub> O <sub>3</sub> | 2.03     | 1.56     | 1.35     | 2.68     | 1.85     |
| MnO                            | .06      | .07      | .04      | .07      | .05      |
| H <sub>2</sub> O               | .67      | 1.59     | .44      | 2.14     | 2.41     |
| CaO                            | 3.20     | 1.85     | 1.70     | 1.21     | .89      |
| Na <sub>2</sub> O              | 2.75     | .67      | 1.50     | .73      | .43      |
| K <sub>2</sub> O               | 2.20     | 4.56     | 2.49     | 4.24     | 3.10     |
| LOI                            | 4.10     | 4.50     | 3.10     | 3.80     | 4.10     |
| TOTAL                          | 98.85    | 98.45    | 100.18   | 97.93    | 98.99    |

## TRACE ELEMENTS (PPM)

|                   |      |     |     |     |      |
|-------------------|------|-----|-----|-----|------|
| Zr                | 129  | 14  | 123 | 149 | 135  |
| Sr                | 235  | 80  | 64  | 129 | 66   |
| Rb                | 54   | 157 | 70  | 154 | 90   |
| Zn                | 34   | 28  | 8   | 14  | 20   |
| Cu                | 1    | 1   | 1   | 1   | 1    |
| Ba                | 2022 | 950 | 930 | 923 | 1431 |
| CuS               | 1    | 1   | 1   | 1   | 1    |
| PbS               | 34   | 24  | 15  | 57  | 21   |
| ZnS               | 9    | 6   | 1   | 6   | 4    |
| BaSO <sub>4</sub> |      |     |     |     |      |
| Pb                | 26   | 10  | 3   | 61  | 6    |
| Ni                | 5    | 2   | 2   | 2   | 5    |
| Co                | 11   | 12  | 38  | 11  | 18   |
| Cr                | 1    | 3   | 0   | 3   | 2    |
| V                 | 20   | 10  | 10  | 10  | 15   |
| Hg                |      |     |     |     |      |
| Ag                |      |     |     |     |      |
| Nb                |      |     |     |     |      |

POOR COPY  
COPIE DE QUALITEE INFÉRIEURE

## LUCKY STRIKE ORE HORIZON SEQUENCE - CLEMENTINE DACITIC TUFFS

## MAJOR ELEMENTS (WT%)

| SAMPLE | 2884 908 | 28851555 | 28861325 | 28861353 | 28971504 |
|--------|----------|----------|----------|----------|----------|
| SiO2   | 75.60    | 71.80    | 61.30    | 71.50    | 71.90    |
| TiO2   | .25      | .27      | .47      | .35      | .20      |
| Al2O3  | 12.06    | 13.35    | 17.03    | 12.91    | 13.12    |
| Fe2O3  | 1.74     | 2.14     | 4.74     | 2.25     | 2.22     |
| MnO    | .03      | .03      | .11      | .07      | .08      |
| MgO    | .19      | .83      | 2.99     | 1.61     | .82      |
| CaO    | .44      | 2.60     | 2.07     | 2.00     | 2.61     |
| Na2O   | 4.36     | 1.86     | 1.92     | .75      | 1.82     |
| K2O    | 3.54     | 2.98     | 4.12     | 3.79     | 3.23     |
| LOI    | .30      | 3.90     | 5.10     | 4.40     | 4.00     |
| TOTAL  | 98.51    | 99.76    | 99.85    | 99.63    | 100.00   |

## TRACE ELEMENTS (PPM)

|       |     |     |     |      |      |
|-------|-----|-----|-----|------|------|
| Zr    | 146 | 166 | 163 | 132  | 135  |
| Sr    | 57  | 214 | 139 | 62   | 171  |
| Rb    | 61  | 126 | 169 | 131  | 87   |
| Zn    | 11  | 27  | 69  | 28   | 25   |
| Cu    | 3   | 1   | 1   | 1    | 4    |
| Mo    | 788 | 791 | 752 | 1031 | 1055 |
| CuS   | 4   | 1   | 1   | 1    | 5    |
| PbS   | 21  | 25  | 25  | 25   | 26   |
| ZnS   | 5   | 8   | 17  | 27   | 5    |
| BaSO4 |     |     |     |      |      |
| Pb    | 6   | 1   | 21  | 25   | 11   |
| Ni    | 3   | 4   | 7   | 2    | 5    |
| Co    | 34  | 11  | 8   | 12   | 22   |
| Cr    | 3   | 3   | 10  | 1    | 2    |
| V     | 25  | 10  | 10  | 15   | 15   |
| Hg    |     |     |     |      |      |
| Ag    |     |     |     |      |      |
| Nb    |     |     |     |      |      |

POOR COPY  
COPIE DE QUALITEE INFERIEURE

## LUCKY STRIKE ORE HORIZON SEQUENCE - CLEMENTINE DACITIC TUFFS

## MAJOR ELEMENTS (WT%)

SAMPLE 2899 909 29001165 2901 753

|                                |       |       |       |
|--------------------------------|-------|-------|-------|
| SiO <sub>2</sub>               | 73.50 | 75.50 | 71.00 |
| TiO <sub>2</sub>               | .25   | .10   | .16   |
| Al <sub>2</sub> O <sub>3</sub> | 12.42 | 11.44 | 12.93 |
| Fe <sub>2</sub> O <sub>3</sub> | 2.06  | 1.57  | 1.70  |
| MnO                            | .05   | .06   | .05   |
| MgO                            | .48   | .83   | .61   |
| CaO                            | 1.78  | 2.33  | 2.93  |
| Na <sub>2</sub> O              | 2.23  | 1.80  | 2.04  |
| K <sub>2</sub> O               | 2.24  | 2.30  | 2.76  |
| LOI                            | 3.50  | 3.60  | 3.80  |

TOTAL 98.51 99.53 98.78

## TRACE ELEMENTS (PPM)

|                   |     |     |      |
|-------------------|-----|-----|------|
| Zr                | 134 | 125 | 138  |
| Sr                | 91  | 161 | 105  |
| Rb                | 43  | 59  | 81   |
| Zn                | 21  | 24  | 17   |
| Cu                | 1   | 1   | 1    |
| Ba                | 431 | 768 | 1766 |
| CuS               | 1   | 1   | 1    |
| PbS               | 19  | 15  | 5    |
| ZnS               | 2   | 5   | 4    |
| BaSO <sub>4</sub> |     |     |      |
| Pb                | 1   | 7   | 1    |
| Ni                | 4   | 4   | 4    |
| Co                | 41  | 19  | 14   |
| Cr                | 4   | 0   | 0    |
| V                 | 20  | 20  | 10   |
| Hg                |     |     |      |
| Ag                |     |     |      |
| Nb                |     |     |      |

POOR COPY  
COPIE DE QUALITEE INFERIEURE

## ORIENTAL ORE HORIZON SEQUENCE - DACITIC TUFFS

## MAJOR ELEMENTS (WT%)

| SAMPLE                         | 1096 300 | 1475  | 1     | 1589 736 | 16061216 | 17301310 |
|--------------------------------|----------|-------|-------|----------|----------|----------|
| SiO <sub>2</sub>               | 68.90    | 68.60 | 68.90 | 67.80    | 67.80    |          |
| TiO <sub>2</sub>               | .36      | .45   | .43   | .50      | .40      |          |
| Al <sub>2</sub> O <sub>3</sub> | 17.36    | 15.24 | 15.24 | 14.03    | 14.03    |          |
| Fe <sub>2</sub> O <sub>3</sub> | 4.61     | 2.07  | 2.63  | 3.44     | 3.28     |          |
| MnO                            | .08      | .04   | .04   | .10      | .07      |          |
| HgO                            | 4.42     | 2.79  | 2.20  | 1.51     | 1.78     |          |
| CaO                            | 2.27     | 2.75  | 1.74  | 2.51     | 2.72     |          |
| Na <sub>2</sub> O              | 1.65     | 2.29  | 1.01  | 2.62     | 1.81     |          |
| K <sub>2</sub> O               | 2.36     | 1.47  | 3.21  | 2.42     | 3.59     |          |
| LOI                            | 5.00     | 2.90  | 4.30  | 3.60     | 4.10     |          |
| TOTAL                          | 99.01    | 98.60 | 99.70 | 98.53    | 99.58    |          |

## TRACE ELEMENTS (PPM)

|                   |      |     |     |      |      |
|-------------------|------|-----|-----|------|------|
| Zr                | 193  | 186 | 196 | 150  | 168  |
| Sr                | 271  | 424 | 226 | 243  | 149  |
| Rb                | 56   | 32  | 75  | 56   | 87   |
| Zn                | 171  | 206 | 55  | 690  | 64   |
| Cu                | 13   | 1   | 1   | 120  | 2    |
| Ba                | 2395 | 865 | 174 | 1934 | 1215 |
| CuS               | 16   | 1   | 1   | 182  | 5    |
| PbS               | 56   | 24  | 21  | 330  | 42   |
| ZnS               | 82   | 111 | 24  | 710  | 37   |
| PbSO <sub>4</sub> |      |     |     |      |      |
| Pb                | 59   | 24  | 10  | 190  | 42   |
| Ni                | 8    | 4   | 4   | 7    | 3    |
| Co                | 15   | 12  | 6   | 20   | 13   |
| Cr                | 16   | 20  | 3   | 17   | 10   |
| V                 | 45   | 20  | 15  | 35   | 20   |
| Hg                |      |     |     |      |      |
| Ag                |      |     |     |      |      |
| Nb                |      |     |     |      |      |

POOR COPY  
COPIE DE QUALITE INFERIEURE

## ORIENTAL ORE HORIZON SEQUENCE - DACITIC TUFFS

## MAJOR ELEMENTS (WT%)

| SAMPLE                         | 1872 166 | 1872 223 | 1872 317 | 1872 393 | 1872 440 |
|--------------------------------|----------|----------|----------|----------|----------|
| SiO <sub>2</sub>               | 69.90    | 70.20    | 64.10    | 66.10    | 66.80    |
| TiO <sub>2</sub>               | .11      | .28      | .21      | .39      | .22      |
| Al <sub>2</sub> O <sub>3</sub> | 15.06    | 15.93    | 17.36    | 16.62    | 15.79    |
| Fe <sub>2</sub> O <sub>3</sub> | 2.01     | 1.84     | 2.49     | 1.90     | 2.09     |
| MnO                            | .09      | .08      | .06      | .06      | .07      |
| MgO                            | 1.89     | 1.43     | 2.52     | 2.88     | 3.41     |
| CaO                            | 1.44     | 1.78     | 1.35     | 1.89     | 1.67     |
| Na <sub>2</sub> O              | 2.10     | 3.63     | .92      | 1.69     | 1.65     |
| K <sub>2</sub> O               | 3.44     | 2.60     | 4.56     | 2.92     | 2.54     |
| LOI                            | 3.00     | 2.30     | 5.10     | 4.10     | 4.30     |
| TOTAL                          | 99.04    | 100.07   | 98.67    | 98.55    | 98.54    |

## TRACE ELEMENTS (PPM)

|                   |     |     |      |      |      |
|-------------------|-----|-----|------|------|------|
| Zr                | 135 | 147 | 151  | 149  | 144  |
| Sr                | 247 | 243 | 138  | 225  | 237  |
| Rb                | 111 | 76  | 152  | 98   | 85   |
| Zn                | 118 | 44  | 42   | 22   | 40   |
| Cu                | 1   | 1   | 1    | 1    | 1    |
| Ba                | 845 | 944 | 1495 | 1194 | 1205 |
| CuS               | 1   | 1   | 1    | 1    | 1    |
| PbS               | 24  | 13  | 19   | 13   | 11   |
| ZnS               | 6   | 4   | 2    | 4    | 5    |
| BaSO <sub>4</sub> |     |     |      |      |      |
| Pb                | 10  | 6   | 8    | 11   | 12   |
| Ni                | 2   | 2   | 1    | 3    | 3    |
| Co                | 19  | 15  | 12   | 18   | 19   |
| Cr                | 1   | 1   | 3    | 1    | 2    |
| V                 | 10  | 10  | 10   | 10   | 10   |
| Hg                |     |     |      |      |      |
| Ag                |     |     |      |      |      |
| Nb                |     |     |      |      |      |

POOR COPY  
COPIE DE QUALITEE INFERIEURE



## ORIENTAL ORE HORIZON SEQUENCE - DACITIC TUFFS

## MAJOR ELEMENTS (WT%)

| SAMPLE                         | 1872 465 | 1872 506 | 1872 649 | 2367 192 | 2367 273 |
|--------------------------------|----------|----------|----------|----------|----------|
| SiO <sub>2</sub>               | 65.90    | 65.10    | 49.70    | 65.30    | 74.00    |
| TiO <sub>2</sub>               | .20      | .33      | .50      | .30      | .27      |
| Al <sub>2</sub> O <sub>3</sub> | 16.00    | 17.36    | 17.86    | 16.93    | 13.74    |
| Fe <sub>2</sub> O <sub>3</sub> | 2.32     | 1.86     | .92      | 2.13     | 1.55     |
| MnO                            | .08      | .10      | .12      | .05      | .06      |
| MgO                            | 3.94     | 3.31     | 2.34     | 2.99     | .90      |
| CaO                            | 1.74     | 2.37     | 6.19     | 1.93     | 1.65     |
| Na <sub>2</sub> O              | 1.37     | 1.48     | 1.66     | 1.15     | 4.30     |
| K <sub>2</sub> O               | 2.43     | 2.75     | 5.10     | 2.65     | 1.28     |
| LOI                            | 4.50     | 4.50     | 10.10    | 5.20     | 1.30     |
| TOTAL                          | 98.48    | 99.16    | 94.49    | 98.63    | 99.05    |

## TRACE ELEMENTS (PPM)

|                   |      |      |      |      |     |
|-------------------|------|------|------|------|-----|
| Zr                | 144  | 157  | 202  | 149  | 132 |
| Sr                | 230  | 289  | 216  | 198  | 317 |
| Rb                | 79   | 75   | 106  | 110  | 33  |
| Zn                | 27   | 25   | 1910 | 31   | 19  |
| Cu                | 1    | 1    | 161  | 1    | 1   |
| Ba                | 1437 | 3876 | 4257 | 1473 | 507 |
| CuS               | 1    | 1    | 180  | 1    | 2   |
| PbS               | 20   | 32   | 1220 | 17   | 5   |
| ZnS               | 3    | 5    | 1950 | 11   | 1   |
| BaSO <sub>4</sub> |      |      |      |      |     |
| Pb                | 13   | 28   | 560  | 19   | 3   |
| Ni                | 4    | 3    | 10   | 5    | 2   |
| Co                | 10   | 4    | 7    | 6    | 19  |
| Cr                | 2    | 2    | 17   | 3    | 3   |
| V                 | 10   | 15   | 20   | 20   | 20  |
| Hg                |      |      |      |      |     |
| Ag                |      |      |      |      |     |
| Nb                |      |      |      |      |     |

POOR COPY  
COPIE DE QUALITE INFERIEURE

## ORIENTAL ORE HORIZON SEQUENCE - DACITIC TUFFS

## MAJOR ELEMENTS (Wt%)

| SAMPLE                         | 2367 394 | 2367 412 | 2837 800 | 2837 900 | 28371050 |
|--------------------------------|----------|----------|----------|----------|----------|
| SiO <sub>2</sub>               | 68.90    | 68.20    | 65.60    | 77.80    | 70.30    |
| TiO <sub>2</sub>               | .29      | .21      | .40      | .20      | .28      |
| Al <sub>2</sub> O <sub>3</sub> | 15.47    | 15.94    | 15.26    | 11.97    | 14.36    |
| Fe <sub>2</sub> O <sub>3</sub> | 1.62     | 1.92     | 4.08     | 1.59     | 2.32     |
| MnO                            | .04      | .03      | .07      | .03      | .03      |
| NiO                            | 4.03     | 4.55     | 2.69     | 1.43     | 2.15     |
| CaO                            | 1.31     | 1.24     | 1.66     | .54      | 1.14     |
| Na <sub>2</sub> O              | 1.70     | 1.66     | 2.42     | 4.30     | 2.78     |
| K <sub>2</sub> O               | 1.29     | 1.22     | 2.07     | 1.26     | 2.68     |
| LOI                            | 4.80     | 4.90     | 3.50     | 1.20     | 2.40     |
| TOTAL                          | 99.45    | 99.87    | 97.75    | 100.32   | 98.44    |

## TRACE ELEMENTS (PPM)

|                   |     |     |     |     |      |
|-------------------|-----|-----|-----|-----|------|
| Zr                | 160 | 162 | 163 | 109 | 132  |
| Sr                | 251 | 262 | 307 | 234 | 306  |
| Rb                | 43  | 41  | 91  | 59  | 90   |
| Zn                | 33  | 39  | 77  | 28  | 26   |
| Cu                | 1   | 2   | 1   | 1   | 1    |
| Ba                | 242 | 919 | 267 | 215 | 1015 |
| CuS               | 1   | 1   | 1   | 1   | 2    |
| PBS               | 27  | 11  | 10  | 10  | 15   |
| ZnS               | 1   | 2   | 11  | 8   | 9    |
| P-SO <sub>4</sub> |     |     | 0   | 0   | 0    |
| Pb                | 21  | 12  | 15  | 3   | 12   |
| Ni                | 4   | 3   | 4   | 2   | 2    |
| Co                | 13  | 12  | 18  | 47  | 24   |
| Cr                | 2   | 2   | 19  | 30  | 41   |
| V                 | 20  | 10  | 10  | 10  | 10   |
| Hg                |     |     |     |     |      |
| Ag                |     |     |     |     |      |
| Pb                |     |     | 0   | 0   | 2    |

POOR COPY  
COPIE DE QUALITEE INFERIEURE

## ORIENTAL ORE HORIZON SEQUENCE - DACITIC TUFFS

## MAJOR ELEMENTS (WT%)

| SAMPLE                         | 2839 529 | 2839 657 | 28391586 | 2853 672 | 2854 708 |
|--------------------------------|----------|----------|----------|----------|----------|
| SiO <sub>2</sub>               | 72.40    | 72.60    | 70.80    | 57.40    | 63.50    |
| TiO <sub>2</sub>               | .24      | .22      | .19      | .47      | .41      |
| Al <sub>2</sub> O <sub>3</sub> | 13.71    | 13.57    | 14.59    | 18.14    | 16.00    |
| Fe <sub>2</sub> O <sub>3</sub> | 2.15     | 1.97     | 2.31     | 6.21     | 5.34     |
| MnO                            | .03      | .03      | .03      | .10      | .06      |
| MgO                            | 2.04     | 2.05     | 1.25     | 3.22     | 2.33     |
| CaO                            | 1.48     | 1.14     | 1.14     | 3.36     | 3.59     |
| Na <sub>2</sub> O              | 2.88     | 2.76     | 3.70     | 1.56     | 1.79     |
| K <sub>2</sub> O               | 2.48     | 2.72     | 3.47     | 4.12     | 2.36     |
| LOI                            | 1.90     | 2.20     | 1.50     | 4.40     | 3.40     |
| TOTAL                          | 99.31    | 99.26    | 98.98    | 98.98    | 98.78    |

## TRACE ELEMENTS (PPM)

|                   |      |      |     |     |     |
|-------------------|------|------|-----|-----|-----|
| Zr                | 111  | 118  | 120 | 203 | 11  |
| Sr                | 246  | 218  | 185 | 272 | 347 |
| Rb                | 76   | 82   | 96  | 161 | 67  |
| Zn                | 23   | 16   | 20  | 42  | 52  |
| Cu                | 1    | 1    | 1   | 11  | 1   |
| Ba                | 1135 | 1339 | 104 | 113 | 109 |
| CdS               | 2    | 1    | 1   | 14  | 1   |
| PbS               | 7    | 11   | 10  | 20  | 16  |
| ZnS               | 3    | 2    | 25  | 25  | 9   |
| BaSO <sub>4</sub> |      |      |     |     |     |
| Pb                | 11   | 8    | 4   | 13  | 10  |
| Ni                | 5    | 2    | 2   | 5   | 6   |
| Co                | 26   | 27   | 17  | 11  | 15  |
| Cr                | 2    | 2    | 5   | 13  | 11  |
| V                 | 15   | 10   | 10  | 20  | 10  |
| Hg                |      |      |     |     |     |
| Ag                |      |      |     |     |     |
| Nb                |      |      |     |     |     |

POOR COPY  
COPIE DE QUALITEE INFERIEURE

## ORIENTAL ORE HORIZON SEQUENCE - DACITIC TUFFS

## MAJOR ELEMENTS (WT%)

SAMPLE 2859 304

|                                |       |
|--------------------------------|-------|
| SiO <sub>2</sub>               | 62.00 |
| TiO <sub>2</sub>               | .41   |
| Al <sub>2</sub> O <sub>3</sub> | 16.97 |
| Fe <sub>2</sub> O <sub>3</sub> | 4.39  |
| MnO                            | .10   |
| H <sub>2</sub> O               | 2.40  |
| CaO                            | 2.76  |
| Na <sub>2</sub> O              | 1.66  |
| K <sub>2</sub> O               | 3.60  |
| LOI                            | 5.50  |

|       |       |
|-------|-------|
| TOTAL | 99.79 |
|-------|-------|

## TRACE ELEMENTS (PPM)

|                   |     |
|-------------------|-----|
| Zr                | 193 |
| Sr                | 206 |
| Rb                | 119 |
| Zn                | 330 |
| Cu                | 1   |
| Ba                | 943 |
| CdS               | 1   |
| PbS               | 27  |
| ZnS               | 67  |
| BaSO <sub>4</sub> |     |
| Pb                | 19  |
| Ni                | 5   |
| Co                | 11  |
| Cr                | 10  |
| V                 | 10  |
| Hg                |     |
| Ag                |     |
| Nb                |     |

POOR COPY  
COPIE DE QUALITEE INFERIEURE

3-96

LAKE SEVEN BASALT

MAJOR ELEMENTS (WT%)

SAMPLE 23491540

|                                |       |
|--------------------------------|-------|
| SiO <sub>2</sub>               | 50.10 |
| TiO <sub>2</sub>               | .52   |
| Al <sub>2</sub> O <sub>3</sub> | 13.67 |
| Fe <sub>2</sub> O <sub>3</sub> | 9.38  |
| MnO                            | .22   |
| H <sub>2</sub> O               | 8.65  |
| CaO                            | 9.87  |
| Na <sub>2</sub> O              | 2.70  |
| K <sub>2</sub> O               | .50   |
| LOI                            | 3.70  |

TOTAL 99.31

TRACE ELEMENTS (PPM)

|                   |     |
|-------------------|-----|
| Zr                | 65  |
| Sr                | 395 |
| Rb                | 9   |
| Zn                | 61  |
| Cu                | 90  |
| Pb                | 309 |
| CuS               | 140 |
| PbS               | 16  |
| ZnS               | 21  |
| BaSO <sub>4</sub> |     |
| Pb                | 14  |
| Ni                | 53  |
| Co                | 25  |
| Cr                | 130 |
| V                 | 145 |
| Hg                |     |
| Ag                |     |
| Nb                |     |

POOR COPY  
COPIE DE QUALITEE INFERIEURE

## LAKE SEVEN BASALT ALTERED FLOWS

## MAJOR ELEMENTS (WT%)

SAMPLE 932 367

|                                |       |
|--------------------------------|-------|
| SiO <sub>2</sub>               | 43.98 |
| TiO <sub>2</sub>               | .88   |
| Al <sub>2</sub> O <sub>3</sub> | 15.48 |
| Fe <sub>2</sub> O <sub>3</sub> | 18.88 |
| MnO                            | .19   |
| MgO                            | 7.72  |
| CaO                            | 12.88 |
| Na <sub>2</sub> O              | 2.58  |
| K <sub>2</sub> O               | .63   |
| LOI                            | 6.18  |

TOTAL 99.24

## TRACE ELEMENTS (PPM)

|                   |     |
|-------------------|-----|
| Zr                | 181 |
| Sr                | 425 |
| Rb                | 9   |
| Zn                | 55  |
| Cu                | 64  |
| Ba                | 288 |
| CuS               | 72  |
| PbS               | 7   |
| ZnS               | 15  |
| BaSO <sub>4</sub> |     |
| Pb                | 18  |
| Ni                | 39  |
| Co                | 24  |
| Cr                | 87  |
| V                 | 148 |
| Hg                |     |
| Ag                |     |
| Nb                |     |

POOR COPY  
COPIE DE QUALITEE INFERIEURE

## SANDY LAKE BASALT FLOWS

## MAJOR ELEMENTS (WT%)

SAMPLE 1352 477 2839 494

|                                |       |       |
|--------------------------------|-------|-------|
| SiO <sub>2</sub>               | 45.60 | 49.80 |
| TiO <sub>2</sub>               | .86   | .60   |
| Al <sub>2</sub> O <sub>3</sub> | 14.99 | 14.70 |
| Fe <sub>2</sub> O <sub>3</sub> | 10.05 | 8.51  |
| MnO                            | .19   | .31   |
| HgO                            | 7.45  | 6.72  |
| CaO                            | 11.40 | 8.76  |
| Na <sub>2</sub> O              | 2.23  | 2.93  |
| K <sub>2</sub> O               | .53   | 1.39  |
| LOI                            | 5.20  | 5.10  |

|       |       |       |
|-------|-------|-------|
| TOTAL | 98.50 | 98.82 |
|-------|-------|-------|

## TRACE ELEMENTS (PPM)

|                   |     |      |
|-------------------|-----|------|
| Zr                | 102 | .68  |
| Sr                | 36  | 334  |
| Rb                | 9   | 29   |
| Zn                | 55  | 48   |
| Cu                | 53  | 62   |
| Ba                | 539 | 2294 |
| CuS               | 76  | 65   |
| PbS               | 15  | 22   |
| ZnS               | 22  | 5    |
| BaSO <sub>4</sub> |     |      |
| Pb                | 21  | 21   |
| Mi                | 102 | 114  |
| Co                | 27  | 36   |
| Cr                | 379 | 429  |
| V                 | 135 | 163  |
| Hg                |     |      |
| Ag                |     |      |
| Pb                |     |      |

POOR COPY  
COPIE DE QUALITEE INFERIEURE

## UPPER BUCHANS SUBGROUP - MAFIC TUFFS

## MAJOR ELEMENTS (WT%)

| SAMPLE                         | 2837 125 | 2837 200 | 2837 350 | 2837 438 | 2837 550 |
|--------------------------------|----------|----------|----------|----------|----------|
| SiO <sub>2</sub>               | 52.20    | 53.60    | 49.50    | 70.00    | 74.10    |
| TiO <sub>2</sub>               | .45      | .53      | .57      | .36      | .27      |
| Al <sub>2</sub> O <sub>3</sub> | 17.48    | 16.04    | 17.88    | 13.25    | 13.11    |
| Fe <sub>2</sub> O <sub>3</sub> | 8.54     | 9.35     | 10.90    | 3.66     | 2.63     |
| MnO                            | .16      | .15      | .17      | .08      | .09      |
| MgO                            | 3.10     | 2.88     | 3.01     | 1.69     | 1.48     |
| CaO                            | 4.47     | 4.54     | 3.69     | 2.87     | 1.42     |
| Na <sub>2</sub> O              | 4.55     | 4.30     | 4.91     | 3.57     | 2.07     |
| K <sub>2</sub> O               | 1.01     | .91      | .05      | 1.39     | 1.76     |
| LOI                            | 6.00     | 5.60     | 6.20     | 3.40     | 3.00     |
| TOTAL                          | 97.96    | 97.90    | 96.88    | 100.27   | 99.93    |

## TRACE ELEMENTS (PPM)

|                   |     |     |     |     |     |
|-------------------|-----|-----|-----|-----|-----|
| Zr                | 56  | 54  | 18  | 169 | 143 |
| Sr                | 142 | 115 | 101 | 135 | 89  |
| Rb                | 20  | 30  | 13  | 36  | 38  |
| Zn                | 68  | 77  | 129 | 45  | 57  |
| Cu                | 65  | 153 | 31  | 8   | 5   |
| Ba                | 74  | 70  | 12  | 137 | 216 |
| CdS               | 70  | 108 | 35  | 10  | 8   |
| PbS               | 10  | 10  | 13  | 13  | 20  |
| ZnS               | 13  | 15  | 17  | 7   | 9   |
| BaSO <sub>4</sub> | 0   | 0   | 0   | 0   | 0   |
| Pb                | 22  | 15  | 25  | 11  | 21  |
| Ri                | 17  | 11  | 18  | 5   | 2   |
| Co                | 27  | 27  | 33  | 18  | 24  |
| Cr                | 61  | 30  | 39  | 15  | 15  |
| V                 | 150 | 190 | 250 | 10  | 10  |
| Hg                |     |     |     |     |     |
| Ag                |     |     |     |     |     |
| Pb                | 0   | 0   | 0   | 0   | 0   |

POOR COPY  
COPIE DE QUALITEE INFERIEURE



## UPPER BUCHANS SUBGROUP - MAFIC TUFFS

## MAJOR ELEMENTS (WT%)

SAMPLE 2837 725

|                                |       |
|--------------------------------|-------|
| SiO <sub>2</sub>               | 69.90 |
| TiO <sub>2</sub>               | .37   |
| Al <sub>2</sub> O <sub>3</sub> | 14.84 |
| Fe <sub>2</sub> O <sub>3</sub> | 3.88  |
| MnO                            | .08   |
| MgO                            | 1.96  |
| CaO                            | .53   |
| Na <sub>2</sub> O              | 3.80  |
| K <sub>2</sub> O               | 1.42  |
| LOI                            | 2.20  |

TOTAL 98.98

## TRACE ELEMENTS (PPM)

|                   |     |
|-------------------|-----|
| Zr                | 143 |
| Sr                | 142 |
| Pb                | 150 |
| Zn                | 68  |
| Cu                | 1   |
| Ba                | 268 |
| ClS               | 1   |
| PbS               | 6   |
| ZnS               | 11  |
| PbSO <sub>4</sub> | 0   |
| Pb                | 12  |
| Pi                | 1   |
| Co                | 20  |
| Cr                | 11  |
| V                 | 10  |
| Hg                |     |
| Ag                |     |
| Sb                |     |

POOR COPY  
COPIE DE QUALITEE INFERIEURE

3-101

## UPPER BUCHANS SUBGROUP - RHYOLITE FLOWS

## MAJOR ELEMENTS (WT%)

| SAMPLE                         | BJ-9-80 | BJ-10-46 | BJ42 447 | BJ42 980 | BJ421470 |
|--------------------------------|---------|----------|----------|----------|----------|
| SiO <sub>2</sub>               | 71.50   | 72.00    | 72.50    | 76.10    | 69.70    |
| TiO <sub>2</sub>               | .51     | .44      | .24      | .04      | .26      |
| Al <sub>2</sub> O <sub>3</sub> | 13.96   | 13.15    | 13.21    | 11.82    | 13.64    |
| Fe <sub>2</sub> O <sub>3</sub> | 3.16    | 3.39     | 2.83     | 1.21     | 3.72     |
| MnO                            | .11     | .11      | .13      | .04      | .23      |
| H <sub>2</sub> O               | 1.74    | 1.51     | .97      | .93      | 1.51     |
| CaO                            | .89     | 1.12     | 1.66     | .47      | 1.59     |
| Na <sub>2</sub> O              | 4.23    | 4.08     | 2.95     | 3.63     | 4.33     |
| K <sub>2</sub> O               | 1.90    | 1.04     | 2.76     | 3.29     | 1.51     |
| LOI                            | 1.70    | 1.40     | 2.80     | 1.40     | 2.70     |
| TOTAL                          | 99.70   | 99.04    | 100.05   | 98.93    | 99.19    |

## TRACE ELEMENTS (PPM)

|                   |      |     |     |     |     |
|-------------------|------|-----|-----|-----|-----|
| Zr                | 174  | 183 | 120 | 111 | 107 |
| Sr                | 151  | 199 | 92  | 110 | 51  |
| Rb                | 36   | 28  | 83  | 67  | 35  |
| Zn                | 126  | 52  | 65  | 18  | 170 |
| Cu                | 3    | 1   | 18  | 10  | 1   |
| Ta                | 1233 | 749 | 610 | 970 | 747 |
| CuS               | 3    | 1   | 19  | 0   | 0   |
| PbS               | 320  | 44  | 9   | 11  | 8   |
| ZnS               | 30   | 15  | 12  | 1   | 21  |
| FeSO <sub>4</sub> | 0    | 0   | 0   | 0   | 0   |
| Pb                | 350  | 46  | 11  | 2   | 9   |
| Ni                | 2    | 5   | 4   | -3  | 2   |
| Co                | 29   | 76  | 12  | 16  | 19  |
| Cr                | 12   | 11  | 10  | 10  | 13  |
| V                 | 10   | 10  | 10  | 10  | 10  |
| Hg                |      |     |     |     |     |
| Ag                |      |     |     |     |     |
| Pb                | 3    | 9   | 9   | 5   | 9   |

POOR COPY  
COPIE DE QUALITEE INFERIEURE

3-102

## UPPER BUCHANS SUBGROUP - RHYOLITE FLOWS

## MAJOR ELEMENTS (WT%)

SAMPLE BJ421658 BJ45 117 BJ46 100

|       |       |       |       |
|-------|-------|-------|-------|
| SiO2  | 77.30 | 69.90 | 67.20 |
| TiO2  | .10   | .15   | .18   |
| Al2O3 | 11.82 | 14.48 | 12.50 |
| Fe2O3 | 1.18  | 1.28  | 2.37  |
| MnO   | .03   | .08   | .05   |
| MgO   | .83   | 1.57  | 2.44  |
| CaO   | .58   | 1.35  | .32   |
| Na2O  | 3.77  | 2.66  | .80   |
| K2O   | 1.72  | 2.68  | 2.86  |
| LOI   | 1.70  | 4.10  | 5.20  |

TOTAL 99.03 98.25 93.42

## TRACE ELEMENTS (PPM)

|       |       |     |     |
|-------|-------|-----|-----|
| Zr    | 111   | 103 | 103 |
| Sr    | 112   | 37  | 9   |
| Rb    | 46    | 50  | 53  |
| Zn    | 27    | 27  | 31  |
| Cu    | 1     | 8   | 5   |
| Ca    | 10.20 | 864 | 795 |
| CuS   | 0     | 11  | 6   |
| PbS   | 11    | 21  | 15  |
| ZnS   | 1     | 18  | 4   |
| PbSO4 | 0     |     |     |
| Pb    | 22    | 25  | 18  |
| Ni    | 1     | 3   | 4   |
| Co    | 15    | 21  | 18  |
| Cr    | 9     | 3   | 11  |
| V     | 10    | 10  | 10  |
| Hg    |       |     |     |
| Ag    |       |     |     |
| Pb    | 9     |     |     |

POOR COPY  
COPIE DE QUALITEE INFERIEURE

## UPPER BUCHANS SUBGROUP - DACITIC TUFFS

## MAJOR ELEMENTS (WT%)

| SAMPLE                         | 1568 640 | 16061070 | 17821036 | 19201726 | 2042 762 |
|--------------------------------|----------|----------|----------|----------|----------|
| SiO <sub>2</sub>               | 61.80    | 70.20    | 66.00    | 72.60    | 68.90    |
| TiO <sub>2</sub>               | .43      | .33      | .28      | .10      | .37      |
| Al <sub>2</sub> O <sub>3</sub> | 17.38    | 13.60    | 15.64    | 12.00    | 13.05    |
| Fe <sub>2</sub> O <sub>3</sub> | 4.14     | 2.89     | 4.64     | 2.01     | 4.88     |
| MnO                            | .07      | .07      | .06      | .04      | .15      |
| MgO                            | 2.88     | 1.47     | 1.53     | .99      | 2.24     |
| CaO                            | 2.11     | 2.14     | 1.60     | 1.72     | 1.19     |
| Na <sub>2</sub> O              | 1.75     | 3.80     | 2.82     | 2.14     | 2.82     |
| K <sub>2</sub> O               | 3.37     | 1.46     | 2.43     | 2.65     | 1.60     |
| LOI                            | 4.00     | 2.50     | 3.80     | 3.40     | 3.70     |
| TOTAL                          | 97.93    | 98.46    | 98.88    | 97.65    | 98.90    |

## TRACE ELEMENTS (PPM)

|                   |      |     |     |     |     |
|-------------------|------|-----|-----|-----|-----|
| Zr                | 221  | 167 | 144 | 97  | 112 |
| Sr                | 203  | 360 | 101 | 68  | 56  |
| Rb                | 112  | 33  | 58  | 58  | 41  |
| Zn                | 62   | 65  | 70  | 57  | 61  |
| Cu                | 1    | 17  | 1   | 200 | 1   |
| Ba                | 1100 | 200 | 240 | 173 | 100 |
| CdS               | 2    | 30  | 1   | 243 | 1   |
| PbS               | 15   | 39  | 15  | 16  | 14  |
| ZnS               | 18   | 31  | 33  | 17  | 11  |
| BaSO <sub>4</sub> |      |     |     |     |     |
| Pb                | 5    | 69  | 16  | 15  | 8   |
| Ni                | 7    | 4   | 3   | 4   | 8   |
| Co                | 11   | 19  | 23  | 19  | 15  |
| Cr                | 10   | 6   | 8   | 0   | 0   |
| V                 | 10   | 20  | 10  | 10  | 10  |
| Hg                |      |     |     |     |     |
| Ag                |      |     |     |     |     |
| Pb                |      |     |     |     |     |

POOR COPY  
COPIE DE QUALITE INFERIEURE

## UPPER BUCHANS SUBGROUP - DACITIC TUFFS

## MAJOR ELEMENTS (WT%)

| SAMPLE                         | BJ42 685 | BJ421975 | BJ422228 | BJ47 344 | BJ48 135 |
|--------------------------------|----------|----------|----------|----------|----------|
| SiO <sub>2</sub>               | 61.80    | 72.20    | 73.50    | 67.30    | 68.30    |
| TiO <sub>2</sub>               | .22      | .18      | .16      | .45      | .38      |
| Al <sub>2</sub> O <sub>3</sub> | 19.92    | 13.01    | 13.07    | 14.93    | 16.97    |
| Fe <sub>2</sub> O <sub>3</sub> | 2.26     | 3.19     | 2.90     | 3.75     | 2.02     |
| MnO                            | .07      | .13      | .08      | .07      | .01      |
| MgO                            | 2.46     | 1.41     | 1.99     | 2.00     | 1.58     |
| CaO                            | .24      | .66      | .39      | 1.04     | 1.19     |
| Na <sub>2</sub> O              | 1.44     | 4.79     | 3.89     | 2.10     | 3.16     |
| K <sub>2</sub> O               | 8.30     | 1.13     | 1.29     | 2.80     | 1.63     |
| LOI                            | 3.50     | 1.80     | 2.00     | 4.10     | 3.50     |
| TOTAL                          | 100.21   | 98.50    | 99.27    | 99.34    | 98.74    |

## TRACE ELEMENTS (PPM)

|                   |      |     |     |     |     |
|-------------------|------|-----|-----|-----|-----|
| Zr                | 128  | 100 | 107 | 108 | 120 |
| Sr                | 68   | 108 | 93  | 45  | 133 |
| Rb                | 216  | 21  | 34  | 58  | 39  |
| Zn                | 26   | 90  | 88  | 52  | 51  |
| Cu                | 1    | 1   | 1   | 11  | 1   |
| Ba                | 1107 | 143 | 5   | 146 | 175 |
| ClS               | 1    | 0   | 1   | 14  | 1   |
| BrS               | 17   | 11  | 11  | 15  | 21  |
| ZnS               | 24   | 12  | 5   | 6   | 9   |
| FeSO <sub>4</sub> | 0    | 0   | 0   |     |     |
| Pb                | 18   | 1   | 11  | 24  | 21  |
| Li                | 5    | 1   | 3   | 7   | 5   |
| Co                | 7    | 14  | 15  | 14  | 14  |
| Cr                | 7    | 10  | 9   | 5   | 2   |
| V                 | 10   | 10  | 10  | 15  | 15  |
| Hg                |      |     | 4   |     |     |
| Ag                |      |     |     |     |     |
| Pb                | 11   | 6   | 8   |     |     |

POOR COPY  
COPIE DE QUALITE INFERIEURE

## UPPER BUCHANS SUBGROUP - DACITIC TUFFS

## MAJOR ELEMENTS (WT%)

| SAMPLE                         | BJ49 263 | 1683 403 | 1730 602 | 2852 674 | 2855 110 |
|--------------------------------|----------|----------|----------|----------|----------|
| SiO <sub>2</sub>               | 70.20    | 65.00    | 76.30    | 71.20    | 71.90    |
| TiO <sub>2</sub>               | .32      | .23      | .18      | .32      | .26      |
| Al <sub>2</sub> O <sub>3</sub> | 13.49    | 14.93    | 10.10    | 14.03    | 13.66    |
| Fe <sub>2</sub> O <sub>3</sub> | 3.73     | 5.80     | .42      | 3.59     | 3.13     |
| MnO                            | .08      | .13      | .02      | .08      | .04      |
| MgO                            | 1.21     | 2.79     | .30      | 1.50     | 1.27     |
| CaO                            | 1.69     | 1.48     | .64      | 2.24     | 2.30     |
| Na <sub>2</sub> O              | 2.03     | 1.96     | .19      | 3.68     | 4.69     |
| K <sub>2</sub> O               | 2.28     | 2.14     | 2.94     | 1.85     | 1.27     |
| LOI                            | 3.70     | 4.20     | 3.70     | 1.70     | 1.40     |
| TOTAL                          | 98.73    | 98.66    | 94.79    | 100.19   | 99.62    |

## TRACE ELEMENTS (PPM)

|       |     |     |     |     |     |
|-------|-----|-----|-----|-----|-----|
| Zr    | 98  | 126 | 110 | 120 | 3   |
| Sr    | 29  | 62  | 18  | 436 | 203 |
| Rb    | 52  | 73  | 45  | 40  | 15  |
| Zn    | 75  | 187 | 76  | 63  | 31  |
| Cu    | 6   | 7   | 49  | 3   | 1   |
| Ca    | 146 | 237 | 200 | 265 | 27  |
| PbS   | 8   | 9   | 67  | 4   | 1   |
| PbS   | 26  | 11  | 18  | 28  | 3   |
| ZnS   | 24  | 29  | 52  | 14  | 12  |
| PbS04 |     |     |     | 0   | 0   |
| Pb    | 20  | 17  | 6   | 21  | 3   |
| Ni    | 4   | 8   | 4   | 5   | 1   |
| Co    | 13  | 15  | 21  | 12  | 21  |
| Cr    | 6   | 15  | 1   | 11  | 13  |
| V     | 30  | 10  | 15  | 30  | 25  |
| Hg    |     |     |     |     |     |
| Ag    |     |     |     |     |     |
| Pb    |     |     |     | 10  | 10  |

POOR COPY  
COPIE DE QUALITE INFERIEURE

3-106

UPPER BUCHANS SUB GROUP - DACITIC TUFFS

MAJOR ELEMENTS (WT%)

SAMPLE 28551476

|                                |       |
|--------------------------------|-------|
| SiO <sub>2</sub>               | 69.90 |
| TiO <sub>2</sub>               | .40   |
| Al <sub>2</sub> O <sub>3</sub> | 14.99 |
| Fe <sub>2</sub> O <sub>3</sub> | 2.99  |
| MnO                            | .07   |
| H <sub>2</sub> O               | 1.43  |
| CaO                            | 2.71  |
| MgO                            | 3.06  |
| K <sub>2</sub> O               | 1.89  |
| LOI                            | 1.60  |

TOTAL 99.04

TRACE ELEMENTS (PPM)

|                   |     |
|-------------------|-----|
| Zr                | 115 |
| Sr                | 17  |
| Rb                | 52  |
| Zn                | 17  |
| Cu                | 2   |
| Li                | 3   |
| Pb                | 3   |
| Cl                | 3   |
| Na                | 8   |
| FeSO <sub>4</sub> | 0   |
| Pb                | 7   |
| Ni                | 3   |
| Co                | 26  |
| Cr                | 16  |
| V                 | 45  |
| Hg                |     |
| Ag                |     |
| As                | 13  |

POOR COPY  
COPIE DE QUALITEE INFERIEURE

## LOW TITANIUM DATABASE

## MAJOR ELEMENTS (WT%)

| SAMPLE | 05421267 | 05421775 | 08691550 | 14970325 | 14930800 |
|--------|----------|----------|----------|----------|----------|
| SiO2   | 45.90    | 39.00    | 54.80    | 48.00    | 40.90    |
| TiO2   | .92      | .70      | .60      | .80      | .20      |
| Al2O3  | 17.49    | 13.40    | 16.00    | 17.00    | 14.95    |
| Fe2O3  | 11.93    | 8.53     | 10.15    | 9.48     | 10.75    |
| MnO    | .12      | .17      | .15      | .15      | .18      |
| MgO    | 6.95     | 6.05     | 5.40     | 7.90     | 9.58     |
| CaO    | 4.31     | 14.79    | 0.33     | 13.54    | 9.76     |
| Na2O   | 2.34     | 2.47     | 2.07     | 2.35     | 1.92     |
| K2O    | 1.25     | .46      | 1.18     | 1.51     | .46      |
| LOI    | 6.90     | 12.00    | 2.30     | 3.00     | 4.00     |
| TOTAL  | 98.11    | 97.08    | 100.98   | 100.31   | 100.00   |

## TRACE ELEMENTS (PPM)

|        |     |     |     |     |     |
|--------|-----|-----|-----|-----|-----|
| Zr     | 185 | 32  | 74  | 94  | 124 |
| Sr     | 354 | 276 | 391 | 317 | 235 |
| Rb     | 33  | 9   | 23  | 25  | 17  |
| Zn     | 93  | 111 | 80  | 56  | 80  |
| Cu     | 69  | 78  | 78  | 77  | 51  |
| Co     | 50  | 103 | 100 | 112 | 13  |
| ClS    | 59  | 29  | 23  | 15  | 34  |
| HS     | 2   | 9   | 7   | 5   | 17  |
| ZnS    | 17  | 2   | 16  | 15  | 13  |
| ZrSiO4 | 0   | 0   |     | 1   | 0   |
| Pb     | 73  | 10  | 73  | 7   | 10  |
| Bi     | 42  | 1   | 12  | 16  | 2   |
| Co     | 39  | 37  | 27  | 37  | 43  |
| Cr     | 170 | 240 | 69  | 213 | 219 |
| V      | 350 | 278 | 275 | 392 | 288 |
| Hg     | 10  | 34  | 13  | 16  | 8   |
| Ag     | 1   | 13  | 2   | 3   | 0   |
| As     |     |     |     |     |     |

POOR COPY  
COPIE DE QUALITEE INFERIEURE



3-108

LOW TITANIUM DAIBASE

MAJOR ELEMENTS (WIZ)

SAMPLE 19170290 26660700 26660975

|       |       |       |       |
|-------|-------|-------|-------|
| SiO2  | 47.30 | 55.20 | 52.90 |
| TiO2  | .71   | .85   | .83   |
| Al2O3 | 14.88 | 16.41 | 16.72 |
| Fe2O3 | 10.48 | 10.48 | 11.06 |
| MnO   | .23   | .22   | .20   |
| MgO   | 3.29  | 4.23  | 5.82  |
| CaO   | 9.46  | 6.08  | 4.67  |
| Na2O  | 1.93  | 3.27  | 2.89  |
| K2O   | .88   | .41   | 1.60  |
| LOI   | 5.90  | 3.60  | 3.90  |

TOTAL 110.06 120.55 120.59

TRACE ELEMENTS (PPM)

|      |      |     |     |
|------|------|-----|-----|
| Zr   | 97   | 71  | 73  |
| Sr   | 278  | 421 | 297 |
| Rb   | 23   | 15  | 21  |
| Zn   | 96   | 40  | 37  |
| Cu   | 62   | 55  | 37  |
| Ba   | 1011 | 274 | 10  |
| Co   | 37   | 67  | 19  |
| Pb   | 9    | 7   | 10  |
| ZnS  | 29   | 17  | 19  |
| FeO4 |      |     |     |
| Pb   | 12   | 6   | 3   |
| Bi   | 78   | 11  | 12  |
| Co   | 33   | 19  | 13  |
| Cr   | 226  | 43  | 45  |
| V    | 256  | 512 | 530 |
| Hg   | 12   | 14  | 10  |
| Ag   | 0    | 0   | 0   |
| As   |      |     |     |

POOR COPY  
COPIE DE QUALITEE INFERIEURE

3-169

FEEDER GRANODIORITE

MAJOR ELEMENTS (WT%)

SAMPLE JGT75227 JGT75228 JGT75229

|       |       |       |       |
|-------|-------|-------|-------|
| SiO2  | 74.50 | 75.20 | 74.30 |
| TiO2  | .24   | .18   | .28   |
| Al2O3 | 12.99 | 12.33 | 12.45 |
| Fe2O3 | 2.19  | 1.86  | 2.36  |
| MnO   | .04   | .03   | .04   |
| MgO   | .50   | .34   | .52   |
| CaO   | 1.42  | 1.39  | 1.32  |
| Na2O  | 4.15  | 3.88  | 4.21  |
| K2O   | 2.40  | 2.69  | 2.44  |
| LOI   | .70   | .60   | .60   |

TOTAL 98.83 98.50 98.52

TRACE ELEMENTS (PPM)

|       |     |     |     |
|-------|-----|-----|-----|
| Zn    | 119 | 97  | 116 |
| Sr    | 127 | 97  | 120 |
| Rb    | 56  | 56  | 49  |
| Zn    | 30  | 14  | 21  |
| Cu    | 1   | 2   | 7   |
| La    | 221 | 113 | 212 |
| Yb    | 1   | 1   | 5   |
| Yb    | 7   | 8   | 8   |
| ZnS   | 2   | 5   | 7   |
| PrSD4 | 3   | 0   | 0   |
| Rb    | 8   | 4   | 10  |
| Li    | 13  | 5   | 1   |
| Co    | 47  | 42  | 57  |
| Cr    | 8   | 9   | 13  |
| V     | 40  | 35  | 40  |
| Hg    |     |     |     |
| Ag    |     |     |     |
| As    |     |     |     |

POOR COPY  
COPIE DE QUALITEE INFERIEURE

3-170

# HIGH TITANIUM DATABASE

## MAJOR ELEMENTS (WT%)

| SAMPLE | JGT75218 | 01750200 | 542 804 | 542 850 | 542 925 |
|--------|----------|----------|---------|---------|---------|
| SiO2   | 48.00    | 43.90    | 45.00   | 44.70   | 45.30   |
| TiO2   | 2.28     | 1.80     | 1.59    | 1.56    | 1.52    |
| Al2O3  | 14.70    | 17.63    | 16.73   | 16.17   | 16.57   |
| Fe2O3  | 12.43    | 10.68    | 11.10   | 10.28   | 10.82   |
| MnO    | 1.20     | .16      | .19     | .17     | .17     |
| HgO    | 6.32     | 6.70     | 7.04    | 5.86    | 6.67    |
| CaO    | 9.40     | 9.32     | 9.45    | 6.66    | 8.67    |
| Na2O   | 2.89     | 3.05     | 2.92    | 5.21    | 4.07    |
| K2O    | .70      | .25      | .24     | .07     | .10     |
| LOI    | 1.60     | 4.10     | 4.10    | 7.10    | 5.00    |
| TOTAL  | 98.52    | 92.59    | 98.36   | 98.78   | 98.89   |

## TRACE ELEMENTS (PPM)

|       |     |     |     |     |     |
|-------|-----|-----|-----|-----|-----|
| Zr    | 142 | 145 | 154 | 108 | 156 |
| Sr    | 24  | 142 | 369 | 361 | 427 |
| Rb    | 16  | 51  | 15  | 14  | 14  |
| Zn    | 42  | 117 | 98  | 92  | 87  |
| Cu    | 57  | 52  | 29  | 27  | 27  |
| Co    | 276 | 755 | 243 | 225 | 147 |
| ClS   | 19  | 40  | 20  | 13  | 12  |
| BrS   | 11  | 75  | 5   | 2   | 3   |
| ZnS   | 10  | 16  | 10  | 13  | 10  |
| PaS04 | 0   | 0   | 0   | 0   | 0   |
| Pb    | 15  | 25  | 42  | 39  | 32  |
| Bi    | 32  | 43  | 16  | 15  | 0   |
| Mo    | 27  | 30  | 41  | 3   | 58  |
| Cr    | 139 | 20  | 87  | 87  | 70  |
| V     | 115 | 255 | 236 | 212 | 250 |
| Hg    |     | 31  | 19  | 17  | 12  |
| Ag    |     | 0   | 17  | 14  | 14  |
| B     | 51  |     |     |     |     |

POOR COPY  
COPIE DE QUALITEE INFERIEURE

## HIGH TITANIUM DATABASE

## MAJOR ELEMENTS (WT%)

| SAMPLE | 07920875 | 07921500 | 08681025 | 08690650 | 08692000 |
|--------|----------|----------|----------|----------|----------|
| SiO2   | 45.00    | 45.60    | 54.80    | 46.20    | 55.20    |
| TiO2   | 2.22     | 1.91     | 1.45     | 2.48     | 1.47     |
| Al2O3  | 15.75    | 17.24    | 15.40    | 15.80    | 15.00    |
| Fe2O3  | 13.90    | 12.33    | 9.70     | 13.39    | 8.18     |
| MnO    | .23      | .21      | .19      | .20      | .13      |
| HgO    | 6.99     | 7.29     | 5.53     | 7.25     | 2.74     |
| CaO    | .96      | 9.40     | 9.17     | 9.54     | 7.83     |
| Na2O   | 2.40     | 2.41     | 2.93     | 3.30     | 4.00     |
| K2O    | .24      | .28      | .80      | .32      | .38      |
| LOI    | 3.60     | 3.00     | 3.10     | 3.20     | 5.80     |
| TOTAL  | 91.29    | 99.95    | 103.07   | 101.68   | 100.73   |

## TRACE ELEMENTS (PPM)

|       |     |     |     |     |     |
|-------|-----|-----|-----|-----|-----|
| Zr    | 167 | 151 | 219 | 156 | 206 |
| Sr    | 325 | 347 | 248 | 332 | 491 |
| Rb    | 17  | 14  | 18  | 17  | 17  |
| Zn    | 126 | 63  | 101 | 110 | 84  |
| Cu    | 62  | 49  | 27  | 58  | 19  |
| Co    | 209 | 208 | 172 | 114 | 114 |
| Cr S  | 61  | 52  | 26  | 62  | 11  |
| Mo S  | 17  | 11  | 12  | 6   | 15  |
| Sn S  | 26  | 11  | 25  | 27  | 22  |
| 27804 |     |     |     | 0   |     |
| Pb    | 23  | 11  | 25  | 23  | 24  |
| Bi    | 57  | 82  | 31  | 61  | 15  |
| Ge    | 45  | 38  | 23  | 41  | 22  |
| Or    | 61  | 58  | 52  | 128 | 45  |
| V     | 354 | 253 | 230 | 330 | 190 |
| Hg    | 21  | 18  | 25  | 54  | 23  |
| Ag    | 0   | 3   | 1   | 5   | 1   |

POOR COPY  
COPIE DE QUALITEE INFERIEURE

## HIGH TITANIUM DAIRASE

## MAJOR ELEMENTS (WT%)

| SAMPLE | 12140025 | 12140790 | 14930630 | 26661950 | 27640125 |
|--------|----------|----------|----------|----------|----------|
| SiO2   | 45.60    | 44.60    | 49.50    | 47.40    | 48.00    |
| TiO2   | 2.80     | 2.22     | 1.82     | 2.34     | 2.77     |
| Al2O3  | 12.10    | 15.57    | 16.10    | 15.46    | 13.24    |
| Fe2O3  | 13.32    | 12.44    | 10.70    | 11.70    | 14.05    |
| MnO    | .22      | .19      | .22      | .20      | .27      |
| H2O    | 3.80     | 7.09     | 6.10     | 6.51     | 4.70     |
| CaO    | 7.41     | 10.28    | 9.76     | 8.97     | 7.53     |
| Na2O   | 3.22     | 2.43     | 2.42     | 2.57     | 4.24     |
| K2O    | .41      | .16      | .82      | .92      | .20      |
| LOI    | 7.10     | 4.80     | 2.90     | 3.10     | 3.80     |
| TOTAL  | 95.08    | 100.58   | 99.94    | 99.37    | 98.10    |

## TRACE ELEMENTS (PPM)

|       |     |     |     |     |     |
|-------|-----|-----|-----|-----|-----|
| Zr    | 192 | 176 | 173 | 174 | 151 |
| Sr    | 310 | 408 | 419 | 421 | 145 |
| Pb    | 16  | 13  | 19  | 19  | 13  |
| Zn    | 161 | 109 | 55  | 77  | 95  |
| Cu    | 15  | 47  | 43  | 41  | 14  |
| Ta    | 476 | 13  | 13  | 77  | 100 |
| CoS   | 12  | 41  | 14  | 19  | 12  |
| MoS   | 11  | 7   | 11  | 8   | 14  |
| ZnS   | 37  | 16  | 19  | 10  | 34  |
| Fe2O4 |     | 0   |     | 0   |     |
| Pb    | 35  | 34  | 14  | 6   | 12  |
| Li    | 9   | 74  | 15  | 3   | 9   |
| Co    | 27  | 13  | 29  | 33  | 2   |
| Cr    | 36  | 170 | 176 | 173 | 13  |
| V     | 312 | 164 | 242 | 178 | 260 |
| Hg    | 49  | 36  | 18  | 7   | 12  |
| Ag    | 12  | 11  | 4   | 3   | 0   |
| S     |     |     |     |     |     |

POOR COPY  
COPIE DE QUALITE INFERIEURE

## HIGH TITANIUM DAIBASE

## MAJOR ELEMENTS (WT%)

SAMPLE 27040350 27040425 27040475

|       |       |       |       |
|-------|-------|-------|-------|
| SiO2  | 44.80 | 45.70 | 41.60 |
| TiO2  | 1.00  | 2.23  | 2.00  |
| Al2O3 | 15.46 | 14.93 | 14.65 |
| Fe2O3 | 11.70 | 13.05 | 11.70 |
| MnO   | .18   | .19   | .20   |
| MgO   | 7.40  | 6.87  | 6.28  |
| CaO   | 9.57  | 9.22  | 9.65  |
| Na2O  | 2.92  | 2.42  | 2.43  |
| K2O   | .37   | .40   | .50   |
| LOI   | 7.00  | 4.30  | 11.60 |

TOTAL 99.50 99.31 100.61

## TRACE ELEMENTS (PPM)

|        |     |     |     |
|--------|-----|-----|-----|
| Zr     | 141 | 136 | 128 |
| Cr     | 432 | 39  | 228 |
| Pb     | 18  | 16  | 22  |
| Zn     | 91  | 105 | 123 |
| Cu     | 49  | 56  | 45  |
| Co     | 107 | 11  | 2   |
| MoS    | 46  | 61  | 48  |
| FeS    | 13  | 13  | 14  |
| ZnS    | 13  | 12  | 19  |
| Fe-SO4 |     |     | 28  |
| Pb     | 13  | 15  | 15  |
| Si     | 04  | 67  | 57  |
| Co     | 5   | 13  | 13  |
| Cr     | 65  | 36  | 28  |
| V      | 310 | 437 | 265 |
| Hg     | 8   | 12  | 14  |
| Ag     | 2   | 0   | 1   |
| Pb     |     |     |     |

POOR COPY  
COPIE DE QUALITEE INFÉRIEURE

ALKALI FELDSPAR GRANITE

MAJOR ELEMENTS (WT%)

| SAMPLE | BJ51 55 | BJ52 60 | JGT/5124 | JGT 124A | JGT 233A |
|--------|---------|---------|----------|----------|----------|
| SiO2   | 74.50   | 73.00   | 74.50    | 76.90    | 75.90    |
| TiO2   | .20     | .28     | .06      | .12      | .10      |
| Al2O3  | 12.99   | 13.53   | 11.10    | 11.53    | 12.05    |
| Fe2O3  | 1.36    | 1.74    | 2.57     | 1.62     | .97      |
| MnO    | 0.00    | .03     | 0.00     | .03      | .01      |
| H2O    | .22     | .43     | .01      | .15      | .01      |
| CaO    | .67     | .71     | .01      | .31      | .27      |
| Na2O   | 4.10    | 4.21    | 3.45     | 3.93     | 3.57     |
| K2O    | 4.33    | 4.25    | 4.12     | 4.40     | 5.03     |
| LOI    | 1.00    | 1.00    | .70      | .50      | .60      |
| TOTAL  | 99.22   | 99.18   | 98.52    | 99.19    | 98.01    |

TRACE ELEMENTS (PPM)

|       |     |     |     |     |     |
|-------|-----|-----|-----|-----|-----|
| Zr    | 145 | 147 | 916 | 204 | 172 |
| Sn    | 72  | 67  | 15  | 14  | 23  |
| Pb    | 126 | 117 | 72  | 148 | 103 |
| Zn    | 14  | 15  | 122 | 65  | 7   |
| Cu    | 1   | 1   | 2   | 1   | 1   |
| B     | 7   | 6   | 36  | 15  | 19  |
| As    | 2   | 4   | 1   | 2   | 1   |
| Sb    | 10  | 11  | 12  | 31  | 2   |
| ZnS   | 1   | 1   | 33  | 37  | 1   |
| FeS04 | 0   | 0   | 0   | 0   | 0   |
| Pb    | 0   | 0   | 20  | 47  | 5   |
| Si    | 3   | 3   | 3   | 3   | 1   |
| Co    | 3   | 41  | 42  | 36  | 07  |
| Cr    | 7   | 77  | 6   | 5   | 6   |
| V     | 10  | 25  | 10  | 10  | 15  |
| Hg    |     |     |     |     |     |
| Ag    |     |     |     |     |     |
| Th    | 13  | 13  | 12  | 12  | 12  |

POOR COPY  
COPIE DE QUALITEE INFERIEURE

## ALKALI FELDSPAR GRANITE

## MAJOR ELEMENTS (WT%)

| SAMPLE                         | JGT25231 | JGT 231A | JGT25276 | JGT25277 | JGT 315A |
|--------------------------------|----------|----------|----------|----------|----------|
| SiO <sub>2</sub>               | 76.50    | 76.90    | 76.70    | 73.10    | 75.60    |
| TiO <sub>2</sub>               | .16      | .12      | .36      | .26      | .10      |
| Al <sub>2</sub> O <sub>3</sub> | 12.38    | 12.05    | 14.83    | 13.22    | 12.70    |
| Fe <sub>2</sub> O <sub>3</sub> | 1.31     | 1.30     | 2.50     | 1.81     | 1.28     |
| MnO                            | .01      | .01      | .04      | .03      | .01      |
| K <sub>2</sub> O               | .02      | .02      | .42      | .35      | .03      |
| CaO                            | .35      | .35      | .49      | .38      | .31      |
| MgO                            | 3.79     | 3.74     | 3.93     | 4.02     | 3.63     |
| Na <sub>2</sub> O              | 4.81     | 4.91     | 5.12     | 5.11     | 5.44     |
| LOI                            | .50      | .50      | 1.10     | .80      | .40      |
| TOTAL                          | 99.83    | 99.90    | 98.69    | 99.33    | 99.58    |

## TRACE ELEMENTS (PPM)

|    |     |     |     |     |     |
|----|-----|-----|-----|-----|-----|
| Cr | 145 | 114 | 263 | 129 | 124 |
| Co | 28  | 32  | 53  | 33  | 34  |
| Pb | 137 | 15  | 133 | 129 | 153 |
| Zn | 28  | 33  | 27  | 31  | 21  |
| Cu | 1   | 1   | 1   | 1   | 1   |
| Fe | 22  | 27  | 31  | 31  | 3   |
| Cl | 1   | 2   | 0   | 3   | 1   |
| Li | 8   | 5   | 1   | 1   | 5   |
| Na | 18  | 15  | 5   | 14  | 2   |
| B  | 0   | 0   | 0   | 0   | 0   |
| Pb | 7   | 12  | 8   | 5   | 11  |
| Bi | 1   | 1   | 1   | 1   | 5   |
| Co | 29  | 38  | 38  | 39  | 52  |
| Cr | 7   | 10  | 8   | 7   | 8   |
| U  | 10  | 10  | 10  | 10  | 10  |
| Th |     |     |     |     |     |
| La | 3   | 3   | 2   | 2   | 3   |

POOR COPY  
COPIE DE QUALITE INFERIEURE



## ALPALI FELDSPAR GRANITE

## MAJOR ELEMENTS (WTZ)

SAMPLE JVN 315B JGT 315C

|                                |       |       |
|--------------------------------|-------|-------|
| SiO <sub>2</sub>               | 74.20 | 72.20 |
| TiO <sub>2</sub>               | .26   | .28   |
| Al <sub>2</sub> O <sub>3</sub> | 13.10 | 13.49 |
| Fe <sub>2</sub> O <sub>3</sub> | 1.59  | 1.83  |
| MnO                            | .02   | .01   |
| H <sub>2</sub> O               | .12   | .20   |
| CaO                            | .49   | .78   |
| Na <sub>2</sub> O              | 3.84  | 3.80  |
| K <sub>2</sub> O               | 5.46  | 5.58  |
| LOI                            | .50   | .50   |

|       |       |       |
|-------|-------|-------|
| TOTAL | 99.58 | 98.47 |
|-------|-------|-------|

## TRACE ELEMENTS (PPM)

|                   |     |     |
|-------------------|-----|-----|
| Zr                | 178 | 165 |
| Sr                | 42  | 77  |
| Rb                | 122 | 147 |
| Zn                | 21  | 29  |
| Cu                | 1   | 1   |
| Ba                | 103 | 9   |
| ClS               | 3   | 2   |
| MS                | 3   | 3   |
| ZnS               | 4   | 18  |
| BaSO <sub>4</sub> | 6   | 6   |
| Pb                | 29  | 21  |
| Ni                | 1   | 1   |
| Co                | 62  | 62  |
| Cr                | 7   | 9   |
| V                 | 10  | 10  |
| Hg                |     |     |
| Ag                |     |     |
| As                | 14  | 15  |

POOR COPY  
COPIE DE QUALITEE INFERIEURE

3-117

HYBIRD GRANITE

MAJOR ELEMENTS (WT%)

SAMPLE JGT75279

|       |       |
|-------|-------|
| SiO2  | 58.48 |
| TiO2  | 1.62  |
| Al2O3 | 13.74 |
| Fe2O3 | 8.68  |
| MnO   | .16   |
| MgO   | 3.36  |
| CaO   | 4.88  |
| Na2O  | 3.89  |
| K2O   | 2.57  |
| LOI   | 1.48  |

TOTAL 98.62

TRACE ELEMENTS (PPM)

|       |     |
|-------|-----|
| Zr    | 288 |
| Sr    | 164 |
| Rb    | 64  |
| Zn    | 56  |
| Cu    | 17  |
| Ba    | 48  |
| CuS   | 15  |
| PbS   | 5   |
| ZnS   | 11  |
| BaSO4 | 0   |
| Pb    | 21  |
| Ni    | 18  |
| Co    | 32  |
| Cr    | 38  |
| V     | 175 |
| Hg    |     |
| Ag    |     |
| Nb    | 20  |

POOR COPY  
COPIE DE QUALITEE INFERIEURE

3-118

## RHYOLITE DYKES

## MAJOR ELEMENTS (WT%)

| SAMPLE | 07920300 | 08681000 | 08690175 | 08691925 |
|--------|----------|----------|----------|----------|
| SiO2   | 71.46    | 73.00    | 65.80    | 67.60    |
| TiO2   | .38      | .13      | .11      | .10      |
| Al2O3  | 13.61    | 11.29    | 15.30    | 13.40    |
| Fe2O3  | 2.88     | 3.24     | 1.27     | 1.45     |
| MnO    | .06      | .11      | .10      | .08      |
| MgO    | .68      | .20      | .62      | .51      |
| CaO    | 1.79     | 1.42     | 5.26     | 5.48     |
| Na2O   | 4.07     | 3.35     | .57      | 4.39     |
| K2O    | 1.91     | 3.49     | 3.59     | 1.87     |
| LOI    | 2.50     | 2.10     | 7.30     | 5.50     |
| TOTAL  | 99.28    | 98.33    | 99.92    | 100.38   |

## TRACE ELEMENTS (PPH)

|       |     |     |     |     |
|-------|-----|-----|-----|-----|
| Zr    | 183 | 562 | 91  | 61  |
| Sr    | 223 | 0   | 211 | 172 |
| Rb    | 46  | 63  | 80  | 52  |
| Zn    | 44  | 129 | 25  | 31  |
| Cu    | 3   | 8   | 5   | 3   |
| Ba    | 873 | 743 | 582 | 425 |
| CuS   | 0   | 5   | 1   | 1   |
| PbS   | 13  | 12  | 22  | 18  |
| ZnS   | 13  | 55  | 11  | 3   |
| BaSO4 | 0   | 0   | 0   | 0   |
| Pb    | 19  | 19  | 20  | 27  |
| Ni    | 2   | 3   | 2   | 1   |
| Co    | 3   | 2   | 1   | 2   |
| Cr    | 34  | 23  | 18  | 21  |
| V     | 4   | 11  | 7   | 13  |
| Hg    | 13  | 82  | 137 | 19  |
| Ag    | 1   | 0   | 2   | 0   |
| Nb    |     |     |     |     |

POOR COPY  
COPIE DE QUALITEE INFERIEURE

## PERALKALINE GRANITE

## MAJOR ELEMENTS (WT%)

| SAMPLE                         | BAFF  | 2     | JBT75265 | JBT75266 | JBT75274 | JBT75275 |
|--------------------------------|-------|-------|----------|----------|----------|----------|
| SiO <sub>2</sub>               | 75.88 | 76.98 | 74.88    | 77.28    | 76.98    |          |
| TiO <sub>2</sub>               | .32   | .14   | .16      | .18      | .12      |          |
| Al <sub>2</sub> O <sub>3</sub> | 11.69 | 11.42 | 11.46    | 11.81    | 11.81    |          |
| Fe <sub>2</sub> O <sub>3</sub> | 2.61  | 2.22  | 2.66     | 2.61     | 1.87     |          |
| MnO                            | .86   | .84   | .88      | 8.88     | .87      |          |
| H <sub>2</sub> O               | .81   | .81   | .81      | .81      | .81      |          |
| CaO                            | .48   | .89   | .49      | .85      | .13      |          |
| Na <sub>2</sub> O              | 4.27  | 3.88  | 3.94     | 3.46     | 3.97     |          |
| K <sub>2</sub> O               | 4.79  | 4.53  | 4.41     | 4.86     | 4.78     |          |
| LOI                            | .38   | .48   | 1.28     | .78      | .38      |          |
| TOTAL                          | 99.45 | 99.63 | 98.41    | 99.28    | 99.88    |          |

## TRACE ELEMENTS (PPM)

|                   |    |     |     |     |     |
|-------------------|----|-----|-----|-----|-----|
| Zr                |    | 458 | 471 | 834 | 167 |
| Sr                | 17 | 17  | 27  | 13  | 19  |
| Rb                | 88 | 94  | 76  | 98  | 198 |
| Zn                | 21 | 57  | 118 | 189 | 29  |
| Cu                | 1  | 7   | 5   | 1   | 1   |
| Ba                |    | 533 | 955 | 326 | 281 |
| CUS               | 1  | 7   | 4   | 2   | 1   |
| PbS               | 24 | 17  | 8   | 11  | 3   |
| ZnS               | 21 | 45  | 187 | 46  | 2   |
| BasO <sub>4</sub> |    | 8   | 8   | 8   | 8   |
| Pb                | 26 | 28  | 19  | 28  | 18  |
| Ni                | 3  | 3   | 4   | 2   | 1   |
| Co                | 53 | 54  | 38  | 32  | 61  |
| Cr                |    | 6   | 18  | 7   | 7   |
| V                 | 18 | 18  | 18  | 18  | 18  |
| Hg                |    |     |     |     |     |
| Ag                |    |     |     |     |     |
| Nb                |    | 31  | 29  | 47  | 92  |

POOR COPY  
COPIE DE QUALITEE INFERIEURE

3-120

SPRINODALÉ GROUP (T) IGNIMBRITE

MAJOR ELEMENTS (WT%)

| SAMPLE | JBT 266A |
|--------|----------|
| S102   | 76.70    |
| T102   | .14      |
| Al2O3  | 12.38    |
| Fe2O3  | 1.18     |
| MnO    | .03      |
| H2O    | .01      |
| CaO    | .05      |
| Na2O   | 5.18     |
| K2O    | 1.58     |
| LOI    | 1.38     |
| TOTAL  | 99.35    |

TRACE ELEMENTS (PPM)

|       |     |
|-------|-----|
| Zr    | 89  |
| Sr    | 40  |
| Rb    | 52  |
| Zn    | 15  |
| Cu    | 1   |
| Ba    | 213 |
| CuS   | 1   |
| PbS   | 2   |
| ZnS   | 1   |
| BaSO4 | 0   |
| Pb    | 12  |
| Ni    | 1   |
| Co    | 45  |
| Cr    | 8   |
| V     | 10  |
| Hg    |     |
| Ag    |     |
| Nb    | 17  |

## HUNGRY MOUNTAIN COMPLEX

## MAJOR ELEMENTS (WT%)

| SAMPLE                         | JGT 317A | 2856 145 | 2856 317 |
|--------------------------------|----------|----------|----------|
| SiO <sub>2</sub>               | 64.18    | 73.98    | 73.88    |
| TiO <sub>2</sub>               | .54      | .14      | .34      |
| Al <sub>2</sub> O <sub>3</sub> | 15.88    | 13.33    | 12.68    |
| Fe <sub>2</sub> O <sub>3</sub> | 5.21     | 1.59     | 2.38     |
| MnO                            | .87      | .83      | .85      |
| MgO                            | 1.63     | .62      | 1.89     |
| CaO                            | 4.84     | 1.88     | 1.86     |
| Na <sub>2</sub> O              | 3.45     | 3.25     | 3.66     |
| K <sub>2</sub> O               | 2.33     | 5.88     | 2.68     |
| LOI                            | 1.28     | 1.88     | 1.28     |
| TOTAL                          | 98.45    | 99.86    | 99.74    |

## TRACE ELEMENTS (PPM)

|                   |      |      |     |
|-------------------|------|------|-----|
| Zr                | 166  | 97   | 86  |
| Sr                | 352  | 218  | 197 |
| Rb                | 68   | 119  | 67  |
| Zn                | 31   | 25   | 28  |
| Cu                | 1    | 6    | 1   |
| Ba                | 1621 | 1858 | 885 |
| CuS               | 1    | 6    | 2   |
| PbS               | 6    | 15   | 22  |
| ZnS               | 3    | 12   | 11  |
| BaSO <sub>4</sub> | 8    | 8    | 8   |
| Pb                | 18   | 15   | 15  |
| Ni                | 1    | 44   | 7   |
| Co                | 38   | 48   | 38  |
| Cr                | 12   | 18   | 18  |
| V                 | 88   | 15   | 38  |
| Hg                |      |      |     |
| Ag                |      |      |     |
| Nb                | 6    | 7    | 5   |

## MISCELLANEOUS SAMPLES

## MAJOR ELEMENTS (WT%)

| SAMPLE                         | 08481612 | 12141315 |
|--------------------------------|----------|----------|
| SiO <sub>2</sub>               | 72.60    | 70.10    |
| TiO <sub>2</sub>               | 0.00     | .35      |
| Al <sub>2</sub> O <sub>3</sub> | 1.82     | 11.43    |
| Fe <sub>2</sub> O <sub>3</sub> | 19.12    | 6.64     |
| MnO                            | .11      | .20      |
| MgO                            | .41      | 2.57     |
| CaO                            | 3.54     | 1.42     |
| Na <sub>2</sub> O              | .05      | 2.60     |
| K <sub>2</sub> O               | .25      | 1.35     |
| LOI                            | 2.60     | 2.70     |
| TOTAL                          | 100.50   | 99.36    |

## TRACE ELEMENTS (PPM)

|                   |     |     |
|-------------------|-----|-----|
| Zr                | 25  | 123 |
| Sr                | 40  | 211 |
| Rb                | 18  | 30  |
| Zn                | 34  | 71  |
| Cu                | 4   | 50  |
| Ba                | 686 | 143 |
| CuS               | 1   | 43  |
| PbS               | 12  | 12  |
| ZnS               | 2   | 13  |
| BaSO <sub>4</sub> |     |     |
| Pb                | 22  | 19  |
| Ni                | 5   | 7   |
| Co                | 9   | 7   |
| Cr                | 34  | 36  |
| V                 | 21  | 48  |
| Hg                | 15  | 245 |
| Ag                | 0   | 1   |
| Nb                |     |     |

## APPENDIX IV

### FACTOR ANALYSIS

R-mode factor analysis used in this study was computed by Newfoundland and Labrador Computer Services using program BMD08M. A general description from their program manual is as follows:

#### General Description, Factor Analysis Program BMD08M

This program performs a factor analysis of up to 198 input variables. The factoring may be done using either covariance or correlation matrices. Initial communality estimates may be squared multiple or they may be specified by the user. If requested, the program will iterate on the initial communality estimates. Three types of rotation are available, all based on the oblimin criterion. In the first, the factors are restricted to be orthogonal which yields, among others, quartimax and varimax rotations. In the second, the criterion are applied to the reference factor structure and the factors are allowed to be oblique which yields the standard oblimin rotations. In the third, the criterion are applied to the primary factor loadings, allowing the factors to be oblique and yielding simple loadings rotations (1). Factor scores may be estimated. Data input may be in the form of raw data, a correlation matrix, a covariance matrix, or a factor loading matrix.



Output from this program includes:

- 1) Means and standard deviations
- 2) Correlation or covariance matrix
- 3) Eigenvalues and cumulative proportion of total variance
- 4) Communalities
- 5) Factor loading matrix before rotation
- 6) Rotated factor loading matrix
- 7) Correlation matrix of the rotated factors
- 8) Factor scores

TOPSAILS GRANITE

JGT.-276

• JGT.-280  
• JGT.-279  
• JGT.-278  
• JGT.-277

JGT.-315

JGT.-1668

H-2622(4)

H 2820

H 2867

H 2528

H 2530

H 2571

H 2769 •  
H 2777 •

H2880

H 2350

H 2882 •

H 2879 •

H2885(2)

H 2886(2)

H 2888

H 2501 (2)

H 256

10F

JGT-124

JGT-317<sup>A</sup>

# HUNGRY MOUNTAIN COMPLEX

H 2856(2)

H 2501(2)

H 2855(2)

H 2852

H 2490

BUCHANS  
LAKE

H 2528

H 2090(2)

H 2858

H 2851(6)

2622(4)

H 2530

H 2571

H 2762

H 1960

H 1940(2)

H 2841(2)

H 2680 H 925

H 2105(3)

H 2880

H 2610

H 2666(40)

H 2043

H 1903

H 1911

H 2348

H 1917 H 2704

H 1493

H 969

H 968

JGT-170

H 1258

H 2769

H 2777

H 2710(2)

H 2785(2)

H 1214

JGT-40-43

H 885

H 1083

H 2800

H 1096

JGT-44

H 1406

H 2771(2)

H 782

H 202

H 75

H 542

H 1908

H 645(7)

H 932

H 2880

H 2901

H 2899

H 2882

H 2900

H 2879

H 2887

H 2889

H 2886(2)

H 2884

H 2350

H 2886(2)

H 2885

H 978(2)

2461

COMPLEX

BUCHANS  
LAKE

BUCHANS  
GROUP

JGT.-113

JGT.-317<sup>A</sup>

H2839 (4)

H 2837 (17)

H 2842

H 1920

H1762

H 1662

H1663

H 1589 (3)

H1568

H1730

H296

H1352

H1062

H972

H158 (2)

H1872

H2367

H1475

H885

H1083

H1096

JGT. 44

JGT.-170

JGT.-40-43

H2800

(2)

JGT.-328

H792

H202

H175

H542

H1908

H 2856 (2)

352

31 (6)

(2)

H1917 B2704

H1493

H2785 (2)

2771 (2)

H932

78 (2)

B.J.-46

B.J.-4

B.J.-47

B

34

TOPSAILS GRANITE

• B.J. 50

• B.J. 51

• B.J. 46  
• B.J. 45  
• B.J. 47  
• B.J. 48  
• B.J. 49

• B.J. 10

BUCHANS  
GROUP

• B.J. 42 (8)

4 of 4

ITE

JGT-265

JGT-266-266<sup>A</sup>

B.J.50

B.J.51


B.J.52

HUNGRY  
MOUNTAIN  
COMPLEX

B.J.-42(8)


5af

(1) JO  
AL  
LO  
(2) NU  
HO  
ON



LEGEND

- (1) JGT. SAMPLES ARE FROM OUTCROP  
ALL OTHER ARE DRILL HOLE SAMPLE  
LOCATIONS.
- (2) NUMBER OF SAMPLES TAKEN FROM DRILL  
HOLES IN BRACKETS, IF GREATER THAN  
ONE.



R.I.-5

JGT-219

JGT-218

60F

JGT-216

JGT.-276

● JGT.-279  
● JGT.-278  
● JGT.-277

H 2622 (4)

H 2530

H 2571

TOPSAILS GRANITE

H 2620

H 2667

H 2769 ●  
H 2777 ●

JGT.-1668

H 2660

JGT.-315

H 2662 ●

H 2679 ●

H 2666 (2)

H 2350

H 2666

H 2666

JGT.-307

H 724 ● JGT.-305

● H 718

JGT.-2271 ● 231

FEEDER GRANODIORITE

JGT.-234

● B.B.257


● B.B.246


● B.B.247

B.B.229 ● B.B.232

70F







1920

32

• B.J.-46  
• B.J.-45  
• B.J.-49  
• B.J.-47  
• B.J.-48

• B.J.10

BUCHANS  
GROUP

• B.J.-42(6)

• JGT.-134

RED INDIAN LAKE

10 of

J.10

B.J.-42(8)

JGT-216

AN LAKE

11 OF

0 1000 2000 3000  
Meters

R.L.-5

JGT.-219

JGT.-218

JGT.-216

120F12

FIGURE 1-2

SAMPLE LOCATION MAP

1" = 5333' OR 1 : 40,000

57°04'

57°00'

48°57'

HIND'S  
LAKE

Sugarloaf  
Mt.  
1250

River

Little  
Hind  
Lake

Orchid  
Lake

Bluff  
Hills

5A<sub>0</sub>?

5A<sub>0</sub>?

1/0F

9a

White  
Hills

Noah's  
Ballast

5A<sub>0</sub>?


8b


Academy  
Bay

5A<sub>0</sub>?

50'

9a









50f



60F

Parts of 12A/10, 12A/11, 12A/14, 12A/15





50°

45°



Lake

Wharf

BM 793

Woodman's Brook

Don River

Tower

Revere Camp

BM 716


RED

INDIAN

GEO

# SYMBOLS

- Geological boundary (defined, approximate, assumed) .....
- Bedding, tops known (inclined, vertical) .....
- Bedding, tops unknown (inclined, vertical) .....
- Pillow bedding, tops known (inclined, vertical) .....
- Flow banding (inclined, vertical) .....
- Schistosity, foliation (inclined, vertical) .....
- Foliation due to thrusting (inclined, vertical) .....
- Anticline (approximate) .....
- Syncline (approximate) .....
- Thrust fault (defined, approximate or assumed) .....
- Fault (defined, approximate, assumed) .....
- In situ and transported siltcliffs, extreme outline projected to surface .....



## CARBONIFEROUS

- Poorly indurated red sandstone and conglomerate

## SILURIAN AND DEVONIAN

- Reddish subaerial ignimbrite, flow-banded rhyolite and pyroclastics. Extrusive equivalents of the Topsails Granite. 10a, subvolcanic quartz-feldspar porphyry


## TOPSAILS GRANITE

- Alkali Feldspar Granite ( $387 \pm 16$  Ma): Fine grained, brick-red granite. Locally approaches syenite. Locally associated bodies of fine grained quartz-feldspar porphyry.
- Mafic Intrusives (associated with Topsails Granite): Fine to coarse grained diabase (9b<sub>1</sub>) and gabbro (9b<sub>2</sub>).
- Peralkaline Granite ( $421 \pm 7$  Ma): Medium to coarse grained, equigranular, white-weathering granite.


Geology compiled by Asarco Inc. and Price Company Limited (now Abitibi-Price Inc.). Geology in remote areas compiled in consultation with Department of Mines and Energy, Government of Newfoundland and Labrador.

Whole rock Rb/Sr isochrons determined by Bell and Richardson using the decay constant  $\lambda = 8.7$

1 90F



# GEOLOGICAL MAP OF BUCHANS, ARE NEWFOUNDLAND



## LEGEND

### ORDOVICIAN - SILURIAN

#### BUCHANS GROUP (447 ± 18 Ma)

##### Upper Buchans Subgroup

indurated red sandstone and conglomerate

### DEVONIAN

sh subaerial ignimbrite, flow-banded rhyolite and pyroclastics. Extrusive  
vents of the Topsails Granite. 10a, subvolcanic quartz-feldspar porphyry.

### QUATERNARY

Feldspar Granite (387 ± 16 Ma): Fine grained, brick-red granite. Locally  
rich in syenite. Locally associated bodies of fine grained quartz-feldspar  
porphyry.

Intrusives (associated with Topsails Granite): Fine to coarse grained  
granite (90%) and gabbro (9%).

Albite Granite, (421 ± 7 Ma): Medium to coarse grained, equigranular,  
weathering granite.

Lake Seven Basalt: Basaltic, amygdaloidal, feldsparphyric, hematitized lava,  
pillow lava, breccia and minor tuff; 8a, arkosic interbeds

##### Lower Buchans Subgroup

Lucky Strike Ore Horizon Sequence: Predominantly felsic sequence of  
pyroclastics, flows, breccias and clastic sediments; 5A, small rhyolite units at  
base, flanked by pyritic siltstone and wacke, followed by 5A, dacitic crystal-vitric  
tuffs gradational with 5A, lithic-rich pyroclastic breccias and in situ massive and  
fragmental transported sulfides; 5A, granite-bearing breccia-conglomerate;  
5A, mafic interbeds.

Intermediate Felsite: Complex assemblage of mafic to felsic pyroclastics,  
breccias, minor flows. Locally altered, mineralized.

10 of


Little Sandy  
Lake

8b

L A K E

# MAP OF BUCHANS AREA NEWFOUNDLAND

Scale 1:50,000 Échelle



## LEGEND

± 18 Ma)

ait: Basaltic, amygdaloidal, feldsparphyric, hematitized lava,  
ie and minor tuff; 6a, eskosic interbeds

re Horizon Sequence: Predominantly felsic sequence of  
n, breccias and clastic sediments; 5A, small rhyolite units at  
pyritic siltstone and wacke, followed by 5A, dacitic crystal-vitric  
with 5A, silic-rich pyroclastic breccias and in situ massive and  
spotted sulfides; 5A, granite-bearing breccia-conglomerate;  
wt.


ault: Complex assemblage of mafic to felsic pyroclastics,  
lows. Locally altered, mineralized.

**8b** Little Sandy Sequence: Interfingering and inter-  
felsic volcanic rocks with the latter predom-  
pillow lava, breccia, pyroclastic flows and  
volcanics; 8b, dacitic breccia and quartz-fel-  
quartz-feldspar crystal tuffs, breccias, tuffs, p-  
massive flows; 8d, dominantly intermediate fra-  
and mineralized.

**7a** Upper Archaean: Resedimented arkosic congl-  
graywacke, siltstone; 7a, interbeds of mafic vo-

**5b** Oriental Ore Horizon Sequence: Complex as-  
breccias, flows and clastic sediments. Similar to  
the Lucky Strike Ore Horizon Sequence; 5b, coe-  
lows, flanked and overlain by 5b, pyritic silt-  
dacitic crystal-vitric tuffs gradational with poly-  
silt massive and fragmental transported tuff-  
with 5b, granite-bearing breccia-conglomerate


**6a** Old Hill Sequence: Stratigraphic equivalent of  
pillow lava, porphyritic and amygdaloidal and




11 of







13DF



Geological boundary (defined, approximate)

Bedding, tops known (inclined, vertical)

Bedding, tops unknown (inclined, vertical)

Pillow bedding, tops known (inclined, vertical)

Flow banding (inclined, vertical)

Schistosity, foliation (inclined, vertical)

Foliation due to thrusting (inclined, vertical)

Anticline (approximate)

Syncline (approximate)

Thrust fault (defined, approximate or assumed)

Fault (defined, approximate, assumed)

In situ and transported sulfides, extrusive

Geology compiled by Asarco Inc. and Provincial  
remote areas compiled in consultation with  
Newfoundland and Labrador.

Whole rock Rb/Sr isochrons determined  
Rb of  $1.42 \times 10^{11}$  yr<sup>-1</sup>

Map accompanies manuscript "Geology of  
Newfoundland", by J. G. Thurlow and  
Thurlow, J. G., eds., The Buchans Orebelt  
Canada Spec. Paper 22, in press.

Geological cartography by Mineral Development  
Government of Newfoundland and Labrador

Base maps at same scale published by  
Mines and Resources, Ottawa

Approximate magnetic declinations, 1960  
annually.

Elevations in feet above mean sea level

Colour separation and printing negatives prepared by Maritime  
Resource Management Service, Amherst, Nova Scotia, 1981

This Provisional Map is equivalent to a standard map  
accuracy of contour


Some names on this map are not yet official. Correct  
names or additions are invited by the Survey and  
Mapping Branch

7405



# SYMBOLS

- Geological boundary (defined, approximate, assumed) .....
- Bedding, tops known (inclined, vertical) .....
- Bedding, tops unknown (inclined, vertical) .....
- Pillow bedding, tops known (inclined, vertical) .....
- Flow bending (inclined, vertical) .....
- Schistosity, foliation (inclined, vertical) .....
- Foliation due to thrusting (inclined, vertical) .....
- Anticline (approximate) .....
- Syncline (approximate) .....
- Thrust fault (defined, approximate or assumed) .....
- Fault (defined, approximate, assumed) .....
- In situ and transported sulfides, extreme outline projected to surface .....



## CARBONIFEROUS

Poorly indurated red sandstone and conglomerate

## SILURIAN AND DEVONIAN

Reddish subaerial ignimbrite, flow-banded rhyolite equivalents of the Topsails Granite 10a, subvolcanic

## TOPSAILS GRANITE

9a Alkali Feldspar Granite (387 ± 16 Ma): Fine grained, approaches syenite. Locally associated bodies of porphyry.

Mafic Intrusives (associated with Topsails Granite): diabase (9b<sub>1</sub>) and gabbro (9b<sub>2</sub>).

9c Peralkaline Granite (421 ± 7 Ma): Medium to coarse grained, white weathering granite.

Geology compiled by Asarco Inc. and Price Company Limited (now Abitibi-Price Inc.). Geology in remote areas compiled in consultation with Department of Mines and Energy, Government of Newfoundland and Labrador.

Whole rock Rb/Sr isochrons determined by Bell and Blenkinsop using the decay-constant for <sup>87</sup>Rb of  $1.42 \times 10^{-11} \text{ yr}^{-1}$

Map accompanies manuscript "Geology and Ore Deposits of the Buchans Area, Central Newfoundland", by J.G. Thurlow and E.A. Swenson in Swenson, E.A., Strong, D.F. and Thurlow, J.G., eds., The Buchans Orebodies: Fifty Years of Geology and Mining: Geol. Assoc. Canada Spec. Paper 22, in press.

Geological cartography by Mineral Development Division, Department of Mines and Energy, Government of Newfoundland and Labrador.

Base maps at same scale published by the Surveys & Mapping Branch, Department of Energy, Mines and Resources, Ottawa.

Approximate magnetic declinations, 1989, for center of map, 28°39'W, decreasing 3.0' annually.

Elevations in feet above mean sea level.

and printing prepared by Bureau of Geology, Department of Energy, Mines and Resources, Ottawa, 1988.

1:50,000 is equivalent to a standard map 1:50,000.

This map is not an official map. Copies are issued by the Surveys and Mapping Branch.

15 of 15

RED

INDIAN

LAKE

# GEOLOGICAL MAP OF BUCHANAN NEWFOUNDLAND

Scale 1:50,000 Échelle



## LEGEND

### ORDOVICIAN - SILURIAN

#### BUCHANAN GROUP (447 ± 18 Ma)

##### Upper Buchans Subgroup

**Lake Seven Basalt:** Basaltic, amygdaloidal, feldsparphyric, hematitized lava, pillow lava, breccia and minor tuff; 6a, arkosic interbeds.

##### Lower Buchans Subgroup

**Lucky Strike Ore Horizon Sequence:** Predominantly felsic sequence of pyroclastics, flows, breccias and clastic sediments; 5A, small rhyolite units at base, flanked by pyritic siltstone and wacke, followed by 5A, dacitic crystal-vitric tuffs gradational with 5A, lithic-rich pyroclastic breccias and in situ massive and fragmental transported sulfides; 5A, granite-bearing breccia-conglomerate; 5A, mafic interbeds.

**Intermediate Footwall:** Complex assemblage of mafic to felsic pyroclastics, breccias, minor flows. Locally altered, mineralized.

**Footwall Arkose:** 3A, Lithic arkose, with local thin, widely separated interbeds of mudstone, siltstone, siliceous graywacke, conglomerate; 3A, sequence of pumiceous felsic pyroclastics and interbedded tuffaceous sediments, underlying lithic arkose east of Buchans; 3A, basaltic lava, pillow lava, pillow breccia, lesser pyroclastics and lenses of bedded chert.

**Footwall Basalt:** 2a, Basaltic lava, pillow lava and pillow breccia, wealdy to strongly amygdaloidal, feldsparphyric and hematitized, with lesser pyroclastics and discontinuous lenses of multicolored bedded chert. Host for Skidder Brook massive sulfide deposit; 2a, felsic flows, breccia and tuff; 2a, rhyolite.

### ORDOVICIAN OR OLDER

**Hungry Mountain Complex (800 ± 70 and 400 ± 60 Ma):** Massive to foliated biotite granodiorite, hornblende diorite, gabbro and amphibolite with less abundant foliated granite.

### CARBONIFEROUS

**Poorly indurated red sandstone and conglomerate**

### SILURIAN AND DEVONIAN

**Reddish subaerial ignimbrite, flow-banded rhyolite and pyroclastics. Extrusive equivalents of the Topsails Granite. 10a, subvolcanic quartz-feldspar porphyry.**

### TOPSAILS GRANITE

**Alkali Feldspar Granite (387 ± 16 Ma):** Fine grained, brick-red granite. Locally approaches syenite. Locally associated bodies of fine grained quartz-feldspar porphyry.

**Mafic Intrusives (associated with Topsails Granite):** Fine to coarse grained diabase (9b<sub>1</sub>) and gabbro (9b<sub>2</sub>).

**Peralkaline Granite (421 ± 7 Ma):** Medium to coarse grained, equigranular, white-weathering granite.

Geology Inc.). Geology of  
Government of

pay-constant for 87

hane Area, Central  
Strong, D.F. and  
Ining: Geol. Assoc.

Mines and Energy

Department of Energy.

V. decreasing 3.0'

16 4

Scale 1:50,000 Échelle

Scale 1:50,000 Échelle

**LEGEND**

**DEVICIAN - SILURIAN**

**HANS GROUP (447 ± 18 Ma)**

Upper Buchans Subgroup

Like Seven Gables: Basaltic, amygdaloidal, feldsparphyric, hematitized lava, pillow lava, breccia and minor tuff; 6a, arkotic interbeds

**Lower Buchans Subgroup**

**5** **lucky series One Herstein Sequence:** Predominantly felsic sequence of pyroclasticites, flows, breccias and clastic sediments; **5a**, small rhyolite units at base, flanked by pyritic alterations and wealds, followed by **5b**, dacitic rhyolite with (a) gradational with **5a**, (b) rhyolite-rich pyroclastic breccias and in situ massive and fragmental transported surfaces; **5c**, granite-bearing breccia-conglomerates; **5d**, mafic interbeds.

**Intermediate Feeder:** Complex assemblage of mafic to felsic pyroclastics, breccias, minor flows. Locally altered, mineralized.

**Footwall Arkose:** 3b. Light arkose, with local thin, widely separated interbeds of mudstones, siltstones, siltyclays, grayclays, conglolites. 3a, sequence of lithomucous talic pyroclastics and interbedded bituminous sediments, underlying the arkose east of Buckhorn. 3c, basaltic lava, pillow lava, pillow breccia, lesser pyroclastics and lenses of bedded chert.

**Footwall Breccia:** 2a, Basaltic lava and pillow breccia, weakly to strongly amygdaloidal, hydrophyritic and hematitized, with lesser pyroclastics and discontinuous lenses of multicolored banded chert. Host for Stibbar Brook massive sulfide deposit; 2b, Mafic flows, breccia and tuff; 3a, mylonite.

**PHYSICIAN OR OTHER**

1 Hungry Mountain Complex (600±70 and 400±50 Ma): Massive to foliated biotite gneissoids, hornblende diorite, gabbro and amphibolite with less abundant foliated granite.

**Wileys' Premium Quartz Sequences:** Stratigraphic equivalent of 3a, felsic pyroclastics characterized by quartz crystals commonly exceeding 10 mm in diameter; 3ay, myofite flows and tuffs; 3bx, acidic pyroclastics; 3bx, interbeds of basaltic lavas; pillow lavas; 3bx, multicolored tuffaceous siltstones, siltstones, mudstones and chert.

Feeder Gneissodiorite (410 ± 80 Ma): Pink, medium to coarse grained, biotite gneissodiorite, probably synchronous with 34g and other, Buchans Group rocks.

**Little Sandy Sequence:** felsic volcanic rocks w/ pillow lava, breccia, & quartz-feldspar crystal massive flows. **Sd.** domes and mineralized.

Upper Arkose: Residual  
graywacke, siltstone. 74  
6.

**Oriental Ore** Mexican breccias, flows and clasts to Lucky Strike Ore horizon, flanked and overlain by dacitic crystal-rich tuffite, silty massive and fragmentary with 50% granitoid debris.

**S&H Min Sequence:** Strata pillow lava, porphyritic coarse pyroclastics, ind overlie in subsurface pumiceous pyroclastics


8-

35

# BUCHANS AREA

3 Miles

4000 Metres



**Little Sandy Sequence:** Interfingering and interbedded mafic, intermediate and felsic volcanic rocks with the latter predominant; 8a, basaltic and andesitic pillow lava, breccia, pyroclastic flows and tuff, minor unseparated felsic volcanics; 8b, dacitic breccia and quartz-feldspar crystal tuff; 8c, rhyolitic quartz-feldspar crystal tuffs, breccias, tuffs, pyroclastic flows, flow-banded to massive flows; 8d, dominantly intermediate fragmental volcanics, locally altered and mineralized.



**Upper Arkose:** Resedimented arkosic conglomerate, lithic arkose, siliceous graywacke, siltstone; 7a, interbeds of mafic volcanics, possibly correlative with 6.



**Oriental Ore Horizon Sequence:** Complex assemblage of felsic pyroclastics, breccias, flows and clastic sediments. Similar lithologically and stratigraphically to Lucky Strike Ore Horizon Sequence; 5B, coalescing rhyolite breccia domes at base, flanked and overlain by 5B, pyritic siltstone and wacke, followed by 5B, dacitic crystal-vitric tuffs gradational with polyolithic breccia-conglomerate and in situ massive and fragmental transported sulfides which in turn are associated with 5B, granitoid-bearing breccia-conglomerate.



**Ski Hill Sequence:** Stratigraphic equivalent of Intermediate Footwall. Basaltic pillow lava, porphyritic and amygdaloidal andesitic flow breccias and fine to coarse pyroclastics. Includes minor polyolithic breccias near stratigraphic top, overlain in subsurface by altered intermediate to felsic flows, breccias and pumiceous pyroclastics (Oriental Intermediate).



**Prominent Quartz Sequence:** Stratigraphic equivalent of 3A: Dacitic (3B<sub>1</sub>) to rhyolitic (3B<sub>2</sub>) pyroclastics; 3B<sub>1</sub> interbedded lenses of graded pumiceous subaqueous pyroclastic flows, cherty mudstone and siltstone; 3B<sub>2</sub> locally important basaltic members.

**Wiley's Prominent Quartz Sequence:** Stratigraphic volcanic equivalent of 3A: felsic pyroclastics characterized by quartz crystals commonly exceeding 10 mm in diameter; 3A<sub>1</sub>, rhyolite flows and tuffs; 3A<sub>2</sub>, dacitic pyroclastics; 3A<sub>3</sub>, interbeds of basaltic lavas; pillow lavas; 3A<sub>4</sub>, multicolored tuffaceous siltstone, siltstone, mudstone and chert.

**Feeder Granodiorite (410 ± 80 Ma):** Pink, medium to coarse grained, biotite granodiorite, probably synchronous with 3A<sub>1</sub> and other Buchans Group rocks.

184/18

FIG. 2-2





