PROTOZOA-CILIATA OF A SMALL POND AT LOGY BAY, NEWFOUNDLAND

CENTRE FOR NEWFOUNDLAND STUDIES

TOTAL OF 10 PAGES ONLY MAY BE XEROXED

(Without Author's Permission)

CHIA-MEI HSU KLEINE



G. 1 385594 17 CENTRE FOR NFLD. STUDIES JCT 6 1975 OF NEWFOUNDLAND

. PROTOZOA-CILIATA OF A SMALL POND AT LOGY BAY, NEWFOUNDLAND.

by.

CHIA-MEI HSU. B.Sc. (Tunghai Univ.).

A thesis submitted in partial fulfilment of the requirements for the degree of Master of Science, Memorial University of Newfoundland.

1971.

ABSTRACT

The ciliate fauna of a small, freshwater pond at Logy Bay, Newfoundland was' studied in 1968-70. A preliminary survey of the seasonal occurrence of the protozoa as affected by ecological paramèters was made.

The study 'pond is quite acidic (pH 4.50-5.15), with high chloride content (99.26 p.p.m.), a wide annual water temperature range (0.5-20.8°C), and a fluctuation of dissolved solid concentration (75-144 p.p.m.).

Twenty-nine ciliate species of 19 genera were determined. Other ciliates belonging to a least 30 taxa were not identified beyond the generic level.

Three groups among the studied ciliates are particularly interesting and demand rediscovery for further study: Tetrahymena uorax for its polymorphic life cycle; Coleps heteracanthus, Aspidisca sp., Condylostoma sp., Dysteria sp., and Trachelocerca sp. which might be new ecotypes of marine ciliates; and Coleps sp., Urotricha sp., Microthorax sp., and Lembadion sp., perhaps new species. TABLE OF CONTENTS .

ACKNOWLEDGEMENTS

ABSTRACT .

INTRODUCTION

HISTORICAL SUMMARY MATERIALS, AND METHODS RESULTS AND DISCUSSION. EXPERIMENTAL

> Physico-chemical factors Cultivation.

Clone cultures. Preparation of living material. Preparation of fixed material. SYSTEMATICS

Coleps octospinus. Coleps heterocanthus. Coleps sp. Urotricha sp. Colpoda cucullus. Microthorax sp. Tetrahymena vorax. Tetrahymena pyriformis. Paramecium bursaria. Cyrtolophosis bursaria. Cyrtolophosis elongata. 3. 5. 11. 11. 11. 18. 19. 20. 20.

23. 32. 36. 38. 43. 46.

49. 55. 65. 71. 78.

ata. 80°

TABLE OF CONTENTS (Cont.).

. · · .	Lembadion sp.	84
	Cristigera sp.	88.
1 4, 1	Cyclidium glaucoma.	90:
•	Vorticella spp.	95.
	Epistylis sp.	/ 101.
	Telotrochidium sp.	105.
	Blepharisma persicinum	108.
	Spirostonum teres.	110.
•	Halteria sp.	113.
•. •	Strombilidium sp.	115.
× • •	Aspidisca sp.	118.
T ,	Additional species examined but not illustrated.	120.
CONCLUSI	CONS.	130.
REFERENC	CES.	132
ADDITION	AL REFERENCES	149.
ADDENDUM		

152.

Carnivorous Tetrahymena vorax and its prey

۰.	PLATES.
I	Location of study pond.
11	Details of study pond. '
III	Temperature of pond.
IN	Dissolved solids concentration of pond
V	Concentration of hydrogen ions.
VI	Coleps octospinus.
VII	Coleps heterocanthus.
VIII	Coleps sp.
IX	Urotricha sp.
x	Colpoda cucullus.
XI	Microthorax sp.
XII	Microthorax sp.
XIII	Microthorax sp.
XIV	Microthorax sp.
XV .	Tetrahymena vorax.
XVI	Tetrahymena vorax.
XVII	Tetrahymena vorax.
XVIII	Tetrahymena pyriformis.
XIX	Tetrahymena pyriformis.
XX	Paramecium bursaria.
XXI	Paramecium bursaria.
XXII	Paramecium bursaria.
XXIII	Cyrtolophosis bursaria.
XXIV	Cyrtolophosis elongata.

13'.

14

15.

33,

37. 39.

44.

47.

50.

51.

52 · :

53.

56.

. 57.

58.

66.

67.

72.

73.

74.

79.

81.

DT ATTO	(Comt
LAILD	loni

85.

89.

91.

96.

98

100

102.

103

10

. 109.

111

-114

116

119.

بير. •

	XXV	Lembadion sp.
	XXVI	Cristigera sp
	XXVII	Cyclidium glaucoma.
	XXVIII	Vorticella p.
	XXIX	Vorticella sp. 1.
	XXX	Vorticella sp. 2.
•	XXXI .	Vorticella nebulifera.
	XXXII (Epistylis sp.
	XXXIII	Epistylis sp.
	XXXIV	Telotrochidium sp.
•	XXXV	Blepharisma persicinum.
	XXXVI	Spirostomum teres.
,	XXXVII	Halteria sp.
	XXXVIII	Strobilidium sp.
	XXXIX	Aspidisca sp.

ACKNOWLEDGEMENTS

I wish to express my sincere thanks for the cooperation of many people who have, in various ways, assisted in this study. I ammost grateful to Memorial University of Newfoundland for the provision of facilities and for the Fellowship which enabled me to undertake this work. I express my appreciation to Drs. C. C. Davis, G. F. Bennett, and Professor J. Phipps, for their ready support and valuable suggestions at various points throughout my programme.

Sincere thanks are also due to Drs. O. A. Olsen, G. R. Brassard, J. K. Pickavance, and S. Frost, and to Mr. H. Heng, for their assistance in the field work that was an indispensable part of this study. I am particularly indebted to Mr. Roy Ficken for invaluable help with photo-

Drs. J. Berger and J. Lackey gave valuable guidance and were sources of inspiration at the outset. Dr. J. Corliss helped greatly, through his cooperation in sending specimens for comparative study. I must thank Dr. D. Steven of McGill University, too, for making library facilities available to me, also Mrs. P. Bennett who typed the thesis with patience and perseverance.

Finally, may I reserve a special acknowledgement for my Supervisor, Dr. M. Laird, Head of the Biology Department, who provided constant advice, guidance and encouragement throughout the project.

To all these and others, I express my sincere appreciation.

Cillate protozoa were first examined from the vicinity of Newfoundland by Ehrenberg in 1854. These were marine species, reported by Kent, 1880-1882, pp. 626-627. Little further work was carried out in this field until Lom and Laird (1969) described some parasitic trichodinids from Newfoundland marine fish. Soon afterwards, Lackey and Lackey (1970) published an extensive list of free-living marine ukicroorganisms, including some cillates, from Logy Bay, near St. John's. However, nobody has previously studied the freshwater cillate fauna of the island of Newfoundland in any depths

Knowledge of the morphology, taxonomy and distribution of Newfoundland's flora and fauna (including the freshwater components) is clearly desirable, as taxonomic and distributional studies of most Newfoundland terrestrial blota are still far from complete. Recognizing the need to be able to make close comparisons with the results of earlier workers, permanent preparations stained by various methods (including "classical" haematoxylin procedures and modern silver impregnation) were employed in the present study to supplement - not to supplant - fresh preparations. The latter were examined by phasecontrast microscopy. In this context it is noted that, to judge from the surprising detail in the figures of such early protozoologists as Ehrenberg, phase-like effects were being obtained about'a century before F. Zernike invented this procedure (in 1935). Assisted by modern photomicrographic techniques, this thesis presents a preliminary survey of a primarily taxonomic study of the free-living ciliates of a

INTRODUCTION

small pond at Logy Bay, Avalon Peninsula,

60.

In addition, some data are presented to show that the appearance and disappearance of particular species of protozoa in a particular locality are usually correlated with seasonal changes (Davis, 1969; Wang, 1928). It was felt worthwhile in this context to make a preliminary survey of the seasonal distribution of protozoa as affected by fluctuations of temperature, hydrogen-ion concentration, and dissolved solids, in a small pond in Newfoundland. Comments on the value of various cultivation techniques are also included.

HISTORICAL SUMMARY

The first protozoan to be discovered was found in standing rain water in 1675. A ciliate of the genus Vorticella, it was described by van Leeuwenhoek (1677). Ledermüller (1763) was the first to introduce the term 'Infusoria' for all the various microscopic "animalicules" (whether bacteria, algae, or protozoa) developing in infusions exposed to air. The first comprehensive monograph on 'Infusoria' was by 0. F. Müller (1773). Bütschli (1880-89) limited the use of 'Infusoria' to Protozoa bearing cilia throughout life, and 'Suctoria' to those with cilia only in the early developmental stages. Doflein (1901) introduced the term 'Ciliophora' to designate a sub-phylum comprising the two classes Ciliata and Suctoria.

Since then, there are two main schemes for classification of protozoa; the pre-war and post-war systems.

The pre-war system was contributed by the pioneer scientists such as Ehrenberg (1832a, b, 1838), Dujardin (1838, 1841), Stein (1854), Claparede and Lachmann (1858-1861), Kent (1880-82), Büţschli (1880-89), Stokes (1888), Schewiakoff (1896).

Shortly before the second world war, Kahl (1930-35) prepared a profusely illustrated, well-documented monograph containing keys to the ciliates of Germany. Kahl's painstaking attention to the comparative morphology and systematics of ciliates at the species leve \mathbf{L}^{2} has not yet been surpassed.

The depth of our knowledge has always been benefitted by the improvement of Technology. It was one of Kahl's contemporaries, Klein, who first applied silver nitrate in demonstrating the silver-line system

of ciliates (Klein, 1926, 1958). In 1936, Chatton-Lwoff's method was introduced into France with the now recognized advantages over Klein's. Under this heading, Bodian (1937) also started the so-called 'Protargo1' technique (Honigberg and Davenport, 1954; Kozloff, 1961; Dragesco, 1962; Tuffrau, 1967). In 1940, Furgason proposed the newly established genus Tetrahymena by applying the silver method to establish systematic relationships on the basis of comparative stomatogenetic data. Studying the life cycle of astomes, de Puytorac (1954, 1959, 1963) revealed basic infraciliary similarities to other groups, e.g., certain members of the order Thigomtrichs. Using morpho-genetic data, Faure-Fremiet (1950) proposed that the Suctorians are very closely allted to the holotrichs and that the pertrichs should be considered closer to the holotrichs than to Spirotrichia. The work of Chatton-Lwoff (1935, 1949, 1950) and Faure-Fremiet (1950, 1953, 1959) validated a phylogenetic hypothesis formalized by Corliss (1956). Based on that, a revised system of classification at higher taxonomic levels within the phylum Protozoa was accepted over the older scheme (Honigberg et al., 1964). This revised scheme has been widely accepted. With the use of the morphogenetic pattern to deduce supposed . affininies among the various key genera involved, Small (1967) established a new order 'Scuticociliatida' which includes all pleuromatine hymenostomes. arhynchodine thigmotrichs, and a number of formerly alleged tetrahymenine hymenostomes. This research stimulates advanced understanding of the systematics and evolution of the holotrichs (Lom, 1968; Corliss, 1968; Nobili, 1969). In Russia, Jankowski (1964a, b, 1967) has constructed a strikingly new taxonomic hierarchy of Ciliophora.

MATERIALS AND METHODS

Habitat

Five hundred metres west of the Marine Sciences Research Laboratory (MSRL) (Fig. 1-A) of Memorial University of Newfoundland at Logy Bay, 8 km. north of St. John's, "There is a flat 400 by 800 m. *Chamaedaphne* and *Sphagnum* marsh in which lie various small permanent and temporary pools and which is crossed by a narrow but relatively deep stream (50 to 100 cm. by 50 to 100 cm.)" (Pickavance <u>et al</u>, 1970). The particular pond of this complex from which all my material was derived is about 50 m. west from the MSRL parking lot, and 10 m. south from the approach foad and the roadside stream (Fig. 1-B) which receives an outflow from it after rain (Station 2, Pickavance, pers. comm.).

The shape and size of the pond are quite variable. Its outline changes in accordance with the volume of its contents, the amount of water held increasing with rain and decreasing during warmer and drier weather. Generally, it is long and rather narrow, but shallow. Its width ranges from 1 m. (Fig. 2C) to nearly 4 m. (Fig. 2A, and 3), and it is about 17 m. long. The shallowest part (Fig. 2C) is 18 cm. deep, the deepest 60 cm. (Fig. 2b). There is an emergent rock at the northern end.

Predominant flora (Fig. 4) around the pool consists of the mosses Sphagnum pulchrum (Braithw.) Warnst and Sphagnum palustre L., Sweet Gale^{*} (Myrica gale L.), a cotton grass (Eriophorum virginicum L.), and Juniperus virginiana L.

* "Gold-withy" is the vernacular name in Newfoundland (Rouleau, 1956).

PLATE I

Figs. 1 and 2, showing location of the pond.

Fig. 1. The pond is located some 500 m. west from the Marine Research Laboratory (A) and 10 m. from the approach road. A roadside stream (B) received outflow from the pond when it rains.

A BAR BAR

Fig. 2. The width of the pond ranges from 1 m. (C) to near 4 m. (A). The shallowest part (C) is 18 cm. deep, the deepest 60 cm. (B).



PLATE II

Fig. 3. Area "A" of Fig. 2, the widest part of the pond, enlarged to show the emergent rock.

Fig. Detail, showing the collecting ladle at the edge of

the pond.



Sampling:

This project was carried out from 1968 to 1970, the most intensive examinations being made during the period from June 1969 to September 1970, when all physico-chemical readings were taken. During the rest of the time, samples were taken at irregular intervals.

Random samples were obtained by using one-litre capped jars as containers. A white-enamelled ladhe was employed in dipping for benthic fauna. Samples were brought back to the laboratory as soon as the physico-chemical conditions had been recorded.

Measurement of physico-chemical factors:

Physico-chemical readings were regularly made in the field from June 1969 to September 1970. Both air and water temperature recordings were made with a mercury thermometer. Concentration of the dissolved solids (D.S.) was measured with a Myron L Dissolved Solids Meter (Model 532 T). The hydrogen ion value was determined using a Porto-Matic pH meter (Model 175). Chloride concentrations were estimated in the laboratory by the Mohr Method (A.P.H.A., 1955).

Preparation of media:

Organisms were concentrated by centrifugation (2500 rpm/ 5 min.). They were then cultured in sterile disposable Petri dishes with 35 ml. of pond water or hay infusion.

The latter was prepared by Turner's method (p. 60, Needham, 1959) modified as follows: to one litre of pond water add 5 gm. of Timothy Hay, *Phleum pratense*, and 10 grains of wheat. Heat to boiling and set aside until cooled. Filter through three layers of gauze into

200 ml. bottles. Each bottle should receive 50 ml. of the hay in-

Four media were used to obtain mixed cultures. (1) Culture with 35 ml. of hay infusion.

(2) Culture with 35 ml. of pond water added, with a surface sprinkling

of Bacto-tryptone (Gruchy, 1955).

(3) Culture with 35 ml. of pond water and four grains of rice.
(4) Culture with 35 ml. of pond water and four grains of wheat.
Cultures were maintained in a brightly lit laboratory at: 70° F.

Clone cultures:

When ciliates were-present, clonal cultures were attempted to serve axenic development (Elliott, 1953 and Gruchy, 1955).

Preparations of living material:

Fresh preparations were dealt with as recommended by Kudo (1966, p. 1073). A 10% solution of methyl cellulose was used to retard the swimming movements of ciliates (Marsland, 1943). For vital staining, the following dyes were dissolved in absolute alcohol (Kudo, 1966): Congo red (1:1,000)--, when, used as an indicator its red colour indicates

alkalinity, a change to blue, weak acidity; ...

Janus green B (1:10,000) - employed for the staining of mitochondria; Methylene blue (1:10,000) - for staining cytoplasmic granules, the

nucleus and cytoplasmic processes;

Neutral red (1:3,000) - when used as an indicator, it becomes yellowish , red (alkaline), cherry red (weak acid); or blue (strong acid);

Lugol solution (made up of potassium iodide 1.5 gm. water 25 cc. and

iodine 1 gm.) - for staining cilia.

Microphotography was undertaken by means of a Carl-Zeiss Photomicroscope I, utilizing both phase-contrast and interference (Nomarski effect) microscopy.

Preparation of fixed material:

Heidenhain's iron haematoxylin was used for nuclear staining (Kudo, 1966).

Silver - impregnation methods.were employed to demonstrate the cortical configuration.

The dry silver method undertaken was that of Klein (1926, 1958).

The Chatton-Lwoff method (Chatton and Lwoff, 1930 and 1936; . Corliss; 1953b;Frankel 1968) was employed when there were

sufficient numbers of ciliates in the culture.

RESULTS 'AND DISCUSSION

EXPERIMENTAL

The physico-chemical factors:

Monthly average records of the pH value, D.S. concentrations and water and air temperatures are given in Table I and Figs. 5, 6, and 7.

In July 1970, the pond dried up at a time of unusual drought and hot weather for Newfoundland (above 80° F). It was replenished on August 10, 1970 after two days' heavy rain. It did not dry up again during the remaining months of the present study. It thus qualifies for the designation of "temporary pond".

Spandl (1926) restricted "temporary pond" to water bodies persisting for no more than 1¹/₂ - 2 months. Others (e.g. Shelford, 1913; Kenk, 1949; Rzoska, 1961; Felton et al, 1967; and Moore, 1970) have so designated any pond drying up completely at least once a year. Laird (1956) sharply differentiated "transient ponds" (primarily rainfilled ones in hollows which do not intersect the water table) from, permanent ones, But followed Bates (1949) in considering "permanent" and "semi-permanent" (= "temporary") ponds as a continuous cline. It is considered that for all practical purposes my study pond can be characterized as a "semi-permanent" pond.

From the end of December 1969 to the end of February 1970, the pond was usually frozen. Records were taken after chopping a hole in the surface ice with an axe. Prior to this time, the strong winds of winter ensured mixing of the epilimnion with hypolimnion .

••	· • 1	1	· ·	
Nonth	Temp.	(°C.)	$D_{2}S_{2}(p_{2},p_{3},m)$	pH.
	Air	Water		
June 1969	14	12.7	-	
July	23.5	20.5	125	-
Aug.	21.2	19.8	112	4.54
Sept.	20	17.9	, 93	4.68
Oct.	18.5	13.3 '	93	4.62
Nov.	7	7.2	144	4.68
Dec.	11.	6.8	^{,75[°]}	4.70
Jan. 1970	Ò .	0.5	106	4.65
Feb.	- 3	0,5	123	4.50
Mar.	3	7.0	102	4.83
Apr.	.6	7':2	85	5.08
May	7.5	, 7.5	85	5.15
June	12.5	10.7	79	5.13
July	30	. –	-	- ^ .
Aug.	22.8	20.8	'100	4.83
Sept.	20.	11	115	4.73

TABLE I

PLATE III

13

Fig. 5. .Temperatures of the pond, June, 1969 to September, 1970.

...



PLATE IV

Fig. 6. Dissolved solids concentration of the pond, July 1969 to Septembér 1970. A and B indicate two peaks of high concentrations of dissolved solids.



PLATE V

Fig. 7. Concentration of hydrogen ions of the

pond, August 1969 to September 1970.



Following the freezing of the pond surface, however, this mixing was inhibited. The epilimnion was depleted of dissolved substances by organisms which, on their death, sank to hypolimnion and decomposed (Macan, 1970). This explains peak B in Fig. 6.

The other peak in Fig. 6,A, reflects a long hot and dry period without much rain from September to November 1969. Usually, the daily readings of the concentration of dissolved solids vary with the amount of rain received. Data missed from the end of June 1970 to the beginning of August 1970 would probably have reached the highest dissolved solids concentration because of the hot weather and unusually long dry period.

Despite the rapid variation of the temperatures and dissolved solid concentrations, the hydrogen-ion concentration of the pond proved very stable, ranging from pH 4.5 in February 1970 to pH 5.15 in May, 1970. It was generally higher in the summer of 1970 and lower in the winter of 1969,

Juday, Frey, and Wilson (1924) followed variations in the hydrogen-ion concentration with the season and depth in Lake Mendota for the period 1919 - 1922. "They observed that in summer particularly, and to a somewhat lesser extent under the ice in winter, the upper waters had a substantially higher pH than the lower waters. They aecribed the summer differences to the photo-synthetic activity of algae towards the surface and to the decomposition of organic matter in the hypolimnion" (Frey, 1966, p. 33). This may serve to explain the slight increase of pH during the summer of 1970, and the slight decrease in the winter of 1969-70, the latter reflecting the decomposition of organic matter in the hypolimnion associated with increases in free $^{CO}_2$ and bound $^{CO}_2$. However, "at pH values below 6.0 acids other than carbonic may be suspected" (Hutchinson, 1957, p. 683). Data from the present pond are closer to those derived by Frey (1949) from some bay lakes of North Carolina (pH 4.34 - 4.92), these figures not being raised by aeration or boiling. Strøm, (1939, 1942, 1944) reported pH values of 8.0 from lakes in drainage basins influenced by Palaeozoic limestones or into which glacial water flowed.

The present data bear comparison with: pH 3.2 - 4.9 reported by Yoshimura (1934) from several Japanese lakes containing strong mineral acid; pH 3.5 - 3.9 recorded by Wehrle (1927) in Sphagnum bog pools near Freiburg in Breisgau, Germany; pH 4.5 - 6.0 from a Sphagnum bog in Iowa (Grant and Thorne, 1955); pH 5.2 - 6.1 in Louisiana ponds (Moore, 1951 and 1955) and one of these temporary ponds was reported pH 4.9 - 5.5 by Margavio (1964). "The source of the acid in such waters is probably twofold. Rain water, which contains sulfate, percolating through peat, tends to lose cations and gain H⁺ by base exchange with the humus of the peat" (Hutchinson, 1957, p. 682).

The dissolved chloride content of the study pond was found to be 99.26 mg/l on January 11, 1970. This figure is considerably higher than the range of 0.08 - 0.51 mg/l of the English Lake District (Mackereth and Heron, 1954).

Prischel (1940) and Conway (1942) state a formula to calculate the chloride concentrations of ponds. These decrease proportionally to the increasing distance from the sea. They show chloride concentrations ranging down from 70-700 mg/1 in ponds 600 m. or less

from the ocean. My study pond is located less than 600 m. from the Atlantic Ocean, its chloride concentration (recorded as 99.26 mg/l on January 11, 1970) falling within the above range.

There is a seasonal fluctuation of chloride content in some localities (Russell and Richards, 1919; Miller, 1914), high winter concentrations doubtless being due to the greater disturbance of the adjacent ocean surface during the winter. Drischel (1940) in fact suggests that a marked seasonal cycle is characteristic of maritime regions. The chloride concentration of the study pond was obtained from the epilimnion, which was covered by ice in January 1970. Regretably, seasonal investigations of the chloride concentrations of the pond were not undertaken.

Aerial chloride transportation can also occur through the wind picking up dry salt from an expanse of arid coastal land from which sea water had evaporated (Holland and Christie, 1909). Typically salt-water ciliates (*Coleps heteracanthus* Noland, and *Trachelocerca* sp.) were discovered in the sample of January 11, 1970. It is possible that wind transportation accounted for the presence of these salt' water.species.

Cultivation:

-

Gruchy (1955) used a surface sprinkling of Bacto-tryptone for mass culture from which *Tetrahymena pyriformis* clonal cultures were axenically established. In companison with other procedures already mentioned, this method was used from the outset as the standard of comparison. Cultures thus fed always formed a thick opaque film on

the top of the medium and smelt very strongly of ammonia. Even after the odour gradually faded, the film was still so opaque that organisms were not observable.

Hay infusion has been very commonly used to grow ciliates. (Turner, Beers, Leray, Jones, in Needham, 1989). However, because the ' sterile culture used lacked prey such as *Chilomonas* sp. for the ciliates to feed upon (Needham, 1959, p. 60), it was generally unsuccessful.

By contrast, media with either four grains of rice or wheat kernels developed flourishing cultures. In those cultures supplied with rice grains, whitish fungi (Saprolegniaceae, as identified by Dr. 0. Olsen) always flourished around the grains. These cultures were always richer in euglenoid phytoflagellates than were those to which wheat kernels were added - a characteristic noted by Brandwein, p. 63 (Needham, 1959). Wheat kernel media always supported more vigorous populations of ciliates than did rice grain media.

Clone cultures:

Ł

Large numbers of a single species were necessary for good preparations destined for permanent staining by the Chatton-Lwoff silver impregantion technique. Stock medium (Elliott and Hayes, 1953; Gruchy, 1955) in a depression slide containing a mixture of penicillin G and streptomycin (250 µg/ml each) was used to obtain axenic cultures: However, it proved that immediately after being transferred into the stock medium, ciliates ruptured. It seems, therefore, that other species of ciliates are not able to tolerate the stock medium designed especially for *Tetrahymena pyriformis* Ehrenberg.

Preparations of living material:

A dissecting microscope was only useful as regards the larger ciliates, such as *Paramecium bursaria* Ehrenberg (140 μ), *Spirostomum teres* Claparède and Lachmann, (200 - 500 μ), and *Stylonychia* sp. (150 μ). Smaller ones, e.g. *Cyclidium glaucoma* 0.F.M. (17.5 - 25 μ), *Urotricha* sp. (27 μ); *Cyrtolophasis elongata* Schewiakoff.(30 μ) and *Halteria* sp. (25 - 50 μ) proved unidentifiable at the low magnifications attainable with this instrument.

Phase microscopy proved indispensable for obtaining comparative data and improving existing taxonomic accounts prepared prior to the availability of this technique. With it, cilia, flagella, fine pseudopodia, stalks and fibrils all showed up to maximum advantage.

As revealed by both the dissecting instrument and high power phase-contrast microscopy, the plankton of the study pond most commonly comprised sarcodinids including Arcella, green algae,

diatoms, blue-green algae, flagellates, rotifers, gastrotrichs and cladocerans in addition to the ciliates reported in the present study. The occurrence of ciliates often coincided with the appearance of *Arcella* and flagellates (e.g. *Chilomonas paramaecium* Ehrenberg) in considerable abundance.

Preparation of fixed material:

Heidenhain's iron haematoxylin was used for nuclear staining when methylene blue had failed to reveal the nucleus. The results were satisfactory both for holotrichs and spirotrichs.

Silver impregnation was indispensible for reveating the
cortical configuration, such structures as basal granules or kinetosomes of the somatic and buccal cilia, the cytoproct, contractile vacuole pores, and the complex fibril network (Chatton and Lwoft, 1930, 1936; Corliss, 1953b, 1956, 1959a, 1961a, 1963; Klein, 1926, 1958; Lom and Laird, 1969; Thompson, 1960, 1963a, and Boggs, 1965). The several modern techniques under this heading furnished a standard means of demonstrating the bases of the entire somatic and buccal ciliature in whole mounts of organisms at various life-history stages.

Klein's dry method and the Chatton-Lwoff procedure for silver impregnation were used in the present study. Although both were invented more than forty years ago (Klein, 1926; Chatton and Lwoff, 1930), the application and preference of these methods are still not universal (Corliss, 1956; Klein, 1958; Raabe, 1959; Lom, 1961; Thompson 1960 and 1963a; and Lom and Laird, 1969).

The Chatton-Lwoff method provided outstanding advantages over Klein's dry method (Corliss, 1956), if sufficient numbers of a single species (e.g. *Tetrahymena pyriformis* Ehrenberg) were available." However, the technique was handicapped when low numbers or a mixture of species were involved. It was thus unsatisfactory for survey studies, although Frankel's (1968) method utilizing centrifugation to wash out Champy's fixative somewhat decreased the chance of losing organisms. Klein's method reveals the argentophilic infraciliature with less such risk. -It is also easier to monitor the mounted organisms by this method, which therefore has much to offer in comparative taxonomy.

With regard to the latter, data concerning the argentophilic infraciliature of ciliates are still very far from complete, being in

fact confined to only a few genera; notably Condylostoma (see Suhama, 1961), Balantidium (see Krascheninnikow and Wenrich, 1958), Cyclidium (Berger and Thompson, 1960), Neobursaridium (see Nilsson, 1962), Paramecium (see Lieberman; 1929), Sathrophilus (see Thompson, 1963a), Spirostomum (see Boggs, 1965), and Tetrahymena (see Corliss, 1952a, 1952b, 1953a, 1957, 1959b, 1960). Tetrahymena is the only genus for which we can now claim virtually complete comparative argentophilic infraciliature data. Application of silver impregnation for purposes of taxonomic identification is handicapped both by the tremendous lack of specific infraciliature "fingerprints" (Thompson, 1960) and the difficulty of ensuring the development of rich, pure cultures. The need of rediscovery and a full redescription of ciliated protozoa in light of modern technique (silver impregnation) for taxanomic data was discussed by Thompson (1963).

It is important to realize, though, that however, sophisticated one's staining procedures may be, fresh preparations still remain indispensable aids to taxonomic identification. As is evident from the photomicrographs reproduced herein, phase microscopy reveals living structural detail scarcely detectable if not altogether invisible by bright-field microscopy, and forms a necessary part of any survey such as this.

SYSTEMATICS

Representatives of 45 ciliate generatives observed from September 1968 to September 1970. Eighteen genera are illustrated by photomicrographs in the following sections. Ecological parameters and data on occurrence for each of the genera and species studied herein are presented in Table II.

23

tina)

Phylum Protozoa

Subphylum

L

Class Ciliata

Subclass I Holotricha '

Ciliophora

Order Gymnostomatidae

Family Colepida

Genus Coleps* . Family Enchelyidae

Genus

Chilophyra Prorodon

. . .

Suborder , Rhabdophorina (=Promstomatina + Pleurostoma-

Microregma

. Platyoph**r**ya

Trachelocera

Urotricha*

Amphileptidae Family Genus *Litonotus* Suborder Cyrtophorina (= Hypostomatina) Dysteriidae Family Dysteria Genus Paratrochilia Chlamydodontidae Family · Chilodonella ~ Genus Family Nassulidae Genus Nassula Order Trichostomatida Family, Colpodidae Colpoda* Genus Microthoracidae Family Genus Microthorax* Hymenostomatida Order Suborder Tetrahymenina Cohnilembidae Family Cohnilembus Genus Urònema Gentus Tetrahymenidae Family Colpidium Genus Tetrahymena*

24

-	, ··	
Subo	der 🕔	Peniculina
Fa	amily	Paramèciidae
, , ,	Genus	Paramecium*
· · Fa	amily,	Frontoniidae
	Genus	Balanonema
	. • • •	Cinetochilum
	•	Cytrolophosis
	: 	Frontonia
	-	Lembadion*
Subo	rder P	leuronematina
· · · · · · · · · · · · · · · · · · ·	amily	Pleuronematidae
· · ·	Genus	Cristigera*
· .	்கு	Cyclidium*.
· · · ·		Pleuronema 🕐
Subclass	Peritr	icha .
Order	Peritr	ichida ,
' Subo	rder .	Seesilina
F	amily	Ophrydiidae

FamilyOphrydiidaeGenusOphrydiumFamilyVorticellidaeGenusVorticella*FamilyEpistylididaeGenusEpistylis*

Tel

Telotrochidium*

·L

Subclass III Spirotricha Order lleterotrichída Suborder Heterotrichina Family Gyrocorythidae Genus^e Metopus Family Spirostomatidae Blcpharisma* Genus Spirestomun* Family Condylostomatidae Condylostoma ' Genus Family Clevelandellidae Genus Paraclevežandia Oligotrichida Order Halteriidae 🤟 Family Genus *Halteria** Strombilidium Hypotrichida Order Aspidiscidae Family Aspidisca* Genus Oxytrichidae Family Gonostomum Genus Steinia Oxytricha Stylonychia Uroleptus Urosoma

ECOLOGICAL	PARAMETERS 'OF	STUDY POND.
		•

b .				· .	
Genus	species	- Date	Temp.(°C.)	D.S.ppm.	pH.
Coleps	octospinus Noland, 1925	Aug. to Oct. '69. March;May,June'70	3 19.5	79 - 113	4.55 - 5.15
<i>n</i>	heteracanthus Noland, 1937	January 14, 1970	0.5	107	4.75 -
11 ₁₁	Sp.	March & June 1970	6.5 & 10.5	78 & 102	4.80 & 5.15
Chilophrya	utahensis Pack, 1919	June to Sept.'69 March & April,'70	6.5 - 20:5	85 - 125	4.5-5.0
Prorodon	discolor (Ehrenberg, 1838)	Aug. & Sept. '69	17.5 - 19.5	93 - 110	, 4. 5 - 4.75
Microregma	sp.	September '69	17.,5	93	4.75
Platyophrya	sp.	September '69	17.5 ; ·	93	4.75 .
Trachelocerca 🗧	sp. '	Jan. & March, 70	0.5 & 7-	102 & 107	4.75 & 4.8 <u>0</u>
Urotricha	sp.	All seasons	0.5 - 20.5	. 78 - 144	4.50 - 5.15
Litonotus	sp. ,	November '69	7.2	144	4.68
Dysteria	sp.	July '69	12.5	126	
۰ ·		1	1.	· ·	•

TABLE II



TABLE , II (Cont.)

Genus	species	Date	Temp.(°C.)	D:S.ppm.	pH.
Paratrochilia	chilodontoides Kahl, 1933	Sept. 6, '69	17.9	93	4.68
Chilodonella	uncinata (Ehrenberg, 1838)	Aug. '69	19.5	110 -	4.50
Nassula ⁴	sp.	Aug. '69	19.5	110	4.50
Colpoda	cucullus O.F.M., 1773	Dec.'69 to Apr:70	0.5 - 8.0	78 - 123	4.75 - 5.00
n	steini Maupas, 1883	Mar.Sept.&Nov.'69	7.5 & 17.5	. 93 & 144	4.75
Microthorax	sp.	July to Oct. '69 March & April '70	7.5 - 20.5	• 85 - 126	4.75 - 5.00
Cohnilembus	sp.	Aug. & Sept. '69	17.5 & 19.5	93 & 110	4.50 & 4.75
Uronéma	sp. ⁹	Sept. '69	17.5	93	4.75
Colpidium	sp.	July:& Aug. '69	12.5 & 19.5	110 & 126	4.50
Tetrahymena ·	vorax (Kidder,Lilly & Claff, 1940)	July 17, 1969	23.0	126	_
11	pyriformis (Ehrenberg, 1830)	All seasons	0.5 - 20.5	78 - 144	4.50 - 5.15
Parameciúm s	bursaria (Ehrenberg, 1833)	All seasons '	0.5 - 20.5	78 - 144	4.50 - 5.15

<			·		•
\ Genus	species	Date	Temp.(°C.)	D.S.ppm.	pH.
Balanonema	biceps (Penard, 1922)	September, 1969	17.5	93	4.75
Cinetochilum	marinum Kahl, 1930	September, 1969	17.5	93	4.75
Cyrtolophosis	bursaria (Schewiakoff, 1893)	April, 1970	7.0	8-5	5.00
<i>II</i>	elongata (Schewiakoff; 1896)	Aug. & Sept. 1969 March 1970	6.5 17.5 19.5	93 - 110	4.55 4.7 4.80
Frontonia	leucas (Ehrenberg, 1838)	September, 1969	17.5	93	4.75
Lembadion	sp.	April to June '70	7.0 - 11.5	72 - 87	5.00 - 5.1
Christigera	sp.	All seasons	0.5 - 20.5	78 - 144	4.50 - 5.1
Cyclidium	glaucoma O.F.M. 1773	Sept.'69-June,'70	0.5 - 18	., 78 - 144	4.75 - 5.1
II 6	citrullus Cohn, 1865	Aug. & Sept. 1969	17.5 & 19.5	93 & 110	4.50 & 4.7
n	elongatum Schewiakoff, 1896	Nov. & Dec. 1969	6.5 & 7.0	78 & 144	4.75
	granulosum Kahl, 1931	September, 1969	17.5	93	4.75
- 17	litomesum Stokes, 1884	August, 1969	19.5	110	4.50
IR .	muscicola Kahl 1931	January, 1970	0.5	106	4.75

(Con T_{4} TT

_ • • •		· (1
Genus	species •	Date	Temp.(°C.) D.S.ppm.	pH.
Pleuronema	marinum Dujardin, 1841	September, 1969	17.5 93	4.75
Ophrydium	sp.	September, 1969	17.5 93	4.75
Vorticella	sp. (Figs. 84 - 85)	April & May, 1969		-
"	microstoma Ehrenberg, 1830	May to August, '69	12.5 - 20.5 113 - 125	4.50
17	sp. (Figs. 87 - 89)	Nov.'69toJan.'70	0.5 - 7.0 78 - 144	4.75 - 4.80
11	nebulifera O.F.M. 1786	Aug. to Oct. 1969	17 - 19.5 93 - 113	4.50 - 4.80
S. W.	sp. (Figs. 90 - 92)	September 1969	17.5 93	4.75
Epistylis	sp. A	January 14, 1970	0.5 106	4.75
Telotrochidium 🎘	sp.	Dec.'69to Jan.'70 May, 1970	0.5 - 6.5 78 - 107	4.75 - 4.80
Metopus	striatus McMurrich, 1884	September 1969	17.5 93	4.75
Blepharisma	persicinum Perty, 1852	Sept. '69; June, '70	12.0 & 17) 5 78 & 93	4.75 & 5.10
Spirostomum	teres Claparède & Lachmann, 1859	All seasons	0.5 - 20.5 78 - 144	4.50 - 5.15
· ·				* 4.

TABLE II (Cont.)

• •

				· -
•	TABLE	TT	(Cont.)	
		~ ~	(001101)	

* * * * * * * * * * * * * * * * *				`·
Genus species	Date	Temp.(°C.)	D.S.ppm.	pH.
Condylostoma sp.	June, 1970	12.0	78	5.15
Paraclevelandia sp. ,	January, 1970	0.5	106	4.75
Halteria sp.	All seasons	0.5 - 20.5	78 - 144	4.50 - 5.15
Strobilidium sp.	August, 1969. Jan. to April,'70	19.5 0.5 - 7.5	85 - 125	4.50 - 5.00
Aspidisca sp.	All seasons	0.5 - 20.5	78 - 144	4.50 - 5.15
Gonostomum sp.	August, 1969	19.5	110	4.50
Steinia sp.	Nov. & Dec. 1969	6.5 & 7.0	. 78 & 144	4.75
Oxytricha sp.	Aug. & Dec. 1969	19.8 & 6.8	112 & 75	4.54 & 4.70
Stylonychia sp.	Aug.'69 to Apr!70	0.5 - 19.5	78 - 144	4.50 - 5.00
Uroleptus longicaudatus Stokes, 1886	August, 1969 May & June, 1970	. 8 12.5 & 19.5	78 86 & 110	4.50 5.00 & 5.15
Urosoma sp.	November, 1969	7.5	144	4.75

Subclass Holotricha Order Gymnostomatida Suborder Rhabdophorina

Family Colepidae

Genus

Coleps (according to Corliss, 1961)

Coleps is distinguishable from related genera of holotrichous ciliates by its characteristic armoured exoskeleton, and by the presence of several spine-like processes at the posterior extremity. Its closest relative (genus *Tiarina*) is also armoured, but lacks any spines at the posterior extremity, which tapers to a point. Therefore, the armour and the spines of the present species dictate allocation to the genus *Coleps*.

Coleps octospinus Noland, 1925. Plate VI, Figs. 8 - 10.

<u>Body</u>: 90 to 115µ in length, 40 - 50µ in diameter. Furrowed longitudinally and transversely, barrel-shaped, armoured with cortical plates. Anterior half slightly flattened dorso-ventrally; nearly circular in medial cross-section. Cytostome apical (Fig. 8). External skeleton transversely bisected by a central groove (Fig. 9,D). Posteriorly, the central section is separated from the terminal one by a groove (Fig. 9,E). No such groove anterforly. Twenty-two to 24 longitudinal rows of plates, in which lattice-like openings allow proto⁴⁴ plasm to extrude (see type 1 of Kahl, 1930, p. 132).

Anterior spines: 4 flat, double-pointed spines on the margins of the cytostome just behind the spinous crown. Their size

PLATE VI

Figs. 8 - 10 Coleps octospinus Noland, 1925

Fig.

8. Showing living example as seen by phase-contrast. (A) thornlike anterior spine; (B) subterminal posterior spine; (C) terminal posterior spine.

Fig. 9. Showing fresh preparation by Nomarski interference-contrast effect. (D) central groove; (E) posterior groove.

Fig. 10. Showing fresh preparation by phase contrast. The arrow "F" indicates the blade-like anterior spine.



variable, the largest ones along the right margin. Both pairs on the lateral margins blade-like (Fig. 10,F) and transparent. Mid-dorsal and mid-ventral sets of spines thorn-like (Fig. 8). A number of smaller, asymmetrically placed spines behind the spinous crown encircling the cytostome.

<u>Posterior spines</u>: four spines symmetrically located at the margin of the posterior extremity (Fig. 8,C), four sub-terminal ones being alternately placed in the penultimate transverse row of plates (Fig. 8,B). Of more uniform size than the anterior spines, and averaging some 9 μ in length.

<u>Ciliation</u>: uniform somatic ciliation throughout, the cilia slender and sparse and gradually decreasing in length anteriorly. Cilia shortest and more numerous around the cytostome (Fig. 9). One caudal cilium.

Movement: swims rapidly, revolving about the long axis in a counter-clockwise direction (viewed from the rear). Occasionally briefly immobile.

Nucleus; Nucleolus at the centre of the round and posteriorly located macronucleus.

<u>Division</u>: dividing examples were observed several times in March, May and June, 1970. The anterior opsterior halves began to separate along the central groove. The posterior half of the anterior daughter and the anterior half of the posterior one remained lightlycoloured - and indeed transparent - during the formation of their new plates and spines.

<u>Contractile vacuole</u>: located at the posterior end of the anterior division product, its contents discharging through the posterior armour by means of a cytopyge. Protoplasm vacuolated and exhibiting ingested algal material.

Food: mainly algae including diatoms, and smaller ciliates.

35

Systematic account: Kahl (1930 and 1932) grouped I4 species of Coleps into three categories on the basis of the structure of their exoskeletal plates. The organism under discussion has plates of Kahl's type 1, which occur in C. hirtus, C. elongatus, C. amphacanthus, C. bicuspis, C. uncinatus and C. octospinus. Noland (1937) described two more species of the genus Coleps. However, both of these differ from the present species in having type 3 plates.

Among the six members of Kahl's type 1, only C. octospinus has more than four posterior spines. There are two posterior spines in C. bicuspis; three in C. hirtus, C. elongatus and C. amphacanthus; and four in C. uncinatus. The Logy Bay species usually has four terminal and four sub-terminal posterior spines. It thus corresponds more closely to C. octospinus than to other species of the genus, despite Geiman's (1931) report that the posterior spines vary from seven to eleven in C. octospinus.

Finally, its large size $(90 - 115\mu \times 40 - 50\mu)$ double-pointed anterior spines and acid marsh habitat are further useful criteria linking it with *C. octospinus*; to which it is accordingly referred by

* °C

it's plate structure (Kahl's type 1), the number (eight) and arrangement (four terminal and four subterminal) of the posterior spines, its double pointed anterior spines and the overall size and shape of its

Coleps heteracanthus Noland, 1937. Plate VII, Fig. 11.

<u>Body</u>: 90μ in length, 40μ in diameter. Furrowed longitudinally and transversely. Barrel-shaped, armoured with cortical plates. Anterior half slightly flattened dorso-ventrally, nearly circular in medial cross-section. Cytostome apical. External skeleton transversely bisected by a central groove. Kahl's type 3 plates (Fig. 11,b).

One flat, double-pointed spine on the margin of the cytostome just behind the spinous crown. A set of thorn-like spines on the other margin of the cytostome. Posterior spines eight in number, arranged as in *Coleps octospinus*. Uniform somatic ciliation throughout, the cilia slender and sparse and gradually decreasing in length anteriorly. Cilia shortest and more numerous around the cytostome. Two'caudal cilia (Fig. 11). Swims rapidly, revolving about the long axis in a counter-clockwise direction as viewed from rear. Two spherical macronuckei, one in each half of the body. Macronucleus (Fig. 11) always with a central red spot in living non-stained material.

Systematic account:

body.

Among the 16 known species of Coleps, group three (e.g. C. incurvus, C. pulcher, C. remanei, C. similis, C. tesselatus, C. spiralis

PLATE VII

Fig. 11. Coleps heteracanthus Noland, 1937.

Showing general appearance and the structure of its plates.

37

(A) Barrel-shaped body with two caudal cilia. (B) Struc-

ture of plates, Kahl's type 3. Ma = Macronucleus.

and C. heteracanthus) bears comparison with the present species in having Kahl's type 3 structure of plates.

C. remanei is immediately separable from the present species bý its giant size (200 - 250 μ), and its numerous posterior spines. C. incurvus is easily distinguishable by virtue of its cylindrical body shape and six posterior spines. C. spiralis differs in exhibiting definite spiral torsion in its longitudinal rows of plates. The number of posterior spines as well as body size, narrow the choice down to C. pulcher (100 μ) and C. heteracanthus (74 - 86 μ), for C. tesselatus (60 - 70 μ) and C. similis 50 - 70 μ) having only three or four posterior spines.

The present species is referable to *C. heteracanthus* rather than *C. pulcher* mainly because its thorns are on only one side anteriorly. The data for the present species fit the description of *C. hateracanthus* (Noland, 1937), which has identical plate structure, thorns on only one side anteriorly, and is of comparable size. However, certain additional characters are evident in the present species. Thus there are a single flat and double-pointed spine in the anterior part of the body, on the side opposite that just mentioned, a spherical macronucleus with a red spot in each half of the unstained body, eight posterior spines, and two caudal cilia.

Coleps sp. (Plate VIII, Figs. 12 - 16). Armoured body (84 µ in length, based on four specimens, 80 -86 µ), with spine-like processes both anteriorly and posteriorly (Figs. 12 - 15). Barrel-shaped, elliptical in medial cross-section, longer

PLATE VIII

39

Figs. 12 - 16. Coleps sp.

Fig. 12. Showing body shape and dorso-ventral ciliation.

Fig. 13. Showing the three posterior spines, and the mid-dorsal and mid-ventral ones.

Fig. 14. Dorsal view, showing the left anterior spines.

Fig. 15. Dorsal view, showing the right anterior spines.

Fig. 16. Plate structure (x 1,200), referable to Kahl's type 1.





diameter, 52.5 μ (Figs. 14 and 15), the shorter (dorso-ventral) one, 43.4 μ (Figs. 12 and 13). Dorsal side more convex than ventral, tapering slightly towards each extremity.

Furrowed longitudinally and transversely with cortical plates of Kahl's type 1 (Fig. 16). Cilia projecting through the interspices between the plates. Ciliation uniformly sparse except for a denser vestiture encircling the cytostome (Fig. 12) and two caudal cilia. Length of somatic cilia, about 11 μ , the caudal ones, 35 μ .

Viewed from the dorsal side, a set of teeth (some 5.2 µ long) projects at each margin anteriorly. The straight set on the left is composed of one single pointed tooth and a wide-angled, double-pointed one (Fig. 14). The right set has a similar single straight tooth anteriorly, but the second and bifurcated one has distinctly closer to one another by comparison with its opposite member on the left (Fig. 15). A few thorn-like mid-dorsal and mid-ventral spines (Fig. 13).

Three incurved claw-like spines evenly spaced about the margin of the rounded posterior extremity. That to the left (the smallest) decidedly incurved, 5.2μ in length. Length of the other two spines, some 8.5 μ (Fig. 13).

Swims leisurely, revolving about the long axis, in a counter-

Systematic account:

This armoured, barrel-shaped ciliate with spines both anteriorly and posteriorly belongs to the genus *Coleps*. Two of the 16

40.

known species of *Coleps*, *C. striatus* and *C. trichotus*, do not merit consideration with it on the Basis of lacking posterior spines.

Plate structure is accepted as one of the chief specific criteria in the other 14 species. The plates of the present species are referable to Kahl's type 1. This narrows the comparison down to the six species bearing Kahl's type 1 plates (C. hirtus, C. elongatus, C. amphacantus, C. bicuspis, C. uncinatus and C. octospinus).

C. uncinatus and C. octospinus are immediately separable from the species under discussion in having more than three posterior spines.

C. hirtus $(40 - 65 \mu)$ with its three spines on the margin of the rounded posterior extremity, is closer to the species under discussion. However, it stands apart in that its armour comprises four girdles and in having single caudal cilium. Furthermore, the present species (84 μ_{s}° long) is very much larger than C. hirtus.

The latter stands closer to C. amphacanthus $(70 - 90 \mu \text{ long})$ in body size, in its plump shape, and in having three posterior spines. Nevertheless, it lacks the major features of C. amphacanthus, which has three posterior spines arranged in almost linear fashion instead of as a triangle, together with a twisted cytostome about sixty degrees clockwise out of line with the main axis of the posterior margin. Furthermore, C. amphacanthus has from four to eight caudal cilia instead of two, as in the present species.

The latter resembles C. bicuspis (55 μ) in shape and anterior spination, differing from it mainly by virtue of its larger size, three posterior spines, and two caudal cilia. As regards the spines of

Q. bicuspis "These pairs of teeth occupy portions at the anterior end that correspond with those occupied by the two spines at the posterior end " (Noland, 1925, p. 6). The two anterior sets of teeth do not "corrspond with the three posterior spines in the present species.

In its specific characters, such as the possession of two caudal cilia, three triangularly arranged posterior spines, and body size, the present species is very close to *C. elongatus* (40 - 55 μ according to Noland, 1925; 65 - 80 μ according to Kahl, 1930). However, the latter is characterized by its slender appearance. The plump shape of the present species (84 μ by 52.5 μ) thus immediately sets it apart from *C. elongatus*. The other major feature distinguishing the present species from *C. elongatus* is its possession of two sets of teeth anteriorly, though differing from other members of *Coleps* in its plate structure, three triangularly arranged posterior spines, two caudal cilia and overall body size and shape, the species described herein requires data based on a sufficiently large population to establish its taxonomic

allocation.

42.

Family Holophryidae

Genus Urotricha Claparede and Lachmann (according to Corliss, 1961).

Urotricha sp. (Plate IX, Figs. 17-19).

Body spherical to slightly oval $(27.3 \ \mu \ x \ 24 \ \mu$, based on 23 specimens, 20 - 34 μ x 17 - 29 μ), posterior third unciliated save for one long caudal cilium (B in Figs. 18 and 19), ciliation otherwise uniform (Figs. 17 - 19). Cytostome round at anterior surface, surrounded by ring of heavier cilia and lacking significant peristome. No trichocysts observed. Contractile vacuole posterior, macronucleus spherical. Progression both by 'leaping' and steady 'swimming'.

Systematic account:

Having its cyclostome at the naterior surface but lacking a peristome, the small ciliate under consideration bears the main characteristics of Urotricha. This genus differs from other members of the Holophryidae in that its rear third is unciliated except for one to several long caudal cilia.

On the basis of body shape, body size, number of caudal cilia and the presence of trichocysts, Kahl (1930) recognized 12 species of Urotricha. The present species differs from U. synuraphaga, U. ovata, U. obliqua, and U. armata in its absence of trichocysts. Its single caudal cilium further distinguishes it from U. pusilla, U. furcata and U. saprophila. On the basis of body shape, the anteriorly trilateral tapering U. agilis, jug-like U. farcta and flask-shaped U. lagenula do not merit comparison with the spherical to oval species under discussion. PLATE IX

Figs. 17 - 19. Urotricha sp.

Fig. 17. Showing spherical to slightly oval forms.

Fig. 18. Showing (A) a ring of heavier cilia around cytostome;

(B) long single caudal cilium.

Fig. 19. Showing (B) long single caudal cilium and truncate posterior extremity.

ارم کلم

The latter is closer to the spherical U. globosa (18 μ) than the cylindrical U. discolor (25 μ). Its body size (27.3 x²24 μ) is closer to the latter than to the former.

Of the known species of Urotricha, none have charcteristics quite comparable to those of the species under consideration. However, the establishment of its systematic allocation requires further study

of its infraciliary morphology.



	Subclass	Holotricha
٠,	•	- ·
	Order	Trichostomatida
2		

Family, Colpodidae

Genus Colpoda

Colpoda cucullus 0. F. M., 1773. Plate X, Figs. 20-23.

Size and shape variable, depending upon the amount of food available. In general, reniform and compressed, overall measurements $42 - 64\mu \ge 31 - 45\mu$ (av. $57.8\mu \ge 39.4\mu$). Right border semi-circular, posterior half of the left border often convex. Oral funnel in the middle of flattened ventral side. Cytostome ventrally located and displaced to the right of the median plane, leading into the peristome cavity and gives rise dorsally to a diagonal groove; its edge bears a 'ciliated area, but no protruding membrane. Right lateral ciliary field, rotated about 60° to assume an anterior position. It appears' as a rather narrow crescent closely adjacent to which lies the anterior 'extremity of the stomatogenous meridian whose basal bodies are 'quite small at this level.

Cilia either single or paired. When present singTy they are confined to the central posterior body region. Being arranged along pellicular grooves they give the distinctive notched appearance to the preoral keel anteriorly. These, which are deepest at the anterior extremity, form a polygonal figure posteriorly. The number of meridians varies from 29 to 34; anterior keel with six to ten indentations (Figs. 20, 21). Macronucleus with a stellate endosome, central Figs. 20 - 23. Colpoda cucullus 0, F. M. 1773. Figs. 20 and 21 from silver impregnation preparation by the Chatton-Lwoff technique, and viewed by phase-contrast microscopy. Figs. 22. and 23, from life, as seen by Nomarski interference-contrast effect.

PLATE X

Fig. 20. Showing the left side.

Fig. 21. Showing the right side.

Fig. 22. Focused on the food vacuoles.

Fig. 23. Pointed to the contractile vacuole, indicated by arrow.



Trichocysts rod-form.

Freshly excysted ciliates exhibit a clear hyaline cytoplasm which rapidly assumes a grayish opacity when food is posthe accumulation of food vacuoles (Fig. 22). Contractile vacuole posteriorly located (Fig. 23).

Systematic account:

On the basis of size, Burt (1940) divided the *Colpoda* species into two groups: (1)—the small forms, *C. steini* and *C. aspera*, averaging about 30μ in length and (2) the large ones, *C. maupasi*, *C. inflata*, and *C. cucullus*, averaging more than 50μ in length. The present species bears comparison with the three whose length ranges from 35μ to 110μ .

The numbers of the pre-oral indentations and the total ciliary meridians of *Colpoda* serve as the significant criteria for accurate species diagnosis. The present species is referable to *C. cucullus* by its size, eight to ten pre-oral indentations and the total number of ciliary meridians. It is easily distinguished from *C. maupasi* (5 pre-oral indentations) and *C. inflata* (6 - 8 pre-oral indentations).

Family Microthoracidae for Genus Microthorax

Microthorax sp.

Plates XI - XIV, Figs. 24-35.

Small (18 - 24µ long, based on 18 specimens), kidney-shaped, compressed laterally; with delicate keeled armour which is more or less pointed anteriorly and rounded posteriorly; ventral armour with three deep ciliary rows (Figs. 29, 31 and 33); oral depression posterior-ventral (Figs. 24 - 34), with a stiff ectoplasmic lip on right side (Figs. 25, 26 and 28), below which there is a small membrane (Figs. 26 - 28, 31 and 34), and with a small tooth on left margin (Figs. 25 - 27, 30 - 34), no cytopharynx; single macronucleus spherical (Fig. 35), two contractile vacuoles.

Systematic account:

The present species bears the characteristics of *Microthorax* which is distinguishable from other genera in Microthoracidae by its posterior-ventrally located oral depression, without cytopharynx, trichocysts, and trichites.

The Logy Bay species differs from Microthorax scutiformis, M. elegans, M. simulans in its frongal half without deep furrow across yentral-dorsal edge. Its rounded posterior extremity separates the present species from M. tridentatus, M. bidentatus, M. glaber, M. ungulatus, M. spiniger, and M. costatus. Among the rest of the species of Microthorax (recognized by Kahl, 1930), Hemicyclium lucidum Eberhard, (Microthorax haliotideus Penard), M. pusillus, M. auricula with.slightly PLATE XI

14'

50

Fig. 24. Microthorax sp._____. Showing small (23.5µ x 12.5µ),

. kidney form with posterior-ventral oral depression.

PLATE XII

Figs. 25-27^f. *Microthorax* sp. Showing a small tooth on left margin of oral depression.
PLATE XIII

52

Figs. 28 - 31. Microthorax sp.

Fig. 28. Showing a stiff ectoplasmic lip on right side of oral depression.

Fig. 29. Showing ventral armour with three deep ciliary rows.

....

d. .

Figs. 30 and, 31. Showing oral depression with membrane.

PLATE XIV

Figs. 32 - 35. Microthorax *p.

;

Figs. 32 - 34. Showing posterior-ventral oral depression with a small

tooth on left margin.

Fig. 35. Showing single spherical macronucleus as stained with methyl green.

ridged ventral side, thus bear no comparison with the deep-ridged present species.

The present species is similar to the remaining two species, M. sulcatus and M. viridis. The latter with zoochlorellae is distinguishable from the present species. Only M. sulcatus resembles the present species in the discussed criteria.

However, the smaller size $(18 - 23 \mu)$ and its slanted oral depression with sigmoid edge along ventral posterior edge do not fit the decsription of *Microthorax sulcatus* Engelmann (1861). Its taxonomfc allocation, however, should only be determined after extensive further study of the infraciliary structure.









Hymenostomatida Order Tetrahymenina Suborder,

Family Tetrahymenidae

Genus Tetrahymena (according-to Corliss 1961a).

Furgason's precise taxonomic work (1940) validated the genëric name "Tetrahymena". This name is derived from the tetrahymenal buccal ciliary apparatus which comprises four membranous structures, one undulating membrane on the right and three membranelles on the left of the buccal cavity. The entire buccal apparatus is typical for the suborder Tetrahymenina (Corliss 1952a, 1953a, 1956). That the direction of the cytostome axis at once distinguishes Tetrahymena from such related genera as Glaucoma, Colpidium, Loxocephalus (Furgason, 1940; Corliss, 1952a, 1953a).

Species of the holotrichous genus *Tetrahymena* Furgason, 1940, are typically of small size and covered with fine cilia arranged in longitudinal rows. Their body is pyriform in at least one stage of the life cycle. They possess a rather inconspicuous oral opening equipped with a tetrahymenal buccal ciliary apparatus and located on the ventral surface near the anterior extremity of the body.

> Tetrahymena voraz (Kidder, Lilly and Claff, 1940) Plates XV - XVII, Figs. 36 - 47.

The marked dimorphism of this ciliate was evident by phase contrast microscopy. Silver impregnation preparations were then made. Figs. 36 - 39. Tetrahymena vorax (Kidder, Lilly and Claff, 1940) from silver impregnation preparation by the Chatton-Lwoff technique, viewed by phase-contrast microscopy.

56

Fig. 36. Showing different stages, for comparison of the size and shape of microstome form with that of macrostome form (pointed one indicated by arrow A).

Fig. 37. Showing microstome stage, both the tailed fusiform and pyriform shapes (indicated by arrows).

Fig. 38. Showing different forms. A pointed one is indicated by arrow B.

'Fig. 39. Showing both tailed fusiform microform and macrostome. The pointed macrostome tomites are indicated by arrow C.

PLATE XV



'PLATE XVI

Figs. 40 - 43. Tetrahymena vorax (Kidder, Lilly, and Claff, 1940) from silver impregantion preparation by the Chatton-Lwoff technique,

viewed by phase-contrast microscopy.

Fig. 40. Showing pyriform microstome from Fig. 38. AZM, adoral zone membranelles; UM, undulating membrane complex; X, unknown structure; POM, post-oral meridians. Notice that the meridians are arranged with uniform regularity-anteriorly, joining irregularly at the posterior extremity (cf. Figs.

Fig. 41. Showing mature macrostome form showing the large pharyngeal pouch and ingested prey.

44 - 47).

Fig. 42. Showing the same mature macrostome as shown in Fig. 36. Notice the size, shape and position of the cytostome and the distance between adjacent cilia increased.

Fig. 43. Showing the same macrostome tomites shown in Fig. 39.



PLATE XVII

Figs. 44 - 47. Tetrahymena vorax (Kidder, Lilly and Claff, 1940) from silver impregnation preparation

by the Chatton-Lwoff technique, viewed by phase-contrast microscopy. 58

Fig. 45. Showing tailed microstome with body ciliation. Notice two

Fig. 46. Illustrating the ciliation around apical-loop (A-P) of microform.

Fig. 47. Showing three contractile vacuole pores (CVP) of microstome stage.

(2° PM)



The following data are based principally upon silver impregnation . material (Chatton-Lwoff method). As necessary, they are supplemented by information derived from live ciliates.

Life cycle.

The changes in body form of the present strain were shown by phase-contrast to be very comparable to Figure 3 of Corliss (1953a). The following descriptions of the pronounced polymorphism exhibited by the present strain are based upon observations of living material as well as mixed staining (neutral red and methylene blue) and (pringipally) silver impregnation.

Tailed microstome stage (Figs. 36-39, 44-47, c.f. Corliss, 1953a, Fig. 3a).

The "thin, tailed organism" reported as the principal form found in nature was seen in both living and stained material. It is a microstome form, i.e. the buccal opening is small. Except for its elongate posterior extremity, slightly larger overall body size, and greater range in number of meridians and contractile vacuole pores, it is practically identical with *T. pyriformis*. Its appearance in silver impregnation proparations is similar to Fig. 3g of Corliss (1953a). These forms are generally very elongate, with long and sharply pointed tails (Figs. 36-38). However, the buccal opening and other morphological features are identical to those described as the "tailed microstome stage" by Corliss (1953a). These forms were probably starved when fixed for silver impregnation.

Pyriform microstome stage. (Figs. 37, 38 and 40, c.f. Figs. 3i, j

of Corliss, 1953a).

This was described by Corliss as "a curious stage in that forms in it are morphologically indistinguishable from *T. pyriformis*". It was observed both in live and stained material, together with tailed microstome forms, from which it is derived. Once all the individuals in a culture have reached this stage, they are irreversibly pyriform, and can only be identified as *T. pyriformis*. 3. Transitional stages from tailed microstome to macrostome form

(Figs. 3b, 3c of Corliss, 1953a).

In life, where both tailed microstome and pyriform microstome stages were observed, the morphology of the former is subject to marked and rapid change. Within a very few minutes, mature macrostome forms result. A condensed account based upon both living and stained material was given by Corliss (1953a).

 Mature macrostome stage (Figs. 36, 39 and 41 - 43, cf. figs. 3d, 3e, and 3f of Corliss, 1953a).

Several rounder, broader forms had eventuated from the transitional stages, resulting in the buccal opening sometimes being broader than long. It may be considered as a true trophont-tomont, the active swimming and hunting phase temporarily over. Division of the tomont, as the macrostome form may be properly considered for a short? time in its life, proceeds normally with approximately equal distribution of food vacuoles and various other cytoplasmic inclusions to the new resulting tomites, the proter and opisthe (cf. figs. 39 and 43 with fig. 3f of Corliss, 1953a). Corliss (1952a, and 1953a) gave detailed figures and investigations; however, superficially this stage is indistinguishable from the macrostome form of *T. patula*. <u>Body size</u>: tailed microstome stage 35 - 80µ long; average overall dimensions, 50 x 20µ. Pyriform microstome stage, 30 - 60µ long; average size, 51 x 35µ. Macrostome stage, 65 - 154µ; average size, 98 x 63µ.

Body shape: Shape varies with stage in life cycle. Tailed microstome stage: elongate; anterior extremity bluntly pointed, posteriorly distinct caudal process variable in shape and length. Body elongation accentuated when starved, tails tapering to very sharp tip. Pyriform microstome stage: pyriform to ovoid to cucumber-shaped, indistinguishable from forms of *T. pyriformis*. Macrostome stage: broadly pyriform with rounded, enlarged posterior extremity due to presence of many large food vacuoles.

<u>Macronucleus</u>: in all microstome forms, identical with that described for *T. pyriformis*. It is ovoid to irregularly spherical, and lies across the central portion of the body, often a little posterior to the midline. In macrostome stage, also similar but of greater size, averaging about 25 x 22µ.

<u>Contractile vacuole</u>: usually one contractile vacuole is posteriorly located.

<u>Food vacuoles</u>: in general similar to those in *T. pyriformis* (for detail see descriptions by Maupas, 1883; Fauré-Fremiet, 1906, 1910; Furgason, 1940; Browning, 1951; Corliss, 1953a and Elliott and Clemmons, 1966). Average size of food vacuoles in bacterized cultures: $3.5 - 4.5\mu$ in diameter. In carnivorous forms, diameters of food *t* vacuoles range up to $35 - 40\mu$; number present in cannibals, generally not over 5; in non-cannibals, 3 - 20, average about 10.

Body ciliation: no apparent differences between ciliary patterns of microstome and macrostome stages. The cilia are arranged in distinct meridians, converging in a uniformly regular manner anteriorly, and joining irregularly at the posterior extremity (Figs. 44 - 47). Distance between adjacent cilia increased in macrostome • forms (Figs. 41 - 43). No "caudal cilium" observed.

<u>Buccal overture</u>: pyriform, right margin slightly more rounded than left; greatly enlarged and relatively broadened in macrostome stage; directly in body axis, ventral (Figs. 40, 42 and 44). Adoral zone of membranelles (AZM). Figs 40 and 44. Three membranelles (M_1, M_2, M_3) attached to left dorsal lateral wall of buccal cavity, M_1 furthest to the left. Composed of cilia considerably longer than those of the body.

Urdulating membrane complex (UM): Undulating membrane, on right-hand border of buccal cavity. See descriptions by Fauré-Fremiet (1948) and Corliss (1953a).

Pharyngeal pouch: present in macrostome form, presumably associated with development of carnivorous habit. Mode of formation briefly described by Kidder et al. (1940).

Apical loop (A-L): identical with that described for T. pyriformis by Corliss (1953a), although proportionately larger and generally open ventrally in macrostome forms.

Ciliary meridians.

<u>Microstome forms</u>: range in number, 17-23; average, 19-21. <u>Macrostome forms</u>: range, 19-26; average 21-23, Two typical unipolar or post-oral meridians (POM's) with anterior extremity

terminating directly at posterior margin of mouth or in case of meridian 1, a very short distance up left side of buccal opening (Figs. 44 and 45). Occasionally three POM's: meridians 1, n and n-1, last stopping half-way up left side of mouth (Fig. 40). Each meridian comprises two main portions, a primary portion (1°PM) and a secondary portion (2°PM) (Fig. 45). The primary portion is much more prominent and is heavily granular as it bears the basal granules of the body cilia; the secondary portion is a strong fibril. Posteriorly, 2°PM's unite with their respective 1°PM's just before irregular convergence of meridians in polar region. Frequently (e.g. in Fig. 41-43) additional 3°PM fibril is in evidence.

Contractile vacuole pores: three contractile vacuole pores (CVP's) are detectable in Fig. 47, they are probably located in meridians 5 - 9 (for details cf. Corliss, 1953a).

Systematic account:

Among the 13 recognized species of Tetrahymena, T. parasitica and T. glaucomaeformis require redescription. So does T. faurei, which has only been reported once.

T. patula, T. vorax and T. paravorax are strikingly distinguishable from other members of the genus by their ability to undergo, a pronounced dimorphism (microstome \leftrightarrow macrostome form) during their life cycle. T. paravorax is readily separated from its two close associates by several characteristics, notably by its possession of a distinct caudal cilium (Gorliss, 1957).

T. vorax and T. patula share one of the taxonomic extremes of the genus Tetrahymena with T. pyriformis. In addition to their ability to undergo microstome \leftrightarrow macrostome transformation, there are , other similarities between T. vorax and T. patula. Both are able to form cysts (Fauré-Fremiet, 1949; Williams, 1961), and the overall anatomy of the body and cytostome of T. vorax is strikingly similar to normal macrostome stage of T. patula.

However, their life cycles are different in certain other *i* respects. The tomont of *T. patula* is able to transform reversibly to either microstome or macrostome tomites. In *T. vorax*, tomites initiate similar reversible transformations. Once the tailed fusiform microstome phase of *T. vorax* is formed from a tomite, it can either transform (reversibly) to macrostome phase or (irreversibly) can become a pyriform microstome. This is not so in *T. patula*. Through the rep-roductive cyst, the mature tomont can develop into a pyriform microstome stage of *T. patula* is commonly so shaped) which may afterwards revert to the macrostome form. On the other hand, the tailed microstome is the form of *T. vorax* commonly found in nature.

The principal differences between *T. vorax* and *T. patula* are based upon the silverline structures. Thus the two POM's, and the location of three CVP's between meridians number five and nine, and the average 19-21 body meridians shown in the present strain are diagpostic of *T. vorax* rather than *T. patula*, which has been reported with four POM's, two to six CVP's (between meridians number 10-15), and an average of 36-42 body meridians (Corliss, 1953a)

of the life cycle, and of the silverline structures of the strain under consideration fit the data described for *T. vorax* (Corliss, 1953a).

However, there are two points which must be considered before a final conclusion is drawn. Firstly, the present strain has a puzzling structure in the buccal cavity (marked with 'X' in the figures). Secondly, the pyriform microstome forms mentioned above are not identifiable with those of *T. vorax*. These pyriform ciliates are certainly not referable to *T. patula*; the microstome phase of which possesses an average of 36-38 meridians. They might either be "*T. pyriformis*" (the prey and stomatin-Buhse, 1966a and 1966b and 1967 - source of *T. vorax*) or iffeversible pyriform microstome forms derived from *T. vorax*. Again, one could be dealing with mixed populations of the two species. This phylogenetic problem with only be solved by future biochemical and serological tests with the strain under discussion, which in the meantime is identified as *Tetrahymena vorax* (Kidder et al., 1940) sensu *latu*.

Tetrahymena pyriformis (Ehrenberg, 1830)

Lwoff, 1947, Plates XVIII, XIX, Figs. 48-57.

Well-fed forms are generally pyriform (Figs. 49-51), starving ones more cylindrical and sometimes becoming extremely attenuated 128-89µ.x 15-32µ). Various modifications (e.g.boomerang and banana forms, Fig. 48; ovoid, Figs. 52, 53) have also been observed. Macronucleus, ovoid to irregularly spherical, and across the central portion of the body, often a little posterior to the midline. No micronucleus. One contractile vacuole posteriorly located (Figs. 49-51). Uniform somatic

PLATE XVIII

Figs. 48 - 53. Tetrahymena pyriformis (Ehrenberg, 1830).

Fig. 48. Showing a variety of shapes. Photomicrograph from silver impregnation preparation by Klein's dry method, viewed by 'phase-contrast microscope.

Fig. 49. Showing typical body shape, ciliation, buccal cavity, and the contractile vacuole. Photomicrograph from life, as seen by Nomarski interference-contrast effect.

Fig. 50. Same as Fig. 49, but focused on food vacuoles for comparison with Figs. 51 - 53. The latter represent phase-contrast views of living examples,

Fig. 51. Showing body shape, ciliation, buccal cavity, and the contracile vacuole.

Fig. 52. Showing well-fed oval individuals with prominent undulating membrane; nucleus (N), food vacuoles and contractile vacuole
 are detectable inside the cells. The dots (b) around the ciliates are the bacterial food of T. pynet, formis.

Fig. 53. Food vacuoles showing clearly after staining (neutral red and methylene blue). Photographed by phase-contrast mic-. roscopy.



PLATE XIX 11

67

Figs. 54 - 57. *Petrahymena pyriformis* (Ehrenberg, 1830) from silver impregnation preparations (Klein's dry method) as seen by phasecontrast microscopy.

Fig. 54. Showing body shape, ciliation, the 'ribs' of undulating membrane complex, and meridian number one-with cytoproct (CYP) and the early stage of stomatogenesis: OA, oral anlage.

Fig. 55. Showing the later stage of stomatogenesis.

Fig. 56. Showing the location of two contractile vacuole pores (CVP's and the arrangement of the primary (1°PM) and secondary

(2°PM) meridians.

Fig: 57. Showing location of the three CVP's.



ciliation (Eilia, 5-8µ in length, Fig. 49).

Tetrahymenal buccal ciliature is evident (Fig. 49-51 and 54) and pyriforma. Its axis parallel to body axis, about one sixth to one eighth of the body length. Adoral zone membranelles are easily demonstrable by staining (Figs. 52, 54 and 55, cf. figs. 40 and 44). Two postoral meridians (Figs. 54 and 55) Ciliary meridians, 17 - 22. 3° PM fibril is not detectable; otherwise similar to *T. vorax*.

New buccal structure (OA, Figs. 54 and 55) originates in a proliferation of granules of meridian 1 out to left. Resulting granular field, located subequatorially, eventually encroaches upon meridian n. (cf. Chatton, Lwoff, Lwoff and Monod, 1931, and Corliss, 1953a).

Cytoproct (CYP, Fig. 54) located in meridian l very near posterior extremity of organism. When detectable, "appears as delicate line, actually a slit, lying more or less between 1°PM and '2°PM.

Two contractile vacuole pores (CVP's, Fig. 56) generally located in meridians 5 and 6, at distance of 6 - 14µ from posterior pole of body. When three, CVP's are present (Fig. 57), 'location, is generally in meridians 5, 6 and 7 in various positions with regard to 1°PM's and 2°PM's of respective meridians.

When this ciliate is held in a medium rich in bacteria, its food vacuoles are readily seen by light microscopy (Figs. 49 - 51). They are still more clearly demonstrable after staining the ciliates in dilute suspensions of methylene blue (1:)10,000) or neutral red (1:30,000) (Fig. 53). Average diameter of food vacuoles in bacterized cultures, 3.5 - 4.5µ (see Maupas, 1883; Fauré-Fremiet, 1906, 1910; Tetrahymena pyriformis was first described by Ehrenberg (1830) as Leucophrys pyriformis. Afterwards it acquired many synonyms. There was thus considerable confusion until Corliss thoroughly analysed and systematically reviewed the earlier literature (Corliss, 1951, 1952a, 1952b, 1953a and 1954).

Furgason, 1940; Browning, 1951; Corliss, 1953a and Elliott and Clemmons

1966).

Systematic account: "

Type species of the genus *Tetrahymena*, *T. pyriformis* is a bacteria-feeder in natural habitats. Characteristically pyriform in shape, it has 17 to 21 meridians and possess strong generic characters - the tetrahymenal buccal apparatus, location of cytoproct on meridian 1, and two contractile vacuole pores. Its approximate size is 50 x 30µ, although there is considerable variation under different conditions: It has a single macronucleus, with or without micronucleus.

The striking similarities between the type species, T. pypiformis, and the strain at Logy Bay strongly suggest that the latter is identical to T. pypiformis. However, a brief and conservative comparison of the strain at Logy Bay with all the members in the genus is stated as follows.

The strain under consideration can be easily distinguished from T. rostrata, T. corlissi, T. setifera and T. paravorax by lacking a caudal cilium and polar basal granule (PBG) - complex.

Its free-living habitat contra-indicates allocation to those species that are parasitic, viz., T. stegomyiac, T. parasitica, T. limacis

T. faurei and T. chironomi.

T. patula and T. vorda are beyond consideration since the present strain does not show dimorphism (microstome \leftrightarrow , macrostome)

during its life cycle.

Redescription of *T_f glaucomaeformis* is needed before any close comparisons can be made.

This situation leads to the suggestion that the present strain should be assigned to T. pyriformis.

In sum, the present species is referable to *T. pyriformis* (Ehrenberg, 1830) by its overall morphology, its lack of a caudal cilium and PBG - complex, its life cycle, its number and arrangement of meridians, the number and position of GVP's, the absence of micronucleus, the free-living habit. The specimens illustrated in the photomicrographs (Figs. 48, 54-57) were double-checked and referred to *T. pyriformis* by Dr. J. O. Corliss (Pers. comm. 1969).

Subclass	Holotricha
Order	Hymenostomat
Suborder	Pèniculina

Oral ciliature characterized by peniculi (Fig. 63) located deep in the buccal cavity; often with ciliated vestibule; body uniformly ciliated.

Family Parameciidae

Cenus

Paramecium (according to Corliss, 1961)

Parameeium bursaria Ehrenberg, 1833, Plates XX-XXII, Figs.58-63.

Body 90 - 150µ foot-shaped. More or less flattened, prominently compressed dorso-ventrally, broadly rounded posteriorly, extending obliquely backwards from left to right to beyond the centre of the body. Cytostome situated at the posterior extremity of the groove, giving entry to a distinct cytopharynx. Buccal cavity with one endoral membrane (Fig. 59) and two peniculi (Fig. 63) Trichocysts numerous (Fig. 60). A contractile vacuole deep in the cytoplasm towards each end of the body (Fig. 58). These vacuoles exhibiting rapid and alternate systole/diastole, each with a short discharge pore and six radiating collecting canals (Fig. 58). A single macronucleus, large and reniform. Micronucleus also single, massive, and lying in the cavity of the macronucleus (Figs. 61 and 62).

Systematic position:

Woodruff (1921) pointed out that species of *Paramecium* fall into two groups according to the shape of the body, namely: (a) The

PLATE XX

Fig. 58. Paramecium bursaria Ehrenberg, 1833, fresh preparation as seen by Nomarski interference-contrast effect. Uniform ciliation without long caudal cilia evident from the figures. 1. trichocyst; 2. contractile vacuoles;

3. zoochlorellae



Figs. 59 - 62. Paramecium bursaria Ehrenberg, 1833. Fig. 59. Showing the endoral membrane around the buccal cavity (by Nomarski interference-contrast effect).

PLATE XXI

Fig. 60. As seen with Nomarski interference-contrast effect, showing abundant development of trichocysts. Beneath the trichocysts the cortical and endoplasmic layers appear green, because of the presence of numerous zoochorellae.
Figs.61,62. Showing the macronucleus (Ma) and the compact micronucleus (Mi), as stained with methyl green and viewed by bright

field.



Fig. 63. Paramecium bursaria Ehrenberg, 1833, impregnated with silver by the Chatton-Lwoff technique, photographed by phase contrast. Showing peniculi located deep in the buccal cavity and body ciliature.

PLATE XXII



"aurelia group" with cigar-shaped bodies, round in cross-section and tapering to a point posteriorly, and (b) the "bursainia group", broadly elliptical in cross-section and rounded posteriorly. In either group two types of micronuclear structure may be found. These are (a) the "caudatum type", in which the micronucleus is a relatively large, rather compact mass, and (b) the "aurelia type", in which the micronuclei are small and distinctly vasicular in organization. On this basis Wenrich (1928) recognized eight well-defined species - P. aurelia, P. caudatum, P. multimicronucleata, P. trichium, P. calkinsi, P.

The organism under discussion is somewhat compressed dorboventrally, rounded posteriorly, and typical of the "bursaria group" in shape. Its body is not cigar-shaped like that of members of the, "aurelia group" (which includes P. aurelia, P. caudatum, and P. multinicronucleata). In the "bursaria group", P. trichium and P. bursaria are distinguished from P. calkinsi, P. polycaryum, and P. woodruffi by the former having a single compact micronucleus (caudatum type), while the latter have small vesicular micronuclei (aurelia type). The length of the present species, $90 - 150\mu$, is closer to that of P. trichium (70 - 90µ). One of the chief features of P. trichium, a convoluted discharge tube connected to the contractile vacuoles, was lacking. However, two contractile vacuoles, each with a short discharge pore and six radiating collecting canals, were clearly evident, especially when the water around the example on the slide was drying out. Rapid cyclosis was meanwhile observed in the ciliate. In sum, these criteria indicate that the organism is referable to P. bursaria rather than to
any other known species. Moreover, the bright green colour. (due to symbiotic zoochlorellae) is itself strongly suggestive of *P. bursaria*, the only well-established member of the genus so characterized.

Two other species, P. putrinum and P. jenningsi (see Kudo, 1966) must, nevertheless, be taken into consideration. In general morphology, the Newfoundland ciliate does not resemble P. jenningsi which is clearly closer to P. aurglia. Though it is certainly closer to another representative of the bursaria group, P. putrinum (80 -150µ long) in size and shape, it can still be identified with confidence as P. bursaria by its two contractile vacuoles and the presence of green zoochlorellae (P. putrinum has only a single contractile vacuole and altogether lacks zoochlorellae).

To summarize, *P. bursaria* can easily be recognized by its size (usually 120 - 160µ in length), rounded posterior end, green colour due to the symbiotic zoochlorellae, and rapid cyclosis. To these characters must be added a highly characteristic feature immediately evident from stained examples - the single large compact micronucleus of the *caudatum*-type, which resolves the question of identity beyond any reasonable doubt.

The Logy Bay ciliate under discussion is typical of *P. bursaria* in every respect, except that its length $(90 - 150\mu)$ is somewhat less than usual $(120 - 160\mu)$. It has two spherical vacuoles, which, depending upon the phase of systole/diastole, show no evidence of radiating canals or appear stellate with radiating canals. The macronucleus is always reniform, the compact micronucleus being situated in its concavity. As is characteristic of *P. bursaria*, the organism is dorso-

ventrally flattened, narrowest and obliquely truncate anteriorly and posteriorly rounded. Its cytoplasm is full of small green algae and cyclosis is rapid.

In the silverline system of *P. bursaria* (Fig. 63), none of the ventral rows of cilia are parallel to the preoral suture in the region anterior to the cytostome. However, they bend towards and terminate against the preoral suture. There are 10.1 rows of cilia in a transverse distance of 12 microns, and 7.0 in a longitudinal distance of 12 microns, there being 70 cilia in an area of 144 square micra on the dorsal surface. These figures are a little higher than those of Lieberman (1929). According to the latter author, *P. bursaria* has 9.75 rows of cilia within a width of 12μ , 6.6 rows within a length of 12μ , and 64.45 in 144 sq. μ .

Family	Frontoniidae			•	
		÷ .			·
Genus	Cyrtolophosis	(according	to	Kudo,	1966)

Ovoid or ellipsoid; with or without mucilaginous envelope in which it lives, but from which it emerges freely; cytostome near anterior extremity with a pocket-forming membrane; on right side a short row of special stiff cilia, bent ventrally; sparse ciliation spiral to posterior-left.

Kahl (1930) considered that the genus Balantiophorus was wrongly established by Schewiakoff, whose proposed species had already described and allocated to the genus Cyrtolophosis by Stokes (1888). Corliss included them as "Unassigned Genera" of tetrahymenine hymenostomes (Corliss, 1961a, p. 146). However, Kudo (1966) allocated Cyrtolophosis as a member of Frontoniidae. Thus the present clas-

> Cyrtolophosis (Balantiophorus) bursaria Schewiakoff, 1893, Plate XXIII, Figs. 64-67.

Body $_{\sigma}(30\mu \times 17\mu)$ transparent, ovoid, rounded at both extremities, dorsal surface more convex than the ventral one (Figs. 64 67). Left, posterior and right margins of the peristome with a baglike undulating membrane (Fig. 66), which can be withdrawn into the peristome. Anterior extremity bearing a fascicle of long, distally curved, vibratile hairs (Figs. 65 - 67). Somatic cilia fine and sparse (Fig. 64). No mucilaginous envelope observed. Contractile vacuole posteriorly (Figs. 64 - 67). Macronucleus ovoid, centrally

PLATE XXIII

Figs. 64 - 67. Cyrtolophosis (Balantiophorus) bursarta Schewiakoff, 1893.

Fig. 64. Showing fine and sparse somatic cilia.

Fig. 65, Showing anterior extremity with a fascicle of long, distally

curved, vibratile hairs.

Fig. 66. Showing ovoid, central located macronucleus.

Fig. 67. Showing posteriorly located contractile vacuole.



located (Figs. 64 - 66). Swims jerkily, revolving about the long axis in a counter-clockwise direction as viewed from the rear.

Systematic account:

síze.

1893.

The above data precisely fit the description of Cyrtolophosis (Balantiophorus) bursaria Schewiakoff, 1893. The present species exhibits the general characters of Cyrtolophosis, being distinct from C. elongata, C. mucicola and C. major in its ovoid body shape. C. bursaria is the only known member of the genus which is ovoid, the other three being elongate-ellipsoid.

Furthermore, the present species (30μ) is distinguishable from *C. elongata* (30μ) in that its contractile vacuole is about onefifth of the body length from the posterior extremity; and from *C. mucicola* (25-28µ) and *C. major* (45µ) in its appreciably different body

For the reasons detailed above, the species under discussion is referable to Cyrtolophosis (Balantiophorus) bursaria Schewiakoff,

Cyrtolophosis (Balantiophorus) elongata

(Schewiakoff, 1896) Plate XXIV, Figs. 68-71.

Body elongate-ellipsoid, 30 x 10μ , the ventro-frontal border obliquely truncate. Anterior cilia longest, about 4μ , the anteriorly placed fascicle of distally vibratile and downwardly curved cilia conspicuous (Fig. 68). Peristome extending from the frontal border for one-third of the length of entire body. Left, posterior and right

PLATE XXIV

Figs. 68 - 71. Cyrtolophosis (Balantiophorus) elongata Schewlakoff, 1896, (phase-contrast).

Fig. 68. Left side of the organism, focused on the elliptical macronucleus and the anterior fascicle of cilia.

Fig. 69. Right side of the organism, showing the right margin of the

peristome, with undulating membrane.

Fig. 70. Extension of the peristome, and oral ciliation.

Fig. 71. Focused on the posterior contractile vacuole.



margins of the peristome with a bag-like undulating membrane (Figs. 69, and 70), which can be withdrawn into the peristome. Contractile vacuole single, spherical, posteriorly located (Fig. 71). Macronucleus ellipsoid, and an adjacent micronucleus. No mucilagious envelope observed. Occasionally swims backwards, revolving about the long axis in a counter-clockwise direction as viewed from the rear.

The available taxonomic data best fit those of *Cyrtolophosis* (*Balantiophorus*) elongata (30µ x 10µ) which is characterized by a posterior contractile vacuole, long, sparse scattered cilia not arranged, in longitudinal rows, and an ellipsoid macronucleus and small micronucleus.

Like *C. elongata* as earlier described it has its contractile vacuole at the posterior extremity - the other three species have it some one-fifth of the body length from this extremity. It is further distinguished from *C. bursaria* by its narrowly elongate body shape, and from *C. major* (45μ) by its smaller size, the lack of diagonal to longitudinal rows of cilia around the body, and an un-notched anterior extremity.

C. mucicola (25-28µ by 9-12µ) is close in size and shape to the Logy Bay ciliate. It is indeed impracticable to differentiate them on these bases or on the location of the contractile vacuole. " However, C. mucicola is distinct from the present species in its denser vestiture of longitudinally arranged cilia, and in having a spherical

macronucleus.

Systematic account:

The Logy Bay species is thus considered referable to C. elongata Stokes on the basis of the location of its contractile vacuole, its body shape and size, its spherical macronucleus, its un-notched anterior extremity and the arrangement of its cilia.

Genus Lembadion Perty (according to Kudo, 1966)

Frontoniidae

Family .

Lembadion sp. Plate XXV, Figs. 72 and 73.

84

Body oval (55 - 62µ x 28 - 32µ based on eleven specimens); dorsal side convex, ventral side slightly flattened; striated longitudinally. Posterior extremity rounded, the anterior one with a small dextral process (Fig. 72, A). Cytostome (47µ) three quarters to four-fifths the body length; voluntarily vibratile membrane to its right, projecting from the anterior extremity as a short, conical, hood-like extension, one side of which is inserted on the anterior left-hand margin, the other (Fig. 72, B) being continued as a conspicuous, lamellate membrane for about two-fifths of cytostome. It is then abruptly narrowed, and descending into the cytostome is continued to the posterior extremity. Two contractile vacuoles, the larger \sim \sim (Fig. 72,C) sub-central and near the dorsal surface with a long posterior tubule opening (D of Figs. 72 and 73) at the right side; the smaller (Fig. 72,F) posteriorly placed somewhat to the left side of the median line. Cilia fine and short anteriorly, bundle of caudal cilia (about 23µ long) (Fig. 72,G) sinistrally directed. Macronucleus, (Fig. 72, E) ellipsoid, subterminally located near the left-hand border. Cell division transversely.

Systematic account:

This holotrichous ciliate exhibits the chief features of *Lembadion*, which is characterized by its huge cytostome. The latter

PLATE XXV Figs. 72.- 73. Lembadion sp. Ventral aspect (phase-contrast)

Dextral anterior process.

Abruptly narrowing right adoral membrane. The larger contractile vacuole (sub-central and near the dorsal surface) with its tubule (D) opening posteriorly at the right side.

E. Macronucleus.

Ά.

Β,

С.

Ø

G.

F. The smaller contractile vacuole.

The sinistrally directed caudal cilia.



occupies at least two thirds of the ventral surface, and while having membranes lacks membrane-bordered pre-oral peristomal furrow.

Kahl (1930) recognized three species of Lembadion¹. The present species differs from all of these. Ranging from 80 - 200µ in length, they are apparently larger than the Logy Bay ciliate (55 - 62µ). Furthermore, the present species has two contractile vacuoles, and a cytostome with abruptly narrowing membrane two-fifths (20µ in Fig. 72) of the body length from its right anterior extremity.

The present species can be further distinguished from L. magnum (100 - 200µ) by the latter's posterior truncation with shorter caudal cilia (one-ninth of the body length). Otherwise the two organisms resemble each other in having a longitudinally striated cuticular surface, and an anteriorly projecting acumination. They both differ from L. bullinum and L. lucens in that these exhibit a transverse stripe all through the cuticular surface and their anterior extremities do not slant to the right.

The description in Kahl's key (Kahl 1930, p. 327) is as follows: 1(2) Pell. nur längsgestreift, nicht gefeldert. Vorderende schräg nach 1. abfallend, Hinterende verjüngt, mit einem Kamm kurzer Caudalwimpern.

Lembadion (Hymenostoma) magnum Stokes, 1887).

- 2(1) Pell. durch Querrippen gefeldert. Vorderende nicht stark nach 1. abgeschrägt. Hinterende breit gerundet, mit einem Büschel langer Caudalwimpern.
- 3(4) Grössere Art über 100µ. Vorn mit deutlicher Einkerbung. Dors. mit 30-35 Längsreihen.

Lembadion bullinum Perty, 1852 (Hymenostoma hymenophora Stokes, 1884).

4(3) Kleinere Art unter 100µ. Die vordere schwache Einkerbung wird durch die Spitze der undulierenden Mbr. verdeckt, so dass das Infusor vorn zugespitzt erscheint. Dors. ca. 15 Längsreihen.

Lembadion (Thurophora) lucens (Maskell, 1887)

Body size is the second criterion used by Kahl (1930) to classify the species of Dembadion. In this respect, Hymenostoma hymenophora Stokes, 1884 should not be the synonym of L. bullinum Perty, 1852. as Kahl considered. Stokes (1884, in Stokes, 1888) pointed out that H. hymenophora differs from other Lembadion in its two contractile vacuoles, the abruptly narrowing adoral membrane, the sinistrally directed caudal cilia and 1/500 inch (50.8µ) body length. In accordance with the International Code of Zoological Nomenclature, H. hymenophora should stand as Lembadion hymenophora (Stokes, 1884), not as a synonym of L. bullinum Perty, 1852 - the former is much the smaller and has distinctive characteristics as outlined.

The species under discussion resembles L. hymenophora in its shape, possession of two contractile vacuoles and its sinistrally directed caudal cilia. However, it differs from the latter mainly in its longitudinally striated cuticular surface, the location of its macronucleus, and its shorter, abruptly narrowing right adoral membrane. Lacking infraciliary data to compare with the descriptions of Dragesco and Tuffrau (1963), the present species should be further studied before its taxonomic allocation is determined.

Subclass	Holotricha			
Order	Hymenostomatida			
Suborder	Pleuronematina			
· ·	• • • •			

Family Pleuronematidae Genus Cristigera

Cristigera sp. Plate XXVI, Figs. 74 - 77.

Small, elliptical (20 x 10µ, Figs. 74 - 77), compressed. Somatic cilia long and sparse (about 10µ); posterior plate without hollow and lacking of cilia except for a single conspicuous caudal cilium (about 20µ, Figs. 74 - 77). Oral aperture close to mid-ventral line with a postoral depression, peristomal furrow continuing back of cytostome to posterior extremity (Figs. 76 - 77). Undulating membrane' on right margin of pre-oral groove forming a pocket around rear cytostome. A single contractile vacuole (Figs. 76, 77), posteriorly

Systematic account:

situated.

The dorsal side of present organism (Fig. 74) is very similar to *Cyclidium*, but it differs from the latter in its dorso-ventral compressed shape and oral aperture with a postoral depression and peristomal furrow continuing back of cytostome to posterior extremity (Figs. 75 - 77). Thus, it fits the description of *Cristigera* Roux, 1901.

PLATE XXVI

89

Figs. 74 - 77. Cristigera sp. (phase-contrast)

Fig. 74. Showing dorsal aspect.

Fig. 75. Showing body ciliation.

÷.,

Fig. 76. Focused on long single caudal cilium.

Fig. 77. Showing oral aperture, peristome and single contractile Br

vacuole.



Family Pleuronematidae

Genus Cyclidium (according to Corliss, 1961)

Cyclidium glaucoma O. F. M., 1773, Plate XXVII, Figs. 78-83.

Length, 17.5 - 25µ; breadth, 10 - 17.5µ; average size, 20 x 12.6µ. Form constant and well-defined; ovoid; circular in cross-section (Fig. 79). Somatic cilia long and sparse, their length equalling or exceeding that of the breadth of the body (Figs. 78 - 83), disposed in even longitudinal lines (Fig. 81). Posterior plate without hollow and lacking cilia except for a single conspicuous caudal cilium, twice at least as long as those of the general surface (Figs. 78 and 80). Ora1 aperture ventral, dominated by the conspicuous undulating membrane, which margins the body and is not thrown into folds. Adoral zone of membranelles hardly recognizable in fresh material. No vestibulum, peristome more or less subequatorial (Figs. 78 and 80). No postoral longitudinal groove. A more or less pronounced area of thigmotactic ciliature is located dorsally (Fig. 81). Macronucleus spheroidal and central (Fig. 78), with an adjacent micronucleus. A single contractile vacuole, posteriorly situated. Cortical layer without trichocysts Examples undergoing transverse binnary fission were commonly observed (Figs. 82 and 83). Free swimming., Under adverse conditions, cysts

are formed after the disappearance of the cilia. Food: bacteria (Fig. 80). While feeding, the organism

remains in one position with spread cilia and beating membranelles.

PLATE XXVII

Figs. 78 - 83. Cyclidium glaucoma O. F. M., 1773, living

examples by phase-contrast microscopy.

Fig. 78. General appearance.

Fig. 79. Equatorial view in apparent transverse section.

Fig. 80. A stage before cyst formation.

Fig. 81. Ciliation.

Fig. 82. Early binary fission.

Fig. 83. Transverse binary fission completed, the filial cells almost separated.

1. Undulating membrane; 2, Macronucleus; 3. Food vacuole;

4. Contractile vacuole; 5. Somatic cilium; 6. Caudal cilium;

7. Indicates the ribs of the undulating membrane complex;

8. Bacteria; 9. Ventral side; 10. Dorsal side.



Systematic account:

The small hymenostome under consideration is easily distinguished from six of the seven genera of Pleuronematidae. Thus Calyptotricha is loricate; Pleuronema is set apart by its giant size (70 - 180μ), and the presence of a semicircular swelling to the left of the peristome; Pleurocoptes is a marine ectocommensal; Ctedoctema has an oblique peristome; the peristome of Cristigera is closer to the midventral line, and there is a postoral depression; while Histiobalantium has a pre-oral area about one quarter of the body length. Those characteristics are absent from the present species, which fits the description of the remaining genus, Cyclidium, in being ovoid and small (15 - 60µ), with an unciliated and truncate frontal plate anteriorly. Its peristome extends backwards from the anterior extremity for up to three-fifths of the length of the body, the cuticular surface is longitudinally ciliated, the somatic cilia are long and sparse, and there is a single caudal cilium. The right peristomial margin has a membrane which forms a pocket around the cytostomal groove, and the left margin bears either free cilia or a membrane which unites with that on the right. There is no semicircular swelling to the left of oral region, a round macronucleus and prominent micronucleus are present, and there is a posterior contractile vacuole.

Kahl (1930) recognized 30 species of Cyclidium. The present one differs from C. brandon, C. paucisetum, C. paradoxum, C. curvatum and C. helgolandicum in having a single caudal cilium. C. litomesum, C. libellus, C. oligotrichum, C. heptatrichum and C. veliferum differ from the Logy Bay species through the reduction of the medial ciliation.

The contractile vacuole of C. mucicola and C. centrale is centrally located (the present species has a contractile vacuole posteriorly). C. opisthostoma, C. lanuginosum and C. flagellatum do not merit consideration by reason of their reduced posterior ciliation. C. obliguum has distinctively oblique ciliation, while C. terricola has cilia on the anterior extremity and an altogether unique shape¹. The long, sparse somatic cilia and the unciliated posterior extremity (save for a single caudal cilium) narrow the choice down to three species, C. glaucoma, C. citrullus and C. elongatum, the remaining ten species² being delicate with short cilia (one half to two-thirds of body C. citrullus is characteristically spindle shaped, C. breadth). elongatum, elongate-ellipsoidal, and C. glaucoma broadly ovoid. However, the total composite of criteria differentiating these protozoa, and the overlap of their sizes and shapes, are such that it is difficult to separate them from one another with complete confidence. terms of body shape, the criteria are so vague that some investigators, e.g. Schewiakoff (Kahl, 1930, p. 376) failed to differentiate them individually. Nevertheless, the present species is distinguishable from C. citrullus not only by body shape but by its smaller size (C. citrullus, 20 - 30µ) and the absence of an invagination at the point of insertion of the caudal cilium.

The description of Kahl's key (Kahl, 1930, p.375) is as follows: 8(9) Moosform ohne abgesetzte Frontalplatte, mit kurzen, derben Trc. (C. terricola).

C. fuscum, C. oblongum,C. singulare, C. gemmuliferum, C. granulosum, C. candens, C. simulans, C. pellucidum, C. instabile, and C. versatile.

Sometimes, the present species is so elongate that it appears closer to the elliptical C. elongatum than the characteristically ovoid C. glaucoma. In addition, its small size $(17.5 - 24\mu, average 20\mu)$ recalls C. elongatum $(16 - 24\mu, never reaching the 25 - 30\mu$ of C. glaucoma - Kahl, 1930). Therefore, its identification as C. elongatum might be understandable. Nevertheless, it is more often ovoid than elliptical. Also, while its peristome is more of less subequatorial, it does not occupy more than three-fifths of the body length. This is held to be the critical feature distinguishing it from C. elongatum, the peristome of which is about three-quarters of the overall length. As regards the body size, many other investigators besides Kahl (1930) have reported that C. glaucoma ranges from 14.7 to 26.1 μ in length (Bhatia, 1936; Berger and Thompson, 1960). It might be pointed out, too, that C. elongatum Schewlakoff was originally considered to be a variety of C. glaucoma (Schewlakoff, 1896).

In sum, the present species is identified as *C. glaucoma* O. F. Müller on the basis of its single caudal cilium, the distribution of its ciliary vestiture, the location of the contractile vacuole, the lack of a posterior invagination and the relative length of its peristome (i.e. up to three-fifths of the overall length).

Subclass	Peri tricha
0rder	Peritrichida
Suborder	Şessilina

Vorticellidae

Family

Genus

s Vorticella, Plates XXVIII - XXXI, Figs. 84 - 94.

Generally, inverted bell form, colorless or yellowish; peristome more or less outwardly extended (Figs. 87-89 and 91-94); colonial or solitary with a contractile stalk (Figs. 84 and 85) attached to submerged objects and aquatic plants (Figs. 84 and 85) or animals; one to two contractile vacuoles (Figs. 87-92). Unbranched stalk with well-developed spasmoneme, thrown into a spiral coil when contractile (Figs. 87-92). Numerous species.

Systematic account:

This group of ciliates (Figs. 84-94) with peristome circular, bordered by membranes running counter-clockwise around it and into a vestibule containing cytostome, cýtopyge, and contractile vacuole pore; body ciliation greatly reduced; all characteristics of Peritrichida.

The sessile form clearly remands this group to Sessilina. The contractile stalk attached to the posterior extremity of the body without lorica, assigns this group to Vorticellidae.

Furthermore, the unbranched stalk with spasmoneme welldeveloped and thrown into a spiral coil when contractile, separates the present group from *Intranstylum* with poorly-developed spasmoneme and from *Zoothamnium* and *Carchesium* with branched stalks.

PLATE XXVIII

Figs. 84 - 86. Vorticella sp.

Figs. 85 & 85. Showing each solitary Vorticella sp. with an unbranched contractile stalk attached to submerged objects.

96

Fig. 85. Vorticella microstoma Ehrenberg, 1830, with

transverse striation.



Thus this group is referable to *Vorticella* Linnaeus. There are more than 220 described species of *Vorticella* (Noland, 1931). The taxonomic data of the four groups from Logy Bay are briefly recorded

as follows:

Vorticella microstoma Ehrenberg 1830, Plate XXVIII, Fig. 8b. Body conical (23 x 20µ) with stalk (18µ); transversely striate. Less twice as long as broad. Peristome narrower than half the body-centre.

Vorticella sp. Plate XXIX, Figs. 87-89.

Surface smooth, broadly campanulate (50 x 34 μ). Width of peristome (34 μ) less than body length. Greenish, stalk (142 μ) more than three times as long as the body.

Vorticella nebulifera O. F. M. 1786, Plate XXXI, Figs. 93,94. var. similis (Stokes, 1887).

Smooth body conical (78 x 48µ), occurring in groups. Translucent, flexible body with granular greyish-white colour bands as shown in Fig. 94 and twists on its stalk in graceful attitudes. Peristome oblique, ciliary disc not elevated while body extended (Fig. 94). The contracted form (Fig. 93) is smoothly ovoid.

PLATE XXIX

Figs. 87 - 89. Vorticella sp. I.

Fig. 87. Showing an enlarged disc-like anterior region with prominent adoral ciliature which winds counter-clockwise to the cytostome in living material, (phase-contrast).

Figs. 88 & 89. Showing smooth surface and broadly companulate shape.

(Nomarski interference-contrast effect).



PLATE XXX -

Figs. 90 - 92. Vorticella sp. 2.

Fig. 90. Showing contractile form.

Fig. 9]. Showing single contractile vacuole centrally located.

" Fig. 92. Showing extended form.



PLATE XXXI

.100

41.

Figs. 93 and 94. Vorticella nebulifera O. F. M. 1786, var. similis (Stokes, 1887).

Fig. 93. Focused on contractile form.

Fig. 94. Showing peristome oblique ciliary disc not elevated while body extended.



Vorticella sp. Plate XXX, Figs. 90 - 92.

Ovoid body (60 x 43μ) with unbranched stalk (70 μ). Transversely striated; single contractile vacuole conspicuously centrally located. Peristome not elevated while body extended (Fig. 92).

Family Epistylidae Genus Epistylis

Epistylis sp. Plates XXXII - XXXIII, Figs. 95-100.

Body longitudinally contractile, fully expanded (300 x 30μ , Fig. 95), contractile (110 x 66μ , Fig. 98). Individuals always on dichotomous non-contractile stalk (Figs. 95 and 97), forming large colonies; pristomial ring prominent; flat cap makes a slight angle with the ring (Fig. 96), gullet with ciliated wall, body full of zoochlorellae(Figs. 99, 100).

Systematic account:

There are six fresh-water genera of Epistylidae (Noland in Edmondson, 1959) which bear comparison with the present organism in having uncontractile stalks

The main feature of its dichotomous non-contractile stalk with one individual at end of each branch of the stalks (Figs. 95 and 97) separates the present organism from *Systylis*; the latter bears clusters of several dozen individuals at the end of each branch.

The peristomial membranes make only a little more than one

PLATE XXXII

102

Figs: 95 and 96. Epistylis sp.

Fig. 95. Showing an expanded and a contracted individual on a

. dichotomous stalk.

Fig. 96. Showing peristomial ring at a slight angle with flat cap.


PLATE XXXIII

103

Figs. 97 - 100. Epistylis sp.

Fig. 97. Showing one individual at end of each branch of the dichotomous non-contractile stalk.

Fig. 98. Showing fully contracted form ...

5

Figs. 99 & 100. Focused on zoochlorellae in body....



turn around the peristome of the present organism, distinguishable from the peristomial membranes that make four to six turns around the peristome of *Campanella*. The remaining three genera *Puxidium*, *Opercularia*, and *Rhabdostyla* with deep peristomial furrow, separating disc and border, bears no comparison with the present organism which has a shallow peristomial furrow, and a disc not set off from border by a deep incision. The data given above fit the description of *Epistylis* Ehrenberg, 1838. Thus, it is referred to *Epistylis*.

104

Family Astylozoonidae Genus Telotrochidium

Telotrochidium sp. Plate XXXIV, Figs. 101-106.

Entirely free-swimming body dontractile from extended campanulate (106 x 60u, Fig. 101) to globose shape (80 x 70u, Fig. 106) without caudal appendage. Two ciliary girdles without body ciliation: one developed at a short distance from the anterior and another from posterior extremity; the anterior one with two parallel rows of adoral zone; cytopyge postero-terminal, tubular, permanently visible (Figs. 102-104). Three contractile vacuoles connected with the cytopharynx (Fig. 101), the largest one subcentral (Figs. 101-103 and 105-106). Macronucleus "L" shape band form (Fig. 102-103 and 105). Movement swift, rotating in alternate directions.

Systematic account:

The free-swimming form, with peristome-bearing extremity forward and without body ciliation, are the characteristics of.

The species under consideration is distinguished from Astylozoon by the latter's conical shape and aboral extremity attenuated, with one or two thigmotactic stiff cilia; from Hastatella by the latter with two to four rings of long conical ectoplasmic processes.

The rounded extremities of the present species without any body ciliation or processes but two ciliary girdles are the main features of *Telotrochidium* which consists of *Telotrochidium crateriforme*.

105

PLATE XXXIV

Figs. 101 - 106. Telotrochidium sp.

Showing a series of body contractile from extended campanulate (Fig. 101) to globose (Fig. 106). The arrows in Figs. 102-104 point to the second, pos-

. 106

terior girdle.



Kent (1881-1882) and Telotrochidium (Opisthonecta) henneguyi (Fauré-Fremiet). Both of them bear a "C" shape sausage form macronucleus which is different from the present species. Thus the "L" shape band form macronucleus suggests the present species as a new member of the genus. However, Dodd (1962) observed some nuclear changes in cysts of *Telotrochidium henneguyi*. Therefore, the species identification of the present organism remains for further investigation.

Subclass	Spirotricha
Order	Heterotrichida
Suborder	Heterotrichina

Spirostomatidae

Blepharisma

Family Genus

Blepharisma persicinum Perty, 1852, Plate XXXV, Figs:107-110.

108

Pyriform (141 x 66µ), somewhat narrowed anteriorly and posterior extremity acuminate; compressed; left peristomal edge sigmoid (Figs. 109 and 110), and is twisted to right at posterior extremity and connected to oral funnel with membrane; in front of cytostome a large well-developed two-layer undulating membrane on right edge (Figs. 107 and 108); peristome about three quarters of body length; ciliary rows longitudinal; ciliation dense; contracile vacuole and cytopyge terminal; macronucleus in six patts; rose coloured.

Systematic account:

Kahl (1932) recognized 19 species of *Blephanisma*. Only five of them with multimacronuclei bear comparison with the present species. Among these five, the present species is distinguishable from *B*. *dileptus*, *B. clarissiumum*, and *B. salinarum* by its rose-coloured smaller (less than 150µ) but rather stumpy body.

This narrows the choice down to *B. musculus* and *B. persicinum.* The former can be easily distinguished from the present species by its more numerous (7-10) macronuclear parts.

PLATE XXXV

Figs. 107 - 110. Blepharisma persicinum Perty, 1852.

Figs. 107 and 108. Showing right edge of cytostome with a large welldeveloped two-layer undulating membrane. Macro-

nucleus in six parts.

Figs. 109 and 110. Showing sigmoid left peristomal edge; contractile vacuole and cytopyge posteriorly, indicated by the arrow in figure 109.

109



The present data fit the descriptions of *B. persicinum* Perty ally which is characteristic pyriform in shape (80 - 120µ long) with accuminated posterior extremity, the range of peristome from one third to three quarters of body length with sigmoid left edge, well-developed undulating membrane on right, macronucleus in three to six parts, and rose-coloured body. The present species is thus referred to *Blepharisma*

persicinum Perty.

Family Spirostomatidae

nus Spirostomum (according to Corliss, 1961)

Spirostomum teres Claparède and Lachmann, 1859, Plate XXXVI, Figs. 111a, 111b, 112.

Body measuring 200 - 500µ by 20 - 354 Slender-elongate, worm-like, 10 - 15 times as long as broad, cylindrical, one side parallel to the other, sometimes slightly swollen medially, rounded anteriorly, usually truncated posteriorly (Figs. IIIa, 111b). Ectoplasm with highly developed and contractile myonemes, arranged lengthwise independent of ciliary rows. Endoplasm clear and transparent. Peristome narrow, occupying about one third of the body, closely lined with a band of well developed cilia or membranelles. Cytostome at the base of the peristome (Fig. 111), leading immediately into a cytopharynx which is very simple and difficult to distinguish. Elliptical macronucleus (Fig. 111) often centrally located below the base of the. peristome,-clearly evident in pressure-immobilized live examples. Micronucleus not observed. Excretory vacuole large and terminal, with Figs. 111 and 112. Spirostomum teres Claparède and Lachmann, 1959. Unstained living ciliate as seen by Nomarski interference-contrast effect.

PLATE XXXVI

Fig. 111a. Ma. oval macronucleus, unlike that of other common members of Spirostomum (except S. filum), which have a moniliform macronucleus. (S) cytostome located at the base of the peristome which extends for one-third of the body length (that of S. filum is only one-fourth of the body length).

111

Fig. 111b. P. truncated posteriorly, a single contractile vacuole occupying the whole of this part of the body and extending forwards as a long canal (C) (S. filum has a tapering, drawn out posterior end).

Fig. 112.

Showing transverse binary fission completed, two filial cells almost separated.



a long dorsal canal (Fig. 111a, 111b). Transverse binary fission (Fig. 112)

Systematic position:

Kahl (1932) recognized seven species of Spirostomum (S. ambiguum, S. minus, S. intermedium, S. inflatum, S. loxodes, S. filum and S. teres). Only in S. teres and S. filum is the macronucleus oval or elliptical in shape. In the other five species, this structure is moniliform. An unidentified but possibly new species referred to by Seshachar and Padmavathi, 1956, has a cylindrical and elongate macronucleus in the vegetative stage. During binary fission, though, this organelle becomes condensed into an oval or polymorphic mass. The macronucleus is always oval in S. teres, and may be clearly observed "either with or without staining in the living animal. S. teres (200 500 μ) is generally longer than S. filum (200 - 300 μ). Anteriorly rounded and posteriorly truncated, S. teres is characterized by one body side parallel to the other. In S. filum, though, there is a tapering, drawn-out posterior extremity. The peristome is about, or more than, one-third of the body length in S. teres, but only onefourth the body length in S. filum. Therefore, the chief features distinguishing S. téres from other members of the genus are the oval shape of the macronucleus, the parallel-sided body shape and the elongate peristome.

Subclass Spirotricha Order Oligotrichida

Family Halteriidae

Halteria

Genus

Halteria sp. Plate XXXVII, Figs. 113 and 114.

Body turbinate (av. 25 x 20 μ ; Figs. 113 and 114); anterior border bears conspicuous adoral zone, unclosed adoral membranelles well-developed; a small membrane on right edge and cirri (Fig. 114,A) on left. Macronucleus oval; a micronucleus. Contractile vacuole left of cytostome. A zone of long cirri or bristles (Fig. 114,B) develops around the equatorial region, the sudden flexure of which appendages, enable the body to progress through the water by a series of leaping movements, in addition to their ordinary swimming motions.

Systematic account:

The present ciliate is easily recognized as *Halteria* by its peculiar movements in the water, which consist of a slow rolling or rotary motion, interrupted at short intervals by a sudden leap backwards or to one side and its turbinate body with equatorial zone of bristles.

However, the present organism clearly bears the main features of Oligotrichida, e.g. the greatly reduced body ciliation and welldeveloped adoral membranelles extended beyond the anterior extremity. There are only two families of Oligatrichida, Halteriidae and Strobiliidae (Kudo, 1966). The present organism is clearly

PLATE XXXVII

114

Figs. 113 and 114. Halteria sp.

Fig. 113. Showing turbinate shape with body ciliation.

Fig. 114. Showing unclosed well-developed membranelles and cirri (A)

around cytostome and a tuft of bristles (B) around

equatorial zone.



distinguishable from Strobiliidae because the latter's cytostome is encircled by adoral membranelles (see Noland in Edmondson, 1966) and allocated to Halteriidae for bearing unclosed adoral membranelles.

Among the three genera of Halteriidae, the present organism is well separated from *Tontonia*, for the latter has well-developed apical collar and a long dytoplasmic (contractile) caudal process; and from *Strombidium* which bears no body bristles or cirri. The organism under consideration bears the generic features of *Halteria* - the greatly reduced body ciliation except equatorial zone of body bristles, the well-developed unclosed adoral membranelles around cytostome. Thus the ciliate under discussion is therefore referable \cdot to *Halteria* Dujardin.

Family Strobilidiidae

Genus Strobilidium

Strobilidium sp. Plate XXXVIII, Figs. 115, 116.

Body turnip-shaped (av. 47 x 34µ, Figs. 115 and 116); adoral membranelles form a spiral crown at the anterior extremity (Fig. 116) without cytopharynx and lateral border with rounded elevation anteriorly; the posterior extremity tapering and truncate; cuticular surface smooth, except at the posterior region (Fig. 115), where there are a few longitudinal ridges which often extend slightly beyond the termination of the body. Horseshoe shaped macronucleus anterior; a micronucleus; a contractile vacuole.

PLATE XXXVIII

116

Figs. 115 and 116. Strobilidium sp.

Fig. 115. Showing truncate posterior extremity and closed adoral membranelles.

Fig. 116. Showing a spiral crown at anterior extremity with rounded .elevation laterally.



Systematic account:

Similar to Halteriidae, the present organism also bears the characteristics of Oligotrichida, the greatly reduced body ciliation nearly absent and the well-developed ador membranelles extended beyond the anterior extremity of the body. However, it is distinguishable from Halteriidae by its closed adoral membranelles form a spiral crown at the anterior extremity while Halteriidae with unclosed adoral membranelles. These discussed criteria fit exactly the description of Strobilidiidae (Kahl, 1930-1935).

There is only one genus of Strobilidiidae, Strobilidium Schewiakoff (see Noland in Edmondson, 1966; Kudo, 1966) of which the present organism bears the generic features - turnip body shape, cytostome at anterior extremity without cytopharynx, with horshshoeshaped macronucleus.

Lateral border-with rounded elevation near anterior extremity, posterior extremity truncate of the present species strongly suggest allocation to *Strombilidium gyrans* Stokes (1885). However, no detailed comparison with other members of *Strombilidium* were made, therefore the species under discussion is left as *Strcmbilidium* sp.

Spirotricha Hypotrichida

Family Aspidicidae

Subclass

Order

Genus Aspidisca

Aspidisca sp. Plate XXXIX, Figs. 117 - 120.

Small (29 x 25µ); ovoid (Figs. 117-120); inflexible; right and dorsal side convex (Figs. 117 and 118), ventral side flattened (Figs. 119 and 120); dorsal surface conspicuously ridged with spines (Fig. 118); adoral zone poorly developed, cirri reduced in number and are limited to frontals and anals; macronucleus horseshoe-shaped; single contractile vacuole; benthic species.

Systematic account:

Reduced zone of adoral membranelles of the present organism differentiates it from all other families of Hypotrichida in that the latter bear well developed conspicuous adoral membranelles. The ovoid body with rudimentary adoral membranelles and seven frontal-ventral, five to twelve anal cirri fits the characteristics of Aspidisca Ehrenberg,

1830.

PLATE XXXIX

119

Figs. 117 - 120. Aspidisca sp.

Figs. 117 and 118. Showing ovoid shape; right and dorsal side convex; spines on ridged dorsal surface.

Figs. 119 and 120. Showing flattened ventral side with poorly developed adoral zone and reduced cirri.

10



The following genera and species were also recovered in samples from the study pond at Logy Bay. The ecological data for each are presented in Table II but illustrations are not included. Systematic positions of the different groups are presented at the beginning of the systematic section.

Chilophrya utahensis Pack, 1919.

Body tube-shaped (56 x $10 - 84 \times 15\mu$); cytostome anteriorly surrounded by protrusible rods; on one side there is a flap process; macronucleus small, central; contractile vacuole terminal; endoplasm with zoochlorellae

Prorodon discolor (Ehrenberg, 1838).

Ovoidal (105 - 130µ x 70 - 85µ); gullet trichites double, conspicuous, their external ends slightly below surface level; contractile vacuole terminal; macronucleus ellipsoid; micronucleus hemispherical.

Microregma sp

Small (23 - 27µ in length), ovoid; dorsal side convex; ventral side flat, with a small slit-like cytostome near anterior extremity, surrounded by longer cilia; single long caudal cilium; contractile vacuole posteriorly.

Matyophrya.sp.

Compressed; elongate ovoid (22.4 - 33µ in length); asymmetrical; dorsal surface convex, ventral surface partly concave; uniform ciliation with spiral striation; cytostome anteriorly; macronucleus round; single contractile vacuole posteriorly.

Trachelocerca sp.

Elongate, flask-shaped; more or less extensible, with drawnout anterior extremity; without any ring-furrow anteriorly; when contracted (84µ in length) pellicular striae not spiral; cytostome anteriorly, surrounded by a ridge containing short trichocysts, cytopharynx with trichocysts; nucleus contains peculiar crystal-like bodies; body full of green coloured zoochlorellae; when extended 120µ in length; single contractile vacuole posteriorly located.

Litonotus, sp.

Elongate flask-shape (48µ in length); with flattened neck and tail, both of which are moderately contractile; neck stout, bent towards the dorsal side; cytostome a long slit (two-fifths of body length), anteriorly located; posterior extremity bluntly rounded; one terminal or several (in one row) contractile vacuoles; two spherical macronuclei between which a micronucleus is located.

Pysteria sp.

Ovate (50 x 30µ); dorsal convex, ventral concave; left ventral non-ciliated; post-oral ciliation is continuation of pre-oral to right of cytostome and parallel to right margin; cytostome in a furrow near right side; posterior stylus conspicuous; two contractile vacuole centrally located.

·Paratrochilia chilodontoides Kahl, 1933.

Rounded anteriorly, narrowed posteriorly $(30 - 35 \times 20 - 22\mu)$; right_side_more_convex_than_left_side; cytostome_anterior_with_cytopharynx and pre-oral membrane; the broad side of funnel-shaped cytopharynx anteriorly; on the left edge two to three bulges; one hem of cilia surrounded the dorsal knoll, otherwise unciliated dorsal surface; two contractile vacuoles in the right side of ventral surface.

Chilodonella uncinata (Ehrenberg, 1838).

Ovoid (48 - 67µ long); dorso-ventrally flattened; dorsal surface convex, ventral surface flat; about eleven ventral ciliary rows; anteriorly flattened dorsal surface with a cross-row of bristles; cytostome round; oral basket conspicuous, protrusible; macronucleus rounded; contractile vacuoles variable in number.

Nassula sp

Ovate (160 x 83µ); ventral flat, dorsal convex; cytostome one-quarter from anterior extremity; body bent to left near cytostome; opening of oral basket deep, in a vestibule with a membrane; macronucleus spherical, central; contractile vacuole large, with accessory vacuoles and opens ventrally through a tubule-pore.

Colpoda steini Maupas, 1883.

Small reniform (15 - 42μ); cytostome about two-fifths from the anterior extremity, and with a bundle of long membranellae; five to six pre-oral ridges; paired and single cilia; one pair of long caudal, cilia; twelve meridians.

Cohnilembus sp.

Slender spindle form (50µ long); peristome from anterior extremity to the middle of body, curved to right, with two membranes on right edge; a few longer caudal cilia; macronucleus oval, central; contractile vacuole posterior.

Uronema sp.

Elongate oval (45µ long); slightly flattened; non-ciliated anteriorly; inconspicuous peristome with ciliated left edge; cytostome in the anterior half, with a small tongue-like membrane; cytopharynx indistinct; macronucleus spherical, central; contractile vacuole terminal.

Colpidium sp.

Elongate reniform $(42 - 70\mu \text{ long})$; 55 - 60 ciliary meridians; triangular cytostome about one-tenth the body length from anterior extremity towards right side; a small ectoplasmic flange along right border of cytostome which shows an undulating membrane on right and three membranellae on left; pre-oral suture curved to left; macronucleus oval, central.

Balanonema biceps (Penard, 1922).

Body ellipsoid (40 - 48µ long); no cilia in the middle region; but with plug-like extremities; cytostome inconspicuous; single long caudal cilium; oval macronucleus posterior to central contractile vacuole.

~ Cinetochilum marinum Kahl, 1930.

Oval (18 - 22µ long); highly flattened; cilia on flat ventral surface only; cytostome right of median line in posterior half, with a membrane on both edges which form a pocket; oblique non-ciliated postoral field leads to left posterior extremity; four caudal cilia; macronucleus spherical, central; contractile vacuole terminal. Frontonia leucas Ehrenberg, 1838.

Ellipsoid (200 - 250µ long); anterior extremity more broadly rounded than posterior extremity; flattened; oral groove in anterior third ventral surface; cytostome with a complex organization (see Kudo, p. 907); a long narrow post-oral groove which is ordinarily nearly closed; cytopharynx with numerous strong fibrils; ciliary rows close and uniform; ectoplasm with numerous trichocysts; feeds on filamentous algae.

. Cyclidium citrullus_Cohn, 1865.

Spindle shape $(20 - 30\mu)$; peristome about two-thirds of body length, dominated by the conspicuous undulating membrane. Spherical macronucleus in the anterior half. Otherwise, very similar to *Cyclidium glaucoma*, 0. F. M.

Cyclidium elongatum Schewiakoff, 1896.

Elongate ellipsoid (16 - 24μ); peristome about three-quarters of body length. Otherwise, very similar to *Cyclidium glaucoma*, O. F. M.

Cyclidium granulosum Kahl, 1931.

More pyriform $(36 - 40\mu)$ than elongate-oval; other characteristics fit the descriptions of Kahl (1931).

125

Cyclidium litomesum Stokes, 1884.

Cylindrical (40µ), the length about two and one-half times the width, the extremities subequally rounded. Cytostome situated slightly behind the centre of the ventral surface. No ciliation in the middle dorsal part of body. The cilia posteriorly situated diverse in length, some of them longer than body length.

Cyclidium muscicola Kahl, 1931.

Small, moss shape (14 - 17μ), similar to Cyclidium glaucoma, 0. F. M., but contractile vacuole in the middle of the body.

Pleuronema marinum Dujardin, 1841.

Elongate-ovoid (62 - 75µ long); peristome in two-thirds body length.anteriorly; a conspicuous membrane at both edges; semicircular swelling to left near oral area; no cytopharynx; close striation longitudinal; single long caudal cilium; trichocysts distinct; macronucleus round; a contractile vacuole.

Ophrydium sp.

Cylindrical (200 - 300µ long) with a contractile neck; posterior end pointed; variable number of individuals in a common mucilaginous mass; pellicle usually cross-striated; with many zoochlorellae.

Metopus striatus McMurrich, 1884.

Body form changeable; when extended oblong or fusiform (64-85µ long); peristome conspicuous, slightly spirally diagonal, beginning at the anterior end and reaching the middle of body; when contracted, peristome much spirally coiled; cytopharynx short; body ciliation uniform, longitudinal spiral; long cilia at extremities; conspicuous contractile vacuole termin**()**; macronucleus oval.

~ Condylostoma sp.

Ellipsoid (132 - 143µ long); truncate anteriorly, rounded posteriorly; slightly flattened; peristome wide anteriorly and Vshaped, peristomal field not ciliated; a large membrane on right edge and adoral zone on left; macronucleus moniliform; several contractile vacuoles with canal; cytopyge posterior.

Paraclevelandia sp.

Oval (22 - 27µ long); body rigid; posterior extremity truncated obliquely to left; peristome posterior without projection; one macronucleus and one micronucleus.

Gonostomum sp.

Flattened, elongate (118 - 127µ long); flexible; eight frontals; one or two oblique ventral rows of short cirri; four or five

anals; two marginal rows; adoral zone well-developed.

Steinia sp

Rounded ellipsoid (148 - $172\mu \ge 80 - 140\mu$); rounded both extremity; right border of peristome curves left, or spirals into a pit in peristome; few frontal cirri, usually eight in three groups; no continuous long rows of cirri; two macronuclei; one contractile vacuale in the middle.

Oxytricha sp.

Ellipsoid (100 - 140µ long); flexible; ventral surface flattened; dorsal surface convex; eight frontals; five ventrals; five anals; short caudals; macronucleus in two parts.

Stylonychia sp.

Ovoid to reniform (100 - 126µ long); not flexible; ventral surface flat; dorsal surface convex; eight frontals, five ventrals; five anals; marginals; three caudals, with short dorsal bristles.

Uroleptus longicaudatus Stokes, 1886.

Elongate body (about 200µ long) drawn out into a tail-like portion; three frontals; two to four rows of ventral cirri; marginals; no anals; sometimes rose or violet coloured. Urosoma sp.

Similar to Oxytricha; but posterior portion drawn out and very narrowed; 160 - 175µ long.

CONCLUSIONS.

"Protozoa inhabit microhabitats, Each microhabitat is an ecological community of producers and consumers with definite relations to each other (Picken, 1937)" said Bamforth (1963). In consideration of the relations between producers and consumers, Felton et al. (1967) suggested that bacteria in the pond ecosystem play a role in the nitrogen, carbon and energy cycles as decomposers and transformers, as a source of nutrients and as members of the food chain. However, "A complete picture of the role of micro-organisms in the natural history of the pond must await further data, particularly about fungi and protozoa" (Felton et al. 1967).

As is well known, ciliated Protozoa may be found wherever there is moisture without deleterious substances. The same species may often be found in both littoral marine waters and inland freshwater lakes, ponds, and pools. Although many species are cosmopolitan, they tend to accumulate in certain places where the environments best suit their needs. Hence, it is necessary to know their natural distribution as well as their ecological parameters.

The pond under study is highly acidic (pH values between 4.50 - 5.15). It also has a high chloride content (99.26 p.p.m.), and a wide annual water temperature range (0.5 - 20.8°C), besides a fluctuation of dissolved solid content (75 - 144 p.p.m.). During the period of regular sampling from June 1969 to September 1970, it dried completely in July 1970, and thus can be defined as a "semi-permanent" pond (Laird, 1956).

â
For free-living ciliates, suitable culture media would

supply sufficient population for their infraciliary analysis. Lacking complete descriptions of infraciliature, many of the taxa presented herein need further study in order to allocate their systematic status.

Forty-five genera of Ciliophora were reported which include many cosmopolitan species.

REFERENCES

AMERIÇAN PUBLIC HEALTH ASSOCIATION. 1955. Standard methods for the examination of water, sewage, and industrial wastes. 10th ed., New York, 522 pp.

BAMFORTH, Ś. S. 1963. Limnetić protozoa of Southeastern Louisiana. Louisiana Academy Sci. XXVI: 120-134.

BATES, M. 1949. The natural history of mosquitoes. New York, Macmillan, XV: 379 pp.

BERGER, J. and J. C. THOMPSON. 1960. A redescription of Cyclidium glaucoma O. F. M., 1786 (Ciliata: Hymenostomatida), with particular attention to the buccal apparatus, J. Protozool. 7: 256-262.

BHATIA, B. L. 1936. Protozoa: Ciliophora. In Sewell, R. B. S. editor, The fauna of British India, Including Ceylon and Burma, pp. 1-493. Taylor and Francis, London.
BOGGS, N. 1965. Comparative studies on Spirostomum: silver impregnation of three species. J. Protozool. 12: 603-606.
BROWNING, I. 1951. Cytoplasmic inclusions of the protozoan, Tetrahymena goleri, Texas Reps. Biol. Med. 9: 3-7.

BUHSE, H. E. 1962. Control of macrostome production in Tetrahymena vorax. J. Protozool. 9, Suppl. 5.

BUHSE, H. E. 1966b. Oral morphogenesis during transformation from microstome to macrostome to microstome in *Tetrohymenaevorax* stain V₂ type S. Trans. Am. Microscop. Soc. 85: 305-313,
<u>1967. Microstome-macrostome transformation in *Tetrahymena* vorax Strain V₂ type S induced by a transforming principle, stomatin, J. Profozool, 14: 608-613.
</u>

BUHSE, H. E. and I. L. CAMERON, 1968. Temporal pattern of macromolecular events during the microstome-macrostome cell transformation of Tetrahymena vorax V₂S. J. Exp. Zool. 169: 229-236.
BUHSE, H. E., J. O. CORLISS, and R. C. HOLSEN. 1970. Tetrahymena vorax: Analysis of stomatogenesis by scanning electron and light microscopy. Trans. Amer. Microsc. Soc. 89: 328-336.
BUTSCHLI, O. 1889. Protozoa. In Bronn, H. G., Klassen und Ordnung des Thiers-Reichs, Pts. 1-64, 2035 pp. C. F. Winter, Leipzig.
CHATTON, E. and A. LWOFF. 1930. Imprégnation, par diffusion argentique, de l'infraciliature des ciliés marins et d'eau douce, après fixation cytologique et sans dessication. Compt. Rend. Soc.

Biol., 104: 834-836.

. 1936. Techniques pour l'étude des protozoaires, spécialement de leurs structures superficielles (cinétomé et argyrome). Bull, Soc. Fr. Micr., 5: 25-39.

CHATTON, E., A. LWOFF, M. LWOFF, and J. L. MONOD. 1931. La formation 1'ébauche buccale posterieure chez les ciliés en division

et ses relations de continuité topographique et génétique avec la bouche anterieure. R. Soc. Biol., Paris, 107: 540-544°. CLAFF, C., L. 1947. Induced morphological changes in Tetrahymena vorax. CLAPAREDE, E. and J. LACHMANN, 1858-1861. Études sur les infusoires et les rhizopodes. Mém. Inst. Nat. Genèvois, 5: 1-260; 6: 261-482; 7: 1-291.

CONWAY, E. J. 1942, Mean geochemical data in relation to oceanic evolution. Proc. R. Irish Acad., 48B: 119-159.
CORBETT, J.J. and J. SWEENEY, 1966. Antigenic differences among some classical *Tetrahymena pyriformis* and *T. vorax* strains.
J. Protozool. 13: 359-366.

CORLISS, J. 0. 1951. Morphology and systematic status of pure culture strains in the Colpidium-Glaucoma-Leucophrys-Tetrahymena group (Abstr.) Proc. Amer. Soc. Protozool. 2: 11. . 1952a. Comparative studies on holotrichous ciliates in the Colpidium-Glaucoma-Leucophyrs-Tetrahymena group. I.

General considerations and history of strains in pure culture. ... Trans. Amer, Micr. Soc. 71: 159-184.

. 1952b. Review of the genus Tetrahymena. Proc. Soc. Protozool. 3:3.

1953a. Comparative studies on holotrichous ciliates in the *Colpidium-Glaucoma-Leuchophrys-Tetrahymena* group. II. Morphology, life cycles and systematic status of strains in pure culture. Parasitology 43: 49-87. CORLISS, J. O. 1953b. Silver impregnation of ciliated protozoa by the Chatton-Lwoff Technique. Stain Tech. 28 : 97-100.

1954. The literature on *Tetrahymena*: its history, growth, and recent trends. J. Protozool. 1 : 156-169.

1956. On the evolution and systematics of ciliated

1957. Tetrahymena paravorax n. sp., the first caudalciliated member of the genus referable to the vorax-patula complex. (Abstr.) J. Protozool. 4 (Suppl.) : 13.

2. 1959a. An illustrated key to the higher groups of the ciliated protozoa with definition of terms. J. Protozool.

1959b. Current composition of the genus Tetrahymena Furgason, 1940. (Abstr.) J. Protozool. 6 (Suppl.) : 24.

1960. Tetrahymena chironomi sp. nov., a ciliate from midge larvae, and the current status of facultative parasitism in the genus Tetrahymena. Parasitology. 50 : 111-153. 1961a. The ciliated protozoa. Pergamon Press, London and New York. 310 pp.

1961b. Natural infection of tropical mosquitoes by ciliated protozoa of the genus *Tetrahymena*. Trans. Roy. Soc. Trop. Med. Hyg. 55 : 149-152.

1963. Application of modern techniques to problems in the systematics of the protozoa. Proc. XVI Internat. Congr. Zool. Washington. 4 : 97-102. CORLISS, J. O. 1965. *Tetrahymena*, a ciliate genus of unusual importance in modern biological research. Acta Protozoologica.

DAVIS, C. C. 1969. Seasonal distribution, constitution, and abundance of zooplankton in Lake Erie. J., Fish. Res. Bk. Canada. 26 : 2459-2476.

DOBELL, C. 1932. Antony van Leeuwenhoek and his "Little Animals". Staples Press, London, vii + 435 pp.

DOFLEIN, F. 1901. Die Protozoen als Parasiten und Krankheitserreger nach biologischen Gesichtsgünkten dargestellt. Jena.

pp. XII + 274, 220 figs.

DRISCHEL, H. 1940. Chlorid-, Sulfat-, und Nitratgehalt der atmosphärischen Niederschlage in Bad Reinerz und Oberschreiberhau in Vergleich zu bisher bekannten Werten anderer Orte. Balneologe. 7 : 321-334.

DUJARDIN, F. 1838. Mémoire sur l'organisation des infusorires. Ann. Sci. nat. (sér. 2, Zool.). 10 : 230-315.

> 1841. Histoire Naturelle des Zoophytes. Infusoires. Páris. 678 pp.

EHRENBERG, C. G. 1830. Abhandl. Akad. Wiss. Berlin, qus dem Jahre 1830 (gedruckt 1832) 1 - 88.

> 1832a. Beiträge zur Kenntnis der Organisation der Infusorien und ihrer geographischen Verbreitung, besonders in Siberien. Abh. Akad. Wiss. Bêrlin, yr. 1831, pp.1-154.

EHRENBERG, C. G. 1832b. Über die Entwicklung und Lebensdauer der Infusionsthiere; "nebst ferneren Beiträgen zu einer Vegleichung ihrer organischen Systeme. Abh. Akad. Wiss. Berlin, yr.1831,

pp. 1-88.

1833. Dritter Beiträgen zur Erkenntniss der grosser Organisation in der Richtung des kleinsten Raumes. Abh. Akad. Wiss. Berlin, yr. 1833. pp. 145-336.

1838. Die Infusionsthierchen als Volkommene Organismen. Leipzig, 612 pp. (Diagnoses given in Latin, French and German). 1854. Nova genera et novae species maris profundi. Monatsberichte der Berliner Akademie, p. 236. In Kent, W. S. (1881-1882), A manual of the Infusoria. II. pp. 626-627. David Bogue, London. 913 pp.

ELLIOTT, A. M. 1959. The Biology of *Tetrahymena*. Ann. Rev. Microbiol. 13: 79-96.

ELLIOTT, A. M., M. A. ADDISON, and S. E. CAREY. 1962. Distribution of .Tetrahymena in Europe. Jg Propozool. 9: 135-141.

ELLIGTT, A. M. and G. L. CLEMMONS. 1966. An ultrastructural study of ingestion and digestion in *Tetrahymena pyriformis*. J.

Protozool. 13 : 311-323.

ELLIOTT, A. M. and D. F. GRUCHY. 1952. The occurrence of mating types in Tetrahymena. Biol. Bull. 103 : 301.

ELLIOTT, A. M. and R. E. HAYES. 1953. Mating types in *Tetrahymena*. Biol. Bull. 105 : 269-284.

ELLIOTT, A. M., M. A. STUDIER, and J. A. WORK. 1964. Tetrahymena pyriformis from several Pacific Islands and Australia. J. Protozool. 11 : 370-378.

ENGELMANN, T. W. 1862. Zur Naturgeschichte der Infusionstiere. Ebenda, 11.

1876. Ueber Entwicklung und Fortpflanzung der Infusorien. Morph. Fb. 1 : 573-635. FAURE-FREMIET, E. 1906. Le *Glaucoma pyriformis* et l'organisation de

la substance vivante. C. R. Assoc. Anat. Session de Bordeaux. 120-127.

1910., Étude sur les mitochondries des protozoaires et des cellules sexuelles. Arch. Anat. Micr. 2 : 457-648. 1948. Les mécanismes de la morphogenèse chez les ciliés. Folia biotheoretica. 3 : 25-58.

1949. Action de lithium sur la stomatogenèse chez les ciliés. Belg-Nederl. Cyto-Embryol. Dagen., Gent.

pp. 100-102.

9 : 183-187.

1951. Écologie des protistes littoraux. Année. Biol.

27 : 437-447.

FELTON, M., J. COONEY and W. MOORE. 1967. A quantitative study of the A bacteria of a temporary pond. J. Gen. Microbiol. 47 : 25-31. FRANKEL, J. and K. HECKMANN. 1968. A simplified Chatton-Lwoff silver impregnation procedure for use in experimental studies with ciliates. Trans. Amer. Microsc. Soc. 87 : 317-321.

FREY, D. G. 1949. Morphometry and hydrography of some natural lakes of the North Carolina coastal plain : the bay lake as a Morphometric type, J. Elisha Mitchell Sci. Soc. 65 : 1-37. 1966. Limnology in North America. Univ. Wisc. Press.

FURGASON, W. H. 1940. The significant cytostomal pattern of the "Glaucoma-Colipidium group", and a proposed new genus and species, Tetrahymena geleii. Arch. Protist. 94 : 224-266.
GELEI, J. von. 1935. Colpidium glaucomaeforme n. sp. (Hymenostomata) und sein Neuronemensystem. Arch. Protistenk. 85 : 289-302.

pp. 734.

GEIMAN, Q. M. 1931. Morphological variations in Colleps octospinus. Tr. Am. Micr. Soc. 50 : 136-143.

GRANT, Mr L. and R. F. THORNE. 1955. Discovery and description of a sphagnum bog in Iowa, with notes on the distribution of bog plants in the State. Proc. Iowa Acad. Sci. 62 : 197-210.

GRUCHY, D. F. 1955. The breeding system and distribution of Tetra-

hymena pyriformis. J. Protozool. 2 : 178-185.

HOLLAND, T. H. and W. A. K. CHRISTIE. 1909. The origin of the salt deposits of Rajputana. Rec. geol. Surv. India. 38 : 154-186.
HOLZ, G. C. Jr. 1955. Loss of the micronucleus and reduction in the number of kinetics in a hymenostome ciliate. (Abstr.) Mol. Bull. 109 : 360.

HURST, D. D. 1957. Occurrence of 'cysts' in Tetrahymena pyriformis. variety 2. (Abstr.) J. Protozool. 4 (Suppl.) : 18. CHINSON, G. E. 1957. A treatise on limnology. I. Geography, Physics, and Chemistry. John Wiley and Sons. pp. 1015.

JUDAY, C., E. A. BIRGE, and V. W. MELOCHE. 1938. Mineral content of northeastern Wisconsin. Trans. Wisc. Acad. Sci. Arts Lett. 31 : 223-276.

UDAY, C., E. B. FRED, and F. C. WILSON. 1924. The hydrogen ion concentration of certain Wisconsin lake waters. Trans. Amer. Micro. Soc. 43 : 177-190.

KAHL, A. 1926. Neue und wenig bekannte formen der holotrichen und heterotrichen Ciliaten. Arch. Protistenk. 55 : 197-438.

> 1930-1935. Urtiere oder Protozoa. I : Wimpertiere oder Ciliata (Infusoria), eine Bearbeitung der freilebenden und ecotocommensalen Infusorien der Erde, unter Ausschluss der marinen Fintinnidae. In Dahl, F., Die Tierwelt Deutschlands, Teil 18²²(yr. 1930), 21 (1931), 25 (1932), 30 (1935) pp. 1-886. G. Fisher, Jena.

KAZUBSKI, S. Ludy 58. Trichia lubomirskii Slos. (Helicidae), a new host of Tetrahymena limacis (Warren; 1932) Kozloff, 1946 (Ciliata) and Zonitoides nitidus MHL, (Zonitidae), a new host of T./ rostrata (Kahl, 1926) Corliss, 1952, in Poland. Bull, Acad. Polonaise Sci. 6 : 247-252.

KEILIN, D. 1921. On a new ciliate, Lambornella stegomyiac; n. g., n. sp., parasitic in the body cavity of the larvae of Stegomyia scutellaris Walker (Diptera, Nematocera, Culicidae) Parasitology. 13 : 216-224.

KENK, R. 1949. The animal life of temporary and permanent ponds in Southern Michigan. Univ. Michigan Misc. Publ. 71 : 1 - 16.
KENT; W. S. 1880-1882. A Manual of the Infusoria. Vols. I - III.
David Bogue, London. 913 pp.

KIDDER, G. W. 1941. Growth studies on ciliates. VII Comparative growth characteristics of four species of sterile ciliates. Biol. Bull. 80 : 50-68.

KIDDER, G. W., D. M. LILLY, and C. L. CLAFF. 1940. Growth studies on ciliates. IV. The influence of food on the structure and growth of *Glaucoma vorax* sp. nov. Biol. Bull. 78 : 9-23.
KLEIN, B. M. 1926. Ergebnisse mit einer Silber-methode bei Ciliation. Arch. Protistenk. 56 : 243-279.

1958. The dry silver method and its proper use. J. Protozool. 5 : 99.

KOZLOFF, E. N. 1946. The morphology and systematic position of a holotrichous ciliate parasitizing *Deroceras agreste* (L). J. Morph. 79 : 445-465.

> 1956a. A comparison of the parasitic phase of Tetrahymena limacis (Warren) with clones in culture, with particular reference to variability in the number of primary ciliary meridians. J. Protozool. 3 : 20-28.

KRASCHENINNIKOW, S. and D. H. WENRICH, 1958. Some observations on the morphology and division of Balantidium coli and Balantidium cavivae (?). J. Protozool. 5: 196-202.
KUDO, R. R. 1966. Protozoology. 5th ed. Charles C. Thomas. 1174 pp.

LACKEY, J. B. and E. W. LACKEY. 1970. A late summer check list of the marine microbiota around Logy Bay, Newfoundland. Can. J.

Zool, 48: 789-795,

LAIRD, M. 1956, Studies of mosquitoes and freshwater ecology in the South Pacific. Roy, Soc. New Zealand Bull. No. 6, 213 pp. LEDERMULLER, M. F. 1763, Mikroskopische Gemüth- und Augen- Ergötzung. Nurnber,

LEEUWENHOEK, A. van, 1677. [Letter No. 18 (9, Oct. 1676) to the Royal Society of London. Partly published in Philosophical Transactions for 1677. Original not seen, information from Dobell, 1932].

LIEBERMAN, P. R. 1929. Ciliary arrangement in different species of Parameciu. Tans. Amer. micro. Soc. 48: 1-LOEFER, J. B., R. D. OWEN, and E. CHRISTENSEN. 1958. Serological

types among thirty-one strains of the ciliated protozoan Tetrahymena pyriformis. J. Protozool. 5: 209-217. LOEFER, J. B., E. B. SMALL, and W. H. FURGASON. 1966. Range of variation in the somatic infraciliature and contractile vacuole pores of Tetrahymena pyriformis. J. Protozool. 13: 90-102: 143

LOM, J. and M. LAIRD. 1969. Parasitic protozoa from marine and euryhaline fish of Newfoundland and New Brunswick. I. Peritrichous ciliates, Can. J. Zoob. 47: 1367-1380.

LWOFF, A. 1947, Some aspects of the problem of growth factors for protozoa. Ann, Rev. Microbiol. 1: 101-114,

MACAN, T. T. 1970, Biological studies of the English lakes. Longman, London, 260 pp.

MACKERETH, F. J. H., and J. HERON. 1954. In Rep. Freshw. biol. Ass. Brit. Emp. 22: 20-21.

MARGAVIO, A. V. 1964, An ecological study of a temporary pond in

southeast Louisiana. M.Sc. Thesis. Loyola University, New

Orleans, La., U.S.A. 62 pp.

MARSLAND, D. A. 1943. Quieting Paramecium for the elementary student. Science, 98: 414.

MAUPAS, E. 1883. Contribution à l'étude morphologique et anatomique des infusoires cilies. Arch. Zool. Exp. Gen. (ser. 2) 1:

427-664.

MAUPAS, E. 1888. Recherches expérimentales sur la multiplication des infusoires Éiliés. Arch. Exp. Gen.

MILLER, N. H. J. 1914. The composition of rain-water collected in the Hebrides and Iceland. With special reference to the amount of nitrogen as ammonia and as nitrates. J. Scot. met. Soc. 16 : 141-158.

MOORE, W. G. 1951. Observations on the biology of *Streptocephalus* seali. Proc. La. Acad. Sci. 14 : 57-65.

1955. The life history of the spiny-tailed fairy shrimp and in Louisiana. Ecology. 36 : 176-184.

. 1970. Limnological studies of temporary ponds in southeastern Louisiana. Southwestern Naturalist. 15 : 83-110.

> Infusorium, Helminthicorum et Testaceorum, non Marinorum, Succincta Historia. Havniae et Lipsiae. 135 pp.

NEEDHAM, J. G., P. S. GALTSOFF, F. E. LUTZ and P. S. WELCH. 1959.

Culture methods for invertebrate animals. Dover, New York.

NILSSON, J: R. 1962. Observations on *Neobursaridium gigas* Balech, 1941 (Ciliata, Heterotrichida). J. Protozool. 9 : 273-276. NOLAND, L. E. 1925a. A review of the genus *Coleps* with descriptions of two new species. Trans. Amer. micr. Soc. 44 : 3.

NOLAND, L. E. 1925b. Factors influencing the distribution of freshwater giliates, Ecology. 6: 437. ; 1937, Observations on marine ciliates of the Gulf coast of Florida, Trans, Amer, micr, Soc, 56: 106-171, 1959, Ciliophora, In Edmondson, W. T., editor, Ward and Shripple's Fresh-Water Biology, 2nd ed., pp. 265-297. Wiley and Sons, New York, - NOLAND, L. E. and H. E. FINLEY, 1931. Studies on the taxonomy of the genus Vortice Ita. In. Am. micr. Soc. 50: 81. PICKAVANCE, J. R., G. F. BENNETT, and J. PHIPPS. 1970. Some mosquitoes and blackflies from Newfoundland, Can. J. Zool. 48: 621-624. RAABE, Z. 1959. Trichodina pédiculus (O. F. Müller, 1786) Ehrenberg 1838 et Trichodina domerguei (Wallengren, 1897). Acta. Parasitol. Po, 7: 189-202. ROULEAU, E, 1956, Some Newfoundland vernacular plant names. Contr. Inst, Bot, Univ: Montreal No. 69, II, : 25-40. RQUX, °J., 1901. Faune Infusorienne des Eaux Stagnantes des Environs de Genève, Kundig, Genève, 148 pp. RUSSELL, E. J. and E. H. RICHARDS, 1919. The amounts and composition of rain and snow falling at Rothamsted. J. agric, Sci. 9: 309-337. RZOSKA, J. 1961) Observations on tropical rainpools and general remarks on temporary waters. Hydrobiologia. 17: 265-286.

SCHEWIAKOFF, W. 1893. Ueber die geographische Verbreitung der Süsswasser-Protozoen. Mém. Acad. impér. Sci. St. Pétersb. (sèr. 7), 41: 1-201. (in Russian).

 1896. The Organization and Systematics of the Infusoria Aspirotricha (Holotricha auctorum). Méta, Acad. impér. Sci. St. Pétersb. (šér. 8), 4: 1-395. (In Russian).
 SESHACHAR, B. R. and P. B. PADMAVATHI. 1956. The cytology of a new species of Spirostomum, J. Protozool, 9: 265.

SHAW, R. F. and N. E. WILLIAMS, 1963. Physiological properties of

Tetrahymena vorax. J. Protozool. 10: 489-491.

SHELFORD, V. 1913. Animal communities in temperate America. Univ. Chicago Press. 368 pp.

SPANDL, H. 1926. Die Tierwelt Vorübergehender Gewässer Mitteleuropas. Arch. Hydrobiol. 16: 74-132.

STEIN, F. 1854. Die Infusionsthiere auf ihre Entwicklungsgeschichte untersucht. Leipzig. 265 pp.

STOKES, A. C. 1884. Hymenostoma hymenophora. Am. Monthly Micros. J.,
V, July, in Stokes A. C. 1888. A preliminary contribution
toward a history of the fresh-water infusoria of the United
States. J. Trention Natural Hist. Soc. I(3): 173-175.
1888. A preliminary contribution toward a history of
the fresh-water infusoria of the United States. J. Trention

Natural Hist, Soc. I(3): 71-344.

- STRØM, K. M. 1939. Conductivity and reaction in Norwegian Lake waters. Int. Rev. Hydrobiol. 38 : 250-258.
- 1942. Hadeland lakes. A limnological outline, Skr. norske 'Vidensk Akad., Mat-nat. Kl. 1941. No. 7. 42 pp.
 - 1944. High mountain limnology. Some observations on stagnant and running waters of the Rondane area. Arh. norske Vidensk Akad., Mat-nat. Kl. 1944. No. 8. 24 pp.
 - SUHAMA, M. 1961. Experimental studies on the morphogenesis of Condylostoma spatiosum Ozaki and Kagui. J. Sci. Hiroshima Univ. (B, 1), 21: 33-82.
- THOMPSON, J. G. Jr. 1955. Morphology of a new species of Tetrahymena. (Abstr.) J. Protozool. 2 (Suppl.), 12.
- 1960. A protozoan "Fingerprint". Turtox News, 38 (10) : 258-259.
 - 1963a. A redescription of the hymenostome ciliate Sathrophilus (saprophilus) mucsorum (Kahl. 1931) Corliss, 1960, with particular attention to the buccal apparatus. Va. J. Sci. 14 : 16-22
 - 1963b. The generic significance of the buccal infraciliature in the family Tetrahymenidae and a proposed new genus and species. Paratetrahymena wassi. Va. J. Sci. 14:126-135

- WANG, C. C. 1928. Ecological studies of the seasonal distribution of protozoa inta fresh water pond. J. Morph. 46 : 431-478.
- WARREN, E. 1932. On a chiliate protozoan inhabiting the liver of a slug. Anh. Natal Mus. 7 : 1-53.
- WEBB, M. G. 1961, The effects of thermal stratification on the distribution of benchic protozoa in Esthwaite water. J. Anim. Ecol. 30 : 137-151.
- WEHRLE, E. 1927. Studien über Wasserstoffionenkonzentrationsverhältnisse und Besiedung an Algenstandorten in der Umgebung von Freiburg im Brisgau. Z. Bot. 19 : 207-287.
- WILLIAMS, N. E. 1961. Polymorphism in Tetrahymena vorax. J. Protozool. 8 : 403-410.
- WOODRUFF, L.L. 1921. The structure, life history and intrageneric relationships of *Parameeium calkinsi*, sp: nov. Biol. Bull. 41 : 171-180.
- YÓSHÍMURA, S. 1934. Kata-numa, a very strong acid-water lake of Volcano, Kata-numa, Miyagi Prefecture, Japan. Arch. Hydrobiol. 26 : 197-202.

ADDITIONAL REFERENCES

- Bodian, D. 1937. The staining of paraffin section of nervous tissues with activated protargol. The role of fixatives. Anat. Rec., 69 : 153-62.
- Burt, R. L. 1940. Specific analysis of the genus *Colpoda* with special reference to the standardization of experimental material. Tr. Am. Micr. Soc., 59 : 414.
- Chatton, E. and A. Lwoff. 1949. Recherches sur les cilies thigmotriches I. Arch. Zool. exptl. gen., 86 : 169 - 253.
 - , 1950. Recherches sur les cilies thigmotriches. II.
 - Corliss, J. O., 1968. The value of ontogenetic data in reconstructing protozoan phylogenies. Trans. A. Micr. Soc., $87 : 1-20_1$
- Dodd, E. E. 1962. A cytochemical investigation of the nuclear changes in the cyst cycle of *Telotrichidium henneguyi*. J., Protozopl., 9:93.
- Dragesco, J. 1960. L'drientation actuelle de la systematique des cilies et la technique d'impregnation au proteinate d'argent. Bull. Micr. Appl., 11: 49-68.
 - 1968. Les genres Pleuronema Dújardin, Schizocalyptra
- nov. gen. et *Histobalantium* Stokes. Protistologica 4 : 85-106. Fraure-Fremiet, E. 1953. La bipartition enantiotrope chez les cilies obligotriches. Arch. Anat. microsc. Morphol. exptly.,
 - 42 : 209-225.
 - .: 1959. La famille des Nassulidae (Ciliatia Gymnostomatidae)

et le genre Wassulopsis n. gen. Compte rend. Acad. Sci., Paris, 249 : 1429-1433.

Honigberg, B. M. and H. A. Davenport. 1954. Staining flagellate protozoa by various silver-protein compounds. Stain Tech., 29:: 241-6.

Honigberg, B. M., W. Balamuth, E. C. Bovee, J.=O. Corliss, M. Gojdics,
R. P. Hall, R. R. Kudo, N. D. Levine, A. R. Loeblich, Jr.,
J. Weiser, and D. H. Weinrich. 1964. A revised classification of the phylum Protozoa. J. Protozool., 11 : 7-20.
Jankowski, A. W. 1964a. Morphology and evolution of Ciliophora. III.

Diagnoses and phylogenesis of 53 sapropebionts, mainly of the order Heterotrichida. Arch. Protistenk., 107 : 185-294.
. 1964b. Morphology and evolution of the Ciliophora. IV. Sapropelebionts of the family Loxocephalidae fam. nova, their
taxonomy and evolutionary history. Acta Protozool., 2 : 33-58.
. 1967. Principles of construction of a new system of ciliate protozoa. Zool. Inst. Acad. Sci. USSR, Sci. Sess., Leningrad, 63: 3-52.

Kozloff, E. N. / 1961. A new genus and two new species of Ancistrocomid Ciliates (Holotricha: Thigmotricha) from Sabellid Polychaeta and from a chiton. J. Protozool., 8 : 60-63.
Lom, J., J. O. Corliss and C. Noirot-Timothee. 1968. Observations of the ultrastructure of the buccal apparatus in Thigmotrich ciliates and their bearing on Thigmotrich-Peritrich affinities.

J. Protozzol., 54": 824-840.

Nobili, R. 1969. Sistematica e filogenesi dei protozoi cliati. Boll. Zool., 36 : 311-326.

Penard, E. 1922. Etudes sur les Infusiores d'Eau Douce. Georg et Cie., Geneve. 331 pp.

Puytorac, P. de. 1954. Contribution a l'etude cytologique et taxonomique des infusibles astomes. Ann. Sci. Nat. Zool., (series 11), 16 : 85-270.

> _____. 1959. Quelque observations sur l'evolution et les origines des cilies Astomatida. Proc. XV Int. Congr. Zool., London, 1958, : 649-651.

. 1963. Contribuțion a l'etude des cilies astomes Haptomain phyridae Cepede, 1903 (cytologie, ultrastructure, taxinomie). Ann. Sci. Nat. Zool., (series 12), 5 : 173-210. Small, E. B. 1967. The Scutiociliatida, a new order of the class

Ciliata (phylum Protozoą, subphylum Ciliophora). Trans. Am.

Tuffrau, M. 1963. L'infraciliature, la morphogenese et la position systematique du genre Lembadion Pertý, 1852 (cilie hymenostome). Ann. Sci. Nat. Zool., (series 12), 5: 479-490.
______. 1967. Perfectionnements et pratique de la technique

d'impregnation au Protargol des Infusoires, Cilies. Protistologica 3: 91-98.

Weinrich, D. H., 1928. Eight well-defined species of Paramecium.

Trans. Am. Micr. Soc., 47 : 275.

ADDENDUM

THE. RELATIONSHIP BETWEEN, CARNIVOROUS Tetrahymena voraz AND THE PREY

On 17 July 1969, T, vorax was collected from the pond water at 23°C., with dissolved solid concentration 126 p.p.m. (see p. 28) and cultured in hay infusion with a surface sprinkling of bacto-tryptone at room temperature (22°C). Tetrahymena pyriformis, Paramecium bursaria, and Urotricha sp. were examined in the same collection.

The marked dimorphism of the cillate under discussion was evident by phase contrast microscopy as stated on page 55. The pyriform microstomes were examined from silver impregnation preparations which also bear the tailed microstomes and the carnivorous macrostomes of T, vorax.

The event of the holotrich eating its prey was observed as follows:

The microstome form of T, vorax started to broaden, and the mouth became larger and more open beneath the microstome oral cavity.

Back of the oral cavity, a pharyngeal pouch appeared which was continuous with the outsides through the mouth opening. The pharyngeal pouch increased in size until it was ready to reach the posterior extremity. At this period, the vacuole occupied the greatest volume of the cell with the protoplasm surrounding it in a thin film.

After this microstome macrostome transformation, the difficult task for the macrostome was to capture and digest its first prey.

As the mouth was open and large, the membranelles created 'strong currents into the "pharyngeal pouch"; eventually a prey was drawn in.

The macrostome immediately became quite active and swam in circles with the mouth directed toward the inner part of the circle. The prey swam about in the vacuole in an entirely normal manner. Eventually, the macrostome protoplasm closed down until the "pharyngeal pouch", with its trapped prey, was cut off from the mouth region.

The enclosed prey continued to swim about, but the fluid content of the vacuole decreased slowly; the protoplasm of the macrostome and prey came to lie close together, and the motion of the latter was restricted. Eventually, the prey lost all activity, and digestion was under way.

As the digestion proceeded, the macrostome increased in size, and afterwards it was able to capture preys much more rapidly.

The event was first described by Kidder et al. (1940), and also reported later by Corliss (1953a), Williams (1961), and Buhse (1966a). Recently, Buhse (1966a & b, 1967, 1968, and 1970) has actively studied the transformation in *T. vorax*. Briefly, Buhse (1967) followed Claff's (1947) study and found that the transformation in *T. vorax* was induced by suspending microstomes in a transforming principle, stomatin, released by the potential prey, *T. pyriformis*. This process occurs in four and a half hours after stomatin addition (Buhse & Carmeron, 1968). Some analysis of stomatogenesis was done by scanning electron and light microscopy (Buhse, Corliss, & Holsen, 1970). This oral replacement is preceded by RNA synthesis and protein synthesis which are stimulated by stomatin (Buhse & Carmeron, 1968). According to Bubge's work (1962), in nature the microstome macrostome transformation of T, vorax is associated only with living T. pyriformis. Once the prey, T. pyriformis, has been exhausted, the reverse transformation, which always involves division of the carnivorous macrostome to form two tailed microstomes, occurs (Kidder et al., 1940; Williams, 1961; and Bubse, 1966b). Evidently, the presence of T. pyriformis in the same collection with T. vorax on 17 July 1969 could explain the occurrence of the carnivorous T. vorax -- T. pyriformis was the source of the microstome + macrostome transforming principle, stomatin, and the prey of the carnivorous macrostome.

Since the irreversible pyriform microstome was described (Kidder et al., 1940; and Corliss, 1953a) as "a curious stage in that forms in it are morphologically indistinguishable from *T. pyriformis*" (quoted on page 60), it had been very uncertain whether the pyriform holotrich is just a stage of *T. vorax* or the true *T. pyriformis* if it is present in the same preparation with the other forms of *T. vorax*. As stated on page 65, "They might either be '*T. pyriformis*' (the prey) and stomatin-...-source of *T. vorax*) or irreversible pyriform microstome forms derived from *T. vorax*." (see the last paragraph of *T. vorax*, p. 65).

The allocation of *T. vorax* strains V_1 and PP which had been reported to be irreversible pyriform microstomes of *T. vorax* (Kidder et al., 1940, and Corliss, 1953) was questioned by Loefer et al. (1958), and Shaw and Williams (1963). Their data from nutritional and serological experiments indicate that the strains V_1 and PP are actually strains of *T. pyriformis*. Later, both the immúnological (Corbett and Sweeney, 1966) and the somatic infraciliature (Loefer et al., 1966)

studies support the view of Shaw and Williams. Consequently, T, vorax strains V, PP and V₁ are proposed to be T, pyriformis (Loefer. et al., 1966). This leaves T, vorax represented by two extant strains, V₂ and Tur, both of which are polymorphic.

Therefore, it is most likely that the non-transformable, pyriform Tetrahymena in the same preparation with T. vorax of the present study is T. pyriformis - the source of stomatin and the prey of T. vorax.

. 5 .

e 5 g



