FISH COMMUNITY CHANGES IN AN EXPLOITED MARINE ECOSYSTEM: NEWFOUNDLAND SOUTHERN GRAND BANK AND ST. PIERRE BANK, 1951–1995

CENTRE FOR NEWFOUNDLAND STUDIES

TOTAL OF 10 PAGES ONLY MAY BE XEROXED

(Without Author's Permission)

JILL M. CASEY







#### INFORMATION TO USERS

This manuscript has been reproduced from the microfilm master. UMI films the text directly from the original or copy submitted. Thus, some thesis and dissertation copies are in typewriter face, while others may be from any type of computer printer.

The quality of this reproduction is dependent upon the quality of the copy submitted. Broken or indistinct print, colored or poor quality illustrations and photographs, print bleedthrough, substandard margins, and improper alignment can adversely affect reproduction.

In the unlikely event that the author did not send UMI a complete manuscript and there are missing pages, these will be noted. Also, if unauthorized copyright material had to be removed, a note will indicate the deletion.

Oversize materials (e.g., maps, drawings, charts) are reproduced by sectioning the original, beginning at the upper left-hand corner and continuing from left to right in equal sections with small overlaps.

Photographs included in the original manuscript have been reproduced xerographically in this copy. Higher quality 6<sup>°</sup> x 9<sup>°</sup> black and white photographic prints are available for any photographs or illustrations appearing in this copy for an additional charge. Contact UMI directly to order.

> Bell & Howell Information and Learning 300 North Zeeb Road, Ann Arbor, MI 48106-1346 USA 800-521-0600

> > UMľ



National Library of Canada

Acquisitions and **Bibliographic Services** 

395 Wellington Street Ottawa ON K1A 0N4 Canada

Bibliothèque nationale Acquisitions et services bibliographiques

395, rue Wellington Ottawa ON K1A 0N4

du Canada

Your file Votre rélérence

Our file Natur réliérence

The author has granted a nonexclusive licence allowing the National Library of Canada to reproduce, loan, distribute or sell copies of this thesis in microform, paper or electronic formats.

The author retains ownership of the copyright in this thesis. Neither the thesis nor substantial extracts from it may be printed or otherwise reproduced without the author's permission.

L'auteur a accordé une licence non exclusive permettant à la Bibliothèque nationale du Canada de reproduire, prêter, distribuer ou vendre des copies de cette thèse sous la forme de microfiche/film, de reproduction sur papier ou sur format électronique.

L'auteur conserve la propriété du droit d'auteur qui protège cette thèse. Ni la thèse ni des extraits substantiels de celle-ci ne doivent être imprimés ou autrement reproduits sans son autorisation

0-612-54866-X



# Fish Community Changes in an Exploited Marine Ecosystem: Newfoundland Southern Grand Bank and St. Pierre Bank, 1951-1995

by

©Jill M. Casey

A thesis submitted to the School of Graduate Studies in partial fulfillment of the requirements for the degree of Master of Science

> Department of Biology Memorial University of Newfoundland April 2000

St. John's

Newfoundland

## Abstract

Longterm fish community changes on southern Grand and St. Pierre Banks were examined from dedicated research survey trawls conducted by the Canadian Department of Fisheries and Oceans from 1951-1995. These time series pre-date the arrival of factory-freezer-trawlers in the 1960's and 1970's, and thus provide insight into changes in the biomass and diversity of an exploited system. The time series were standardized by applying conversion factors for changes in vessel, diel changes in catchability, and relative catchability of selected species. Total biomass in the 1990's was reduced to 11% and 9% of that observed in the 1950's on southern Grand Bank and St. Pierre Bank, respectively, and largely resulted from the decline of the haddock population. Compensatory responses to this decline were visible with the flatfish on southern Grand Bank and skate on St. Pierre Bank but continued fisheries for flatfish and bycatch of skate ensured that total species biomass would remain at low levels. This study shows the importance of examining data on as long a time-scale as possible. Failure to examine such historical data has resulted in the largest skate in the northwest Atlantic, the barndoor skate, being driven to near extinction without anyone noticing.

#### Acknowledgements

I would like to thank Dr. R. A. Myers for giving the opportunity to work on such an interesting and challenging Masters project, and for having such confidence in my ability to get the job done. I would also like to thank Dr. R. L. Haedrich, Dr. P. Sneigrove, and Dr. R. Knoechel for provided valuable criticism on this thesis.

The Canadian Department of Fisheries and Oceans Groundfish Division, St. John's, Newfoundland is thanked for providing data for this analysis. B. Atkinson, G. Lily, and S. Walsh provided much appreciated help in interpreting the data. Noel Cadigan is thanked for suggesting the form of model 2 in Chapter 1.

The Canadian Department of Fisheries and Oceans, Halifax, and the National Marine Fisheries Service at Woods Hole, MA are thanked for providing the data for Chapter 3. Also, B. Atkinson, K. Frank, J. E. Simon, M. Fogarty, S. Murawski and J. Hutchings are thanked for their helpful discussions of Chapter 3.

And last, but not least, I would like to thank my husband, Gavin Crutcher, for his patience and emotional support. I would also like to thank our son, Andrew, who showed me the benefits of applauding your own accomplishments no matter how small they might be. The story of this population is a very good example of how a fishery can disappear while being watched carefully by fisheries biologists and an international regulatory body (Templeman 1978 - in reference to the haddock population on the southern Grand Bank).

# Contents

	Abs	tract
	Ack	nowledgementsiii
	List	of Tables vii
	List	of Figures viii
	Intr	oduction and Overview 1
	Co-	authorship Statement 5
1	Die	variation in trawl catchability: Is it as clear as day and
	nigl	t? 6
	1.1	Abstract
	1.2	Introduction
	1.3	Methods
		1.3.1 Data
		1.3.2 Model 10
		1.3.3 Regional and Seasonal Differences in Diel Catchability 12
		1.3.4 Randomization Tests
		1.3.5 Depth Effects
	1.4	Results
	1.5	Discussion

2	Fish	a Community	Cha	nges	in	1 a	m	E	xp	lo	it	ed	1	M	ar	in	е	E	cc	s	ys	te	em:	
	Nev	vfoundland So	uthe	rn G	rai	ıd	B	an	k	an	d	St		P	ie	rr	e I	Ba	m	k,	1	9	51-	
	199	5																						26
	2.1	Abstract									•													26
	2.2	Introduction .							•															27
	2.3	Methods							•	•	•		•			•					•	•		28
	2.4	Results				•	•		•	•	•					•						•		38
	2.5	Discussion				•		-	•	•	•									•	•			47
3	Nea	r Extinction o	of a I	arge	, V	Vie	de	ly	D	is	tri	ibı	ıt	ed	1	Fis	h							<b>52</b>
	3.1	Abstract				•									•							•		52
	3.2	Introduction .				•							•							•				53
	3.3	Methods			• •		•		•							•				·	•	•		55
	3.4	Results		• • •	• •	•	•	• •	•	•	• •	• •	•	•	•	•	•	•	•	•	•	•	÷	56
	3.5	Discussion		• • •	• •	•	•	• •		•	•	• •	•	•	•	•		•	•	•	•	•	•	57
Summary								61																
Bibliography								63																
A Estimation in SAS								73																
в	B Research Survey Tows Per Stratum								75															
С	Sou	thern Grand I	Bank	Dat	a I	Jse	ed	in	0	Ch	aŗ	ote	r	2										78
D	D St. Pierre Bank Data Used in Chapter 2								96															

# List of Tables

1.1	Comparison of species catchability during day and night trawls	17
1.2	Influence of depth on diel catchability. $\ldots$	22
2.1	Number of research survey tows in which each species was caught.	32
2.2	Comparison of research survey vessel and gear characteristics	34
2.3	Conversion factors for diel catchability and the relative catchability	
	of species	35
A.1	Sample data set to be used with Model 2 (Chapter 1). $\hdots$ .	74
B.1	Number of tows per stratum completed on the southern Grand	
	Bank, 1951-1995	76
B.2	Number of tows per stratum completed on the St. Pierre Bank,	
	1951-1995	77
C.1	Southern Grand Bank Data	95
D.1	St. Pierre Bank Data.	113

# List of Figures

1.1	Map of NAFO areas.	9
1.2	Frequency of the proportion of nominally significant tests. $\hdots$ .	14
1.3	Estimates of the day/night effect for all region and season combi-	
	nations (as in Table 1.1)	20
1.4	Comparison of the parameter estimates combined over all regions	
	and seasons for Model 1 and Model 2	21
2.1	Map of research survey trawl locations	29
2.2	Estimate of total biomass from southern Grand Bank $\hdots$	40
2.3	Estimate of total biomass from St. Pierre Bank $\ldots$	41
2.4	Estimates of biomass for species of the southern Grand Bank	42
2.5	Estimates of biomass for species of the St. Pierre Bank $\ \ldots \ \ldots$	43
2.6	Species diversity	45
2.7	Cumulative species biomass	46
2.8	Compensation by pelagic species	49
3.1	Map of the Northwest Atlantic Fisheries Organization	54
3.2	Estimates of absolute biomass for barndoor skate	58

#### Introduction and Overview

Fish populations have been exploited for centuries, with the expansion from simple food fisheries to extensive commercial fisheries (Sahrhage and Lundbeck 1992). At a time when many fish stocks around the world have been overexploited (Hilborn and Walters 1992; Larkin 1977) and the world's fishery harvest has reached one million metric tonnes (Pauly 1996), serious consideration must be given to the effect exploitation has, not only on commercial species, but on the marine community as a whole.

The natural variability observed in many fish populations (Caddy and Gulland 1983) may result from fishing pressure, changing environmental conditions, or from species interactions, such as competition and predation (Sissenwine 1984). The importance of species interactions are becoming better recognized, with multispecies studies rapidly replacing the single-species approach to fishery assessment (Kerr and Ryder 1989). Fisheries harvest more than the targeted species, as evidenced in by-catch statistics (Alverson et al. 1994), and thus are multispecies in nature. In the North Sea, however, a recent study has shown that non-targeted groundfish communities have remained stable in spite of intensive fishing (Greenstreet and Hall 1996). A similar result was observed for non-targeted deep-water fish assemblages in the Northwest Atlantic (Haedrich 1995).

Changes in species composition may occur as a result of competitive or predatorprey relationships which are precipitated by changes in environmental conditions or fishing pressure. One species may be better adapted to changing environmental conditions and thrive, thus out-competing the other species for food and space. In the Northeast Pacific, the sardine (Sardinops caerulea) was replaced by a competitor, the northern anchovy (Engraulis mordax), as a result of changes in the oceanographic environment that favoured the anchovy (?). A similar interaction was observed between the Atlantic herring (*Clupea harengus*) and Atlantic mackerel (*Scomber scombrus*) populations in the Northwest Atlantic (Skud 1982).

Fisheries tend to target the largest, most abundant species which are higher up the food chain (Deimiling and Liss 1994). Removal of these larger species enables smaller species to expand their populations. Analysis of the Great Lakes' fish populations has shown a shift from large, long-lived species with higher commercial value, to small, short-lived species which are less preferred (Smith 1968; Steedman and Regier 1990). Intensive exploitation in the North Sea has resulted in the decrease in the number of fish from the larger length classes (Anon 1996). A similar result was observed on Georges Bank following the heavy exploitation of the 1960's and 1970's. Recently the proportion of fish from larger length classes has increased, due primarily to the increase in elasmobranchs (Murawski and Idoine 1992).

The distinction between the effects of exploitation and the environment is not easily made (Harris and Poiner 1991), and often effects due to changing environmental conditions may be further exacerbated by exploitation (Ludwig et al. 1993). Furthermore, environmental conditions, while being difficult to predict, cannot be controlled (Walters and Collie 1988). Alternatively, fishing effort is easily predicted and controlled.

In the Northwest Atlantic, commercial fisheries date back to the 1400's and over the following 500 years the fishery expanded from inshore to offshore. The Inshore fishery had been more successful than the offshore until the mid-1950's when factory-freezer-trawlers first arrived on the Newfoundland Grand Banks (Hutchings and Myers 1995). The incredible increase in fishing effort, due to the power and storage capabilities of the factory freezer trawlers, was evidenced by the fact that the total amount of fish caught in two thirty-minute trawls was equal to the total amount of fish harvested in one year by the French in the 1500's (Warner 1983). The number of factory freezer trawlers fishing off the coast of Newfoundland increased into the 1960's and 1970's. In 1977, fishing by foreign fleets was reduced with the establishment of Canada's 200 mile limit (Pinhorn and Halliday 1990). This decrease in foreign fishing effort, however, was not matched by a decrease in effort from Canadian vessels, and in 1992 a moratorium was imposed in response to drastically depleted cod stocks off the coast of Newfoundland (Bishop et al. 1993).

Recent studies in the eastern North Atlantic (NAFO Divisions 2J3K) suggest that there has been a gradual decline in species abundance since 1978 (Gomes et al. 1995; Haedrich 1995; Villagarcia 1995). Total species abundance has remained fairly constant on the Newfoundland Grand Banks (Gomes et al. 1992) and Georges Bank (Fogarty and Murawski 1998) since 1971 and 1963, respectively. On Georges Bank this stability has resulted from an increase in the population size of elasmobranchs (Murawski and Idoine 1992). Both study periods began after the arrival the factory-freezer-trawlers, and thus may have missed the population maxima during the low exploitation period.

The main purpose of this study is to reconstruct the population history of fish species from the Southern Grand Bank and St. Pierre Bank back to the early 1950's, prior to the arrival of factory-freezer-trawlers. Research vessel survey data from the Department of Fisheries and Oceans will be examined to initially determine if the change in the timing of the surveys, from daylight hours to twentyfour hours a day, significantly effects the catchability of fish species (Chapter 1). Prior to 1972, research surveys in this region were primarily conducted during daylight hours. Without this correction, estimated abundance of species with variable patterns of diel catchability may be inaccurate.

In Chapter 2 this information will be used to calculate estimates of absolute abundance of groundfish species to determine if relative species abundance has been altered over the time period. The structure of the groundfish community will be examined in terms of changes in species diversity and dominance throughout this period of intense exploitation.

Finally, Chapter 3 concentrates on one species, the barndoor skate, and the effect that the fishery has had on this non-commercial species. This chapter shows the importance of assembling data on as long a time-scale as possible and as wide a spatial scale as possible. Failure to examine such historical data has allowed this species to disappear without anyone noticing.

## **Co-authorship Statement**

Chapter 1 was published in the Canadian Journal of Fisheries and Aquatic Science (1998) 55: 2329-2340 and Chapter 3 was published in Science (1998) 281: 690-692. Both studies were co-authored by Dr. R. A. Myers who suggested the proper analyses for each chapter. I researched the literature, manipulated the data, completed the analyses and wrote the text under Dr. Myers direct supervision.

# Chapter 1

# Diel variation in trawl catchability: Is it as clear as day and night?

## 1.1 Abstract

Diel variation in the catchability of over 50 species was examined using research vessel surveys conducted off the coast of Newfoundland and Labrador from 1972 to 1995. Catchability during the day and night was estimated for several seasons and geographic areas in the northwest Atlantic using two generalized linear models. In general, species exhibiting diel vertical migrations, such as redfish, northern sand lance and haddock, were caught in higher proportions during the day. Nonmigrating species, such as flatfish, skate and scuplin, which rely on visibility of the traw as a means of escapement, were caught in higher proportions during the day, night. Analysis of the effect of depth indicated that catchability during the day, relative to the night, increased significantly with depth for 21 species. This study demonstrates that the accepted methods of estimating standard errors for generalized linear models are not valid for survey trawl data and suggest alternative methods.

## 1.2 Introduction

Many countries conduct bottom trawl surveys for the purpose of estimating abundance of commercial species. Determining the efficiency of survey gear, however, is a key component of estimating abundance and interpreting these survey results. The survey vessel, time of year and time of day affect the survey trawl efficiency. Fish behaviour in response to physical factors, such as light intensity, has been shown to affect the efficiency of the survey gear over a 24 hour period (Walsh 1991). Abundance of species exhibiting diel vertical migration would be overestimated during surveys conducted during the day as compared with surveys conducted at night, when these species would be out of the vertical range of the trawl (Michalsen et al. 1996). Increased visibility of the trawl during the day (Glass and Wardle 1989) could result in lower catches of non-migrating species and thus abundance would be underestimated. Consequently, estimated abundance of species with variable patterns of diel catchability may be inaccurate. Analysis must be carried out to determine if die varbility in catchability exists, and if so, to correct for this difference.

Several studies of diel catchability have made use of designed experiments in which a predetermined number of tows was carried out under controlled conditions (Walsh 1991; Walsh and Hickey 1993; Engås and Soldal 1996). While important information on fish behaviour in the vicinity of survey gear has been obtained, the number of tows completed has often been too few to give precise estimates. Furthermore, the results may not take into account regional or seasonal changes in catchability.

The purpose of this paper is to present a simple method that allows the relative efficiency of fishing gear during the day and night to be rigorously estimated using data routinely collected during research surveys. This approach combines data from many years and several geographic areas.

A further motivation for this work is to provide correction factors for older research surveys of the Canadian Department of Fisheries and Oceans, many of which were conducted almost exclusively during daylight hours. As data collected from commercial vessels are thought to be unreliable, more emphasis is being placed on research vessel surveys. Correction factors for diel variation in trawl catchability are essential in determining absolute abundances from these surveys.

### 1.3 Methods

#### 1.3.1 Data

Research surveys have been conducted by the Canadian Department of Fisheries and Oceans off the coast of Newfoundland since 1946 (Templeman 1966). Prior to 1972, however, surveys were primarily conducted during daylight hours. We examined stratified random surveys conducted around the Island of Newfoundland and the coast of Labrador (Pitt et al. 1981) from 1972 (when both day and night surveys were conducted in roughly equal proportion) to 1995 during which time approximately 20,000 research tows were successfully made. Only tows with a duration of 30 minutes were selected for this analysis. Data from seven Northwest Atlantic Fisheries Organization (NAFO) subdivisions (Figure 1.1), each with different bottom topographies, were included in this study. Seasons of the surveys are given in Table 1.1. Species were selected for analysis if they occurred in more than 100 tows, and only strata in which individual species were caught in more than five tows were included. Sunrise and sunset were determined using the day of year and latitude of the tows (Brock 1981). Tows which occurred within one hour of sunrise and sunset were excluded so that successive tows would not be compared (e.g. the last tow of the night and the first tow of the day).



Figure 1.1: Map of NAFO areas around Newfoundland. The 100 (dotted line), 200 (narrow dash), and 300 (wide dash) meter isobaths are given for reference.

#### 1.3.2 Model

#### Model 1

Consider a survey in year y, in which  $C_{y,s,l}$  fish are caught in stratum s during the time of day (i.e. day or night) l. The catchability associated with time of day l is  $S_l$ , where the catchability will be scaled so that the catchability during the night will be 1. We construct a simple model in which an equal proportion of fish will be in each stratum in all years. The expected value of the catch,  $E[C_{y,s,l}]$ , is

$$E[C_{y,s,l}] = N_y P_s S_l$$

where  $N_y$  is the number of fish in the population in year y,  $P_s$  is the proportion of fish stratum s, and  $S_l$  is the combined availability and vulnerability of each species to the survey gear (or catchability) associated with time of day, l. The simplest model for the probability of catching  $C_{y,s,l}$  fish is a Poisson distribution. This is not a realistic model, however, because fish usually aggregate (i.e. in schools) and, as such, are not captured independently. Also, habitat within a stratum is not equally suitable. An over-dispersed, i.e. extra-Poisson model, is preferred in which over-dispersion is modeled using a scale factor for the variances (McCullagh and Neider 1989). The scale factor only affects the variance, but not the parameter estimates. The data can be analyzed in terms of a generalized linear model (GLM) with a log link.

The main assumption of this model is that an equal proportion of fish will be in each stratum in all years. This may not be a valid assumption, however, as it is unlikely that the distribution of fish will remain constant from year to year.

#### Model 2

To remove the assumption of a constant stratum and year effect, the data considered will be restricted to strata within years in which at least one day and night tow has occurred.

Let  $G_{ysd}$  be the total catch in numbers during the day in a particular year and stratum, and  $G_{ys}$  be the total catch for the day and the night in the same year and stratum.  $T_{ysd}$  and  $T_{ysn}$  will be the corresponding number of day and night tows, respectively. Let the probability that a fish caught in a year, y, and stratum, s, is caught during the day be  $p_{ys}$ , and  $1 - p_{ys}$  for night catches. If there is no difference between day and night catchability, then we would expect

$$\frac{p_{ys}}{1 - p_{ys}} = \frac{T_{ysd}}{T_{ysn}}$$

We are interested, however, in whether or not a different proportion of fish are caught during the day and night, and so the catchability term,  $S_d$ , is multiplied to the right hand side of the above equation. After a log transformation we have

$$\log\left(\frac{p_{ys}}{1-p_{ys}}\right) = \log(S_d) + \log\left(\frac{T_{ysd}}{T_{ysn}}\right)$$

The left side of the equation represents the logit transformation  $p_{\mu\nu}$ . The term on the far right is the offset (McCullagh and Neider 1989), a known quantitative variate, which will account for the number of day and night tows. The log of the catchability during the day,  $S_d$ , is the intercept and will be estimated. Positive estimates indicate higher daytime catchabilities, while negative estimates indicate higher catchabilities at night.

If fish of a given species are captured independently, and if the probability of catching a fish during the day is constant for individuals of that species, then the probability of catching a fish during the day in a particular year and stratum  $(C_{yyd})$ , given the total number of fish caught in that year and stratum  $(C_{yy})$ , is binomial. Over-dispersion must also be taken into account with Model 2. An extra-binomial model is preferred in which over-dispersion is modeled using a scale factor for the variances (McCullagh and Nelder 1989). A GLIM with a logit link and an offset is used to analyze the data. The method of programming is given in the Appendix A.

This model also makes a potentially unrealistic assumption that the same proportion of fish will remain in a stratum during the survey period for a given year. Many species make seasonal migrations which take them into and out of the arbitrarily assigned strata, and as such would not be expected to restrict their distribution during the year.

It is not clear that either of the above models is superior as both make assumptions that may not be valid for research survey data. For simplicity, Model 2 was chosen for this analysis, although reference will be made to the results of Model 1 as a method of comparison.

# 1.3.3 Regional and Seasonal Differences in Diel Catchability

Differences in bottom topographies associated with each NAFO area could possibly affect diel catchability. Banks within the study area range from 150-300m in depth surrounded by troughs of 450-500m in the north, to depths of 100m with 150-200m deep troughs in the south (Litvin and Rvachev 1963).

Seasonal changes in diel catchability have been suggested for various groundfish species (Beamish 1965). Since the degree of vertical migration within the water column may vary seasonally, research surveys conducted in different seasons were examined separately.

This analysis examines nine region and season combinations over which estimates of diel catchability will be compared.

#### 1.3.4 Randomization Tests

The reliability of confidence limits and significance tests for the day/night effect was assessed using a randomization test (Manly 1991). For each area/season combination studied (Table 1.1), the catchability of day tows was estimated using Model 2 and the original data, with half of the data randomly assigned to the day and half to the night. Significance was determined by calculating the 95% confidence interval based on the standard errors of the GLIM estimates. The proportion of significant tests from 100 randomizations was then cletermined.

The randomization tests showed that the standard significance levels and standard errors of the parameter estimates were not reliable. Figure 1.2 shows the proportion of significant tests from the 100 randomizations for each species and region/season combination. With a significance level of 5%, the proportion of nominally significant tests should have been 0.05. For only 33% of the species and region/season combinations, however, was the proportion of significant tests less than or equal to 0.05.

Fish species which are known to school, such as capelin, redfish, Atlantic cod and arctic cod, tended to have a higher proportion of significant tests, while with solitary species, such as wolffish, the proportion of significant tests was closer to 0.05. In subdivision 3M, where the number of species and number of observations were the lowest for all subdivisions, the proportion of significant tests tended to be lower. In subdivisions with many observations, such as 3L and 3N O, the opposite was true.



Figure 1.2: Frequency of the proportion of nominally significant tests obtained from the randomizations for all species and region/season combinations (as in Table 1.1). The frequency is given as the percentage of species and region/season combinations (n=266).

Alternative estimates of reliability were constructed as the standard deviation of the mean of the parameter estimates from the 100 randomizations. These estimates of reliability in Table 1.1 represent the standard errors under the null hypothesis (i.e. no difference between day and night catchability), and are used in hypothesis testing (Manly 1991). It was decided that 100 randomizations were sufficient as this study deals with many region/season combinations. The main interest of this study is to examine trends across all region and season combinations rather than the reliability of a single estimate.

#### 1.3.5 Depth Effects

Depth, which has been thought to contribute to the variability in diel catchability (Pitt 1967), was examined using Model 2. Tows were categorized into approximately 100m depth classes that corresponded with the average depth of the strata included in this study (Bishop 1994). The model was run initially as a regression analysis with depth as a covariate. Since the standard significance tests are considered to be liberal, any species for which depth was not significant was eliminated. For the remaining species, estimates of reliability for the depth effect were then determined with the depth categories randomly assigned in the same proportion as would normally be observed for each species. For example, it would be unlikely that a species normally found at shallow depths would be caught in equal proportions at the greatest depths. The standard deviation of the estimates from 100 randomizations were used in hypothesis testing. Estimates of diel catchability in each depth class were then determined using Model 2.

## 1.4 Results

Estimates of diel catchability were obtained for all region/season combinations (Table 1.1 and Figure 1.3). Negative estimates indicated higher catchabilities at night. The summary refers to estimates determined over all regions and seasons. Of the 32 species with significant differences over all regions and seasons, 12 were caught significantly more during the day and 20 were caught significantly more at night.

Estimates for the family Gadidae differed among species. Arctic cod was caught significantly more during the day in subdivision 3L in both the fall and spring surveys, while the estimate for subdivision 3Ps indicated a higher night catchability. The estimates for Atlantic cod were also variable with higher day catchabilities in subdivisions 3M and 3NO and a higher night catchability in subdivision 2J. Haddock was caught significantly more during the day, while the opposite was true for longfin hake. Only longfin hake had significant results across all regions and seasons.

All of the flatfish species included in this analysis had significant differences in diel catchability, although no consistent pattern among the species could be found. Witch flounder was caught significantly more during the night in 3L (summer only) and 3NO. Higher daytime catchability for witch founder was found in subdivision 2J only. Atlantic halibut and Greenland halibut were caught significantly more during the day, while yellowtail flounder was caught significant more at night. Estimates for American plaice differed regionally with catchability in 3NO and 3Ps spring surveys higher at night. In the north, catches were higher during the day.

and a second descent of the second second	and the second second									
Species	2J Fall	3K Fall	3L. Fall	31L Spring	3L Summer	3M Spring	3NO Fali	3NO Spring	3Ps Spring	Summary
Herring								2.02(0.53)	0.81(0.22)	0.91(0.22
Aretic ond	0.24(0.23)	-0.25/0.16)	0.5(0.14)	0.38(0.17)	-0.04(0.53)		0.95(1.22)		-1.12(0.24)	0.27(0.08
Atlantic oud	0.23/0 08	-0 16/0 14	-0 15(0 1)	1000000	0.09/0.14	1.16(0.43)	0.28(0.11)	0.23(0.09)	-0.26(0.15)	0.05/0.05
Campbed and	100.000	firmon o	devolution.	for alara.	(unalored)	(manharer)	()	0.0000000000000000000000000000000000000	for other on	CV 0/86 0
Greenland cod	-0.28(0.38)						-			-0.28(0.42
Haddock			-1.29(1.08)		1(0.97)		-1.32(0.83)	0.87(0.25)	0.34(0.17)	0.51(0.2
Silver hake							-0.99(0.66)	0.02(0.24)	0.37(0.2)	0.26(0.18
Pollock								0.99(0.75)	0.11(0.36)	0.2(0.32)
Longfin hake						-0.95(0.16)	-1.55(0.73)	-1.01(0.26)	-0.42(0.11)	-0.58(0.09
White hake						100000	-0.18(0.46)	0.24(0.15)	0.01(0.09)	0.04(0.07
Roundnose grenndier	-0.9(0.65)	-1.38(0.59)								-1.3(0.49
Roughlead greandier	0.08(0.06)	0.09(0.05)	0.41(0.13)	-0.12(0.09)	0.35(0.14)	0.1(0.11)	0.87(0.48)	0.15(0.16)		0.19(0.04
Marlin-spike	0.21(0.24)	-0.14(0.1)	0.11(0.16)	0.39(0.26)	-0.1(0.24)	-0.27(0.1)	-0.5(0.31)	-0.39(0.18)	0.04(0.05)	-0.02(0.04
Blue hake	-0.72(0.42)	-0.43(0.32)				-0.14(0.18)				-0.27(0.14
Monkfish								1.12(1.33)	0.19(0.12)	0.14(0.12
Lanternfishes	0.12(0.22)	0.78(0.49)	-0.25(0.27)	3.56(0.69)	1.84(3.94)	-0.4(0.43)		-1.22(1.07)	-0.6(0.43)	-0.18(0.18
Barracudina	-0.35(0.54)	0.09(0.3)	1.07(0.43)	0.4(0.25)	0.05(0.26)				0.43(0.3)	0.48(0.16
Longnose cel	-0.16(0.44)	-0.28(0.5)								-0.24(0.37
Atlantic hagfish									-0.65(0.18)	-0.65(0.2
Spiny cels		-0.74(0.38)	0.75(0.79)	0.11(1.17)	1.47(0.98)	-0.58(0.29)				-0.19(0.24
Northern alligatorfish	-0.34(0.14)	-0.15(0.11)	-0.44(0.15)	-1(0.2)	-0.2(0.6)					-0.56(0.11
Common alligatorfish	-2.69(0.93)		-1.06(0.27)	-0.81(0.48)			-2.38(0.98)			-1.01(0.21
Northern sand lance			1.09(0.3)	1.78(0.42)	1.91(2.34)		2.43(0.38)	2.7(0.4)	3.58(2)	1.59(0.19
Broadhead wolffish	-0.2(0.05)	-0.02(0.08)	0.16(0.12)	-0.11(0.23)	0.01(0.18)	-0.14(0.19)				-0.11(0.04
Stripod wolffish	-0.09(0.08)	-0(0.05)	0.20(0.11)	-0.03(0.1)	-0.12(0.16)	-0.87(0.09)	-0.82(0.32)	-0.71(0.12)	-1.19(0.19)	-0.29(0.04
Spotted wolffish	-0.12(0.09)	0.32(0.15)	-0.17(0.14)	0.07(0.13)	0.09(0.25)	-0.63(0.12)		2(0.77)		-0.09(0.06
Hookear sculpin	-1.73(0.59)	-0.89(0.13)	-2.13(0.24)	-1.87(0.55)						-2.02(0.2
Arctic Deepsea sculpin	-0.63(0.23)	0.23(0.16)	-0.14(0.78)	0.56(0.44)						0.01(0.11
Sen Raven				0.25(0.3)			-1.08(0.26)	-0.52(0.08)	-0.51(0.08)	-0.56(0.06

Table 11 Comparison of species exclusibility during day and night travels. The number in parentheses (roles to the standard errors compared using a randomization procedure (see eac). Estimates for all regions and assess (Summary) are provided with standards errors. Positive estimates indicate higher excluses aftering the dury relative to the right. Significant estimates are given in hold. For a list

Species	2.J Fall	3K Fall	3L Full	3L Spring	3L Summer	3M Spring	3NO Fall	3NO Spring	3Ps Spring	Summary
Longhorn sculpin Shorthorn sculpin			-3.55(1.06)	-0.56(0.55)			-1.33(0.24)	-1.2(0.31)	-1.84(0.44)	-1.44(0.19
Mailed sculpin	-1.96(0.38)	-1.39(0.63)	-0.83(0.17)	-0.78(0.25)	-1.82(0.29)		-0.52(0.27)	-1.12(0.29)	-1.13(0.33)	-0.78(0.14)
Common lumpfish	1.29(0.42)	0.95(0.49)	1.84(0.67)	0.45(0.27)				1.21(0.24)	0.56(0.2)	0.57(0.14)
Spiny lumpfish	-1.52(0.23)	100	-1.21(0.12)	-2(0.2)	-2.13(0.64)		-1.95(0.54)	-1.83(0.33)	-2.68(0.28)	-1.54(0.08)
Seasnails	-1.17(0.87)	-1.96(0.65)	-1.28(0.16)	-0.95(0.25)			-3.75(1.59)			-1.31(0.14)
Rodfish	0.65(0.26)	0.28(0.38)	0.59(0.23)	0.2(0.16)	1.01(0.37)	0.94(0.15)	0.99(0.3)	1.24(0.23)	0.63(0.08)	0.8(0.07)
Fourline snakeblenny			-1.37(0.7)							-1.37(0.62)
Snake blenny			0.51(0.28)				0.36(0.44)			0.43(0.27)
Shanny			0.28(0.32)	-1.05(1.05)						0.12(0.25)
Arctic celpout	-0.26(0.09)	-0.46(0.12)	-0.7(0.11)	-0.38(0.08)	-0.44(0.09)		-0.12(0.39)	-0.53(0.15)	-0.27(0.18)	-0.54(0.06)
Eclpout (Lycodes sp.)	0.28(0.21)	0.19(0.29)	-0.86(0.68)	0.13(0.07)				-0.35(0.56)		0.1(0.07
Vahi's colpout	-0.18(0.13)	-0.24(0.06)	-0.19(0.1)	-0.1(0.11)	-0.58(0.16)	-0.06(0.16)	-1.79(1.48)	0.42(0.27)		-0.21(0.04
Witch flounder	0(0.1)	0.25(0.1)	0.19(0.16)	0(0.11)	-0.32(0.1)	0.28(0.16)	-0.59(0.14)	-0.62(0.13)	-0.09(0.08)	-0.1(0.05
American plaice	0.3(0.09)	0.16(0.05)	0.38(0.08)	0.39(0.05)	0.23(0.08)	0.21(0.13)	0.03(0.07)	-0.38(0.07)	-0.28(0.11)	0.2(0.03
Atlantic halibut								0.26(0.13)	0.41(0.16)	0.34(0.12)
Yellowtail flounder			-0.76(0.19)	-0.68(0.23)	-0.62(0.28)		-1.02(0.17)	-1.06(0.11)	-0.32(0.22)	-1(0.1
Greenland halibut	-0.01(0.06)	-0.07(0.05)	0.28(0.07)	0.32(0.08)	-0.08(0.1)	-0.05(0.09)	0.78(0.15)	0.1(0.11)	0.02(0.08)	0.02(0.03
Thorny skate	-1.07(0.12)	-0.64(0.06)	-0.8(0.08)	-0.75(0.09)	-0.77(0.12)	-1.19(0.13)	-0.74(0.08)	-0.98(0.12)	-1.15(0.13)	-0.87(0.04
Smooth skate	-0.64(0.24)	-0.61(0.1)	-0.8(0.36)	-0.13(0.29)		-1.49(0.43)	-1.54(0.64)	-1.03(0.36)	-0.43(0.1)	-0.62(0.07
Atlantic argentine								2.03(0.64)	0.83(0.2)	0.93(0.19
Capelin	0.59(1.21)	0.32(0.19)	0.23(0.13)	1.82(0.2)	1.33(0.62)		1.16(0.78)	0.34(0.36)	1.4(0.74)	0.95(0.12
Black dogfish	-0.03(0.25)	-0.66(0.85)				-0.47(0.47)			0.11(0.06)	0.1(0.06
Spiny dogfish						2 2		1.39(3.65)	-0.34(0.51)	-0.33(0.49
and the second s								A DESCRIPTION OF A DESC		

Table 1.1 Continued

Higher catchabilities during the day were found for several deepwater species, such as Atlantic argentine and roughhead grenadier, and pelagic species, such as capelin and herring. Redfish and northern sand lance, which have higher daytime catchabilities, have a demersal existence during the day, but become pelagic feeders at night (Scott and Scott 1988).

Estimates for each species of lumpfish were consistent for all regions and seasons. Common lumpfish catches, however, were significantly higher during the day, while spiny lumpfish had a significantly higher catchability at night across all regions and seasons.

For species of wolffish, sculpin, skate and eelpout, catches were higher at night. Both striped and spotted wolffish, however, also exhibited higher day catchabilities in subdivision 3L (fall only) and 3K surveys, respectively.

Parameter estimates for catchability during the day were generally not different for Models 1 and 2 (Figure 1.4). The only exceptions were roundnose grenadier and northern sand lance. The fact that two models, using two variations on the same data set (Model 2 used a much smaller proportion of the Model 1 data set), would have such similar results, suggests that the results of this analysis are believable.

Significant estimates for the effect of depth on diel catchability were obtained for 24 species (Table 1.2). For the majority of these species, depth was positively associated with higher catchabilities during the day relative to the night. At shallow depths, these species were more likely to caught in greater proportions during night tows, but at greater depths, catchability was likely to be higher during the day. Only three species had a negative association with depth.

Catchability was not affected by the type of vessel used over the time period. Estimates of diel catchability obtained for each region, season, and vessel combination did not differ significantly from those obtained using only region and season combinations.



Figure 1.3: Estimates of the day/night effect for all region and season combinations (as in Table 1.1). Negative results indicate higher catchability at night. Significant (+) and non-significant (O) results are differentiated and a reference line at zero is provided for comparison.



Model 1 Estimates (log(Sd))

Figure 1.4: Comparison of the parameter estimates combined over all regions and seasons for Model 1 and Model 2. Each point represents an individual species. A one-to-one reference line is also given.

Table 1.2: Influence of depth on diel catchability. The second column is the estimate of the slope from the regression analysis, where positive estimates indicate higher catches during the day relative to the night as depth increases. The number in parentheses refers to the standard error under the null hypothesis (as in Table 1.1). Only those species with significant estimates are aboven. Estimates of diel catchability in each depth zone, which correspond to average depths of the strata, are given for comparison. Significant estimates are given in bold.

			Dept	th Zone (in me	ters)	
Species	Estimate of the depth effect	< 91	92-183	184-274	275-366	367-549
American plaice	0.39(0.03)	-0.28(0.06)	0.42(0.06)	0.51(0.1)	0.24(0.11)	0.28(0.28)
Arctic cod	0.29(0.12)	-0.47(0.32)	0.35(0.15)	0.22(0.18)	0.93(0.43)	
Arctic eelpout	0.38(0.09)	-1.09(0.13)	-0.58(0.11)	-0.14(0.15)	-0.38(0.15)	
Atlantic hagfish	0.99(0.35)		-3.33(19.1)	-1.05(0.48)	-0.13(0.4)	0.21(21.91)
Barracudina	0.56(0.25)			-0.16(0.43)	0.46(0.31)	0.96(0.26)
Capelin	-0.44(0.13)	1.41(0.44)	1.12(0.2)	0.2(0.21)	0.72(0.43)	0.49(9.24)
Common alligatorfish	1.06(0.29)	-3.41(0.69)	-1.25(0.3)	-0.05(0.39)	-0.11(0.53)	
Common lumpfish	-0.52(0.21)	1.28(0.36)	0.34(0.31)	0.87(0.67)	-0.1(0.51)	0.66(0.34)
Hookear sculpin	0.84(0.22)	-2.38(0.75)	-2.18(0.27)	-0.55(0.3)	-0.34(0.54)	
Longfin hake	0.46(0.14)			-0.96(0.32)	-0.82(0.16)	-0.12(0.12)
Longhorn sculpin	0.89(0.37)	-1.48(0.22)	-0.39(0.47)	-1.42(5.33)		
Longnose cel	-1.87(0.52)				1.11(17.02)	-0.76(0.7)
Marlin-spike	0.15(0.07)		-0.09(14.34)	0.02(0.14)	-0.27(0.14)	0.07(0.07)
Northern alligatorfish	0.67(0.22)	-1.52(3.84)	-1.17(0.21)	-0.39(0.31)	0.15(0.28)	
Sea raven	0.39(0.11)	-0.72(0.08)	-0.35(0.14)	0.13(0.47)		
Seasnails	0.79(0.23)	-4.05(0.35)	-1.22(0.21)	-0.84(0.2)	-0.28(0.27)	
Shorthorn sculpin	0.94(0.29)	-0.22(0.17)	0.72(0.22)			
Smooth skate	0.32(0.09)	-1.33(9.36)	-0.99(0.12)	-0.59(0.13)	-0.3(0.18)	-0.06(0.23)
Spiny eels	1.79(0.35)				-1.46(0.71)	0.33(0.29)
Spiny lumpfish	0.99(0.18)	-2.51(0.18)	-1.43(0.12)	-0.55(0.41)		
Striped wolffish	0.22(0.05)	-0.77(0.12)	-0.37(0.08)	-0.16(0.07)	0.01(0.09)	-0.78(0.56)
Thorny skate	0.14(0.04)	-0.93(0.1)	-0.99(0.07)	-0.82(0.08)	-0.44(0.11)	-0.13(0.17)
Vahl's eelpout	0.16(0.06)	0.09(6.55)	-0.49(0.17)	-0.21(0.08)	-0.06(0.1)	-0.06(0.14)
Witch flounder	0.22(0.06)	-0.96(0.17)	-0.24(0.17)	-0.11(0.1)	0.1(0.13)	0.17(0.11)
# 1.5 Discussion

The observed diel variation in catchability may be attributed to patterns of vertical migration, and escapement associated with visibility of the trawl. Redfish and northern sand lance are known to migrate vertically at night in search of food high in the water column (Scott and Scott 1988). The high catchability during the day for both species is consistent with this type of diel vertical migration. Atkinson (1989) and Beamish (1965) found similar results for redfish in the northwest Atlantic.

Previous studies of Atlantic cod and haddock have shown that catches were higher during the day (Engås and Soldal 1996; Walsh and Hickey 1993; Michalsen et al. 1996). Using acoustic techniques Beamish (1965) found that both species were found in higher concentrations on the bottom during the day. Our results for haddock are consistent with these studies. For Atlantic cod, however, diel catchabilities were variable among regions. The previous studies considered surveys conducted over a small geographic area and a short period of time and thus may not represent the variability of these species over their entire range.

Visibility is important to the ability to escape the trawl during daylight hours for some species (Glass and Wardle 1989), e.g. underwater video cameras have shown that some flatfish are able to avoid the trawl during the day (Walsh 1988; Walsh and Hickey 1993). In this study, however, not all flatfish showed higher catchabilities during night tows. Atlantic halibut, a deepwater species, was caught significantly more during the day indicating that this species may be migrating up into the water column at night, a previously undocumented phenomenon. The same may be true for American plaice found in deep water. Walsh (1991) carried out experiments on the Grand Banks and found higher catchabilities of American plaice at night, while Beamish (1965), who examined data from the Scotian shelf and the Gulf of St. Lawrence, found the opposite. These regional differences, however, may be confounded by the effects of depth on diel catchability. In our analysis, estimates of diel catchability for American place were higher during the day in the northern regions which also have the deepest strata. In the south, where strata are shallower, catchabilities were higher at night.

Analysis of the influence of depth indicated that catchability during the day, relative to the night, increased significantly with depth for 21 species. When diel catchability was examined for each depth zone, the greatest differences were between the shallowest depth (< 91m) and the rest of the zones. If light does not penetrate farther than the shallowest depth, then the diel catchability should not necessarily differ beyond this depth. This increase in catchability during the day with depth may be the result of diel vertical migrations.

This study has shown that there are clear differences between day and night catchabilities for many species in the northwest Atlantic, and that depth and geography are also important factors. Size and abundance of species, although not considered in this analysis, may also be important in determining the diel variation in trawl catchability (Korsbrekke and Nakken 1997). These differences should be taken into account in the assessment of fish stocks to improve estimates of abundance.

Stratified random surveys are conducted annually for the purpose of estimating abundance of commercial species. Our analysis has shown that these data can be used to augment other studies which require the use of specified controlled experiments. Research surveys provide many replicates of data necessary for rigorous hypothesis testing and should not be overlooked as a source of valuable information. Our results will allow the time series of groundfish research surveys to be extended back 45 years which was previously not possible because the earlier surveys occurred mostly during the day. Such an analysis is crucial for resolving major ecological issues (Greenstreet and Hall 1996).

An important general statistical issue has been raised as a result of this analysis. The standard methods for dealing with extra-Poisson and extra-binomial variation in generalized linear models (GLIM's) do not appear to provide an adequate basis for inference and hypothesis testing when dealing with stratified random research survey data. Our randomization testing clearly demonstrated that the variation in both models could not be explained by the scale factor, which inflates the binomial and Poisson variances in GLIM's. We suggest that for research survey data, and perhaps for other highly variable fisheries data, that randomization testing be undertaken to demonstrate the reliability of inferences. We have demonstrated, however, that by examining data from many geographical areas, the reliability of inferences can often be determined.

# Chapter 2

Fish Community Changes in an Exploited Marine Ecosystem: Newfoundland Southern Grand Bank and St. Pierre Bank, 1951-1995

# 2.1 Abstract

Longterm fish community changes on southern Grand and St. Pierre Banks were examined from dedicated research survey trawls conducted by the Canadian Department of Fisheries and Oceans from 1951-1995. These time series pre-date the arrival of factory-freezer-trawlers in the 1960's and 1970's, and thus provide insight into changes in the biomass and diversity of a heavily exploited system. The time series were standardized by applying conversion factors for changes in vessel, diel changes in catchability, and relative catchability of selected species. Total biomass in the 1990's was reduced to 11% and 9% of that observed in the 1950's on southern Grand Bank and St. Pierre Bank, respectively, and largely resulted from the decline of the haddock population. Compensatory responses to this decline were visible with the flatfish on southern Grand Bank and skate on St. Pierre Bank but continued fisheries for flatfish and bycatch of skate kept total species biomass at low levels.

#### 2.2 Introduction

At a time when many fish stocks around the world have been overexploited serious consideration must be given to the role that fisheries play in the structuring of marine communities. Fisheries often target the largest and most abundant species which are higher up on the food chain (Deimling and Liss 1994). The removal of these top predators enables species at lower trophic levels to compensate for this loss of fish biomass. Fisheries, however, have become quite adaptive and efforts have often concentrated on the next trophic level when the catch per unit effort at the highest trophic level declines. This phenomenon has recently been referred to as "fishing down marine food webs" (Pauly et al. 1998).

The concept that total fish biomass remains constant over time, through species compensation, is over 50 years old (Allee et al. 1949). Two widely known examples have lent credibility to this idea. In the Great Lacks, the total weight of the catch remained relatively constant over the last 150 years (Regier and Hartman 1973). This is a case of species compensation where large predators, such as lake trout, are replaced by smaller, often introduced, species. This finding, lowever, is based upon commercial catches, and not true abundance. Consequently, trends in absolute biomass are difficult to interpret. The second is the observation that the total biomass of demersal fish on Georges Bank remained relatively constant from 1963 to 1990, with sharks and skates compensating for the loss of gadoid and flatfish biomass (Fogarty and Murawski 1998). By 1963, however, the biomass of the most desirable species on Georges Bank, such as halibut, haddock and cod, had already been greatly reduced by fishing (Smith 1994).

A unique data set from two regions in the northwest Atlantic is examined and allows the abundance of groundfish species to be estimated from the early 1950's, before the arrival of factory-freezer-trawlers, to 1995. The structure of the groundfish community will be examined in terms of changes in species diversity and dominance throughout this period of intense exploitation.

## 2.3 Methods

#### Data

From 1947 to 1970 a "fixed" location survey along line transects, designed to sample a range of depths, was used to estimate groundfish abundance off the coast of Newfoundland (Pitt et al. 1981). Stratified random surveys began in 1971 and 1972, on the southern Grand Bank and St. Pierre Bank, respectively. The data collected prior to 1971 was converted to the new stratification scheme using the latitude, longitude and depth of the survey locations (Figure 1). Strata were included in the analysis if fewer than 10 years of data were missing, which would allow for a more complete time series. Fourteen strata from the southern Grand Bank, with depths ranging from 57-183 meters, and eleven strata from the St. Pierre Bank, with depths ranging from 56-274 meters, were selected for analysis (see Appendix B). Six years of St. Pierre Bank data were excluded due to the stratum constraints.



Figure 2.1: Map of research survey trawl locations on the southern Grand Bank and the St. Pierre Bank. Each (•) represents one trawl. The area of the southern Grand Bank used in this analysis is outlined and extends down to a depth of 183m. The Canadian 200 mile limit is provided as a reference on the southern Grand Bank. St. Pierre Bank strata considered are located on the bank and extend down to a depth of 366m west of 56° latitude. Years shown were selected arbitrarily.

Annual research surveys of the Southern Grand and St. Pierre Banks were conducted in the spring, but often extended into the summer and fall. For the purpose of this analysis, data was selected if the timing of the trawl occurred between January 1st and June 30th. Changes in the timing of the survey has consequences for species such as lumpfish (*Cyclopterus lumpus*) which migrate inshore of the survey area in the spring to spawn (Scott and Scott 1988). Also, spiny dogfish (*Squalus acanthias*), which migrate from the south into the study area in the spring (Templeman 1963), may not be surveyed by trawls which occurred before the month of May. These species will not be considered.

The number of tows in which each species was caught was determined, and the common species were selected for analysis (Table 2.1). All redfish (*Sebastes* sp.) species were grouped because in the 1950's the distinction among species was not made as redfish species identification requires a dissection. Rare species and fish that were only identified to genus or family, but which occurred in the same family as the common species, were also selected and will be used to estimate biomass at the family level. Individual estimates of biomass are calculated for fish identified to species (see Appendices C and D).

#### Standardization of the Data

#### Vessels and Survey Gear

Dedicated groundfish research surveys were conducted using otter trawls with 29 or 30 mm mesh liners in all years (Table 2.2) in the study area. The *Investigator II* conducted research surveys from 1951-1960 and from 1963-1965 using a Yanke 36 otter trawl, while the *A. T. Cameron*, which replaced the *Investigator II*, conducted surveys from 1959-1982 using a Yankee 41.5 otter trawl. The *Wilfred Templeman*, which subsequently replaced the *A.T. Cameron*, conducted surveys from 1983present using the Engels 145 high-lift otter trawl.

Catches were standardized using correction factors determined from the area surveyed by each gear type relative to that of the A.T. Cameron (Table 2.2). The area surveyed was calculated as wingspread multiplied by the towing speed (Rijnsdorp et al. 1996). Paired tow comparisons, which were carried out in 1959-1960 between the Investigator II and the A.T. Cameron (Templeman et al. 1978), and in 1983 between the A.T. Cameron and the Wilfred Templeman (Gavaris and Brodie 1984), yielded similar results. It should be noted, however, that neither of the correction factors is large and is unlikely to have a profound effect on the observed trends in biomass.

#### Diel variability in trawl catches

Prior to 1972, research surveys were primarily conducted during daylight hours. This may cause early estimates of abundance to be either underestimated, due to the increased visibility of the trawl during the day (Glass and Wardle 1989), or overestimated, if species exhibit diel vertical migrations that take them out of the vertical range of the trawl at night (Michalsen et al. 1996).

Conversion factors for the variation in diel catchability were calculated for over 50 species in the northwest Atlantic (Casey and Myers 1998a). Since the majority of the early research surveys were conducted during daylight hours, night catches for the entire period were adjusted by multiplying the weight by the appropriate conversion factor (Table 2.3). Table 2.1: Number of research survey tows on St. Pierre Bank (3Fe) and southern Grand Bank (3NO) in which each species was caught. The ordering is alphabetical for the order, family and species. "Other" refers to any fish that was identified to family only. The habitat of each species, and the reported range relative to the study area are given for reference. D-demersal, P=pelagic, M=mesopelagic, C=-coastal, DI=species found locally but usually at depths greater than that surveyed in the study area, L=local, LO=local species that migrates out of the area seasonally. MS=migrates into the area seasonally from the south, N=north of the study area, S=south of the study area, W=west of the study area.

Species	3Ps (n=1595)	3NO (n=2678)	Habitat	Range
CLUREIEORMES	124	41		
Clupeldae	124	41		
Alosa nseudoharenaus (Alawifa)	3	-	P	S
Chupen harenous (Herring)	123	41	P	I.
CADIFORMES	1249	2238		
Gadidae	1247	2237		
Boreogadus saida (Arctic cod)	8	10	P	N
Enchelyopus cimbrius (Fourbeard rockling)	63	12	D	L
Gadus mortug (Atlantic cod)	1060	2108	D	I.
Gadus ogac (Greenland cod)	1		D	N
Gaidronsarus sp. (Threeheard rockling)	10	4	D	DL
Melanogrammus aeglefinus (Haddock)	585	851	D	L
Merluccius bilinearis (Silver hake)	385	135	D	L
Micromesistius poutassou (Blue whiting)	2	14	M	S
Pollachius virms (Pollock)	173	99	P	L
Urophucis chesteri (Longfin hake)	295	51	D	DL
Urophucis tenuis (White hake)	468	387	D	L
Macrouridae	294	76		
Comphaenoides rupestris (Roundnose grenadier)	1		D	DW
Nezomia bairdi (Marlin-spike)	294	76	D	S
LOPHIIFORMES	375	128		
Lophiidae	375	128		
Lophius americanus (Monkfish)	375	128	D	L
MYCTOPHIFORMES	65	13		
Paralepididae	65	13		
Notolepis rissoi (White barracudina)	9		M	DL
Notolepis sp. (Barracudina)	56	13		
MYXINIFORMES	91	3		
Myxinidae	91	3		
Myzine glutinosa (Atlantic hagfish)	91	3	D	L
PERCIFORMES	1547	1891		
Ammodytidae	95	492		
Ammodytes dubius (Northern sand lance)	95	492	D	L
Anarhichadidae	268	577		
Anarhichas denticulatus (Broadhead wolffish)	39	28	D	N
Anarhichas lupus (Striped wolffish)	241	538	D	L
Anarhichas minor (Spotted wolffish)	17	23	D	N
Cottidae	1018	1091		
Artediellus sp. (Hookear sculpin)	17	42		
Cottunculus microps (Arctic deepsea sculpin)	2		D	w
Cottunculus sp. (Deepsea sculpin)	2			
Gymnocanthus tricuspis (Arctic staghorn sculpin)	1		D	N
Hemitripterus americanus (Sea raven)	658	718	D	L

Table 2.1 (continued).

Species	3Ps (n=1595)	3NO (n=2678)	Habitat	Range
PERCIFORMES				
Cottidae				
Muorocenhalus octodecemaninosus (Longhorn sculpin)	654	656	D	L
Muoroenhalus madricornis (Fourborn sculpin)	1	000	D	N
Muorocenhalus scornius (Shorthorn sculpin)	213	35	D	N
Trielons an (Mailed sculpin)	87	130	-	
Cyclopteridae	682	124		
Cuclonterus lumpus (Common lumpfish)	655	59	D	LO
Eumierntremus minasus (Spiny lumpfish)	98	66	D	N
Scorpaenidae	631	491		
Sebastes en (Redfish)	631	491	D	T.
Zoarcidae	28	107		
Cumnelus viridis (Green ocean nout)	2	3	D	N
Lucenchelus en (Wolf eel)	5	1		
Lucenchalus verrilli (Verrill's wolf cel)	2		D	S
(ucodes esmerki (Esmark's celpout)	12	1	D	DL
Lucodes landlesi (Land's ealpout)	1	Å	D	I.
(ucodes reticulatus (Arctic celpout)	27	94	D	N
(usedes on (Falcout)	7	74	5	
(ucodes ushlii (Vahl's calpout)	8	26	D	DI.
Management an anisonus (Common comp pout)	-	27	D	T
Melanastiama altentiaum (Soft exhout)	3	1	D	DI
Other (Feleeute)	2	2		00
DI EURONECTIEORMES	1297	2650		
Playmontides	1207	2650		
Charleson halve annealassus (Witch floweder)	654	045	D	r
Biggeodeprintus cynogossus (Which hounder)	1224	2576	D	ř
Rippoglossotales platessotales (American platee)	100	280	Ď	DI
Limondo fermicinas (Velloutail Bounder)	492	1743	D	I.
Donanda Jerruginae (Tellowial Hounder)	452	1	D	I.C.
Pseudoptueronectes unterretards (Winter Houtder)	174	959	D	DI
DA HEODMER	1977	2222	0	00
Balidae	1977	2232		
Raia fullas (Despurator skato)	1	2202	D	DI.
Raja Igniae (Deepwater skate)	22	18	D	L
Raja coellata (Winter skate)	28	10	Ď	S
Raja mediata (Winter state)	1228	2210	D	ř.
Raja radiata (Chordy Skate)	242	192	D	ĩ
Paia minimuda (Colontail slate)	1	100	D	DI
SAT MONTEOPMES	320	670	0	55
Argentinidae	242	90		
Amenting silve (Atlantic amenting)	242	90	D	DL
Ormonidan	88	585	5	JL
Mallatus millours (Canalin)	88	585	P	1.
COLLAT IEODMES	208	25	-	L
Pauelidea	208	25		
Controscullium fabricii (Black dorfich)	81	-0	P	DI.
Contrologianting (Colour deaffish)	247	25	P	MS

	Investigator II <sup>1</sup>	A.T.Cameron <sup>2</sup>	W. Templeman <sup>3</sup>
Vessel type	Stern trawler	Side trawler	Stern trawler
Tonnage	124	753	925
Length (m)	24.6	53	50
Speed of tow	4 knots	3.5 knots	3.5 knots
Trawl	Yankee 36	Yankee 41.5	Engels 145
Footrope (m)	24.4	30.5	44.2
Headrope (m)	18.3	24.4	29.3
Doors			
Weight (kg)	362.9	589.7	1247.4
Net opening (m)			
Wing spread	10.7	13.4	18.3
Headline height	2.4-3.4	2.4-3.4	3.7-5.5
Mesh size (mm)			
Codend liner	294	30 <sup>3</sup>	30
Area swept $(1000 \text{ m}^{2}\text{h}^{-1})$	79.2	86.8	118.5
Relative catchability	1.1	1	0.73

Table 2.2: Comparison of research survey vessel and gear characteristics of the Investigator II, the A.T. Cameron and the W. Templeman.

<sup>2</sup>(Gavaris and Brodie 1984) <sup>3</sup>(Walsh and McCallum 1995) <sup>4</sup>(Templeman et al. 1978)

Table 2.3: Conversion factors for diel catchability and the relative catchability of species. Conversions for diel catchability are applied to all night catches to make them comparable to that of the day. Conversions for relative species catchability are applied to biomass estimates to obtain estimates of absolute abundance (Edwards 1968).

	Diel Catchability		Relative Species	
Species	3Ps	3NO	Catchability	
American plaice	0.75	0.68	0.28	
Atlantic cod	1.00	1.26	0.28	
Atlantic halibut	1.40	1.40	0.38	
Barndoor skate	0.48	0.48	0.10	
Broadhead wolffish	0.90	0.90	0.16	
Haddock	1.40	2.39	0.48	
Longfin hake	0.65	0.36	0.42	
Longhorn sculpin	0.24	0.24	0.42	
Monkfish	1.00	1.00	0.16	
Pollock	1.00	1.00	0.08	
Redfish	2.23	2.23	0.27	
Sea raven	0.57	0.57	0.90	
Shorthorn sculpin	1.00	1.00	0.42	
Silver hake	1.00	1.00	0.04	
Smooth skate	0.54	0.54	0.10	
Spotted wolffish	0.91	0.91	0.16	
Striped wolffish	0.75	0.75	0.16	
Thorny skate	0.42	0.42	0.10	
White hake	1.00	1.00	0.51	
Witch flounder	0.90	0.90	0.49	
Yellowtail flounder	0.36	0.36	0.39	

#### **Biomass Estimates**

The survey area is primarily stratified along depth zones. For the purpose of this analysis, strata of equal depths were combined within each region and then biomass estimates were calculated (Pinhorn 1971). All strata were not surveyed in all years and therefore by combining same-depth strata more years could be included in the analysis.

Biomass estimates, with standard errors, were determined for individual species using the standard calculations for stratified random methods (Cochran 1977). The estimation of standard error was modified when there was only one sample per stratum: the sample variances in the same stratum from the previous and following years were averaged. This correction was only used in one year for the southern Grand Bank data and in four years for the St. Pierre Bank data.

All species are not sampled equally by the survey gear, as the survey trawl used may be more efficient at catching certain species. With the use of research submarines, underwater cameras and acoustic surveys, Edwards (1968) determined the availability and vulnerability of species to the survey gear. More rigorous methods employed by Clark and Brown (1977), which considered commercial and research vessel surveys and weight at age data, yielded catchability coefficients comparable to those of Edwards' (1968). Correction factors for the relative catchability of species used in this analysis are listed in Table 2.3. Absolute indices of abundance are calculated by dividing the biomass estimates by these correction factors.

In order to visualize changes within the grounfish community, estimates of biomass were calculated at the family level. A "losss" was then used to graphically display the data (Cleveland 1979). This is a robust, local smooth using locally linear fits. A window, dependent on the fraction of the data selected to be analyzed, is placed around each x value; points inside the window are weighted so that nearby points get the most weight. A fraction equal to 0.25 was used for all families, except the Gadidae in the southern Grand Bank for which 0.45 of the data was used. The smoothing is used for graphical purposes only.

#### **Species Diversity**

Two measures of species diversity were examined to identify changes within the groundfish community. Species richness is a simple measure of the total number of species in an given area (Gotelli and Graves 1996) and can provide information regarding the appearance or disappearance of species. Although changes in individual species are not identified with this summary measure, any changes in species richness over time can be further investigated. Species richness was calculated as the average number of species per tow in each year.

Evenness, a measure of the distribution of species abundances within the community, was chosen to examine changes in dominance within the groundfish community. Species evenness is negatively related to proportional abundances (Hill 1973). That is, low species evenness indicates that proportional abundances are high, and necessarily, that the community is dominated by only a few species.

Species evenness (E) was defined as  $E = \frac{ex(H)}{N_e}$ , where H is the Shannon-Wiener function and  $N_o$  is the corresponding species richness for each year (Hill 1973).

As a complement to the diversity indices, the cumulative biomass was calculated for each year. That is, biomass estimates were ranked from lowest to highest in each year and then the cumulative biomass was plotted against rank. A steeper rank abundance curve denotes dominance by only a few species.

### 2.4 Results

Large changes in the groundfish community occurred on both the southern Grand Bank (Figure 2.2) and St. Pierre Bank (Figure 2.3) where total biomass decreased by 70% and 80%, respectively, from the 1950's to the 1970's, and was largely representative of the family Gadidae. During the decrease in gadoid biomass on the southern Grand Bank, flatfish biomass increased and dominated through the 1960's, 1970's and into the 1980's, but was not nearly sufficient to compensate for the decrease in gadoid biomass. Flatfish biomass also increased on the St. Pierre Bank over the same time period, but the total biomas in this region was dominated by skate species which also increased as gadoids declined. Although gadoid biomass increased in both areas in the 1980's, by the 1990's the biomass of all species had decreased to its lowest level in at least 43 years. Only wolffish continued to increased in 1990's on the southern Grand Bank.

Biomass of Atlantic cod, haddock and white hake, was greatest in the 1950's (Figures 2.4 and 2.5). While the biomass of cod peaked again in the mid 1980's, the biomass of the haddock and white hake remained low for the rest of the time period. On the southern Grand Bank, both cod and haddock had been equally abundant, while on the St. Pierre Bank, haddock had been twice as abundant as cod. Silver hake was abundant on the St. Pierre Bank where biomass was the greatest in the 1950's and peaked again in the mid 1980's.

Redfish biomass in both areas contributed to the high estimate of total biomass in the 1950's. Since redfish inhabit a range of depths from about 90m to over 350m (Templeman 1959), and the survey area being investigated only covers 50-80% of that range, these estimates represent only that portion of the population found at the upper limit of its vertical range. Redfish biomass, however, has declined since the 1950's, except for an increase on the southern Grand Bank in the 1980's. Biomass estimates of American plaice, the dominant flatfish in this region, were greatest in the 1960's but gradually declined to the early 1990's. Yellowtail flounder biomass did not peak until the early 1970's and has decreased steadily since that time. The trends in biomass of Atlantic halibut and witch flounder more closely resemble that of the family Gadidae, with maximum estimates obtained in the 1950's and 1960's, and estimates remaining low since that time.

Skate biomass remained fairly stable over the time period. Biomass estimates of thorny skate, the dominant skate species in both regions, was greatest in the 1950's on the St. Pierre Bank thorny skate biomass idi not peak until the mid 1960's and has been decreasing since that time. Smooth skate, one of the smallest skate species in the northwest Atlantic, was highly variable on the southern Grand Bank, but on the St. Pierre Bank, smooth skate biomass had been increasing until the 1980's. Barndoor skate and winter skate biomass had been increasing until the 1980's. Barndoor skate and winter skate on the St. Pierre Bank had opposite trends in biomass over the time period. Barndoor skate biomass was greatest in the 1950's but decreased into the mid 1960's where it has remained at low levels ever since. Winter skate, however, was rarely caught until the mid 1980's when biomass estimates doubled. This may have resulted from a shift in distribution from the south.

Trends in biomass for wolffish, dominated by striped wolffish, differed for the southern Grand Bank and the St. Pierre Bank. On the St. Pierre Bank, wolffish biomass had been increasing until the mid 1970's when it peaked and subsequently declined. Estimates of wolffish biomass on the southern Grand Bank, however, have been increasing steadily.



Figure 2.2: Estimate of total biomass from selected southern Grand Bank strata from 1952-1995. Locas curves are used to depict changes in family composition during the time series.



Figure 2.3: Estimate of total biomass from selected St. Pierre Bank strata from 1951-1995. Loess curves are used to depict changes in family composition during the time series.



Figure 2.4: Estimates of biomass (with standard errors) for species in the selected region of the southern Grand Bank for the years 1952-1995.



Figure 2.5: Estimates of biomass (with standard errors) for species in the selected region of the St. Pierre Bank for the years 1951-1995.

Other species, such as those in the family Cottidae, increased in biomass over time. Sea raven biomass peaked in the late 1960's on the St. Pierre Bank and the 1970's on the southern Grand Bank, and has since decreased in both areas. Trends in biomass for longhorn sculpin were similar to that of the sea raven, except on the St. Pierre Bank where estimates of biomass were highly variable.

Species richness (Figure 2.6) increased on both the southern Grand Bank and the St. Pierre Bank, although the results for the southern Grand Bank were more variable. On St. Pierre Bank, species richness increased sharply from about 18 species in the 1950's to about 24 species in the 1970's, at a time when the total number of individuals was actually in decline. To ensure that this result was not due to the research vessel, survey gear or survey location, species richness was recalculated with only one vessel and gear type, and a subset of survey locations. Each subset produced similar trends in species richness over that time period. Most of the new species encountered were deepwater species, such as Greenland halibut and edepout, that had moved into shallower depths.

Species evenness (Figure 2.6) on the southern Grand bank increased from the 1950's to the 1970's and then remained fairly stable through the 1990's. On the St. Pierre Bank, however, species evenness continued to increase from the 1950's through the 1990's. These patterns are shown by the plot of rank abundance where the steepest curves for the southern Grand Bank (Figure 2.7), were found in the 1950's and 1960's, while on the St. Pierre Bank (Figure 2.7), the steepness of the rank abundance curves decreased gradually over the time period. The steep curve for 1995, however, is the result of a large catch of redfish on the St. Pierre Bank.



Figure 2.6: Species diversity. Two measures of species diversity, species evenness (Hill's  $N_i$ /species richenss) and species richness (mean number of species) are given for the southern Grand and St. Pierre Banks.



Figure 2.7: Cumulative species biomass for the southern Grand Bank from 1952-1995 and the St. Pierre Bank from 1951-1995.

#### 2.5 Discussion

This study represents one of the first attempts to obtain longterm, absolute estimates of biomass without relying on data from commercial fish catches. Commercial catch data, which can produce absolute indices of abundance through virtual population analysis, can only give insight into commercially important species and thus fish community changes may not be detected (Gulland and Garcia 1984). Although attempts have been made to examine historical fisheries data (Greenstreet and Hall 1996), such studies have not assessed changes in survey design and research vessel, or diel variability of research surveys conducted 24 hours a day.

The reduction in the biomass of major species, namely cod and haddock, fundamentally changed the groundfish community structure and reduced total species biomass by 90% from the 1950's to the 1990's. The largest biomass decrease of an individual species was experienced by the haddock population. In the 1950's, haddock had been the most abundant species on the southern Grand and St. Pierre Banks. Intensive fishing effort, both foreign and domestic, combined with an unprecedented discard rate (50-80% of the haddock catch was discarded due to the small size of the fish (Templeman et al. 1978; Templeman and Bishop 1979)), drastically depleted the haddock population by the 1960's. Haddock biomass has not recovered since that time is likely due to the fact that it has been taken as by-catch in the Atlantic cod fishery.

The findings of this study do not support the concept that total fish biomass will remain constant over a period of intense exploitation through the mechanism of species compensation. Increases in flatfish biomass on the southern Grand Bank and flatfish and skate biomass on the St. Pierre Bank were not great enough to compensate for the decline of the gadolids in the 1950's and 1960's. The increase that did occur was probably due to functional complementarity within the groundfish community. The function of the gadoids, flatfish and skates can be considered in terms of prey preferences. Both haddock and yellowtail flounder feed primarily on crustaceans and polychaetes, while cod and American plaice share capelln as the preferred prey (Methven 1999). Thoray skate, the most abundant skate on the St. Pierre Bank, prey on crustaceans and polychaetes (Templeman 1982) and would have been able to increase in biomass in this region where yellowtail flounder was less abundant. American plaice and yellowtail flounder became the target of fisheries, however, preventing either species from attaining their potential maximum biomass level. Skate species, although not taken commercially until 1994, were taken as bycatch in major fisheries.

Whether or not pelagic fish species would have compensated for this loss of biomass is uncertain as surprisingly little published information is available for pelagies during this time period. Pelagic species, such as capelin and northern sand lance, are not well sampled in groundfish surveys as they tend to swim above the range of the trawl. For this reason no attempt was made in this study to compare estimates of biomass between demersal and pelagic species. For comparison, the proportion of trawls in which each pelagic species was caught (?) was calculated as an index of biomass (Figure 2.8). Trawls that were directed capelin surveys were excluded. It is assumed that biomass is high when the species is caught in a high proportion of tows. Only the southern Grand Bank data were used for this calculation as neither species is abundant on the St. Pierre Bank.



Figure 2.8: Proportion of tows in which capelin and northern sand lance were caught in research survey trawls of the southern Grand Bank, 1952-1995.

Pelagics were only caught in 20% of the tows in the 1950's, but with the decrease in gadid biomass into the 1960's, 80% of the tows contained forage fish. The increase in gadid biomass in the early 1980's was coincident with a decrease to about 30% of the tows containing forage fish. In the early 1990's, when most of the demersal species had decreased in biomass, the proportion of tows in which forage fish were caught again increased to over 60%.

The increase in species evenness and richness with fishing pressure was not expected but can be explained using the intermediate disturbance hypothesis (Huston 1979). This hypothesis states that in the absence of environmental stress, such as fishing pressure, species diversity would be low as one or more species dominate the community. If the biomass of the dominant species is reduced (i.e. intermediate disturbance), more resources would be made available to other species thus increasing species diversity. In this case, the fishery acts as a keystone predator. Such predators actually facilitate the increase in species diversity by reducing competitive exclusion through species removal (Hacker and Gaines 1990). The removal of cod and haddock allowed flatfish on the southern Grand Bank, and flatfish and skate on the St. Pierre Bank to increase in biomass thus increasing species evenness. The release of resources into the groundfish community on the St. Pierre Bank may have allowed deepwater species, such as Greenland halibut and eelpout, to move into shallower depths, contributing to the increase in species richness.

This study differs dramatically from the analysis by Greenstreet and Hall (1996) of northern North Sea data from 1923 to 1993. They did not show this decline in absolute abundance and increase in diversity. The North Sea, however, was perhaps fully exploited at the time that survey began. Hislop (1996) showed that fishing mortality on haddock had already reached very high levels when the first world war began. By the 1930's hundreds of trawlers were operating in the North Sea, while on the Grand Banks, only about 50 trawlers operated using large mesh nets to capture mainly large cod (Thompson 1939).

This is the only study of a demersal fish community that makes estimates of the long term impact of trawling on the biomass and diversity before the introduction of intensive industrial trawling. There are two other studies that examined changes in species composition and began close to the onset of industrial trawling. Pauly (1988) describes bottom trawl surveys from 1963 to 1982 in the Gulf of Thailand when trawling effort increased by a factor of 8, and virtually all species groups declined except squid. Jin and Tang (1996) describe demersal surveys from 1959 to 1986 in the Yellow Sea where the dominant demersal fish species were replaced by small pelagic fish species.

Examination of historical fisheries data allows current species biomass to be put into perspective. Recent estimates of species biomass are often considered to be normal and are used in comparison only with estimates of future years. This has been referred to by Pauly (1995) as the 'shifting baseline syndrome'. In these instances, current biomass estimates are often a fraction of that suggested by historical data. Such was the case with the barndoor skate in the northwest Atlantic (Casey and Myers 1998b). The historical data for this species was readily available but not examined and so this species was able to disappear without anyone taking notice. Haddock populations on the southern Grand and St. Pierre Banks, which are thought to be genetically distinct from other stocks in the northwest Atlantic (Zwanenburg et al. 1992), could be facing the same fate.

# Chapter 3

# Near Extinction of a Large, Widely Distributed Fish

# 3.1 Abstract

Are extinctions of marine vertebrates as rare and unlikely as current data indicate? Longterm research surveys on the continental shelf between the Grand Banks of Newfoundland and Southern New England reveal that one of the largest skates in the northwest Atlantic, the barndoor skate (*Raja laevis*), is close to extinction. Forty-five years ago, research surveys on St. Pierre Bank (off southern Newfoundland) recorded barndoor skates in 10% of their tows; in the last 20 years none has been caught and this pattern of decline is similar throughout the range of the species.

## 3.2 Introduction

Elasmobranchs tend to be very susceptible to the effects of fishing because they grow slowly, mature late in life and produce few offspring (Holden 1973). Species of the family Rajidae, while the most fecund of the elasmobranchs (Holden 1974), are known to experience varying degrees of resilience to exploitation (Walker and Hislop 1988) due to the large range in life history characteristics within this family. While shark fisheries usually cause a sharp decline in species abundance (Anderson 1990), dogfish and skate on Georges Bank have increased in biomass following the depletion of groundfish stocks (Murawski and Idoine 1992). It has been suggested that the energy released into the ecosystem by these depleted stocks provided resources for elasmobranch populations to increase. The recent introduction of a directed fishery for dogfish and skate on Georges Bank, however, has resulted in a marked decline of these species (Fogarty and Murawski 1998).

The barndoor skate, Raja Laevis, is one of the largest skates in the northwest Atlantic (Bigelow and Schroeder 1953b), ranging from Cape Hatteras to the Grand Banks of Newfoundland (Figure 3.1). In May 1929 in the southern New England region, barndoor skates were captured at an average rate of 7 per tow (Bigelow and Schroeder 1953b). On Georges Eank in 1951 the average capture rate for barndoor skates was as high as 21 per tow (Bigelow and Schroeder 1953a). Once common, this distinctive species with a maximum body width of just over one meter, now appears to be near extinction. Although the extinction of marine species is thought to be rare (Huntsman 1994), the closely related "common" skate in the northeast Atlantic, Raja batis, was shown to be locally extinct in the Irish Sea (Brander 1981). If current population trends continue, however, the barndoor skate could become the first well-documented example of extinction in a marine fish species.



Figure 3.1: Map of the Northwest Atlantic Fisheries Organization. Subdivisions in which populations were assessed in this analysis are shown. The 300-m isobath (dotted line) is given for reference. 3N and 3O, southern Grand Bank; 3Ps, St. Pierre Bank; 4Vn, Sydney Bight; 4Vs, Banquereau Bank; 4W, Sable Island Bank; 4X, Browns Bank; 5Y, Gulf of Maine; 5Ze, Georges Bank; 5Zw, southern New England. The numbers on the axes are degrees of latitude and longitude.

# 3.3 Methods

Biomass (kg/km<sup>2</sup>) of the barndoor skate was determined from research vessel survey data assembled from the southern Grank Bank, at the northern limit of the range, to southern New England. From 1971 in Canadian waters and 1963 in the United States, annual research surveys have been conducted using a stratified random survey design (Doubleday and Rivard 1981). These are systematic surveys, covering a range of depths from 50 to 400 m, which are designed to provide unbiased estimates of abundance.

Prior to 1971, a fixed location survey along line transects was used off the coasts of Newfoundland and Nova Scotia. These surveys were converted to the stratification scheme using the latitude, longitude and depth of the tow. The earlier surveys were conducted primarily during the day and, as skates are caught in significantly higher proportions at night, day catches would underestimate the true abundance (Casey and Myers 1998a). To account for this difference in diel catchability, half of the catches prior to 1970 were converted to the night using a factor of 2.08. These surveys would then be comparable to those after 1970 in which day and night tows were conducted in roughly equal proportion.

Biomass estimates are based upon the stratified random design (Cochran 1977). Estimates of absolute abundance were obtained by dividing the biomass estimates by a factor of 0.15, if 10% of the skates in the path of the trawl are caught during the day (Edwards 1968) and 20% of the population would be caught at night. Neither of these conversions affects the observed biomass trends. An exponential decay curve  $(Ne^{-6})$  was fit to the data using non-linear least squares, where N is the population size in the first year of the surveys, and t is the time since the first year.

If barndoor skate are rarely caught in recent surveys, then the statistical power

of detecting individuals must be determined. On St. Pierre Bank, 504 of 1075 research survey tows were conducted over the past 25 years at depths where, historically, barndoor skate caught in one tow has a negative binomial distribution, with a probability of  $\frac{(k+a)}{(k+a)} \frac{\mu^{a}k^{k}}{\mu^{k}}$ , where  $\mu$  is the mean catch per tow, and the constant, k, is the inverse of aggregation. Then, with a theoretical low mean abundance of 500 individuals in a 7368 km<sup>2</sup> area (with a tow sample area of 0.05 km<sup>2</sup>, and the probability of ratching a barndoor skate if encountered is 0.15), and k=0.5, the probability of not detecting a barndoor skate in 504 tows is 0.77. Greater aggregation (that is, lower k) results in only small changes in this number. If 1000 individuals remained, the probability of not detecting a barndoor skate decreases to 0.6.

#### 3.4 Results

The population trend of the barndoor skate is similar for all regions (Figure 3.2), with biomass decreasing into the early 1970's, after which barndoor skate were caught only on Browns Bank and nearby Georges Bank. The estimated rate of population decline  $(\hat{\delta})$  was lowest in the northern and highest in southern regions. If only data since 1960 are considered, the population decline on St. Pierre Bank, Sydney Bight and Banquereau Bank is similar to that in southernmost regions (that is, Gulf of Maine, Georges Bank, and southern New England).

The longest time-series available is for southern Grand Bank, at the species' northern limit on the continental shelf, and St. Pierre Bank, where barndoor skate were once commonly found. Compared with other skate species on St. Pierre Bank, the barndoor skate had been one of the most numerous skates, second in abundance only to the thorny skate *Raje radiata* (Templeman 1966). Considering the mean biomass of the barndoor skate in each decade and the corresponding mean weight of individuals on St. Pierre Bank, the average number of barndoor skates in the 1950's would have been on the order of 0.6 million. That number would decrease to about 0.2 million individuals in the 1960's, and to less than an estimated 500 individuals in the 1970's. The other smaller skate species, namely thorny skate and smooth skate (*Raja senta*), were actually increasing in biomass over this time period (see Figure 2.5 in the previous chapter).

#### 3.5 Discussion

Direct biological information on skates in the northwest Atlantic is scarce. Sufficient comparative information, however, is available to estimate the mortality required to drive this species to extinction. The closest relative of the barndoor skate in the north Atlantic, the common skate (Bigelow and Schroeder 1953b), matures at approximately 11 years (Brander 1981). We should expect a similar age at maturity on Georges Bank, which has a similar temperature regime (Myers et al. 1997). Maximum egg production, which can be estimated from the inverse relationship with the weight of the young at hatching (The relationship between the weight of the young at hatching, w, and the maximum estimate of fecundity is 10.9m<sup>-0.46</sup> (Holden 1973)) is approximately 47 eggs/year.

Considering the age at maturity and the annual fecundity of the barndoor skate, the instantaneous mortality rate required to drive this species to extinction is approximately 0.4, assuming that mature and immature mortalities are equal.



Figure 3.2: Estimates of absolute biomass for barndoor skate (*Raja laevis*) from the southern Grand Bank to southern New England. Open circles are zero catches. The estimated rate of population decline,  $\delta$ , is provided with the standard error (SE).
In order for the barndoor skate population to be self-sustaining, the survival to age  $x, l_x$ , must be sufficient such that  $\sum l_x m_x = 1$ , where  $m_x$  is the number of eggs developing into females from individuals at age x. If survival, P, is constant such that  $l_x = P^x$ , then  $P^a m_o \sum_{i=0}^{\infty} P^i = 1$ , where the survival to the age of maturity a, is  $P^a$ , and the fecundity in each year is  $m_o$ . Summing the geometric series,  $P^a m_o(\frac{1}{1+p}) = 1$ . The instantaneous mortality rate is  $\log(P)$ . If the mortality on juveniles is twice that of adults, the instantaneous mortality required to drive the barndoor skate to extinction is 0.45.

At the northern limit of the barndoor skate's range, where the bottom temperature averages 2.5°C, about 6°C colder than on Georges Bank (Myers et al. 1997), the age at maturity should be approximately double and the instantaneous mortality required to drive the species to extinction would be expected to be closer to 0.2. This is based on the relationship between somatic growth rate and temperature, and a comparison of age at maturity of cod which occurs at the same locations (Myers et al. 1997).

The fishing mortalities for cod (*Cadus morhus*) have been above 0.4, and often much greater, in these regions for more than 30 years (Myers et al. 1996). It is therefore not surprising that the barndoor skate population would be reduced to such extreme levels when taken as by-catch in major fisheries, for example cod and redfish (*Sebastes* sp.). For a species that matures at such a late age (at least 11 years), there is a greater probability that individuals will be caught before they are able to reproduce. This situation is amplified for populations at the northern limit of the range.

The only recent research survey catches of barndoor skate in the Newfoundland area have been very deep (greater than 1000 m) and in an area north of the reported range of this species, where surveys were conducted in support of a new fishery for Greenland halibut (*Reinharditus hippoglossoides*). Whether or not barndoor skate were commonly found at this depth over the entire range is not known, as research surveys were rarely conducted at such depths. These great depths, however, may have represented one of the last refuges for this species.

Barndoor skate have survived on Browns Bank and Georges Bank probably because of a faster growth rate combined with the seasonal closure of parts of these banks to trawling (Shackell and Lien 1995). Perhaps the only hope for the longterm survival of this species is to designate an area protected from trawling on all the banks that is sufficiently large to allow for a self-sustaining population. A protected area would also provide a simple and effective means to conserve other species.

Failure to examine historical data has resulted in the largest skate in the northwest Atlantic being driven to near extinction without anyone noticing. If such a large, easily identified species, has been allowed to disappear in an area that is well surveyed, the fate of little known species is likely to be worse. This study shows the importance of assembling data on as long a time-scale as possible and as wide a spatial scale as possible; otherwise, the near extinction of a very apparent species may be missed.

A directed fishery for skate was initiated off the coasts of Newfoundland and Nova Scotia in 1994 as an attempt to utilize non-traditional groundfish species. Elasmobranchs experience low natural mortality when compared with teleost fish species, and, at low population levels, are not able to compensate, for example, with increased fecundity or reduced natural mortality (Anderson 1990). Consequently, the end result of increased mortality due to fishing pressure could be extinction. Consideration should be given to fact that species targeted in a commercial fishery are less likely to be driven to extinction for economic reasons. Non-target species, however, would continue to be taken as by-catch regardless of their commercial importance. As such, the extinction rate for non-target species, like the barndoor skate, would be far greater.

### Summary

Examination of large data sets presents many challenges due to the fact that the method of data collection changes over time with the introduction of new technology. Prior to 1972, research surveys of the southern Grand and St. Pierre Banks were conducted during daylight hours. This is a concern when determining absolute estimates of abundance. Species exhibiting diel vertical migrations, such as redfish and haddock, would be caught in greater proportions during the day. Non-migrating species, such as flatfish and skate, which rely on the visibility of the trawl as a means of escapement, would be caught in higher proportions during the night. Correction factors were determined for over 50 species and research surveys conducted at night were adjusted appropriately.

Total groundfish biomass in the 1990's was reduced to 11% and 9% of that observed in the 1950's on the southern Grand Bank and St. Pierre Bank, respectively. This trend largely resulted from the decline of the haddock population. Compensatory responses to this decline were visible with the flatfish on southern Grand Bank and skate on St. Pierre Bank. Continued fisheries for flatfish and bycatch of skate, however, ensured that total species biomass would remain at low levels.

Failure to examine such historical data has resulted in the largest skate in the northwest Atlantic, the barndoor skate, being driven to near extinction without anyone noticing. If such a large, easily identified species, has been allowed to disappear in an area that is well surveyed, the fate of little known species is likely to be worse. This study shows the importance of assembling data on as long a time-scale as possible and as wide a spatial scale as possible; otherwise, the near extinction of a very apparent species may be missed.

### Bibliography

- Allee, W. C., A. E. Emerson, O. Park, T. Park, and K. P. Schmidt (1949). Principles of Animal Ecology. Philadelphia, PA.: W.B. Saunders.
- Alverson, D. L., M. H. Freeberg, S. A. Murawski, and J. G. Pope (1994). A global assessment of fisheries bycatch and discards. *FAO Fisheries Technical Paper (339)*.
- Anderson, E. D. (1990). Fishery models as applied to elasmobranch fisheries. In H. L. Pratt, S. H. Gruber, and T. Taniuchi (Eds.), Elasmobranchs as Living Resources: Advances in Biology, Ecology, Systematics, and the Status of the Fisheries, pp. 475–484.
- Anon (1996). Report of the working group on ecosystem effects of fishing activities. ICES CM 1996/Assess/Env:1.
- Atkinson, D. B. (1989). Diel movements of beaked redfish and the implications of these for stratified random bottom trawl estimates of biomass and abundance. N. Am. J. Fish. Manage. 9, 163–170.
- Beamish, F. W. H. (1965). Vertical migration by demersal fish in the Northwest Atlantic. J. Fish. Res. Bd. Can. 23, 109–139.
- Bigelow, H. B. and W. C. Schroeder (1953a). Fishes of the Gulf of Maine. Fish. Bull. 53(74), 577.
- Bigelow, H. B. and W. C. Schroeder (1953b). Fishes of the Western North Atlantic, Part 2. New Haven, CT: Sears Foundation for Marine Research.
- Bishop, C. A. (1994). Revisions and additions to stratification schemes used during research vessel surveys in NAFO Subareas 2 and 3. NAFO SCR Document 94/43.
- Bishop, C. A., E. F. Murphy, M. B. Davis, J. W. Baird, and G. A. Rose (1993). An assessment of the cod stock in NAFO Divisions 2J+3KL. NAFO Sci. Counc. Rep. 93/86, Department of Fisheries and Oceans, Canada.

- Brander, K. (1981). Disappearance of common skate, Raia batis, from Irish Sea. Nature 290, 48–49.
- Brock, T. D. (1981). Calculating solar radiation for ecological studies. Ecol. Modell. 14, 1–19.
- Caddy, J. F. and J. A. Gulland (1983). Historical patterns of fish stocks. Marine Policy 7, 267–278.
- Casey, J. M. and R. A. Myers (1998a). Diel variation in trawl catchability: Is it as clear as day and night? Can. J. Fish. Aquat. Sci. 55, 2329–2340.
- Casey, J. M. and R. A. Myers (1998b). Near extinction of a large, widely distributed fish. *Science 281*, 690–692.
- Clark, S. H. and B. E. Brown (1977). Changes in biomass of finfish and squids from the Gulf of Maine to Cape Hatteras, 1963-1974, as determined from research vessel survey data. Fish. Bull. 75, 1–21.
- Cleveland, W. S. (1979). Robust locally weighted regression and smoothing scatterplots. J. Am. Stat. Assoc. 74, 829–836.
- Cochran, W. J. (1977). Sampling Techniques. New York, N. Y.: John Wiley & Sons.
- Deimling, E. A. and W. J. Liss (1994). Fishery development in the eastern North Pacific: a natural-cultural system perspective 1888-1976. *Fish. Oceanogr. 3*, 60-77.
- Doubleday, W. G. and D. Rivard (Eds.) (1981). Bottom Trawl Surveys. Ottawa, Canada: Government of Canada Fisheries and Oceans.
- Edwards, R. L. (1968). Fishery resources of the North Atlantic area. In D. W. Gilbert (Ed.), *The Future of the Fishing Industry of the United States*, pp. 52-60. University of Washington.
- Engås, A. and A. V. Soldal (1996). Diurnal variations in bottom trawl catch rates of cod and haddock and their influence on abundance indices. *ICES J. Mar. Sci.* 49, 89–95.
- Fogarty, M. J. and S. A. Murawski (1998). Large-scale disturbance and the structure of marine systems: fishery impacts on Georges Bank. *Ecol. Appl. 8, Supplement 1*, 56–522.
- Gavaris, S. and W. B. Brodie (1984). Results of comparative fishing between the A. T. Cameron and the Wilfred Templeman during July-August 1983. CAFSAC Research Document 84/41.

- Glass, C. W. and C. S. Wardle (1989). Comparison of the reactions of fish to a trawl gear, at high and low light intensities. *Fish. Res.* 7, 249–266.
- Gomes, M. C., R. L. Haedrich, and J. C. Rice (1992). Biogeography of groundfish assemblages on the Grand Bank. J. Northwest Atl. Fish. Sci. 14, 13–27.
- Gomes, M. C., R. L. Haedrich, and M. G. Villagarcia (1995). Spatial and temporal changes in the groundfish assemblages on the north-east Newfoundland/Labrador shelf, north-west Atlantic, 1978-1991. *Fish. Oceanogr.* 4, 85– 101.
- Gotelli, N. J. and G. R. Graves (1996). Null Models in Ecology. Washington: Smithsonian Institution Press.
- Greenstreet, S. P. R. and S. J. Hall (1996). Fishing and the ground-fish assemblage structure in the north-western North Sea: an analysis of long-term and spatial trends. J. Anim. Ecol. 65, 577–598.
- Gulland, J. A. and S. Garcia (1984). Observed patterns in multispecies fisheries. In R. M. May (Ed.), *Exploitation of Marine Communities*, pp. 155–190. Springer-Verlag.
- Hacker, S. D. and S. D. Gaines (1990). Some implications of direct positive interactions for community species diversity. *Ecology* 78(7), 1990–2003.
- Haedrich, R. L. (1995). Structure over time of an exploited deep-water fish assemblage. In A. G. Hopper (Ed.), Deep-Water Fisheries of the North Atlantic Oceanic Slope, pp. 27–50. Netherlands: Kluwer Academic Publishers.
- Harris, A. N. and I. R. Poiner (1991). Changes in species composition of demersal fish fauna of Southeast Gulf of Carpentaria, Australia, after 20 year of fishing. *Marine Biology* 111, 503–519.
- Hilborn, R. and C. J. Walters (1992). Quantitative Fisheries Stock Assessment: Choice, Dynamics and Uncertainty. New York: Chapman and Hall.
- Hill, M. O. (1973). Diversity and evenness: A unifying notation and its consequences. Ecology 54, 360–367.
- Hislop, J. R. G. (1996). Changes in North Sea gadoid stocks. ICES J. Mar. Sci. 53, 1146–1156.
- Holden, M. J. (1973). Are long-term sustainable fisheries for elasmobranchs possible? Rapp. P.-v. Réun. Cons. int. Explor. Mer 164, 427-432.
- Holden, M. J. (1974). Problems in the rational exploitation of elasmobranch populations and some suggested solutions. In F. R. H. Jones (Ed.), Sea Fisheries Research, pp. 117–137. Elek Science, London.

- Huntsman, G. R. (1994). Endangered marine finfish: Neglected resources or beasts of fiction? *Fisheries 19*(7), 8–15.
- Huston, M. A. (1979). A general hypothesis of species diversity. American Naturalist 113, 81–101.
- Hutchings, J. A. and R. A. Myers (1995). The biological collapse of Atlantic cod off Newfoundland and Labrador: an exploration of historical changes in exploitation, harvesting technology and management. In R. Arnason and L. Felt (Eds.), *The North Atlantic Fisheries: Successes, Failures and Challenges*, pp. 38–93. Charlottetown, Prince Edward Island: The Institute of Island Studies.
- Jin, X. and Q. Tang (1996). Changes in fish species diversity and dominant species composition in the Yellow Sea. Fish. Res. 26, 1996.
- Kerr, S. R. and R. A. Ryder (1989). Current approaches to multispecies analysis of marine fisheries. Can. J. Fish. Aquat. Sci. 46, 528–534.
- Korsbrekke, K. and O. Nakken (1997). Length and species dependant diurnal variation in catch rates in the Norwegian Barents Sea bottom trawl surveys. *ICES C.M. W:20.*
- Larkin, P. A. (1977). An epitaph for the concept of maximum sustained yield. Trans. Am. Fish. Soc. 106, 1–11.
- Litvin, V. M. and V. D. Rvachev (1963). The bottom topography and sediments of the Labrador and Newfoundland fishing areas. In Y. Y. Marti (Ed.), Soviet Fisheries Investigations in the Northwest Atlantic, Jerusalem. Israel Program for Scientific Translantions.
- Ludwig, D., R. Hilborn, and C. Walters (1993). Uncertainty, resource exploitation, and conservation: lessons from history. Science 260, 17–36.
- Manly, B. F. (1991). Randomization and Monte Carlo Methods in Biology. New York: Chapman and Hall.
- McCullagh, P. and J. A. Nelder (1989). Generalized Linear Models. Monographs on Statistics and Applied Probability. London: Chapman & Hall.
- Methven, D. A. (1999). Annotated bibliography of demersal fish feeding with emphasis on selected studies from the Scotian Shelf and Grand Banks of the Northwestern Atlantic. Technical report, Department of Fisheries and Oceans.

- Michalsen, K., O. R. Godø, and A. Fernö (1996). Diel variation in the catchability of gadoids and its influence on the reliability of abundance indices. *ICES J. Mar. Sci.* 53, 389–395.
- Murawski, S. A. and J. S. Idoine (1992). Multispecies size compositon: a conservative property of exploited fishery systems? J. Northwest Atl. Fish. Sci. 14, 79–85.
- Myers, R. A., J. A. Hutchings, and N. J. Barrowman (1996). Hypotheses for the decline of cod in the North Atlantic. Mar. Ecol. Prog. Ser. 138, 293–308.
- Myers, R. A., G. Mertz, and S. Fowlow (1997). Maximum population growth rates and recovery times of Atlantic cod, *Gadus morhua. Fish. Bull.* 95, 762–772.
- Pauly, D. (1988). Fisheries research and the demersal fisheries of Southeast Asia. In J. A. Gulland (Ed.), Fish Population Dynamics, 2nd edition, pp. 329–348. New York: Wiley.
- Pauly, D. (1995). Anecdotes and the shifting baseline syndrome of fisheries. Trends. Ecol. Evol. 10, Trends in Ecology and Evolution, 1995, v.10, n.10, p-1994.
- Pauly, D. (1996). One hundred million tonnes of fish, and fisheries research. Fish. Res. 25, 25–38.
- Pauly, D., V. Christensen, J. Dalsgaard, R. Forese, and F. Torres (1998). Fishing down marine food webs. *Science* 279, 860–862.
- Pinhorn, A. T. (1971). Accuracy of abundance indices for cod from St. Pierre Bank (ICNAF Div. 3Ps) based on Canada (Newfoundland) research vessel surveys in terms of commercial abundance indices. In International Commission for the Northwest Atlantic Fisheries Redbook, Part III, pp. 239–248. Government of Canada.
- Pinhorn, A. T. and R. G. Halliday (1990). Canadian versus international regulation of Northwest Atlantic Fisheries: Management practices, fishery yields, and resource trends, 1960-1986. N. Am. J. Fish. Manage. 10, 154–174.
- Pitt, T. K. (1967). Diurnal variation in the catches of American plaice, *Hip-poglossoides platessoides* Fabr., from the Grand Bank. *ICNAF Res. Bull.* 4, 53-58.
- Pitt, T. K., R. Wells, and W. D. Mckone (1981). A critique of research vessel otter trawls by the St. John's Research and Resource Services. In W. G.

Doubleday and D. Rivard (Eds.), *Bottom Survey Trawls*, pp. 42–61. Government of Canada Fisheries and Oceans.

- Regier, H. A. and W. L. Hartman (1973). Lake Erie's fish community: 150 years of cultural stresses. *Science* 180, 1248–1255.
- Rijnsdorp, Q. D., P. I. van Leeuwen, N. Daan, and H. J. L. Heessen (1996). Changes in abundance of demersal fish species in the North Sea between 1906-1909 and 1990-1995. *ICES J. Mar. Sci.* 53, 1054–1062.
- Sahrhage, D. and J. Lundbeck (1992). A History of Fishing. Berlin: Springer-Verlag.
- Scott, W. B. and M. G. Scott (1988). Atlantic Fishes of Canada, Volume No. 219 of Can. Bull. Fish. Aquat. Sci. University of Toronto Press.
- Shackell, N. and J. Lien (1995). An under-utilized conservation option for fishery managers: marine protected areas in the northwest Atlantic. In N. L. Shackell and J. H. M. Willison (Eds.), Marine Protected Areas and Sustainable Fisheries, pp. 21–30. Wolfville, Nova Scotia: SAMPAA, Centre for Conservation Biology, Acadia University.
- Sissenwine, M. P. (1984). Why do fish populations vary? In R. M. May (Ed.), Exploitation of Marine Communities, pp. 59–94. Springer-Verlag, Berlin.
- Skud, B. E. (1982). Dominance in fishes: the relation between environment and abundance. Science 216, 144–149.
- Smith, S. H. (1968). Species succession and fishery exploitation in the Great Lakes. Fisheries Research Board of Canada 25, 667–693.
- Smith, T. D. (1994). Scaling Fisheries: The Science of Measuring the Effects of Fishing, 1855-1955. Great Britain: Cambridge University Press.
- Steedman, R. J. and H. A. Regier (1990). Ecological bases for an understanding of ecosystem integrity in the Great Lakes basin. In C. J. Edwards and H. A. Regier (Eds.), An Ecosystem Approach to the Integrity of the Great Lakes in Turbulent Times, pp. 254–270. Great Lakes Fishery Commission, Ann Arbor, Michigan.
- Templeman, W. (1959). Redfish Distribution in the North Atlantic. Bulletin No. 120. Ottawa: Fisheries Research Board of Canada.
- Templeman, W. (1963). Distribution of Sharks in the Canadian Atlantic. Bulletin No. 140. Ottawa: Fisheries Research Board of Canada.
- Templeman, W. (1966). Marine Resources of Newfoundland. Bulletin No. 154. Ottawa: Fisheries Research Board of Canada.

- Templeman, W. (1982). Stomach contents of the thorny skate, raja radiata, from the Northwest Atlantic. J. Northwest Atl. Fish. Sci. 3, 123–126.
- Templeman, W. and C. A. Bishop (1979). Age, growth, year-class strength, and mortality of haddock, *Melanogrammus aeglefinus*, on St. Pierre Bank in 1948-1975 and their relation to the haddock fishery in this area. *ICNAF Res. Bull. 14*, 85–99.
- Templeman, W., V. M. Hodder, and R. Wells (1978). Age, growth, year-class strength and mortality of the haddock *Melanogrammus aeglefinus*, on the southern Grand Bank and their relation to the haddock fishery of the area. *ICNAF Res. Bull.* 13, 31–52.
- Thompson, H. (1939). The occurrence and biological features of haddock in the Newfoundland area. Research Bulletin, Department of Natural Resources, Newfoundland Government 6, 1–30.
- Villagarcia, M. G. (1995). Structure and distribution of demersal fish assemblages on the northeast Newfoundland and Labrador shelf. Master's thesis, Memorial University of Newfoundland, St. John's, Newfoundland.
- Walker, P. A. and J. R. G. Hislop (1988). Sensitive skates or resilient rays? Spatial and temporal shifts in ray species composition in the central and north-western North Sea between 1930 and the present day. *ICES J. Mar. Sci.* 55, 392–402.
- Walsh, S. J. (1988). Diel variability in the trawl catches of juvenile and adult yellowtail flounder on the Grand Banks and the effect on resource assessment. N. Am. J. Fish. Manage. 8, 1261–1272.
- Walsh, S. J. (1991). Diel variation in availability and vulnerability of fish to a survey trawl. J. Appl. Ichthyol. 7, 147–159.
- Walsh, S. J. and W. M. Hickey (1993). Behavioural reactions of demersal fish to bottom travls at various light conditions. *ICES Mar. Sci. Symp.* 196, 68–76.
- Walsh, S. J. and B. R. McCallum (1995). Survey trawl mensuration using acoustic trawl instrumentation. ICES CM 1995/B:26.
- Walters, C. J. and J. S. Collie (1988). Is research on environmental factors useful to fisheries management? Can. J. Fish. Aquat. Sci. 45, 1848–1854.
- Warner, W. W. (1983). Distant Water: The Fate of the North Atlantic Fisherman. Boston: Little, Brown & Co.

Zwanenburg, K. C. T., P. Bentzen, and J. M. Wright (1992). Mitochondrial DNA differentiation in western North Atlantic populations of haddock (Melanogrammus acqlefinus). Car. J. Fish. Ayuat. Sci. 49, 2527–2537.

### Appendix A

### Estimation in SAS

This appendix demonstrates how to fit the proposed Model 2 (Chapter 1) to catch data from stratified random surveys.

Create a data set with seven variables per observation: species, region and season, number caught during the day (Cysd) in a particular year(y) and stratum (s), total number caught (Cys) in a particular year and stratum, number of tows during the day (Tysd), number of tows during the night (Tysn), and time of day (daytime). A sample of what the data set should look like is given in Table A.1.

The following is a brief description of the method of programming in SAS:

```
data dl;

infile survey;

input species season Cysd Cys Tysd Tysn daytime;

offset=Tysd/Tysn

proc sort; by species season;

proc genmod; by species season;

class daytime;

model Cysd/Cys = daytime / link=logit dist=binomial offset=offset

dscale noint;
```

Table A.1: Sample data set to be used with Model 2 (Chapter 1). The data represent Atlantic cod in subdivision 3NO from the spring research surveys.

y	8	$C_{ysd}$	$C_{ys}$	$T_{ysd}$	$T_{ysn}$	Offset
72	383	62	65	1	1	0
72	359	73	78	1	2	-0.69
72	382	154	358	2	2	0
72	381	83	228	2	2	0
73	374	8	10	3	1	1.1
73	376	1	1	1	2	-0.69
73	354	15	22	1	2	-0.69
73	359	13	59	1	2	-0.69
74	378	129	484	1	2	-0.69
74	381	5	23	1	3	-1.1
74	379	32	296	1	2	-0.69
74	380	8	16	1	1	0
75	376	17	23	1	1	0
75	377	81	110	1	1	0
75	333	169	169	1	1	0
76	354	32	60	2	1	0.69
76	333	15	15	1	1	0
76	334	0	2	1	1	0

Appendix B

# Research Survey Tows Per Stratum

	Stratum Number													
	336	337	338	351	352	353	354	359	360	361	362	374	375	376
1951	0	8	4	0	9	2	3	0	0	0	0	0	0	0
1952	5	6	3	3	3	3	7	6	6	3	1	0	3	1
1953	2	9	3	0	7	3	6	3	4	2	3	1	2	0
1954	0	3	5	1	9	1	2	0	5	S	4	2	2	1
1955	0	6	6	3	9	1	3	4	11	6	3	2	2	1
1956	0	4	4	2	3	0	1	0	5	2	2	2	1	1
1957	0	4	2	2	4	1	2	2	7	2	2	0	2	0
1958	0	4	2	3	4	0	2	2	7	1	0	2	1	0
1959	4	8	4	2	4	2	9	10	5	4	1	2	1	2
960	8	6	6	6	3	1	3	3	5	7	1	2	0	4
961	3	10	5	1	6	3	5	6	8	4	0	1	1	2
962	13	14	7	3	3	2	12	6	5	3	n.	0	2	3
963	11	14	10	5	10	4	11	4	10	9	2	0	7	5
964	11	22	13	3	8	4	10	3	3	3	1	0	3	6
965	9	15	7	1	6	4	9	5	7	4	0	2	0	4
966	7	8	4	0	5	2	10	3	5	1	0	ō	1	4
967	3	8	1	0	1	1	3	7	8	1	0	3	4	9
968	5	4	3	2	3	1	3	4	4	3	2	1	2	4
969	3	1	2	0	0	0	0	2	1	1	2	0	2	0
970	1	3	0	2	0	0	3	1	4	õ	6	2	4	4
971	1	5	1	ō	0	1	3	5	4	2	2	2	5	2
972	0	0	0	2	0	0	0	3	10	8	4	5	7	5
973	3	3	5	5	5	3	3	3	1	4	s	4	3	3
974	0	0	0	0	0	0	0	0	ō	4	4	2	3	0
975	2	3	2	4	4	3	0	0	4	4	3	2	3	2
976	2	2	3	4	4	2	3	3	4	5	5	0	1	3
977	2	2	4	5	5	3	2	2	4	3	5	3	4	3
978	3	6	7	9	8	5	2	4	12	7	6	5	8	7
979	4	4	7	11	12	5	4	3	9	8	12	4	s	4
980	2	3	5	10	11	4	3	4	11	6	11	3	4	3
981	4	5	2	4	0	1	8	6	14	6	13	7	9	7
982	3	3	5	9	7	3	2	2	7	6	8	4	5	7
984	2	2	5	6	7	2	2	2	7	5	7	2	s	4
985	2	5	9	8	11	6	3	2	16	6	81	4	7	7
986	2	5	9	14	14	7	3	2	13	10	14	6	8	8
987	2	6	9	13	13	6	2	2	15	8	13	5	8	8
988	2	4	8	10	11	5	2	2	12	7	10	5	6	6
989	2	5	10	15	13	7	2	2	15	9	13	5	8	8
990	5	10	16	22	23	8	4	4	28	17	21	10	14	13
991	4	9	15	22	24	11	5	8	20	14	20	7	10	12
992	4	6	10	16	14	7	4	4	20	12	14	5	6	10
993	4	5	11	16	14	7	4	4	20	13	16	6	10	10
994	3	5	10	16	14	6	4	4	16	11	15	5	9	8
005	-	-				-	- C					<b>1</b>	-	-

Table B.1 Number of research survey tows completed in selected strata from the Southern Grand Bank during the years of 1951-1995.

					Str	atum ?	fumber				_
	311	312	313	314	315	316	317	320	321	705	706
1951	3	0	2	6	2	2	4	9	0	2	0
1952	3	1	0	4	4	0	0	0	1	0	0
1953	3	3	0	3	5	1	2	8	3	1	0
1954	2	0	0	4	4	1	2	8	3	0	0
1955	1	1	1	3	5	1	2	10	2	0	0
1956	2	3	1	1	3	2	4	7	3	1	2
1957	2	1	1	4	3	2	3	6	4	1	2
1958	2	3	1	3	5	2	4	6	3	1	2
1959	2	0	0	6	2	4	4	6	4	0	0
1960	2	4	5	8	4	11	7	15	6	1	1
1962	0	0	0	3	1	2	2	3	0	0	1
1963	2	1	3	5	4	6	4	7	4	0	0
1964	2	2	1	3	3	4	5	6	1	0	1
1965	3	1	5	3	4	6	4	5	2	3	0
1967	5	3	6	2	2	6	4	2	0	2	1
1968	2	1	3	0	2	6	4	2	0	0	0
1969	3	1	6	1	1	6	4	2	0	0	0
1970	4	-	4	1	2	6	7	0	0	1	1
1972	á.	2	2	2	2	2	4	0	2	2	2
1973	8	1	2	0	2	3	7	2	2	2	2
1974	8	2	12	2	2	14	8	1	1	11	15
1975	4	3	3	0	0	1	4	0	0	2	1
1976	6	5	6	2	2	4	4	1	2	5	4
1977	4	4	10	4	4	6	4	0	0	4	4
1978	8	2	5	3	0	6	4	0	3	5	2
1979	4	3	5	1	3	3	3	1	0	4	3
1980	2	1	2	1	4	2	2	6	5	2	2
1981	2	2	2	4	5	7	5	2	2	2	4
1982	3	2	2	5	3	1	3	4	4	2	4
1983	3	3	3	7	8	4	3	13	10	3	5
1984	2	2	2	4	S	2	2	7	6	2	2
1985	3	2	2	7	7	3	2	5	7	2	4
1986	3	2	2	7	6	2	2	9	10	2	4
1987	3	2	2	4	8	3	3	9	10	2	4
1988	4	2	2	7	6	3	2	11	11	2	4
1989	3	3	2	9	7	2	2	10	9	2	4
1990	3	2	2	6	1	2	2	5	8	2	4
1991	3	2	2	7	7	2	2	12	11	2	4
1992	2	2	2	5	7	2	2	8	9	2	5
1993	5	4	4	13	10	4	4	13	14	4	8
1994	4	2	2	9	6	2	2	8	10	3	4
1005		3	2	7	6	2	2	12	11	2	4

Table B.2 Number of research survey tows completed in selected strata from the St. Pierre Bank during the years of 1951-1995.

# Appendix C

## Southern Grand Bank Data Used in Chapter 2

Table	B.1	Total	weight	for	each	specie	es and	year	surveyed	on
southe	rn G	rand H	Bank. ]	The .	orderin	g of t	he spec	ies is	alphabeti	cal.
The to	otal n	umber	of tows	5 (N	) is giv	en for	referen	ice.		

Species	Year	Total Weight (kg)	N	Year	Total Weight (kg)	N
American plaice	1951	103.31	26	1973	1870.63	50
American plaice	1952	1859.69	50	1974	478.99	13
American plaice	1953	504.83	45	1975	1368.03	36
American plaice	1954	317.05	40	1976	1791.72	41
American plaice	1955	179.97	57	1977	2045.76	47
American plaice	1956	431.22	27	1978	3943.79	89
American plaice	1957	594.35	30	1979	2999.55	92
American plaice	1958	418.56	28	1980	3731	80
American plaice	1959	1613.44	52	1981	2579.5	49
American plaice	1960	930.32	54	1982	2255.8	71
American plaice	1961	3123.02	55	1984	2474.64	58
American plaice	1962	4743.28	64	1985	11741.18	167
American plaice	1963	2922.42	56	1986	3829.29	115
American plaice	1964	3763.52	90	1987	3794.14	110
American plaice	1965	6138.58	66	1988	2550.84	90
American plaice	1966	2670.32	50	1989	3033.01	112
American plaice	1967	851.87	19	1990	2946.01	104
American plaice	1968	1191.6	40	1991	1753.6	103
American plaice	1969	406.42	8	1992	871.67	89
American plaice	1970	1394.12	28	1993	1090.75	77
American plaice	1971	2245.56	30	1994	487.52	64
American plaice	1972	1076.84	21	1995	426.5	85
Arctic cod	1951	0	26	1973	0	50
Arctic cod	1952	0	50	1974	0	13
Arctic cod	1953	0	45	1975	0	36
Arctic cod	1954	0	40	1976	0	41
Arctic cod	1955	0	57	1977	0	47
Arctic cod	1956	0	27	1978	0	89

		Total			Total	
Species	Year	Weight (kg)	N	Year	Weight (kg)	N
Arctic cod	1957	0	30	1979	0	92
Arctic cod	1958	0	28	1980	0	80
Arctic cod	1959	0	52	1981	0	49
Arctic cod	1960	0	54	1982	0	71
Arctic cod	1961	ő	55	1984	0	58
Arctic cod	1962	0	64	1985	0	167
Arctic cod	1963	0	56	1986	0.04	115
Anotic cod	1064	0	00	1087	0.04	110
Arctic cod	1965	0	66	1988	0.01	90
Aretic cod	1966	0	50	1090	0.04	112
Arctic cod	1967	0	19	1990	0.01	104
Aretic cod	1069	0	40	1001	0.02	102
Arctic cou	1060	0	0	1000	0.02	80
Aretic cod	1909	0	20	1002	0.02	77
Arctic cou	1071	0	20	1004	0.02	64
Arelie cou	1971	0	30	1005	0.01	04
Arctic cod	1972	0	21	1995	0	60
Arctic eeipout	1951	0	20	1973	0	30
Arctic esipout	1952	0	30	1974	0	13
Arctic eeipout	1953	0	45	1975	0	30
Arctic eelpout	1954	0	40	1976	0	41
Arctic ecipout	1955	0	57	1977	0	4/
Arctic eelpout	1956	0	27	1978	0	89
Arctic eelpout	1957	0	30	1979	0	92
Arctic eelpout	1958	0	28	1980	0	80
Arctic eelpout	1959	0	52	1981	10.75	49
Arctic eelpout	1960	0	54	1982	12.4	71
Arctic eelpout	1961	0	55	1984	1.91	58
Arctic eelpout	1962	0	64	1985	28.1	167
Arctic eelpout	1963	0	56	1986	12.12	115
Arctic eelpout	1964	0	90	1987	10	110
Arctic eelpout	1965	0	66	1988	3.07	90
Arctic eelpout	1966	0	50	1989	7.39	112
Arctic eelpout	1967	0	19	1990	16.7	104
Arctic eelpout	1968	0	40	1991	9.45	103
Arctic eelpout	1969	0	8	1992	4.56	89
Arctic eelpout	1970	0	28	1993	2.95	77
Arctic eelpout	1971	0	30	1994	1.11	64
Arctic eelpout	1972	5.45	21	1995	2.81	85
Atlantic argentine	1951	0	52	1973	0.23	100
Atlantic argentine	1952	6.88	100	1974	0	26
Atlantic argentine	1953	1.81	90	1975	2.49	72
Atlantic argentine	1954	0	80	1976	0.05	82
Atlantic argentine	1955	0	114	1977	0.23	94
Atlantic argentine	1956	0	54	1978	6.43	178
Atlantic argentine	1957	0	60	1979	23.16	184
Atlantic argentine	1958	0	56	1980	0.7	160
Atlantic argentine	1959	5.13	104	1981	0	98
Atlantic argentine	1960	52.03	108	1982	o	142
Atlantic argentine	1961	2.18	110	1984	1.33	116
Atlantic argentine	1962	4.54	128	1985	0.42	334
Atlantic argentine	1963	313.59	112	1986	29.46	230
Atlantic argentine	1964	34.18	180	1987	0	220

Table B.1 Total weight for each species and year surveyed on southern Grand Bank. The ordering of the species is alphabetical. The total number of tows (N) is given for reference.

	201	Total			Total	
Species	Year	Weight (kg)	N	Year	Weight (kg)	N
Atlantic argentine	1965	4.27	132	1988	0	180
Atlantic argentine	1966	59.64	100	1989	0	224
Atlantic argentine	1967	0	38	1990	0.35	208
Atlantic argentine	1968	70.76	80	1991	0	206
Atlantic argentine	1969	0	16	1992	1.25	178
Atlantic argentine	1970	0	56	1993	0.98	154
Atlantic argentine	1971	0	60	1994	0.18	128
Atlantic argentine	1972	0	42	1995	1.41	170
Atlantic cod	1951	1621.82	26	1973	714.44	50
Atlantic cod	1952	4227.76	50	1974	81.18	13
Atlantic cod	1953	3277.31	45	1975	587.29	36
Atlantic cod	1954	7118.99	40	1976	367.13	41
Atlantic cod	1955	6849.32	57	1977	1510	47
Atlantic cod	1956	7912.11	27	1978	2506.76	89
Atlantic cod	1957	1530.45	30	1979	3937.41	92
Atlantic cod	1958	1295.98	28	1980	1549.15	80
Atlantic cod	1959	5912.23	52	1981	1577.7	49
Atlantic cod	1960	4008.15	54	1982	1639.32	71
Atlantic cod	1961	5091.08	55	1984	5111.14	58
Atlantic cod	1962	3598.3	64	1985	17672.87	167
Atlantic cod	1963	3865.55	56	1986	9854.82	115
Atlantic cod	1964	8020.83	90	1987	13897.56	110
Atlantic cod	1965	5070.13	66	1988	6272.14	90
Atlantic cod	1966	2237.81	50	1989	6613.15	112
Atlantic cod	1967	1285.36	19	1990	6693.52	104
Atlantic cod	1968	2013.44	40	1991	2730.1	103
Atlantic cod	1969	1094.8	8	1992	1736.21	89
Atlantic cod	1970	1253.99	28	1993	1024	77
Atlantic cod	1971	2687.79	30	1994	136.47	64
Atlantic cod	1972	652.02	21	1995	217.7	85
Atlantic harfish	1951	0	26	1973	0	50
Atlantic hagfish	1952	0	50	1974	0	13
Atlantic harfish	1953	0	45	1975	0	36
Atlantic harfish	1954	0	40	1976	0	41
Atlantic harfish	1955	0	57	1977	0	47
Atlantic hagfish	1956	0	27	1978	0	89
Atlantic harfish	1957	0	30	1979	0	92
Atlantic harfish	1958	0	28	1980	0	80
Atlantic harfish	1959	0	52	1981	0	49
Atlantic hagfish	1960	0	54	1982	0	71
Atlantic hagfish	1961	0.46	55	1984	0	58
Atlantic harfish	1962	0.23	64	1985	0	167
Atlantic hagfish	1963	0	56	1986	0	115
Atlantic harfish	1964	0	90	1987	0	110
Atlantic hagfish	1965	0	66	1988	0	90
Atlantic harfish	1966	0.23	50	1989	0	112
Atlantic hagfish	1967	0	19	1990	0	104
Atlantic bagfish	1968	0	40	1991	0	103
Atlantic hagfish	1969	0	8	1992	0	89
Atlantic harfish	1970	0	28	1993	0	77
Atlantic hagfish	1971	0	30	1994	0	64
Atlantic hagfish	1972	ő	21	1995	ō	85

Table B.1 Total weight for each species and year surveyed on southern Grand Bank. The ordering of the species is alphabetical. The total number of tows  $\left(N\right)$  is given for reference.

Species	Vear	Total Weight (kg)	N	Vear	Total Weight (kg)	N
openio	tout	mentine (nt)		TOUL	mergine (wg)	
Atlantic halibut	1951	0	26	1973	15.42	50
Atlantic halibut	1952	48.47	50	1974	0	13
Atlantic halibut	1953	41.91	45	1975	7.94	36
Atlantic halibut	1954	32.33	40	1976	68.04	41
Atlantic halibut	1955	30.93	57	1977	43.55	47
Atlantic halibut	1956	21.45	27	1978	42.25	89
Atlantic halibut	1957	59.86	30	1979	30.42	92
Atlantic halibut	1958	4.99	28	1980	245.8	80
Atlantic halibut	1959	173.54	52	1981	81	49
Atlantic halibut	1960	372.74	54	1982	140.9	71
Atlantic halibut	1961	541.36	55	1984	92.96	58
Atlantic halibut	1962	588.03	64	1985	283.44	167
Atlantic halibut	1963	150.88	56	1986	239.46	115
Atlantic halibut	1964	173.08	90	1987	220.32	110
Atlantic halibut	1965	335.59	66	1988	23.22	90
Atlantic halibut	1966	115.53	50	1989	132.8	112
Atlantic halibut	1967	0	19	1990	20.43	104
Atlantic halibut	1968	29.48	40	1991	81.13	103
Atlantic halibut	1969	41.28	8	1992	58.1	89
Atlantic halibut	1970	8.62	28	1993	8.01	77
Atlantic halibut	1971	38.1	30	1994	3.78	64
Atlantic halibut	1972	0	21	1995	15.98	85
Barndoor skate	1951	0	26	1973	0	50
Barndoor skate	1952	15.54	50	1974	0	13
Barndoor skate	1953	44.91	45	1975	0	36
Barndoor skate	1954	31.08	40	1976	õ	41
Barndoor skate	1955	0	57	1977	0	47
Barndoor skate	1956	0	27	1978	0	89
Barndoor skate	1957	0	30	1979	0	92
Barndoor skate	1958	15.54	28	1980	ő	80
Barndoor skate	1959	10.04	52	1981	0	49
Barndoor skate	1960	110.68	54	1982	0	71
Barndoor skate	1961	90.72	55	1984	0	58
Daradoor skate	1962	61.22	64	1085	0	167
Darndoor shate	1062	01.20	56	1086	0	115
Darnuoor shate	1064	0.08	00	1087	0	110
Barndoor skate	1965	9.90	66	1088	0	90
Barndoor skate	1965	27.21	50	1080	0	112
Barndoor shave	1067	27.21	10	1000	0	104
Darndoor skille	1000	0	10	1001	0	102
Barndoor skate	1968	0	40	1991	0	103
Barndoor skate	1969	0		1992	0	09
Barndoor skate	1970	0	28	1993	0	64
Darndoor skate	1971	0	30	1005	0	04
Darngoor skate	1972	0	21	1995	0	60
Blue whiting	1951	0	26	1973	0	30
Blue whiting	1952	0	50	1974	0	13
Blue whiting	1953	0	45	1975	0	36
Blue whiting	1954	0	40	1976	0	41
Blue whiting	1955	0	57	1977	0	47
Blue whiting	1956	0	27	1978	0.2	89
Blue whiting	1957	0	30	1979	0	92

Table B.1 Total weight for each species and year surveyed on southern Grand Bank. The ordering of the species is alphabetical. The total number of tows (N) is given for reference.

		Total			Total	
Species	Year	Weight (kg)	N	Year	Weight (kg)	N
Blue whiting	1959	0	52	1981	0	49
Blue whiting	1960	0.18	54	1982	0	71
Blue whiting	1961	0	55	1984	0	58
Blue whiting	1962	0	64	1985	0.83	167
Blue whiting	1963	0.79	56	1986	0.17	115
Blue whiting	1964	0.54	90	1987	0	110
Blue whiting	1965	0.18	66	1988	0	90
Blue whiting	1966	0	50	1989	0	112
Blue whiting	1967	0	19	1990	0	104
Blue whiting	1968	4.31	40	1991	0	103
Blue whiting	1969	0	8	1992	0	89
Blue whiting	1970	0	28	1993	0	77
Blue whiting	1971	0	30	1994	0	64
Blue whiting	1972	0	21	1995	0	85
Broadhead wolffish	1951	0	26	1973	0	50
Broadhead wolffish	1952	8.66	50	1974	0	13
Broadhead wolffish	1953	0	45	1975	0	36
Broadhead wolffish	1954	0	40	1976	0	41
Broadhead wolffish	1955	0	57	1977	7.26	47
Broadhead wolffish	1956	0	27	1978	15.5	89
Broadhead wolffish	1957	0	30	1979	35.41	92
Broadhead wolffish	1958	0	28	1980	15.1	80
Broadhead wolffish	1959	4 99	52	1981	73	49
Broadhead wolffish	1960	10.89	54	1982	0	71
Broadhead wolffish	1961	0	55	1984	0	58
Broadhead molffish	1962	0	64	1985	4.15	167
Broadhead wolffish	1963	0	56	1986	12.45	115
Broadhead wolffish	1964	0	90	1987	0	110
Broadhead wolffish	1965	44.97	66	1988	ő	90
Broadhead wolffish	1966	21.33	50	1989	0	112
Broadhead wolffish	1967	0	19	1990	0	104
Broadhead wolffish	1968	0	40	1991	0	103
Broadhead wolffish	1969	19.05	8	1992	15.85	89
Broadhead wolffish	1970	22.68	28	1993	24.61	77
Broadhead wolffish	1971	13.61	30	1994	0	64
Broadhead wolffish	1972	0	21	1995	0	85
Capelin	1951	15.27	26	1973	1.96	50
Capelin	1952	1.3	50	1974	0	13
Capelin	1953	24.9	45	1975	1500.59	36
Capelin	1954	2.65	40	1976	0.28	41
Capelin	1955	4.9	57	1977	42.63	47
Capelin	1956	0.3	27	1978	16.61	89
Capelin	1957	0.05	30	1979	211.14	92
Capelin	1958	0.55	28	1980	0.7	80
Capelin	1959	31.64	52	1981	147	49
Capelin	1960	6.54	54	1982	31.65	71
Capelin	1961	12.61	55	1984	2.24	58
Capelin	1962	265.32	64	1985	52.23	167
Capelin	1963	1511.98	56	1986	12.54	115
Capelin	1964	811.66	90	1987	325.77	110
Capalin	1965	636 23	66	1988	5.98	90
Capelin	1966	96.74	50	1989	43.94	112

Table B.1 Total weight for each species and year surveyed on southern Grand Bank. The ordering of the species is alphabetical. The total number of tows  $\left(N\right)$  is given for reference.

		Total			Total	
Species	Year	Weight (kg)	N	Year	Weight (kg)	N
Capelin	1967	249.8	19	1990	28	104
Capelin	1968	381.73	40	1991	84.5	103
Capelin	1969	30.09	8	1992	16.4	89
Capelin	1970	220.24	28	1993	408.9	77
Capelin	1971	11.37	30	1994	9.42	64
Capelin	1972	60.88	21	1995	35.6	85
Common ocean pout	1951	0	26	1973	16.33	50
Common ocean pout	1952	0	50	1974	0	13
Common ocean pout	1953	0	45	1975	6.8	36
Common ocean pout	1954	0	40	1976	0	41
Common ocean pout	1955	0	57	1977	0	47
Common ocean pout	1956	0	27	1978	0	89
Common ocean pout	1957	0	30	1979	0	92
Common ocean pout	1958	0	28	1980	0.8	80
Common ocean pout	1959	1.54	52	1981	4.25	49
Common ocean pout	1960	0.75	54	1982	6.4	71
Common ocean pout	1961	0	55	1984	4.81	58
Common ocean pout	1962	0.91	64	1985	5.6	167
Common ocean pout	1963	0	56	1986	0	115
Common ocean pout	1964	0.45	90	1987	2.49	110
Common ocean pout	1965	1.54	66	1988	1	90
Common ocean pout	1966	0.91	50	1080	1.83	112
Common ocean pout	1967	0	19	1990	7.74	104
Common ocean pout	1068	0	40	1001	0	103
Common ocean pout	1969	0	8	1992	0	89
Common ocean pout	1970	0	28	1993	0	77
Common ocean pout	1071	0	20	1004	0	64
Common ocean pour	1072	22.26	21	1005	0	95
Deepses couloin	1051	40.00	26	1973	0	50
Deepsea sculpin	1052	0	50	1074	0	12
Deepsea sculpin	1952	0	45	1975	0	36
Deepses sculpin	1054	0	40	1076	0	41
Deepses sculpin	1055	0	57	1077	0	47
Deepsea sculpin	1056	0	27	1978	0	80
Deepees sculpin	1057	0	20	1070	0	03
Deepsea sculpin	1059	0	28	1980	0	80
Deepsea scuipin	1050	0	20	1960	0	40
Deepsea scuipin	1000	0	54	1082	0	71
Deepses scuipin	1900	0	04	1984	0	70
Deepsea scutpin	1961	0	00	1964	0	167
Deepsea scuipin	1902	0	04	1960	0	115
Deepsea sculpin	1903	0	00	1980	0	110
Deepsea sculpin	1964	0	90	1089	0	110
Deepsea scutpin	1903	0	00	1966	0	110
Deepsea scutpin	1906	0	30	1909	0	104
Deepsea sculpin	1967	0	19	1990	0	102
Deepsea sculpin	1968	0	40	1991	0	103
Deepsea sculpin	1969	0	8	1992	0	69
Deepsea sculpin	1970	0	28	1993	0	
Deepsea sculpin	1971	0	30	1994	0	04
Deepsea sculpin	1972	0	21	1995	0	60
Exelpont	1951	0	26	1973	3.17	20
P. ANTINGUES	1 10/10/2		- CH2	17769	1.30	

Table B.1 Total weight for each species and year surveyed on southern Grand Bank. The ordering of the species is alphabetical. The total number of tows (N) is given for reference.

		Treat			Travel	
Species	Vear	Weight (kg)	N	Vear	Weight (kg)	N
Species	1.004	meigne (wg)		rear	weight (kg)	14
Eelpout	1953	0	45	1975	4.08	36
Eelpout	1954	0	40	1976	3.18	41
Eelpout	1955	0	57	1977	33.34	47
Eelpout	1956	0	27	1978	50.53	89
Feloout	1957	0	30	1979	12 93	92
Felpout	1958	0	28	1980	12.85	80
Eelpout	1959	0	52	1981	0	49
Felpout	1960	0	54	1982	0	71
Felpout	1961	19 74	55	1984	0	58
Eelpout	1962	0	64	1985	0	167
Felpout	1963	18 14	56	1986	0	115
Felpout	1964	0.9	90	1987	0	110
Eelpout	1965	14 51	66	1088	0	90
Eelpout	1966	0	50	1989	0	112
Eeloout	1967	24.02	10	1000	0	104
Edgout	1069	04.02	40	1001	0	102
Esloout	1060	0	-	1002	0	203
Eelpout	1970	7.94	28	1992	0	77
Ealpoint	1071	0	20	1004	0	64
Estpour	1072	0	30	1005	0	05
Estpour	1972	0	00	1990	0	50
Eamark's elipout	1951	0	20	1973	0	12
Esmark's elipout	1052	0	45	1075	0	20
Esmark's elipout	1903	0	40	1970	0	30
Esmark's elipout	1954	0	40	1970	0	41
Esmark's elipout	1933	0	07	1078	0	80
Esmark's elipout	1956	0	21	1978	0	09
Esmark's elipout	1050	0	30	1000	0	54
Esmark's eelpout	1908	0	20	1960	0	00
Esmark's eeipout	1959	0	52	1981	0	49
Esmark's eelpout	1960	0	04	1962	0	71
Esmark's ecipout	1961	0	33	1984	0	08
Esmark's eelpout	1962	0	04	1985	0	167
Esmark's eelpout	1963	0	36	1986	0	115
Esmark's eeipout	1904	0	90	1987	0	110
Esmark's eelpout	1965	0	00	1988	0	90
Esmark's eelpout	1966	0	50	1989	0	112
Esmark's ecipout	1967	0	19	1990	0	104
Esmark's eelpout	1968	1.13	40	1991	U	103
Esmark's eelpout	1969	0	8	1992	0	89
Esmark's ecipout	1970	0	28	1993	0	
Esmark's eelpout	1971	0	30	1994	0	04
Esmark's eelpout	1972	0	21	1995	0	85
Fourbeard rockling	1951	0	26	1973	0	50
Fourbeard rockling	1952	0	50	1974	0	13
Fourbeard rockling	1953	0	45	1975	0	36
Fourbeard rockling	1954	0	40	1976	0.05	41
Fourbeard rockling	1955	0	57	1977	0	47
Fourbeard rockling	1956	0	27	1978	0.14	89
Fourbeard rockling	1957	0	30	1979	0	92
Fourbeard rockling	1958	0	28	1980	0.02	80
Fourbeard rockling	1959	0.14	52	1981	0	49
Fourbeard rockling	1960	0	54	1982	0	71

Table B.1 Total weight for each species and year surveyed on southern Grand Bank. The ordering of the species is alphabetical. The total number of tows  $\left(N\right)$  is given for reference.

Table B.1	Total	weight f	or each	species	and	year	surveyed	on
southern ( The total	Grand E	ank. The	e orderi N) is gi	ng of the ven for re	e spec	cies is ice.	alphabeti	cal.

		Total			Total	
Species	Year	Weight (kg)	N	Year	Weight (kg)	N
Fourbeard rockling	1961	0.14	55	1984	0	58
Fourbeard rockling	1962	0	64	1985	0	167
Fourbeard rockling	1963	0	56	1986	0	115
Fourbeard rockling	1964	0.28	90	1987	0	110
Fourbeard rockling	1965	0.28	66	1988	0	90
Fourbeard rockling	1966	0	50	1989	0	112
Fourbeard rockling	1967	0	19	1990	0	104
Fourbeard rockling	1968	0.14	40	1991	0	103
Fourbeard rockling	1969	0	8	1992	0	89
Fourbeard rockling	1970	0.23	28	1993	0	77
Fourheard rockling	1971	0	30	1994	0	64
Fourbeard rockling	1972	0	21	1995	0	85
Greenland halibut	1951	0	26	1973	1.05	50
Greenland halibut	1952	0	50	1974	0	13
Greenland halibut	1953	0	45	1975	0.78	36
Greenland halibut	1954	0	40	1976	3.09	41
Greenland halibut	1955	0	57	1977	6.79	47
Greenland halibut	1956	0	27	1978	10.62	89
Greenland halibut	1957	0.67	30	1979	42.62	92
Greenland halibut	1958	2.67	28	1980	14.6	80
Greenland halibut	1959	4 98	52	1981	11.5	49
Greenland halibut	1960	1.87	54	1982	3.8	71
Greenland halibut	1961	3.63	55	1984	0.17	58
Greenland halibut	1962	2.97	64	1985	0.54	167
Greenland halibut	1963	22.44	56	1986	4.52	115
Greenland halibut	1964	33 32	90	1987	4.96	110
Greenland halibut	1965	46	66	1988	3.78	90
Greenland halibut	1966	20.41	50	1989	1.18	112
Greenland halibut	1967	53.98	19	1990	3.69	104
Greenland halibut	1968	4.41	40	1991	4.43	103
Greenland halibut	1969	0	8	1992	3.71	89
Greenland halibut	1970	3.77	28	1993	7.29	77
Greenland halibut	1971	4.98	30	1994	1.42	64
Greenland halibut	1972	0	21	1995	12.62	85
Haddock	1951	6886.19	26	1973	29.4	50
Taddock	1952	5295.16	50	1974	0	13
Taddock	1953	11176.23	45	1975	6.35	36
Haddock	1954	11628.28	40	1976	6.8	41
Haddock	1955	11543.03	57	1977	11.1	47
Haddock	1956	12396.57	27	1978	31.27	89
Haddock	1957	5506.08	30	1979	35.19	92
Taddock	1958	4217.07	28	1980	76	80
Haddock	1959	8398.19	52	1981	119.95	49
Taddock	1960	11649.92	54	1982	502.38	71
Taddock	1961	7773.31	55	1984	1179.6	58
Haddock	1962	5762.97	64	1985	1063.25	167
Haddock	1963	1507.25	56	1986	265.85	115
Haddock	1964	1856.81	90	1987	1289.07	110
Haddock	1965	644.56	66	1988	1739.51	90
Haddock	1966	1017.64	50	1989	142.26	112
Haddock	1967	0.45	19	1990	170.94	104
Haddock	1968	183.39	40	1991	102.61	103

		Treat			Treat	
Sourcies	Vear	Weight (kg)	N	Vear	Weight (kg)	N
opecies	1 COR	and Bue (ng)			troibus (nD)	
Haddock	1969	1.82	8	1992	27.35	89
Haddock	1970	7.39	28	1993	75.07	77
Haddock	1971	187.1	30	1994	68.81	64
Haddock	1972	5.21	21	1995	37.06	85
Hookear sculpin	1951	0	26	1973	0	50
Hookear sculpin	1952	0	50	1974	0	13
Hookear sculpin	1953	0	45	1975	0.14	36
Hookear sculpin	1954	0	40	1976	0.05	41
Hookear sculpin	1955	0	57	1977	0.23	47
Hookear sculpin	1956	0	27	1978	0.12	89
Hookear sculpin	1957	0	30	1979	0.06	92
Hookear sculpin	1958	0	28	1980	0	80
Hookear sculpin	1959	0.07	52	1981	0	49
Hookear sculpin	1960	0.06	54	1982	0	71
Hookear sculpin	1961	0.14	55	1984	0	58
Hookear sculpin	1962	0.03	64	1985	0	167
Hookear sculpin	1963	0.01	56	1986	0	115
Hookear sculpin	1964	0.01	90	1987	0.01	110
Hookear sculpin	1965	0.35	66	1988	0	90
Hookear sculpin	1966	0.1	50	1989	0	112
Hookear sculpin	1967	0	19	1990	0.01	104
Hookear sculpin	1968	0	40	1991	0.01	103
Hookear sculpin	1969	0	8	1992	0.02	89
Hookear sculpin	1970	0	28	1993	0	77
Hookear sculpin	1971	0	30	1994	0.02	64
Hookear sculpin	1972	0	21	1995	0.01	85
Longfin bake	1951	0	26	1973	0	50
Longfin hake	1952	16.77	50	1974	0	13
Longfin bake	1953	0	45	1975	1.14	36
Longfin hake	1954	0	40	1976	0.96	41
Longfin hake	1955	0	57	1977	0.68	47
Longfin hake	1956	0	27	1978	17.3	89
Longfin hake	1957	0	30	1979	5.72	92
Longfin bake	1958	0	28	1980	4	80
Longfin hake	1959	0.9	52	1981	0	49
Longfin bake	1960	2.74	54	1982	0.1	71
Longfin hake	1961	0	55	1984	0.66	58
Longfin hake	1962	0	64	1985	0.83	167
Longfin bake	1963	3.05	56	1986	0	115
Longfin hake	1964	29.49	90	1987	0.5	110
Longfin hake	1965	2.72	66	1988	0	90
Longfin hake	1966	6.94	50	1989	0.66	112
Longfin hake	1967	0	19	1990	0	104
Longfin hake	1968	1.81	40	1991	0.11	103
Longfin hake	1969	0	8	1992	0.12	89
Longfin hake	1970	0	28	1993	0	77
Longfin hake	1971	0	30	1994	0.05	64
Longfin hake	1972	0	21	1995	0.00	85
Longhorn sculpin	1951	0	26	1973	72.63	50
Longhorn sculpin	1952	1.77	50	1974	74.02	13
Longhorn sculpin	1953	1.06	45	1975	28.34	36
Longhorn sculpin	1954	2.48	40	1976	97.09	41

Table B.1 Total weight for each species and year surveyed on southern Grand Bank. The ordering of the species is alphabetical. The total number of tows  $\left(N\right)$  is given for reference.

		Total			Total	
Species	Year	Weight (kg)	N	Year	Weight (kg)	N
Longhorn sculpin	1955	0	57	1977	23.8	47
Longhorn sculpin	1956	0.79	27	1978	33.7	89
Longhorn sculpin	1957	2.83	30	1979	117.51	92
Longhorn sculpin	1958	3.95	28	1980	108.25	80
Longhorn sculpin	1959	11.32	52	1981	16.1	49
Longhorn sculpin	1960	3.66	54	1982	114.5	71
Longhorn sculpin	1961	15.09	55	1984	32.29	58
Longhorn sculpin	1962	11.17	64	1985	314.57	167
Longhorn sculpin	1963	9.48	56	1986	83.37	115
Longhorn sculpin	1964	27.23	90	1987	100.1	110
Longhorn sculpin	1965	14.12	66	1988	20.92	90
Longhorn sculpin	1966	3.05	50	1989	24.27	112
Longhorn sculpin	1967	1.49	19	1990	29.68	104
Longhorn sculpin	1968	1.83	40	1991	26.39	103
Longhorn sculpin	1969	1.99	8	1992	3.82	89
Longhorn sculpin	1970	1.13	28	1993	12.08	77
Longhorn sculpin	1971	9.31	30	1994	2.33	64
Longhorn sculpin	1972	125.41	21	1995	4.67	85
Mailed sculpin	1951	0	52	1973	0.1	100
Mailed sculpin	1952	0	100	1974	0.24	26
Mailed sculpin	1953	0	90	1975	0.43	72
Mailed sculpin	1954	0	80	1976	0.43	82
Mailed sculpin	1955	0	114	1977	0.09	94
Mailed sculpin	1956	0.03	54	1978	1.29	178
Mailed sculpin	1957	0	60	1979	0.56	184
Mailed sculpin	1958	0	56	1980	0.3	160
Mailed sculpin	1959	0.1	104	1981	1.6	98
Mailed sculpin	1960	0.09	108	1982	0.01	142
Mailed eculoin	1961	0.3	110	1984	0	116
Mailed sculpin	1962	0.0	128	1985	0.02	334
Mailed couloin	1967	0.30	112	1986	0.08	230
Mailed sculpin	1964	0.25	180	1987	0.12	220
Mailed couloin	1065	0.7	132	1988	0.03	180
Mailed sculpin	1066	0.07	100	1020	0.22	224
Mailed sculpin	1967	0.05	28	1990	0.16	208
Mailed sculpin	1069	0.02	80	1001	0.22	206
Mailed sculpin	1908	0.05	16	1002	0.57	178
Mailed sculpin	1070	0.02	56	1007	0.02	154
Mailed sculpin	1970	0.00	60	1004	0.02	128
Mailed sculpin	1072	0.69	42	1005	0.01	170
Marlin-spike	1051	0.00	26	1973	0.09	50
Marlin miles	1052	9.79	50	1074	0.00	13
Martin-spike	1052	0.20	45	1075	0.27	26
Marin-spike	1903	0	40	1076	0.37	41
Marlin mike	1055	0	57	1977	0.1	47
Marlin spike	1056	0	27	1078	0.45	80
Marlin spike	1057	0	30	1979	0.23	92
Marlin spike	1059	0	28	1080	1.4	80
Marlin spike	1050	5.43	52	1981	1.4	49
Marin-spike	1909	5.43	54	1082	0.1	-19
Martin-spike	1960	0.9	55	1984	0.08	58
INTERACIO-NOIME	1301	0.08	00	1004	0.08	00

Table B.1 Total weight for each species and year surveyed on southern Grand Bank. The ordering of the species is alphabetical. The total number of tows  $({\rm N})$  is given for reference.

		Total			Total	
Species	Year	Weight (kg)	N	Year	Weight (kg)	N
Marlin-enike	1963	40.1	56	1986	0	115
Marlin-spike	1964	44.03	90	1987	0.59	110
Marlin-spike	1965	16.02	66	1988	0.17	90
Marlin-spike	1966	12.64	50	1989	0.07	112
Marlin-spike	1967	0	19	1990	0	104
Marlin-spike	1968	3.4	40	1991	0.27	103
Marlin-spike	1969	0	8	1992	0	89
Marlin-spike	1970	0	28	1993	0	77
Marlin-spike	1971	0	30	1994	0	64
Marlin-spike	1972	0	21	1995	0	85
Monkfish	1951	0	26	1973	27.44	50
Monkfish	1952	52.46	50	1974	0	13
Monkfish	1953	22.08	45	1975	5.44	36
Monkfish	1954	6.56	40	1976	13.61	41
Monkfish	1955	5.98	57	1977	0.95	47
Monkfish	1956	1	27	1978	22.13	89
Monkfish	1957	0.5	30	1979	2.95	92
Monkfish	1958	13.97	28	1980	45	80
Monkfish	1959	89.36	52	1981	22	49
Monkfish	1960	44.58	54	1982	2.95	71
Monkfish	1961	142.87	55	1984	0	58
Monkfish	1962	148.66	64	1985	36.94	167
Monkfish	1963	68.6	56	1986	25.73	115
Monkfieh	1964	39.47	90	1987	47.72	110
Monkfish	1965	39.82	66	1988	46.48	90
Monkfish	1966	23 59	50	1989	12.95	112
Monkfish	1967	0	19	1990	13.07	104
Monkfish	1968	13.6	40	1991	18.9	103
Monkfish	1969	0	8	1992	0.62	89
Monkfish	1970	0	28	1993	4.65	77
Monkfish	1971	0	30	1994	9.54	64
Monkfish	1972	0	21	1995	0	85
Northern sand lance	1951	0	26	1973	41.29	50
Northern sand lance	1952	1.05	50	1974	34.02	13
Northern sand lance	1953	0.7	45	1975	252.19	36
Northern sand lance	1954	0.22	40	1976	26.07	41
Northern sand lance	1955	7.48	57	1977	70.99	47
Northern sand lance	1956	0.62	27	1978	483.3	89
Northern sand lance	1957	1.05	30	1979	469.83	92
Northern sand lance	1958	52.88	28	1980	528.55	80
Northern sand lance	1959	1.14	52	1981	89.15	49
Northern sand lance	1960	27.39	54	1982	182.36	71
Northern sand lance	1961	1.2	55	1984	1.66	58
Northern sand lance	1962	0.12	64	1985	3.54	167
Northern sand lance	1963	0.18	56	1986	0.24	115
Northern sand lance	1964	3.15	90	1987	1.19	110
Northern sand lance	1965	1.85	66	1988	0.8	90
Northern sand lance	1966	1.02	50	1989	3.57	112
Northern sand lance	1967	2.66	19	1990	0.42	104
Northern sand lance	1968	0.32	40	1991	1.45	103
Northern sand lance	1969	3.18	8	1992	0.58	89
Northern sand lance	1970	91.56	28	1993	0.75	77

Table B.1 Total weight for each species and year surveyed on southern Grand Bank. The ordering of the species is alphabetical. The total number of tows (N) is given for reference.

		Total			Total	
Species	Year	Weight (kg)	N	Year	Weight (kg)	N
Northern sand lance	1971	36.35	30	1994	0.89	64
Northern sand lance	1972	35.38	21	1995	0.47	85
Pollock	1951	66.53	26	1973	0	50
Pollock	1952	123.03	50	1974	0	13
Pollock	1953	12.47	45	1975	0	36
Pollock	1954	14.46	40	1976	0.91	41
Pollock	1955	10.48	57	1977	0	47
Pollock	1956	602.49	27	1978	1.82	89
Pollock	1957	7.48	30	1979	7.49	92
Pollock	1958	3.99	28	1980	0	80
Pollock	1959	47.66	52	1981	0	49
Pollock	1960	219.07	54	1982	0.7	71
Pollock	1961	20.41	55	1984	0	58
Pollock	1962	19.05	64	1985	1.7	167
Pollock	1963	156.66	56	1986	1.25	115
Pollock	1964	120.21	90	1987	16.43	110
Pollock	1965	13.39	66	1988	0.5	90
Pollock	1966	37.87	50	1989	13.7	112
Pollock	1967	0	19	1990	15.13	104
Pollock	1968	0	40	1991	15.44	103
Pollock	1969	0	8	1992	0	89
Pollock	1970	0	28	1993	0	77
Pollock	1971	103.65	30	1994	0	64
Pollock	1972	0	21	1995	0	85
Redfish	1951	174.89	78	1973	639.61	150
Redfish	1952	18936.93	150	1974	0	39
Redfish	1953	10132.08	135	1975	694.78	108
Redfish	1954	2116.6	120	1976	176.41	123
Redfish	1955	1896.92	171	1977	3683.8	141
Redfish	1956	3016.57	81	1978	126.7	267
Redfish	1957	7910.48	90	1979	1222.65	276
Redfish	1958	2247.04	84	1980	194.36	240
Redfish	1959	6552.77	156	1981	241	147
Redfish	1960	19059.76	162	1982	4340.5	213
Redfish	1961	4061.4	165	1984	299.55	174
Redfish	1962	7739.65	192	1985	563.49	501
Redfish	1963	3376.8	168	1986	4024.09	345
Redfish	1964	5108.69	270	1987	463.85	330
Redfish	1965	2704.55	198	1988	651.53	270
Redfish	1966	3679.19	150	1989	368.85	336
Redfish	1967	69.85	57	1990	1513.05	312
Bedfish	1968	6609.13	120	1991	29.9	309
Redfish	1969	0	24	1992	107.14	267
Redfish	1970	375.82	84	1993	1295.73	231
Redfish	1971	460.4	90	1994	1162.21	192
Redfish	1972	0.6	63	1995	9282.72	255
Sea raven	1951	0	26	1973	157.39	50
Sea raven	1952	8.73	50	1974	62.58	13
Sea raven	1953	8.73	45	1975	78.71	36
Sea raven	1954	15.71	40	1976	158.98	41
Sea raven	1955	4.99	57	1977	115.65	47
See rasion	1956	12.48	27	1978	189.4	89

Table B.1 Total weight for each species and year surveyed on southern Grand Bank. The ordering of the species is alphabetical. The total number of tows  $({\rm N})$  is given for reference.

		Total			Total	
Species	Year	Weight (kg)	Ν	Year	Weight (kg)	N
Sea raven	1957	3.49	30	1979	176.42	92
Sea raven	1958	13.48	28	1980	238.35	80
Sea raven	1959	17.45	52	1981	89.5	49
Sea raven	1960	0	54	1982	279	71
Sea raven	1961	47.21	55	1984	177.04	58
Sea raven	1962	31.98	64	1985	392.96	167
Sea raven	1963	16.25	56	1986	370.68	115
Sea raven	1964	45.29	90	1987	259 22	110
Sea raven	1965	35.57	66	1988	115.83	90
Sea raven	1966	17.9	50	1989	137.2	112
Sea raven	1967	13.61	19	1990	159.15	104
Sea raven	1968	29.94	40	1991	126.05	103
Sea raven	1969	26.31	8	1992	89.34	89
Sea raven	1970	23.14	28	1993	126.21	77
Sea raven	1971	30.38	30	1994	59.76	64
Sea raven	1972	26.08	21	1995	41.63	85
Shorthorn sculpin	1951	0	26	1973	0	50
Shorthorn sculpin	1952	0.75	50	1974	0.91	13
Shorthorn sculpin	1953	1.5	45	1975	0.23	36
Shortborn sculpin	1954	0	40	1976	0	41
Shorthorn sculpin	1955	0	57	1977	0.32	47
Shorthorn sculpin	1956	0	27	1978	0.91	89
Shorthorn sculpin	1957	0	30	1979	0.45	92
Shorthorn sculpin	1958	0	28	1980	0	80
Shorthorn sculpin	1959	0	52	1981	0	49
Shorthorn sculpin	1960	0	54	1982	0.9	71
Shorthorn sculpin	1961	0	55	1984	2.49	58
Shorthorn sculpin	1962	0.68	64	1985	2.32	167
Shorthorn sculpin	1963	0	56	1986	0	115
Shorthorn sculpin	1964	0	90	1987	2.57	110
Shorthorn sculpin	1965	ō	66	1988	0.25	90
Shorthorn sculpin	1966	0	50	1989	1	112
Shorthorn sculpin	1967	0	19	1990	3.29	104
Shorthorn sculpin	1968	0	40	1991	0.84	103
Shorthorn sculpin	1969	0	8	1992	5.38	89
Shorthorn sculpin	1970	0	28	1993	1.09	77
Shorthorn sculpin	1971	0	30	1994	0.33	64
Shorthorn sculpin	1972	0	21	1995	0	85
Silver hake	1951	0.94	26	1973	0	50
Silver hake	1952	3.78	50	1974	0	13
Silver hake	1953	2334.62	45	1975	1.36	36
Silver hake	1954	14.64	40	1976	1.59	41
Silver hake	1955	14.34	57	1977	0.45	47
Silver hake	1956	6.24	27	1978	56.56	89
Silver hake	1957	1	30	1979	29.78	92
Silver hake	1958	4.88	28	1980	10.1	80
Silver hake	1959	2.05	52	1981	318.55	49
Silver hake	1960	34.38	54	1982	0.1	71
Silver hake	1961	31.97	55	1984	0	58
Silver hake	1962	9.07	64	1985	35.28	167
Silver hake	1963	0	56	1986	123.75	115
Silver hake	1964	0	90	1987	25.07	110

Table B.1 Total weight for each species and year surveyed on southern Grand Bank. The ordering of the species is alphabetical. The total number of tows (N) is given for reference.

		Total			Total	
Species	Year	Weight (kg)	N	Year	Weight (kg)	N
20 I I	10.05	0.03	~	1088	0.49	00
Silver hake	1965	0.23	50	1090	1.09	112
Suver hake	1900	0	10	1000	1.45	104
Silver hake	1967	0.00	19	1990	1.95	102
Silver hake	1968	0.43	40	1002	0.01	103
Suver nake	1969	1.00	~	1002	0.00	
Sliver have	1970	1.36	20	1004	1.04	64
Sliver nake	1971	0	30	1005		95
Suver hake	1972	0.75	20	1990	4.98	50
Smooth skate	1991	10.75	50	1074		12
Smooth skate	1902	12.10	30	1075	0.45	26
Smooth skate	1903	0.75	40	1975	0.45	41
Smooth skate	1904	1.5	57	1077	2.26	47
Smooth skate	1965	4.74	27	1078	37.13	89
Smooth skate	1900	0.00	20	1070	217	02
Smooth skate	1987	0.99	30	1000	5.11	80
Smooth skate	1998	10.58	20	1081		49
Smooth skate	1909	10.56	52	1081		45
Smooth skate	1960	9.00		1084		59
Smooth skate	1961	11.57	00	1005	0.84	167
Smooth skate	1962	11.01		1000	2.0	115
Smooth skate	1903	14.21	30	1980	0.00	110
Smooth skate	1964	31.09	90	1987	2.99	00
Smooth skate	1965	13.06	00	1966	2.00	110
Smooth skate	1966	13.04	10	1000	9.12	104
Smooth skate	1907	0.00	19	1001	0.13	102
Smooth skate	1968	0.25	40	1002	4.47	80
Smooth skate	1969	0	~	1002	0.97	
Smooth skate	1970	6.26	20	1004	0.03	64
Smooth skate	1971	0.30	30	1005	1.00	er
Smooth skate	19/2	0	21	1073	1.00	100
Spiny tumptish	1901	0	100	1074	0	26
Spiny tumptish	1962	0	200	1075	0.14	79
Spiny lumphsh	1953	0	90	1975	0.14	82
Spiny lumpish	1904	0	114	1077	0	0.4
Spiny lumphsn	1900	0		1079	0.08	179
Spiny lumpian	1900	0	60	1070	0.2	184
Spiny lumpasa	1967	0	50	1090	0.11	160
Spiny lumptish	1968	0	104	1981	0.11	98
Spiny lumpian	1909	0	108	1082	0.15	142
Spiny lumptish	1960	0.06	110	1984	0.10	116
Spiny lumplish	1001	0.00	120	1085	0	334
Spiny tumpish	1962	0.04	112	1096	0.12	230
Spiny tumplish	1064	0.04	190	1087	0.2	220
Spiny lumpian	1004	0.12	122	1000	0.12	180
Spiny tumption	1066	0	100	1989	0.26	224
Solay lumpfish	1960	0	38	1990	0.24	208
Spiny lumpfish	1967	0	80	1991	0.06	206
Colory tumptish	1960	0	16	1992	0.66	178
Spiny humplish	1970	0	56	1993	0.25	154
Colory tumpfish	1071	0	60	1994	0.19	128
Spiny lumpfish	1972	0.23	42	1995	0.57	170
APPROX AND ADDRESS AND ADDRESS						

Table B.1 Total weight for each species and year surveyed on southern Grand Bank. The ordering of the species is alphabetical. The total number of tows (N) is given for reference.

		Total			Total	
Species	Year	Weight (kg)	N	Year	Weight (kg)	N
		0.100				
Spotted wolffish	1951	0	26	1973	0	50
Spotted wolffish	1952	0	50	1974	0	13
Spotted wolffish	1953	0	45	1975	0	36
Spotted wolffish	1954	0	40	1976	0	41
Spotted wolffish	1955	0	57	1977	0	47
Spotted wolffish	1956	0	27	1978	16.86	89
Spotted wolffish	1957	0	30	1979	0	92
Spotted wolffish	1958	0	28	1980	0	80
Spotted wolffish	1959	0	52	1981	7.5	49
Spotted wolffish	1960	0	54	1982	4.1	71
Spotted wolffish	1961	3.63	55	1984	17.02	58
Spotted wolffish	1962	7.71	64	1985	0	167
Spotted wolffish	1963	7.53	56	1986	0	115
Spotted wolffish	1964	13.61	90	1987	0	110
Spotted wolffish	1965	15.25	66	1988	11.62	90
Spotted wolffish	1966	0	50	1989	0	112
Spotted wolffish	1967	0	19	1990	0	104
Spotted wolffish	1968	30.84	-40	1991	9.34	103
Spotted wolffish	1969	0	8	1992	0	89
Spotted wolffish	1970	9.53	28	1993	0	77
Spotted wolffish	1971	0	30	1994	0	64
Spotted wolffish	1972	0	21	1995	0	85
Striped wolffish	1951	6.95	26	1973	180.99	50
Strined wolfEsh	1952	39.91	50	1974	87.55	13
Striped wolffish	1953	25.86	45	1975	186.66	36
Striped wolffish	1954	37.42	40	1976	195.52	41
Strined wolffish	1955	14.96	57	1977	86.18	47
Strined wolffish	1956	7.99	27	1978	228.47	89
Striped wolffish	1957	66.36	30	1979	442.27	92
Strined wolffish	1958	1.99	28	1980	462.65	80
Stringd wolffish	1959	61.28	52	1981	65	49
Striped wolffieh	1960	157.3	54	1982	630.5	71
Striped wolffich	1961	64.41	55	1984	305.02	58
Striped wolffich	1062	151.94	64	1085	575 10	167
Striped wolffish	1963	55.42	56	1986	751 15	115
Striped wolffich	1064	154.24	90	1987	739.7	110
Striped wolffich	1065	50.68	66	1000	407.67	90
Striped wollfish	1066	192.00	50	1080	247 55	112
Scriped wollish	1990	10 02	10	1000	607.04	104
Striped wolffish	1069	19.36	40	1990	288.82	103
Striped wolmsn	1908	15.99	40	1002	300.03	80
Striped wollinsh	1070	1.20	20	1002	204 65	77
Striped wollinsh	1970	1.30	20	1004	407.10	84
Striped wolmsn	1971	0.44	30	1994	430.12	04
Striped wolmsn	1972	30.29	21	1995	133.57	80
Thorny skate	1951	104.43	20	1973	1394.59	30
1 norny skate	1952	1094.72	00	1974	172.51	1.5
Thorny skate	1953	1328.45	45	1975	a16.29	30
Thorny skate	1954	1351.48	40	1976	1264.39	41
Thorny skate	1955	239.15	57	1977	1164.86	47
Thorny skate	1956	177.87	27	1978	906.59	09
Thorny skate	1957	651.52	30	1979	939.82	92
Thorny skate	1958	421.18	28	1980	1395.6	80

Table B.1 Total weight for each species and year surveyed on southern Grand Bank. The ordering of the species is alphabetical. The total number of tows (N) is given for reference.

		Total			Total	
Species	Year	Weight (kg)	N	Year	Weight (kg)	N
Thorny skate	1959	2275.78	52	1981	493.4	49
Thorny skate	1960	1491.5	54	1982	1233.6	71
Thorny skate	1961	1600.89	55	1984	926.86	58
Thorny skate	1962	2659.96	64	1985	3740.85	167
Thorny skate	1963	2065.85	56	1986	2631.39	115
Thorny skate	1964	2380.83	90	1987	1379.21	110
Thorny skate	1965	1709.92	66	1988	1346.97	90
Thorny skate	1966	805.15	50	1989	1261.72	112
Thorny skate	1967	655.9	19	1990	1656.61	104
Thorny skate	1968	467.45	40	1991	670.69	103
Thorny skate	1969	238.15	8	1992	786.32	89
Thorny skate	1970	940.78	28	1993	455.89	77
Thorny skate	1971	550.88	30	1994	234.05	64
Thorny skate	1972	555.88	21	1995	370.47	85
Threebeard rockling	1951	0	26	1973	0	50
Threebeard rockling	1952	0	50	1974	0	13
Threebeard rockling	1953	0	45	1975	0	36
Threebeard rockling	1954	0	40	1976	0.05	41
Threebeard rockling	1955	0	57	1977	0.09	47
Threebeard rockling	1956	0	27	1978	0.1	89
Threebeard rockling	1957	0	30	1979	0	92
Threebeard rockling	1958	0	28	1980	0	80
Threebeard rockling	1959	0	52	1981	0	49
Threebeard rockling	1960	0	54	1982	0	71
Threebeard rockling	1961	0.09	55	1984	0	58
Threebeard rockling	1962	0	64	1985	0	167
Threebeard rockling	1963	0	56	1986	0	115
Threebeard rockling	1964	0	90	1987	0	110
Threebeard rockling	1965	0	66	1988	0	90
Threebeard rockling	1966	0	50	1989	0	112
Threebeard rockling	1967	0	19	1990	0	104
Threebeard rockling	1968	0	40	1991	0	103
Threebeard rockling	1969	0	8	1992	0	89
Threebeard rockling	1970	0	28	1993	0	77
Threebeard rockling	1971	0	30	1994	0	64
Threebeard rockling	1972	0	21	1995	0	85
Vahl's eelpout	1951	0	26	1973	0.23	50
Vahl's eelpout	1952	0	50	1974	0	13
Vahl's eelpout	1953	0	45	1975	0.18	36
Vahl's eelpout	1954	0	40	1976	0.45	41
Vahl's eelpout	1955	0	57	1977	0	47
Vahl's eelpout	1956	0	27	1978	25.56	89
Vahl's eelpout	1957	0	30	1979	1.13	92
Vahl's eelpout	1958	0	28	1980	0.5	80
Vahl's eelpout	1959	0	52	1981	0	49
Vahl's eelpout	1960	0	54	1982	0	71
Vahl's eelpout	1961	0	55	1984	0	58
Vahl's eelpout	1962	1.29	64	1985	0.66	167
Vahl's celpout	1963	0	56	1986	0	115
Vahl's eelpout	1964	2.26	90	1987	0	110
Vahl's eelpout	1965	0.23	66	1988	0	90
Vahl's eelpout	1966	0.42	50	1989	0	112

Table B.1 Total weight for each species and year surveyed on southern Grand Bank. The ordering of the species is alphabetical. The total number of tows (N) is given for reference.

		Total	N	Year	Total	
Species	Year	Weight (kg)	N	Year	Weight (kg)	N
Vahl's eelpout	1967	7.71	19	1990	0.04	104
Vahl's eelpout	1968	0	40	1991	0	103
Vahl's eelpout	1969	0	8	1992	0	89
Vahl's celpout	1970	0	28	1993	0	77
Vahl's eelpout	1971	0	30	1994	0	64
Vahl's eelpout	1972	0	21	1995	0	85
White hake	1951	608.73	26	1973	116.35	50
White hake	1952	1635.18	50	1974	0	13
White hake	1953	1139.15	45	1975	24.04	36
White hake	1954	328.44	40	1976	95.93	41
White hake	1955	59.39	57	1977	24.94	47
White hake	1956	88.81	27	1978	472.67	89
White hake	1957	86.81	30	1979	220.65	92
White hake	1958	51.85	28	1980	57.5	80
White hake	1959	696.8	52	1981	331.3	49
White hake	1960	1079.53	54	1982	0.1	71
White hake	1961	1741.57	55	1984	28.64	58
White hake	1962	1936.75	64	1985	61.42	167
White hake	1963	1484.39	56	1986	21.58	115
White hake	1964	7725.97	90	1987	157.66	110
White hake	1965	116.68	66	1988	126.12	90
White hake	1966	949.07	50	1989	28.3	112
White hake	1967	0	19	1990	6.89	104
White hake	1968	181.89	40	1991	15.27	103
White hake	1969	0	8	1992	10.79	89
White hake	1970	31.98	28	1993	18.05	77
White hake	1971	43.37	30	1994	52.37	64
White hake	1972	16.33	21	1995	2.41	85
Witch flounder	1951	144.21	26	1973	286.23	50
Witch flounder	1952	205.21	50	1974	0	13
Witch flounder	1953	352.15	45	1975	58.96	36
Witch flounder	1954	241.99	40	1976	278.64	41
Witch flounder	1955	78.88	57	1977	74.62	47
Witch flounder	1956	132.37	27	1978	77.08	89
Witch flounder	1957	91.3	30	1979	73.33	92
Witch flounder	1958	135.23	28	1980	173.5	80
Witch flounder	1959	814.13	52	1981	105	49
Witch flounder	1960	163.9	54	1982	167	71
Witch flounder	1961	329.99	55	1984	111.05	58
Witch flounder	1962	1161.2	64	1985	489.37	167
Witch flounder	1963	342.87	56	1986	229	115
Witch flounder	1964	841.63	90	1987	250.7	110
Witch flounder	1965	286.03	66	1988	419.4	90
Witch flounder	1966	517.23	50	1989	265.56	112
Witch flounder	1967	4.54	19	1990	192.54	104
Witch flounder	1968	136.76	40	1991	99.99	103
Witch flounder	1969	7.48	8	1992	154.16	89
Witch flounder	1970	8.75	28	1993	50.7	77
Witch flounder	1971	18.15	30	1994	85.74	64
Witch flounder	1972	25.85	21	1995	29.63	85
Yellowtail flounder	1951	2.99	26	1973	1906.37	50
Yellowtail flounder	1952	75.49	51	1974	1077.28	13

Table B.1 Total weight for each species and year surveyed on southern Grand Bank. The ordering of the species is alphabetical. The total number of tows (N) is given for reference.

Section	Vere	Total	N	Veen	Total	N	
opecies	rear	weight (kg)	14	rear	weight (ng)	14	
Yellowtail flounder	1953	108.78	45	1975	1606.39	36	
Yellowtail flounder	1954	135.37	40	1976	2094.24	41	
Yellowtail flounder	1955	34.33	57	1977	2542.19	47	
Yellowtail flounder	1956	49.7	27	1978	4463.43	89	
Yellowtail flounder	1957	121.69	30	1979	4790.51	92	
Yellowtail flounder	1958	257.3	28	1980	4651.87	80	
Yellowtail flounder	1959	241.27	52	1981	2114.5	49	
Yellowtail flounder	1960	144.23	54	1982	2780.4	71	
Yellowtail flounder	1961	298.25	55	1984	3214.53	58	
Yellowtail flounder	1962	743.44	64	1985	6815.42	167	
Yellowtail flounder	1963	684.93	56	1986	4435.6	115	
Yellowtail flounder	1964	1677.53	90	1987	3800.03	110	
Yellowtail flounder	1965	1377.25	66	1988	1968.05	90	
Yellowtail flounder	1966	1278.44	50	1989	2337.58	112	
Yellowtail flounder	1967	493.73	19	1990	2991.73	104	
Yellowtail flounder	1968	1104.52	40	1991	2563.03	103	
Yellowtail flounder	1969	365.15	8	1992	1533.83	89	
Yellowtail flounder	1970	1089.1	28	1993	2217.12	77	
Yellowtail flounder	1971	728.24	30	1994	1093.41	64	
Yellowtail flounder	1972	1494.15	21	1995	1894.17	85	

Table B.1 Total weight for each species and year surveyed on southern Grand Bank. The ordering of the species is alphabetical. The total number of tows  $\left(N\right)$  is given for reference.

## Appendix D

# St. Pierre Bank Data Used in Chapter 2

Table C.1 Total weight for each species and year surveyed on southern Grand Bank. The ordering of the species is alphabetical. The total number of tows (N) is given for reference.

Species	Year	Total Weight (kg)	N	Year	Total Weight (kg)	N
						_
American plaice	1951	103.31	26	1973	1870.63	50
American plaice	1952	1859.69	50	1974	478.99	13
American plaice	1953	504.83	45	1975	1368.03	36
American plaice	1954	317.05	40	1976	1791.72	41
American plaice	1955	179.97	57	1977	2045.76	47
American plaice	1956	431.22	27	1978	3943.79	89
American plaice	1957	594.35	30	1979	2999.55	92
American plaice	1958	418.56	28	1980	3731	80
American plaice	1959	1613.44	52	1981	2579.5	49
American plaice	1960	930.32	54	1982	2255.8	71
American plaice	1961	3123.02	55	1984	2474.64	58
American plaice	1962	4743.28	64	1985	11741.18	167
American plaice	1963	2922.42	56	1986	3829.29	115
American plaice	1964	3763.52	90	1987	3794.14	110
American plaice	1965	6138.58	66	1988	2550.84	90
American plaice	1966	2670.32	50	1989	3033.01	112
American plaice	1967	851.87	19	1990	2946.01	104
American plaice	1968	1191.6	40	1991	1753.6	103
American plaice	1969	406.42	8	1992	871.67	89
American plaice	1970	1394.12	28	1993	1090.75	77
American plaice	1971	2245.56	30	1994	487.52	64
American plaice	1972	1076.84	21	1995	426.5	85
Arctic cod	1951	0	26	1973	0	50
Arctic cod	1952	0	50	1974	0	13
Arctic cod	1953	0	45	1975	0	36
Arctic cod	1954	0	40	1976	0	41
Arctic cod	1955	0	57	1977	0	47
Arctic cod	1956	0	27	1978	0	89
		Total			Total	
--------------------	------	-------------	-----	------	-------------	-----
Species	Year	Weight (kg)	N	Year	Weight (kg)	N
Arctic cod	1957	0	30	1979	0	92
Arctic cod	1958	0	28	1980	0	80
Arctic cod	1959	0	52	1981	0	49
Arctic cod	1960	0	54	1982	0	71
Arctic cod	1961	0	55	1984	0	58
Arctic cod	1962	0	64	1985	0	167
Arctic cod	1963	0	56	1986	0.04	115
Arctic cod	1964	0	90	1987	0.08	110
Arctic cod	1965	0	66	1988	0.01	90
Arctic cod	1966	0	50	1989	0.04	112
Arctic cod	1967	0	19	1990	0.01	104
Arctic cod	1968	0	40	1991	0.02	103
Arctic cod	1969	0	8	1992	0	89
Arctic cod	1970	0	28	1993	0.02	77
Arctic cod	1971	0	30	1994	0.01	64
Arctic cod	1972	0	21	1995	0	85
Arctic eelpout	1951	0	26	1973	0	50
Arctic eelpout	1952	0	50	1974	0	13
Arctic eelpout	1953	0	45	1975	0	36
Arctic eelpout	1954	0	40	1976	0	41
Arctic eelpout	1955	0	57	1977	0	47
Arctic eelpout	1956	0	27	1978	0	89
Arctic eelpout	1957	0	30	1979	0	92
Arctic eelpout	1958	0	28	1980	0	80
Arctic eelpout	1959	0	52	1981	10.75	49
Arctic eelpout	1960	0	54	1982	12.4	71
Arctic eelpout	1961	0	55	1984	1.91	58
Arctic eelpout	1962	0	64	1985	28.1	167
Arctic eelpout	1963	0	56	1986	12.12	115
Arctic eelpout	1964	0	90	1987	10	110
Arctic eelpout	1965	0	66	1988	3.07	90
Arctic eelpout	1966	0	50	1989	7.39	112
Arctic eelpout	1967	0	19	1990	16.7	104
Arctic eelpout	1968	0	40	1991	9.45	103
Arctic eelpout	1969	0	8	1992	4.56	89
Arctic eelpout	1970	0	28	1993	2.95	77
Arctic eelpout	1971	0	30	1994	1.11	64
Arctic eelpout	1972	5.45	21	1995	2.81	85
Atlantic argentine	1951	0	52	1973	0.23	100
Atlantic argentine	1952	6.88	100	1974	0	26
Atlantic argentine	1953	1.81	90	1975	2.49	72
Atlantic argentine	1954	0	80	1976	0.05	82
Atlantic argentine	1955	0	114	1977	0.23	94
Atlantic argentine	1956	0	54	1978	6.43	178
Atlantic argentine	1957	0	60	1979	23.16	184
Atlantic argentine	1958	0	56	1980	0.7	160
Atlantic argentine	1959	5.13	104	1981	0	38
Atlantic argentine	1960	52.03	108	1982	0	142
Atlantic argentine	1961	2.18	110	1984	1.33	116
Atlantic argentine	1962	4.54	128	1985	0.42	334
Atlantic argentine	1963	313.59	112	1986	29.46	230
Atlantic argentine	1964	34.18	180	1987	0	220

Table C.1 Total weight for each species and year surveyed on southern Grand Bank. The ordering of the species is alphabetical. The total number of tows (N) is given for reference.

Species	Year	Total Weight (kg)	N	Year	Total Weight (kg)	N
Atlantic argentine	1965	4.27	132	1988	0	180
Atlantic argentine	1966	59.64	100	1989	0	224
Atlantic arrentine	1967	0	38	1990	0.35	208
Atlantic argentine	1968	70.76	80	1991	0	206
Atlantic argentine	1969	0	16	1992	1.25	178
Atlantic argentine	1970	0	56	1993	0.98	154
Atlantic argentine	1971	0	60	1994	0.18	128
Atlantic argentine	1972	0	42	1995	1.41	170
Atlantic cod	1951	1621.82	26	1973	714.44	50
Atlantic cod	1952	4227.76	50	1974	81.18	13
Atlantic cod	1953	3277.31	45	1975	587.29	36
Atlantic cod	1954	7118.99	40	1976	367.13	41
Atlantic cod	1955	6849.32	57	1977	1510	47
Atlantic cod	1956	7912 11	27	1978	2506.76	89
Atlantic cod	1957	1530.45	30	1979	3937.41	92
Atlantic cod	1958	1295 98	28	1980	1549.15	80
Atlantic cod	1959	5912 23 -	52	1981	1577.7	49
Atlantic cod	1960	4008 15	54	1982	1639 32	71
Atlantic cod	1961	5091.08	55	1984	5111.14	58
Atlantic cod	1962	3508 3	64	1985	17672 87	167
Atlantic cod	1062	2865 55	56	1086	9854 82	115
Atlantic cod	1064	8020.82	00	1087	12607.56	110
Atlantic cod	1965	5070 13	66	1988	6272 14	90
Atlantic cod	1066	2227 81	50	1080	6612.15	112
Atlantic cod	1067	1995 26	10	1000	6602 52	104
Actancic cou	1068	2012.44	40	1001	2720.1	102
Atlantic cod	1060	1004 9	40	1002	1776 21	80
Atlantic cou	1070	1252.00	28	1002	1024	
Atlantic cod	1970	1203.55	20	1004	120.47	64
Atlantic cod	1971	2007.79	30	1005	130.47	0.4
Atlantic cod	1051	652.02	21	1072	217.7	80
Atlantic hagisa	1901	0	20	1074	0	12
Atlantic nagisa	1902	0	45	1075	0	1.5
Atlantic hagist	1054	0	40	1076	0	41
Atlantic hagish	1904	0	40	1077	0	47
Atiantic nagian	1900	0	07	1078	0	
Atlantic nagnan	1956	0	21	1978	0	89
Atlantic hagtish	1957	0	30	1080	0	92
Atlantic nagnan	1908	0	20	1081	0	00
Atlantic nagrish	1939	0	02	1981	0	-19
Atlantic nagnan	1900	0	24	1982	0	11
Atlantic haghsh	1901	0.45	20	1984	0	38
Atlantic nagnan	1902	0.23	04	1960	0	107
Atlantic hagnsh	1903	U	30	1980	0	115
Atlantic naghsh	1964	0	90	1088	0	110
Atlantic nagfish	1965	0 00	50	1088	0	30
Atlantic nagnsh	1900	0.23	30	1000	0	104
Atlantic nagfish	1967	0	19	1001	0	104
Atlantic nagfish	1968	0	40	1991	0	103
Atlantic hagfish	1969	0	8	1992	0	89
Atlantic nagfish	1970	0	28	1993	0	
Atlantic hagfish	1971	0	30	1994	0	04
Atlantic hagfish	1972	0	21	1995	0	85

Table C.1 Total weight for each species and year surveyed on southern Grand Bank. The ordering of the species is alphabetical. The total number of tows (N) is given for reference.

				Total		
Species	Year	Weight (kg)	N	Year	Weight (kg)	N
Atlantic halibut	1051		26	1077	15.42	50
Atlantic halibut	1951	48.47	50	1973	10.42	13
Atlantic halibut	1053	41.01	45	1975	7.94	36
Atlantic halibut	1954	32 33	40	1976	68.04	41
Atlantic halibut	1055	10.03	57	1977	43.55	47
Atlantia halibut	1056	21.45	27	1078	42.25	80
Atlantic hallout	1057	50.96	20	1070	20.42	02
Atlantic hallbut	1059	4.00	28	1080	245.8	80
Atlantic halibut	1050	172 54	50	1091	81	40
Atlantic hallbut	1960	272 74	5.4	1082	140.9	71
Atlantic halibut	1061	541.26	55	1084	02.06	59
Atlantic hallbut	1062	592.02	64	1085	283 44	167
Atlantic hallout	1902	150.00	5-2	1086	220.44	115
Atlantic hallout	1903	100.00	30	1087	235.40	110
Atlantic hallout	1904	173.08	90	1000	220.32	00
Achancie nanout	1903	333.09	50	1080	120.0	110
Atlantic hallout	1966	110.03	10	1989	20.42	104
Atlantic hallbut	1967	00 48	19	1990	20.43	104
Atlantic hallout	1908	29.40	40	1000	61.13	103
Atlantic hallout	1969	41.28	0	1992	8.01	09
Atlantic halibut	1970	8.62	28	1993	3.01	
Atlantic hallout	1971	36.1	30	1994	3.70	04
Atlantic halibut	1972	0	21	1995	10.96	60
Barndoor skate	1951	10.00	20	1973	0	10
Barndoor skate	1952	15.54	30	1974	0	13
Barndoor skate	1953	44.91	45	1975	0	30
Barndoor skate	1904	31.08	40	1910	0	41
Barndoor skate	1955	0	57	1977	0	97
Barndoor skate	1956	0	27	1978	0	89
Barndoor skate	1957	0	30	19/9	0	92
Barndoor skate	1958	15.54	28	1980	0	80
Barndoor skate	1959	0	52	1981	0	49
Barndoor skate	1960	110.68	54	1982	0	11
Barndoor skate	1961	90.72	55	1984	0	38
Barndoor skate	1962	61.23	04	1985	0	107
Barndoor skate	1963	0	56	1980	0	115
Barndoor skate	1964	9.98	90	1987	0	110
Barndoor skate	1965	0	66	1988	0	90
Barndoor skate	1966	27.21	50	1989	0	112
Barndoor skate	1967	0	19	1990	0	104
Barndoor skate	1968	0	40	1991	0	103
Barndoor skate	1969	0	8	1992	0	89
Barndoor skate	1970	0	28	1993	0	11
Barndoor skate	1971	0	30	1994	0	64
Barndoor skate	1972	0	21	1995	0	85
Blue whiting	1951	0	26	1973	0	50
Blue whiting	1952	0	50	1974	0	13
Blue whiting	1953	0	45	1975	0	36
Blue whiting	1954	0	40	1976	0	41
Blue whiting	1955	0	57	1977	0	47
Blue whiting	1956	0	27	1978	0.2	89
Blue whiting	1957	0	30	1979	0	92
Blue whiting	1059	0	78	1980	0	80

Table C.1 Total weight for each species and year surveyed on southern Grand Bank. The ordering of the species is alphabetical. The total number of tows (N) is given for reference.

Species  Yar  Weight (bg)  N  Yar  Weight (bg)  N    Bins whiting  1991			These			Tread	
Bits  Bits <th< th=""><th>Species</th><th>Year</th><th>Weight (kg)</th><th>N</th><th>Year</th><th>Weight (kg)</th><th>N</th></th<>	Species	Year	Weight (kg)	N	Year	Weight (kg)	N
Blue whiting 1591 0.6 2.1 812 0.6 41 Blue whiting 1591 0.6 51 1848 0.6 41 Blue whiting 1592 0.7 64 1859 0.5 17 Blue whiting 1592 0.7 64 1859 0.7 1.0 110 Blue whiting 1593 0.1 56 1858 0.0 00 Blue whiting 1594 0.5 18 66 1858 0.0 00 Blue whiting 1595 0.1 56 1858 0.0 10 Blue whiting 1597 0.5 18 66 1858 0.0 10 Blue whiting 1597 0.5 18 66 1959 0.0 110 Blue whiting 1597 0.0 20 1992 0.0 84 Blue whiting 1597 0.0 21 1992 0.0 84 Blue whiting 1597 0.0 30 1994 0.0 13 Blue whiting 1597 0.0 30 1994 0.0 13 Blue whiting 1597 0.0 30 1997 0.0 33 Breadhaut wolfinh 1533 0.4 54 1973 0.0 33 Breadhaut wolfinh 1558 0.4 27 1978 1.5 40 Breadhaut wolfinh 1598 0.4 90 1979 0.1 34 Breadhaut wolfinh 1598 0.4 90 1979 0.1 34 Breadhaut wolfinh 1598 0.0 49 1977 0.0 34 Breadhaut wolfinh 1598 0.0 49 1978 0.5 40 Breadhaut wolfinh 1598 0.0 49 1978 0.5 40 Breadhaut wolfinh 1598 0.0 49 1978 0.0 10 Breadhaut wolfinh 1598 0.0 49 1978 0.0 10 Breadhaut wolfinh 1598 0.0 198 0.0 110 Breadhaut wolfinh 1598 0.0 199 0.0 110 Breadhaut wolfinh 1598 0.0 199 0.0 100 Breadhaut wolfinh 1598 0.0 199 0.0 100 Breadhaut wolfinh 1597 0.0 30 1989 0.0 110 Breadhaut wolfinh 1597 0.0 199 1990 0.0 110 Breadhaut wolfinh 1597 0.0 199 1990 0.0 110 Breadhaut wolfinh 1597 0.1 35 21 1978 0.0 10 Breadhaut wolfinh 1597 1.1 35 21 1978 0.0 45 Breadhaut wolfinh 1597 1.1 35 21 1978 0.0 10 Breadhaut wolfinh 1597 1.1 35 21 1978 0.0 45 Breadhaut wolfinh 1597 1.1 35 21 1978 0.0 110 Breadhaut wolfinh 1597 1.1 35 21 1978 0.0 110 Breadhaut wolfinh 1597 1.1 35 21 1977 1.2 40 Breadhaut wolfinh 1597 1.1 35 20 1977 1.1 48 0.0 10 Breadhaut wolfinh 1597 1.1 35 20 1977 1.1 48 0.0 10 Breadhaut wolfinh 1597 1.1 35 20 1977 1.2 41.4 40 Capelin 1550 0.5 24 0.1 1977 1.2 4.3 4.4 11.5 77 Breadha			0 ( 0)				
	Blue whiting	1959	0	52	1981	0	49
	Blue whiting	1960	0.18	54	1982	0	71
	Blue whiting	1961	0	55	1984	0	58
	Blue whiting	1962	0	64	1985	0.83	167
	Blue whiting	1963	0.79	56	1986	0.17	115
	Blue whiting	1964	0.54	90	1987	0	110
	Blue whiting	1965	0.18	66	1988	0	90
Bise white the set of the set	Blue whiting	1966	0	50	1989	0	112
	Blue whiting	1967	0	19	1990	0	104
Bise whiting 1969 0 \$ \$ 1920 0 \$ \$ 1920 0 \$ \$ 74 \$ \$ 1920 0 \$ \$ 74 \$ \$ 1920 0 \$ \$ 74 \$ \$ 1920 0 \$ \$ 74 \$ \$ 1920 0 \$ \$ 74 \$ \$ 1920 0 \$ \$ 74 \$ \$ 1920 0 \$ \$ 10 \$ \$ 100 0 \$ \$ 1	Blue whiting	1968	4.31	40	1991	0	103
Bisse white \$\$ 1571 0 \$ 38 \$ 1932 \$ 0 \$ 76 \$ 45 \$ 1935 \$ 0 \$ 76 \$ 45 \$ 1975 \$ 0 \$ 21 \$ 1956 \$ 0 \$ 65 \$ 1975 \$ 0 \$ 25 \$ 1975 \$ 0 \$ 25 \$ 1975 \$ 0 \$ 25 \$ 1975 \$ 0 \$ 25 \$ 1975 \$ 0 \$ 25 \$ 1975 \$ 0 \$ 25 \$ 1975 \$ 0 \$ 25 \$ 1975 \$ 0 \$ 25 \$ 1975 \$ 0 \$ 25 \$ 1975 \$ 0 \$ 25 \$ 1975 \$ 0 \$ 25 \$ 1975 \$ 0 \$ 25 \$ 1975 \$ 0 \$ 25 \$ 1975 \$ 15.5 \$ 46 \$ 1975 \$ 0 \$ 25 \$ 1975 \$ 1.5 \$ 3 \$ 46 \$ 1975 \$ 0 \$ 25 \$ 1975 \$ 1.5 \$ 3 \$ 46 \$ 1975 \$ 0 \$ 25 \$ 1975 \$ 1.5 \$ 3 \$ 46 \$ 1975 \$ 0 \$ 25 \$ 1975 \$ 1.5 \$ 3 \$ 46 \$ 1975 \$ 0 \$ 35 \$ 1975 \$ 1.5 \$ 3 \$ 46 \$ 1975 \$ 1.5 \$ 3 \$ 46 \$ 1975 \$ 1.5 \$ 3 \$ 46 \$ 1975 \$ 1.5 \$ 3 \$ 46 \$ 1975 \$ 1.5 \$ 3 \$ 46 \$ 1975 \$ 1.5 \$ 3 \$ 46 \$ 1975 \$ 1.5 \$ 3 \$ 46 \$ 1975 \$ 1.5 \$ 3 \$ 46 \$ 1975 \$ 1.5 \$ 3 \$ 46 \$ 1975 \$ 1.5 \$ 3 \$ 46 \$ 1975 \$ 1.5 \$ 3 \$ 46 \$ 1975 \$ 1.5 \$ 10 \$ 10 \$ 10 \$ 10 \$ 10 \$ 10 \$ 10 \$ 1	Blue whiting	1969	0	8	1992	0	89
	Blue whiting	1970	0	28	1993	0	77
Bise white the set of the set	Blue whiting	1971	0	30	1994	0	64
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	Blue whiting	1972	0	21	1995	0	85
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	Broadhead wolffish	1951	0	26	1973	0	50
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	Broadhead wolffish	1952	8.66	50	1974	0	13
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	Broadhead wolffish	1953	0	45	1975	0	36
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	Broadhead wolffish	1954	0	40	1976	0	41
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	Broadhead wolffish	1955	0	57	1977	7.26	47
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	Broadhead wolffish	1956	0	27	1978	15.5	89
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	Broadhead wolffish	1957	0	30	1979	35.41	92
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	Broadhead wolffish	1958	0	28	1980	15.1	80
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	Broadhead wolffish	1959	4.99	52	1981	73	49
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	Broadhead wolffish	1960	10.89	54	1982	0	71
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	Broadhead wolffish	1961	0	55	1984	0	58
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	Broadhead wolffish	1962	0	64	1985	4.15	167
	Broadhead wolffish	1963	0	56	1986	12.45	115
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	Broadhead wolffish	1964	0	90	1987	0	110
Broadhad wolfini  1968  21.33  60  101  102    Broadhad wolfini  1967  0.40  190  0.0  103    Broadhad wolfini  1969  10.40  190  0.0  103    Broadhad wolfini  1969  10.60  8  1992  1.68  104    Broadhad wolfini  1970  12.82  20  1.98  0  163    Broadhad wolfini  1971  12.82  20  1.985  0  85    Broadhad wolfini  1971  1.32  20  1.975  1.07  1.06  20    Capelin  1951  1.57  20  1.975  1.06  50  1.06  1.02  1.04  50  1.06  1.02  1.06  50  1.075  6.28.1  41  1.06  1.02  1.06  50  1.075  6.28.1  41  1.06  1.06  1.06  1.06  1.06  1.06  1.06  1.06  1.06  1.06  1.06  1.06 <td>Broadhead wolffish</td> <td>1965</td> <td>44.27</td> <td>66</td> <td>1988</td> <td>0</td> <td>90</td>	Broadhead wolffish	1965	44.27	66	1988	0	90
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	Broadhead wolffish	1966	21.33	50	1989	0	112
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	Broadhead wolffish	1967	0	19	1990	0	104
Broadhad wolfini  1980  10.06  8  1992  15.85  89    Broadhad wolfini  1970  12.82  20  10.46  7  6    Broadhad wolfini  1971  13.02  21  1985  0  85    Broadhad wolfini  1971  13.02  21  1985  0  85    Broadhad wolfini  1972  13.17  26  1971  L66  0  10    Capelin  1631  1.51  5  40  1977  L26  0  10  23  10  0.28  41  7  0.28  41  7  0.28  41  7  0.28  41  7  0.28  41  7  0.28  41  7  0.28  41  7  0.28  41  7  0.28  41  7  0.28  41  7  0.28  41  7  0.28  41  7  0.28  41  7  0.28  41  1.25  41  <	Broadhead wolffish	1968	0	40	1991	0	103
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	Broadhead wolffish	1969	19.05	8	1992	15.85	89
Broadbade wolffah  1971  13.61  30  1994  0  64    Broadbade wolffah  1972  0  21  1965  0  85    Capelin  1972  0  21  1973  0.0  85    Capelin  1953  2.43  45  1977  0.0  85    Capelin  1953  2.43  45  1976  0.028  41    Capelin  1955  4.0  77  107  42.83  47    Capelin  1955  0.3  70  1977  42.83  47    Capelin  1955  0.3  57  1977  42.83  47    Capelin  1958  0.35  28  1980  6.7  80    Capelin  1959  3.64  82  188  1.47  49    Capelin  1960  6.54  84  1845  1.55  78  60    Capelin  1960  155.85  28  18.55	Broadbead wolffish	1970	22.68	28	1993	24.61	77
Broadhead wolffab  1972  0  21  1985  0  85    Capelin  1811  115.7  20  197.1  L0  93    Capelin  1981  115.7  20  197.1  L0  93    Capelin  1983  24.3  45  197.5  150.0  83    Capelin  1564  24.3  40  197.7  0.28.4  1    Capelin  1564  24.3  40  197.7  0.28.4  1    Capelin  1564  6.3  97  197.8  16.01.8  41    Capelin  1567  0.05  20  197.7  10.41.8  42    Capelin  1567  0.05  20  187.2  21.1.4  92    Capelin  1560  16.4  1812  1.4.6  41  92    Capelin  1560  15.4  182  1.4.6  71  10.04.1  10.04  10.04  10.04  10.04  10.04  10.04 <td>Broadhead wolffish</td> <td>1971</td> <td>13.61</td> <td>30</td> <td>1994</td> <td>0</td> <td>64</td>	Broadhead wolffish	1971	13.61	30	1994	0	64
Capelin  1931  15.77  26  1973  1.06  50    Capelin  1982  1.3  50  1377  100  101    Capelin  1982  1.3  50  1377  100  100    Capelin  1982  1.3  50  1377  100  100  101    Capelin  1985  4.4  87  1976  100  101  100  101    Capelin  1965  4.3  87  1978  16.61  49  100  11.1  90  10.4  40  100 </td <td>Broadbead wolffish</td> <td>1972</td> <td>0</td> <td>21</td> <td>1995</td> <td>0</td> <td>85</td>	Broadbead wolffish	1972	0	21	1995	0	85
Capelini  1922  1.3  60  1974  0  13    Capelini  1964  2.04  61  1976  80.38  41    Capelini  1954  2.04  61  1977  42.63  47    Capelini  1955  4.0  57  1977  42.63  47    Capelini  1955  4.0  57  1977  42.63  47    Capelini  1956  0.3  57  1977  42.63  47    Capelini  1956  0.3  57  1977  42.64  49    Capelini  1969  0.54  57  197  42.64  49    Capelini  1969  1.44  51  1984  1.57  47    Capelini  1969  1.24  57  1894  2.54  167    Capelini  1962  1.63  46  1894  2.54  197    Capelini  1962  1.53  64  1894  2.54 <td>Capelin</td> <td>1951</td> <td>15.27</td> <td>26</td> <td>1973</td> <td>1.96</td> <td>50</td>	Capelin	1951	15.27	26	1973	1.96	50
Operation  1933  24.0  45  1975  126.08  36    Coperini  1184.4  2.2.8  40  1977  0.2.8  41    Coperini  1184.4  2.2.8  40  1977  0.2.8  41    Coperini  1186.4  0.3  27  1978  16.61  89    Coperini  1587  0.0.8  30  1977  21.1.44  92    Caperini  1587  0.0.8  30  1979  21.1.14  92    Caperini  1580  0.4.5  30  1989  0.4.7  80    Caperini  1560  1.6.4  41  182  31.65  71    Caperini  1560  1.6.4  41  182  31.65  71    Caperini  1560  1.6.4  41  182  32.4.2  31.65    Caperini  1560  1.6.4  41  46  11.5  54  11.5    Caperini  1563  52.3.2  <	Capelin	1952	1.3	50	1974	0	13
Capelin  1954  2.85  40  1976  0.28  41    Capelin  1955  4.0  67  1977  42.83  47    Capelin  1955  4.0  77  1977  16.81  49    Capelin  1956  0.3  77  1977  16.81  49    Capelin  1958  0.3  23  1980  0.7  80    Capelin  1959  3.6.4  52  1881  147  49    Capelin  1960  1.6.4  41  81.85  7.8  7.8    Capelin  1960  1.6.4  41  81.85  7.8  7.8    Capelin  1960  1.6.4  81  81.85  7.8  7.8    Capelin  1960  1.6.4  81  1.85  7.8  7.8    Capelin  1963  1.85  5.8  8.1  8.2  1.15    Capelin  1963  1.86  5.9  19.8  3.5.8	Capelin	1953	24.9	45	1975	1500.59	36
Capelin  1955  -4.9  57  1977  42.83  47    Capelin  1956  0.3  27  1978  16.61  89    Capelin  1957  0.05  30  1979  211.14  92    Capelin  1968  0.05  28  1890  0.77  80    Capelin  1968  1.54  22  1840  0.77  80    Capelin  1968  1.54  21  1898  0.17  80    Capelin  1961  1.24  21  184  2.4  58    Capelin  1961  1.54  51  54  2.24  58    Capelin  1963  15.18  56  185  52.23  167    Capelin  1963  15.18  56  186  2.24  58    Capelin  1963  15.18  60  198  12.54  115    Capelin  1964  81.68  80  198  5.88	Capelin	1954	2.65	40	1976	0.28	41
Capelin  1988  0.3  27  1978  16.61  89    Capelin  1867  0.05  30  1579  21.1.1.4  90    Capelin  1897  0.05  30  1579  21.1.3.4  90    Capelin  1590  0.5.4  31.851  1.47  40    Capelin  1560  6.5.4  51  189.2  31.65  71    Capelin  1560  6.5.4  51  189.2  31.65  71    Capelin  1560  1.5.4  51  189.2  2.34.2  50    Capelin  1560  1.5.4  51  189.4  2.34.2  50    Capelin  1563  51.16.8  60  19.64  12.5.4  115    Capelin  1964  81.1.6.8  90  19.87  3.5.97  110    Capelin  1964  81.6.6  90  1897  3.5.37  100    Capelin  1964  81.6.2  2.5.98  109	Capelin	1955	4.9	57	1977	42.63	47
Capelin  1957  0.05  39  1979  211.14  92    Capelin  1988  0.55  28  1890  0.7  80    Capelin  1969  0.46  82  1981  1.47  49    Logefin  1960  6.45  181  3.1.26  78    Gapelin  1962  26.5.32  64  184  3.1.26  78    Gapelin  1962  26.5.32  64  1985  82.2.34  167    Capelin  1963  15.11.68  59  185  82.2.34  115    Capelin  1963  15.11.68  59  1987  3.5.37  110    Capelin  1964  81.1.68  59  1987  3.5.36  69    Capelin  1964  81.1.68  50  1987  3.5.84  69	Capelin	1956	0.3	27	1978	16.61	89
Capatin  1558  0.35  28  1990  0.7  80    Capatin  1599  31.64  32  1981  147  49    Capatin  1969  6.54  48  1892  31.65  71    Capatin  1960  6.54  48  1892  31.65  71    Capatin  1961  12.61  55  184  2.24  58    Capatin  1962  265.32  64  1985  82.23  167    Capatin  1963  151.18  50  186  2.24  58    Capatin  1963  151.18  50  186  12.54  115    Capatin  1963  151.18  50  1987  313.77  110    Capatin  1964  81.1.86  50  197  333.77  110    Capatin  1964  81.35  60  1386  5.52  90  90	Capelin	1957	0.05	30	1979	211.14	92
Capelin  1999  31.64  52  1981  147  49    Capelin  1990  6.54  44  1982  31.65  71    Capelin  1990  6.54  44  1982  31.65  71    Capelin  1940  1.51  86  1982  3.65  71    Capelin  1940  1.51  86  1983  2.53  87    Capelin  1943  51.68  69  1984  13.7  710    Capelin  1944  81.66  90  1987  325.77  110    Capelin  1965  65.53  69  1987  326.77  10	Capelin	1958	0.55	28	1980	0.7	80
Capelin  1990  6.5.4  54  1982  31.65  71    Capelin  1961  12.61  55  1984  2.24  58    Capelin  1962  265.32  64  1985  52.23  167    Capelin  1962  265.32  64  1985  52.24  18    Capelin  1964  811.68  60  1987  325.77  110    Capelin  1964  63.52  66  1985  53.69  50	Capelin	1959	31.64	52	1981	147	49
Capellin  1961  12.81  P5  1984  2.24  58    Capellin  1962  255.22  64  1985  52.23  167    Capellin  1963  151.96  56  1986  12.54  115    Capellin  1963  151.96  56  1986  12.54  115    Capellin  1963  151.96  90  1987  325.77  110    Capellin  1965  62.32  90  1987  325.97  110	Capelin	1960	6 54	54	1982	31.65	71
Capelin  1967  265.2  50  1969  2.1.3  107    Capelin  1962  265.3  64  1985  52.23  167    Capelin  1963  1511.98  56  1986  12.54  115    Capelin  1964  811.66  90  1987  325.77  110    Capelin  1965  636.23  66  1988  5.98  90    Capelin  1965  637.23  60  1983  5.98  90	Capalin	1961	12.61	55	1984	2.24	58
Origonia  1963  1511.98  56  1986  12.54  115    Capelin  1963  1511.98  56  1986  12.54  115    Capelin  1963  651.05  90  100  100  100    Capelin  1965  662.23  66  1986  43.04  112	Capelin	1962	265 32	64	1985	52.23	167
Capelin 1964 811.66 90 1987 325.77 110 Capelin 1965 636.23 66 1988 5.98 90 Carelin 1966 674 50 1980 43 34 112	Capalin	1963	1511 98	56	1986	12.54	115
Capelin 1965 636.23 66 1988 5.98 90 Capelin 1966 96.74 50 1989 43.94 112	Capelin	1964	811.66	90	1987	325.77	110
Capella 100 0624 50 1960 030 10	Capalin	1065	636 23	66	1988	5.98	90
	Capelin	1966	96 74	50	1989	43.94	112

Table C.1 Total weight for each species and year surveyed on southern Grand Bank. The ordering of the species is alphabetical. The total number of tows (N) is given for reference.

Section	Vere	Total	N	Year	Total Weight (kg)	N
Species	LOW	weight (kg)	14	1.004	meißur (wE)	
Capelin	1967	249.8	19	1990	28	104
Capelin	1968	381.73	40	1991	84.5	103
Capelin	1969	30.09	8	1992	16.4	89
Capelin	1970	220.24	28	1993	408.9	77
Capelin	1971	11.37	30	1994	9.42	64
Capelin	1972	60.88	21	1995	35.6	85
Common ocean pout	1951	0	26	1973	16.33	50
Common ocean pout	1952	0	50	1974	0	13
Common ocean pout	1953	0	45	1975	6.8	36
Common ocean pout	1954	0	40	1976	0	41
Common ocean pout	1955	0	57	1977	0	47
Common ocean pout	1956	0	27	1978	0	89
Common ocean pout	1957	0	30	1979	0	92
Common ocean pout	1958	0	28	1980	0.8	80
Common ocean pout	1959	1.54	52	1981	4.25	49
Common ocean pout	1960	0.75	54	1982	6.4	71
Common ocean pout	1961	0	55	1984	4.81	58
Common ocean pout	1962	0.91	64	1985	5.6	167
Common ocean pout	1963	0	56	1986	0	115
Common ocean pout	1964	0.45	90	1987	2.49	110
Common ocean pout	1965	1.54	66	1988	1	90
Common ocean pout	1966	0.91	50	1989	1.83	112
Common ocean pout	1967	0	19	1990	7.74	104
Common ocean pout	1968	0	40	1991	0	103
Common ocean pout	1969	0	8	1992	0	89
Common ocean pout	1970	õ	28	1993	0	77
Common ocean pout	1971	0	30	1994	0	64
Common ocean pout	1972	23.36	21	1995	0	85
Deepsea sculpin	1951	0	26	1973	0	50
Deepsea sculpin	1952	ō	50	1974	0	13
Deepsea sculpin	1953	0	-45	1975	0	36
Deepsea sculpin	1954	0	40	1976	0	41
Deepsea sculpin	1955	0	57	1977	0	47
Deepsea sculpin	1956	0	27	1978	0	89
Deepsea sculpin	1957	0	30	1979	0	92
Deepsea sculpin	1958	0	28	1980	0	80
Deepsea sculpin	1959	0	52	1981	0	49
Deepsea sculpin	1960	0	54	1982	0	71
Deepsea sculpin	1961	0	55	1984	0	58
Deepsea sculpin	1962	ō	64	1985	0	167
Deepsea sculpin	1963	0	56	1986	0	115
Deepsea sculpin	1964	0	90	1987	0	110
Deepsea sculpin	1965	0	66	1988	0	90
Deepsea sculpin	1966	0	50	1989	0	112
Deepsea sculpin	1967	0	19	1990	0	104
Deepsea sculpin	1968	0	40	1991	0	103
Deepsea sculpin	1969	0	8	1992	0	89
Deepsea sculpin	1970	0	28	1993	0	77
Deepsea sculpin	1971	ō	30	1994	0	64
Deepsea sculpin	1972	ő	21	1995	õ	85
Eelpout	1951	0	26	1973	3.17	50
Eelpout	1952	ő	50	1974	1.36	13

Table C.1 Total weight for each species and year surveyed on southern Grand Bank. The ordering of the species is alphabetical. The total number of tows (N) is given for reference.

		Total			Total	
Species	Year	Weight (kg)	N	Year	Weight (kg)	N
Eelpout	1953	0	45	1975	4.08	36
Eelpout	1954	0	40	1976	3.18	41
Eelpout	1955	0	57	1977	33.34	47
Felpout	1956	0	27	1978	50.53	89
Felnout	1957	0	30	1979	12.93	92
Eelpout	1958	0	28	1980	12.85	80
Felnout	1959	0	52	1981	0	49
Felnout	1960	0	54	1982	0	71
Feloout	1961	19 74	55	1984	0	58
Eelpout	1962	0	64	1985	0	167
Felnout	1963	18.14	56	1986	0	115
Falcout	1964	0.9	90	1987	0	110
Felnout	1065	14.51	66	1088	0	90
Ealpout	1066		50	1090	0	112
Felnout	1967	34 02	19	1990	0	104
Eelpout	1069	34.02	40	1001	0	102
Ealpout	1060	0	0	1002	0	80
Eelpout	1070	7.04	00	1002	0	27
Eerpoor	1970	7.54	20	1004	0	
Eelpout	1971	0	30	1005	0	04
Capacitic astrong	1072	0	21	1990	0	50
Esmark's elipout	1951	0	20	1973	0	30
Esmark's ecipout	1952	0	50	1974	0	13
Esmark's eelpout	1953	0	40	1975	0	30
Esmark s eeipout	1954	0	40	1970	0	41
Esmark's ecipout	1955	0	57	1977	0	91
Esmark's ecipout	1999	U	21	1978	0	89
Esmark's eelpout	1957	0	30	1979	0	92
Esmark's eeipout	1958	0	28	1980	0	80
Esmark's ecipout	1959	0	52	1981	0	49
Esmark's eelpout	1960	0	54	1982	0	71
Esmark's eelpout	1961	0	55	1984	0	58
Esmark's eelpout	1962	0	64	1985	0	167
Esmark's eelpout	1963	U	56	1980	U	115
Esmark's eelpout	1964	0	90	1987	0	110
Esmark's eelpout	1965	0	66	1988	0	90
Esmark's eelpout	1966	0	50	1989	0	112
Esmark's eelpout	1967	0	19	1990	0	104
Esmark's eelpout	1968	1.13	40	1991	0	103
Esmark's eelpout	1969	0	8	1992	0	89
Esmark's eelpout	1970	0	28	1993	0	77
Esmark's eelpout	1971	0	30	1994	0	64
Esmark's eelpout	1972	0	21	1995	0	85
Fourbeard rockling	1951	0	26	1973	0	50
Fourbeard rockling	1952	0	50	1974	0	13
Fourbeard rockling	1953	0	45	1975	0	36
Fourbeard rockling	1954	0	40	1976	0.05	41
Fourbeard rockling	1955	0	57	1977	0	47
Fourbeard rockling	1956	0	27	1978	0.14	89
Fourbeard rockling	1957	0	30	1979	0	92
Fourbeard rockling	1958	0	28	1980	0.02	80
Fourbeard rockling	1959	0.14	52	1981	0	49
Fourbeard rockling	1960	0	54	1982	0	71

Table C.1 Total weight for each species and year surveyed on southern Grand Bank. The ordering of the species is alphabetical. The total number of tows  $({\rm N})$  is given for reference.

Table (	C.1	Total	weight	for	each	species	and	year	surveyed	on
souther	n G	rand E	Bank. T	he e	orderin	ng of the	spec	ies is	alphabeti	cal.
The tot	tal n	umber	of tows	(N)	) is giv	en for re	feren	ice.		

		Total			Total	
Species	Year	Weight (kg)	N	Year	Weight (kg)	N
Fourbeard rockling	1961	0.14	55	1984	0	58
Fourbeard rockling	1962	0	64	1985	0	167
Fourbeard rockling	1963	0	56	1986	0	115
Fourbeard rockling	1964	0.28	90	1987	0	110
Fourbeard rockling	1965	0.28	66	1988	0	90
Fourbeard rockling	1966	0	50	1989	0	112
Fourbeard rockling	1967	0	19	1990	0	104
Fourbeard rockling	1968	0.14	-40	1991	0	103
Fourbeard rockling	1969	0	8	1992	0	89
Fourbeard rockling	1970	0.23	28	1993	0	77
Fourbeard rockling	1971	0	30	1994	0	64
Fourbeard rockling	1972	0	21	1995	0	85
Greenland halibut	1951	0	26	1973	1.05	50
Greenland halibut	1952	0	50	1974	0	13
Greenland halibut	1953	0	45	1975	0.78	36
Greenland halibut	1954	0	-40	1976	3.09	41
Greenland halibut	1955	0	57	1977	6.79	47
Greenland halibut	1956	0	27	1978	10.62	89
Greenland halibut	1957	0.67	30	1979	42.52	92
Greenland halibut	1958	2.67	28	1980	14.6	80
Greenland halibut	1959	4.98	52	1981	11.5	49
Greenland halibut	1960	1.87	54	1982	3.8	71
Greenland halibut	1961	3.63	55	1984	0.17	58
Greenland halibut	1962	2.97	64	1985	0.54	167
Greenland halibut	1963	22.44	30	1985	4.52	115
Greenland halibut	1964	33.32	90	1987	4.90	110
Greenland halibut	1960	40	00	1988	3.78	90
Greenland halibut	1966	20.41	50	1989	1.18	112
Greenland halibut	1967	33.98	19	1990	3.69	102
Greenland hallout	1900		-	1991	9.43	203
Greenland halibut	1909	2	20	1992	7.20	77
Greenland halibut	1970	4.00	20	1004	1.42	64
Greenland hallbut	1971	4.30	30	1005	12.62	85
Greenland natious	1051	6996 10	26	1973	29.4	50
Tiaddock .	1052	5205 16	50	1074	0	13
Haddock	1952	11176 22	45	1975	6 35	36
Haddock	1054	11020 20	40	1076	6.8	41
Waddock	1055	11543.03	57	1977	11.1	47
Haddock	1056	12396 57	27	1978	31 27	89
Haddock	1957	5506.08	30	1979	35.19	92
Haddock	1958	4217.07	28	1980	76	80
Haddock	1959	8398.19	52	1981	119.95	49
Waddock	1960	11649.92	54	1982	502.38	71
Haddock	1961	7773.31	55	1984	1179.6	58
Haddock	1962	5762.97	64	1985	1063.25	167
Haddock	1963	1507.25	56	1986	265.85	115
Haddock	1964	1856.81	90	1987	1289.07	110
Haddock	1965	644.56	66	1988	1739.51	90
Haddock	1966	1017.64	50	1989	142.26	112
Haddock	1967	0.45	19	1990	170.94	104
Haddock	1968	183.39	40	1991	102.61	103

		Track			Tread	
Snacias	Vear	Weight (kg)	N	Year	Weight (kg)	N
operio	1004					-
Haddock	1969	1.82	8	1992	27.35	89
Haddock	1970	7.39	28	1993	75.07	77
Haddock	1971	187.1	30	1994	68.81	64
Haddock	1972	5.21	21	1995	37.06	85
Hookear sculpin	1951	0	26	1973	0	50
Hookear sculpin	1952	0	50	1974	0	13
Hookear sculpin	1953	0	45	1975	0.14	36
Hookear sculpin	1954	0	40	1976	0.05	41
Hookear sculpin	1955	0	57	1977	0.23	47
Hookear sculpin	1956	0	27	1978	0.12	89
Hookear sculpin	1957	0	30	1979	0.06	92
Hookear sculpin	1958	0	28	1980	0	80
Hookear sculpin	1959	0.07	52	1981	0	49
Hookear sculpin	1960	0.06	54	1982	0	71
Hookear sculpin	1961	0.14	55	1984	0	58
Hookear sculpin	1062	0.03	64	1985	0	167
Hookear sculpin	1963	0.01	56	1986	0	115
Hookean soulpin	1064	0.01	90	1987	0.01	110
Hookear sculpin	1065	0.75	66	1088	0	90
Hookear sculpin	1066	0.00	50	1989	0	112
Hookear sculpin	1067	0.1	10	1000	0.01	104
Hookear sculpin	1000	0	10	1001	0.01	107
Hookear sculpin	1060	0		1002	0.02	80
Hookear scuipin	1909	0	20	1002	0.02	77
Hookear sculpin	1071		20	1004	0.02	64
Hookear sculpin	1971	0	01	1005	0.01	95
hookear sculpin	1972	0	21	1077	0.01	50
Longhn nake	1901	10.00	20	1074	0	12
Longin hake	1052	10.77	10	1075	1.14	26
Longfin nake	1953	0	40	1076	0.96	41
Longhn hake	1954	0	40	1077	0.50	41
Longha hake	1935	0	07	1070	177.9	20
Longhn hake	1999	0	21	1910	17.5	00
Longfin hake	1957	0	30	1979	0.72	92
Longhn hake	1958	0	20	1980	4	00
Longfin hake	1959	0.9	52	1981	01	-19
Longhn hake	1960	2.14	34	1964	0.1	71
Longfin hake	1951	0	00	1984	0.00	167
Longhn hake	1962	0	04	1960	0.03	107
Longfin hake	1963	3.05	00	1980	0.5	115
Longfin hake	1964	29.49	90	1987	0.5	110
Longfin hake	1965	2.72	66	1988	0	90
Longfin hake	1966	6.94	50	1989	0.00	112
Longfin hake	1967	0	19	1990	0	104
Longfin hake	1968	1.81	40	1991	0.11	103
Longfin hake	1969	0	8	1992	0.12	89
Longfin hake	1970	0	28	1993	0	<i>a</i>
Longfin hake	1971	0	30	1994	0.05	64
Longfin hake	1972	0	21	1995	0	85
Longhorn sculpin	1951	0	26	1973	72.63	50
Longhorn sculpin	1952	1.77	50	1974	74.02	13
Longhorn sculpin	1953	1.06	45	1975	28.34	36
Longhorn sculpin	1954	2.48	40	1976	97.09	41

Table C.1 Total weight for each species and year surveyed on southern Grand Bank. The ordering of the species is alphabetical. The total number of tows (N) is given for reference.

Canadian	Vere	Total	N	Vere	Total	\$7
opecies	tear.	weight (kg)	14	reat	weight (kg)	14
Longhorn sculpin	1955	0	57	1977	23.8	47
Longhorn sculpin	1956	0.79	27	1978	33.7	89
Longhorn sculpin	1957	2.83	30	1979	117.51	92
Longhorn sculpin	1958	3.95	28	1980	108.25	80
Longhorn sculpin	1959	11.32	52	1981	16.1	49
Longhorn sculpin	1960	3.66	54	1982	114.5	71
Longhorn sculpin	1961	15.09	55	1984	32.29	58
Longhorn sculpin	1962	11.17	64	1985	314.57	167
Longhorn sculpin	1963	9.48	56	1986	83.37	115
Longhorn sculpin	1964	27.23	90	1987	100.1	110
Longborn sculpin	1965	14.12	66	1988	20.92	90
Longhorn sculpin	1966	3.05	50	1989	24.27	112
Longborn sculpin	1967	1.49	19	1990	29.68	104
Longhorn sculpin	1968	1.83	40	1991	26.39	103
Longhorn sculpin	1969	1.99	8	1992	3.82	89
Longhorn sculpin	1970	1.13	28	1993	12.08	77
Longhorn sculpin	1971	9.31	30	1994	2 33	64
Longhorn sculpin	1972	125.41	21	1995	4.67	85
Mailed eculoin	1951	0	52	1973	0.1	100
Mailed coulpin	1052	0	100	1974	0.24	26
Mailed couloin	1052	0	00	1975	0.43	79
Mailed sculpin	1054	0	80	1976	0.43	82
Mailed coulpin	1055	0	114	1977	0.09	04
Mailed sculpin	1056	0.02	54	1079	1 20	179
Mailed sculpin	1950	0.05	60	1979	0.56	184
Mailed sculpin	1907	0	56	1980	0.30	160
Mailed sculpin	1050	0.1	104	1081	1.6	08
Mailed sculpin	1939	0.00	108	1002	0.01	149
Mailed sculpin	1960	0.05	110	1064	0.01	116
Mailed sculpin	1001	0.0	120	1095	0.02	224
Mailed sculpin	1902	0.20	112	1986	0.02	220
Mailed sculpin	1064	0.35	190	1087	0.12	220
Mailed sculpin	1004	0.20	122	1000	0.02	190
Malled scupin	1966	0.02	100	1080	0.03	224
Mailed sculpin	1960	0.03	29	1909	0.16	209
Mailed sculpin	1007	0.02	80	1001	0.20	200
Mailed sculpin	1000	0.03	16	1991	0.57	178
Mailed sculpin	1070	0.02	56	1007	0.07	154
Mailed sculpin	1970	0.03	60	1004	0.02	100
Mailed sculpin	1971	0.69	42	1005	0.07	120
Maalla sella	1072	0.00	26	1072	0.00	50
Martin-spile	1931	8.00	20	1074	0.05	12
Mariin-spike	1952	0.10	45	1075	0.27	13
Marin-spike	1933	0	40	1070	0.37	41
Marin-spike	1954	0	40	1077	0.1	41
Mariin-spike	1955	0	07	1079	0.45	47
Marin-spice	1900	0	20	1070	0.40	00
Marile apile	1059	0	28	1090	0.23	80
Martin-spike	1908	5 42	±0	1091	1.4	40
Marin-spike	1909	0.43	52	1080	0.1	
Maalla miles	1001	0.9	55	1094	0.09	59
Marlin spike	1961	7.12	64	1085	0.08	167
IN CALLED TO A CARE	1.002	6.14			0.02	494

Table C.1 Total weight for each species and year surveyed on southern Grand Bank. The ordering of the species is alphabetical. The total number of tows (N) is given for reference.

		Total			Total	
Species	Year	Weight (kg)	N	Year	Weight (kg)	N
Marlin-spike	1963	40.1	56	1986	0	115
Marlin-spike	1964	44.03	90	1987	0.59	110
Marlin-spike	1965	16.02	66	1988	0.17	90
Marlin-spike	1966	12.64	50	1989	0.07	112
Marlin-spike	1967	0	19	1990	0	104
Marlin-spike	1968	34	40	1991	0.27	103
Marlin-spike	1969	0	8	1992	0	89
Marlin-spike	1970	0	28	1993	0	77
Marlin-spike	1971	0	30	1994	0	64
Marlin-spike	1972	0	21	1995	0	85
Monkfieh	1951	0	26	1973	27.44	50
Monkfish	1952	52.46	50	1974	0	13
Monkfish	1953	22.08	45	1975	5.44	36
Monkfish	1954	6.56	40	1976	13.61	41
Monkfish	1955	5.98	57	1977	0.95	47
Monkfieh	1956	1	27	1978	22.13	89
Monkfish	1957	0.5	30	1979	2.95	92
Monkfish	1958	13.97	28	1980	45	80
Monkfieh	1959	89.36	52	1981	22	49
Monkfish	1960	44.58	54	1982	2.95	71
Monkfish	1961	142.87	55	1984	0	58
Monkfieh	1967	148.66	64	1985	36.94	167
Monkfish	1963	68.6	56	1986	25.73	115
Monkfish	1964	39.47	90	1987	47.72	110
Monkfieh	1965	39.82	66	1988	46.48	90
Monkfish	1966	23.59	50	1989	12.95	112
Monkfish	1967	0	19	1990	13.07	104
Monkfish	1968	13.6	40	1991	18.9	103
Monkfish	1969	0	8	1992	0.62	89
Monkfish	1970	0	28	1993	4.65	77
Monkfish	1971	0	30	1994	9.54	64
Monkfish	1972	0	21	1995	0	85
Northern sand lance	1951	0	26	1973	41.29	50
Northern sand lance	1952	1.05	50	1974	34.02	13
Northern sand lance	1953	0.7	45	1975	252.19	36
Northern sand lance	1954	0.22	40	1976	26.07	41
Northern sand lance	1955	7.48	57	1977	70.99	47
Northern and lance	1956	0.62	27	1978	483.3	89
Northern sand lance	1957	1.05	30	1979	469.83	92
Northern sand lance	1958	52.88	28	1980	528.55	80
Northern sand lance	1959	1.14	52	1981	89.15	49
Northern sand lance	1960	27.39	54	1982	182.36	71
Northern sand lance	1961	1.2	55	1984	1.66	58
Northern sand lance	1962	0.12	64	1985	3.54	167
Northern sand lance	1963	0.18	56	1986	0.24	115
Northern sand lance	1964	3.15	90	1987	1.19	110
Northern sand lance	1965	1.85	66	1988	0.8	90
Northern sand lance	1966	1.02	50	1989	3.57	112
Northern sand lance	1967	2.66	19	1990	0.42	104
Northern sand lance	1968	0.32	40	1991	1.45	103
Northern sand lance	1969	3.18	8	1992	0.58	89
Northern sand lance	1970	91.56	28	1993	0.75	77

Table C.1 Total weight for each species and year surveyed on southern Grand Bank. The ordering of the species is alphabetical. The total number of tows  $\left(N\right)$  is given for reference.

Species	Year	Total Weight (kg)	N	Year	Total Weight (kg)	N
Northern cand lance	1071	26.25	20	1004	0.90	64
Northern sand lance	1072	25 28	21	1005	0.47	85
Pollock	1051	66 52	26	1072	0.47	50
Pollock	1052	122.02	50	1973	0	12
Pollock	1052	123.03	30	1075	0	26
Pollock	1903	14.40	40	1973	0.01	30
Pollock	1055	14.40	40	1976	0.91	41
Pollock	1056	602.40	37	1079	1.92	80
Pollock	1057	7.49	20	1070	7.40	02
Pollock	1059	2.00	28	1080	1.45	80
Dellock	1050	17 66	50	1091	0	40
Pollock	1969	210.07	54	1981	07	71
Dellock	1001	20.41		1084	0.1	59
Pollock	1062	10.41	6.1	1005	17	167
Delleek	1062	150.00	50	1086	1.05	115
Pollock	1064	130.00	00	1087	16.42	110
Dellesk	1005	12 20	00	1000	0.5	00
Pollock	1066	27.97	50	1090	12.7	112
Pollock	1067	31.81	10	1000	15.7	104
Pollock	1907	0	19	1990	10.13	102
Pollock	1908	0	40	1000	10.44	103
Pollock	1909	0	0	1992	0	09
Pollock	1970	102.05	28	1004	0	84
Pollock	1971	103.65	30	1005	0	04
Pollock	1972	174.00	21	1995	e20 e1	150
Dedish	1931	10000.00	10	1973	039.01	100
Rearish	1952	18930.93	150	1974	001 00	39
Redhsh	1953	10132.08	135	1975	694.78	108
Rednish	1994	2116.6	120	1970	1/0.41	123
Redrish	1955	1890.92	1/1	1977	3003.0	141
Redhsh	1955	3016.57	81	1978	120.7	207
reeansh	1957	7910.48	90	1979	1222.00	210
Redhsh	1958	2247.04	84	1980	194.30	240
Reansh	1909	0002.77	130	1981	1010 5	147
Redhsh	1960	19059.76	162	1982	4340.5	213
Rednan	1901	P.100P	105	1964	239.00	1/4
Redhsh	1962	7739.65	192	1985	563.49	301
Rednsn	1903	33/0.8	108	1990	4024.09	340
Redfish	1964	5108.69	270	1987	463.85	330
Reansn	1965	2/04.55	198	1988	001.03	270
Redhsh	1966	3679.19	150	1989	308.80	330
Rednsn	1907	09.80	54	1990	1313.00	312
Redhsh	1968	6609.13	120	1991	29.9	309
Redfish	1969	0	24	1992	107.14	267
Redhsh	1970	375.82	84	1993	1295.73	231
Redlish	1971	460.4	90	1994	1162.21	192
Redhsh	1972	0.6	63	1995	9282.72	200
Sea raven	1951	0	26	1973	157.39	30
Sea raven	1952	8.73	50	1974	62.58	13
Sea raven	1953	8.73	45	1975	78.71	30
Sea raven	1954	15.71	40	1976	158.98	41
Sea raven	1955	4.99	57	1977	115.65	
Sea raven	1956	12.48	27	1978	189.4	69

Table C.1 Total weight for each species and year surveyed on southern Grand Bank. The ordering of the species is alphabetical. The total number of tows (N) is given for reference.

Species		Total		10.02	Total	
	Year	Weight (kg)	N	Year	Weight (kg)	N
Sea raven	1957	3.49	30	1979	176.42	92
Sea raven	1958	13.48	28	1980	238.35	80
Sea raven	1959	17.45	52	1981	89.5	49
Sea raven	1960	0	54	1982	279	71
Sea caven	1961	47.21	55	1984	177.04	58
Sea raven	1962	31.98	64	1985	392.96	167
Sea raven	1963	16.25	56	1986	370.68	115
Sea caven	1964	45.29	90	1987	259.22	110
Sea raven	1965	35.57	66	1988	115.83	90
Sea raven	1966	17.9	50	1989	137.2	112
Sea raven	1967	13.61	19	1990	159.15	104
Sea raven	1968	29.94	40	1991	126.05	103
Sea raven	1969	26.31	8	1992	89.34	89
Sea raven	1970	23.14	28	1993	126.21	77
Sea raven	1971	30.38	30	1994	59,76	64
Sea raven	1972	26.08	21	1995	41.63	85
Shorthorn sculpin	1951	0	26	1973	0	50
Shorthorn sculpin	1952	0.75	50	1974	0.91	13
Shorthorn sculpin	1953	1.5	45	1975	0.23	36
Shorthorn sculpin	1054	0	40	1976	0	41
Shorthorn sculpin	1955	0	57	1977	0.32	47
Shorthorn couloin	1056	0	27	1978	0.91	89
Chesthern coulpin	1057	0	20	1070	0.45	92
Shorthorn sculpin	1058	0	28	1980	0.40	80
Shorthorn sculpin	1050	0	52	1981	0	49
Shorthorn sculpin	1960	0	54	1982	0.9	71
Charthorn couloin	1061	0	55	1064	2.49	58
Shorthorn sculpin	1001	0.68	64	1085	2 22	167
Chesthere couloin	1062	0.00	56	1086	0	115
Shorthorn Sculpin	1064	0	00	1097	2 57	110
Chosthern coulpin	1065	0	66	1088	0.25	90
Shorthorn sculpin	1066	0	50	1080	1	112
Chosthorn couloin	1067	0	10	1000	3 20	104
Shorthorn sculpin	1068	0	40	1991	0.84	103
Shorthorn sculpin	1908	0	8	1002	5 18	80
Shorthorn sculpin	1070	0	20	1002	1.00	77
Shorthorn sculpin	1970	0	20	1004	0.33	64
Charthan sculpin	1070	0	21	1005	0.00	95
Shorthorn scuipin	1051	0.04	26	1973	0	50
Cilum hale	1052	2.79	50	1974	0	12
Silver have	1902	0124 60	45	1075	1 26	26
Cilcum hales	1054	14.64	40	1076	1 50	41
Suver have	1904	14.04	67	1077	0.45	47
Suver nake	1900	6 24	27	1078	56 56	80
Cilum halo	1950	0.24	20	1979	29.78	92
Cilver hale	1059	4.00	28	1080	10.1	80
Suver nake	1908	4.00	40	1001	219 55	40
Sliver naké	1959	2.00	54	1082	318.00	-49
Sliver nake	1960	34.38	04	1982	0.1	11
Suver nake	1961	31.97	00	1984	27.00	36
Silver hake	1962	9.07	04	1985	35.28	107
Suver nake	1963	0	00	1980	123.75	115

Table C.1 Total weight for each species and year surveyed on southern Grand Bank. The ordering of the species is alphabetical. The total number of tows (N) is given for reference.

		Total			Total		
Species	Year	Weight (kg)	N	Year	Weight (kg)	N	
Silver hake	1965	0.23	66	1988	0.42	90	
Silver hake	1966	0	50	1989	1.08	112	
Silver hake	1967	Ő.	19	1990	1.45	104	
Silver hake	1968	0.43	40	1991	1.35	103	
Silver hake	1969	0	8	1992	0.91	89	
Silver hake	1970	1.36	28	1993	0.09	77	
Silver hake	1971	0	30	1994	1.04	64	
Silver hake	1972	0	21	1995	0	85	
Smooth skate	1951	0.75	26	1973	4.98	50	
Smooth skate	1952	12.75	50	1974	0	13	
Smooth skate	1953	3.75	45	1975	0.45	36	
Smooth skate	1954	9.75	40	1976	0	41	
Smooth skate	1955	1.5	57	1977	2.26	47	
Smooth skate	1956	4 74	27	1978	37.13	89	
Smooth skate	1957	0.99	30	1979	3.17	92	
Smooth skate	1958	2.66	28	1980	5	80	
Smooth skate	1959	10.58	52	1981	8	49	
Smooth skate	1960	9.66	54	1982	4	71	
Smooth skate	1961	2.27	55	1984	1.41	58	
Smooth skate	1962	11.57	64	1985	9.84	167	
Smooth skate	1963	14.21	56	1986	2.9	115	
Smooth ekate	1964	37.69	90	1987	2.99	110	
Smooth skate	1965	13.06	66	1988	2.08	90	
Smooth skate	1966	13.84	50	1989	0.66	112	
Smooth skate	1967	0	19	1990	8.13	104	
Smooth skate	1968	0.23	40	1991	9.94	103	
Smooth skate	1969	0	8	1992	4.47	89	
Smooth skate	1970	0	28	1993	0.83	77	
Smooth skate	1971	6.36	30	1994	0.79	64	
Smooth skate	1972	0	21	1995	1.99	85	
Sping lumpfich	1951	0	52	1973	0	100	
Spiny lumpfish	1952	0	100	1974	0	26	
Spiny lumpfish	1953	0	90	1975	0.14	72	
Solny lumpfich	1054	0	80	1976	0	82	
Spiny lumpfieh	1955	0	114	1977	0	94	
Spiny lumpfieh	1956	0	54	1978	0.08	178	
Spiny lumpfish	1957	0	60	1979	0.2	184	
Spiny lumpfieh	1958	0	56	1980	0.11	160	
Spiny lumofieh	1959	0	104	1981	0	98	
Spiny lumpfish	1960	0	108	1982	0.15	142	
Spiny lumpfieh	1961	0.08	110	1984	0	116	
Spiny lumofeh	1962	0	128	1985	0	334	
Spiny lumpfish	1963	0.04	112	1986	0.12	230	
Spiny lumpfish	1964	0.04	180	1987	0.2	220	
Coiny humpfish	1065	0.12	132	1988	0.12	180	
Spiny lumpfieh	1966	0.12	100	1989	0.26	224	
Spiny lumpfish	1967	0	38	1990	0.24	208	
Spiny lumpfish	1968	0	80	1991	0.06	206	
Spiny lumpfish	1969	0	16	1992	0.66	178	
Spiny lumpfish	1970	0	56	1993	0.25	154	
Spiny lumphed	1971	0	60	1994	0.19	128	
Spiny lumpfish	1972	0.23	42	1995	0.57	170	

Table C.1 Total weight for each species and year surveyed on southern Grand Bank. The ordering of the species is alphabetical. The total number of tows  $({\rm N})$  is given for reference.

		Total			Total	
Species	Year	Weight (kg)	N	Year	Weight (kg)	N
		0 . 0				
Spotted wolffish	1951	0	26	1973	0	50
Spotted wolffish	1952	0	50	1974	0	13
Spotted wolffish	1953	0	45	1975	0	36
Spotted wolffish	1954	0	40	1976	0	41
Spotted wolffish	1955	0	57	1977	0	47
Spotted wolffish	1956	0	27	1978	16.86	89
Spotted wolffish	1957	0	30	1979	0	92
Spotted wolffish	1958	0	28	1980	0	80
Spotted wolffish	1959	0	52	1981	7.5	49
Spotted wolffish	1960	0	54	1982	4.1	71
Spotted wolffish	1961	3.63	55	1984	17.02	58
Spotted wolffish	1962	7.71	64	1985	0	167
Spotted wolffish	1963	7.53	56	1986	0	115
Spotted wolffish	1964	13.61	90	1987	0	110
Spotted wolffish	1965	15.25	66	1988	11.62	90
Spotted wolffish	1966	0	50	1989	0	112
Spotted wolffish	1967	0	19	1990	0	104
Spotted wolffish	1968	30.84	40	1991	9.34	103
Spotted wolffish	1969	0	8	1992	0	89
Spotted wolffish	1970	9.53	28	1993	0	77
Spotted wolffish	1971	0	30	1994	0	64
Spotted wolffish	1972	0	21	1995	0	85
Stringd wolffish	1951	6.95	26	1973	180.99	50
Stringd wolffieh	1952	39.91	50	1974	87.55	13
Strined wolffish	1953	25.86	45	1975	186.66	36
Striped wolffish	1954	37.42	40	1976	195.52	41
Striped wolffieh	1055	14.96	57	1977	86.18	47
Striped wolffish	1956	7.99	27	1978	228.47	89
Striped wolffish	1957	66.36	30	1979	442.27	92
Carined wolffich	1059	1 99	28	1980	462.65	80
Carlored wollman	1050	61.98	52	1081	65	49
Stained wolffich	1060	157.3	54	1982	630.5	71
Striped wolldish	1061	64.41	55	1984	305.02	58
Striped wolfish	1062	151.04	64	1095	575 19	167
Striped wolffish	1062	55.42	55	1986	751 15	115
Striped wolfish	1064	154.94	00	1087	730 7	110
Striped wollinsh	1004	50.69	66	1099	497.67	90
Striped wolfnah	1966	282.26	50	1080	547 55	112
Striped wolmsn	1900	10 02	10	1000	697.04	104
Striped wormsn	1000	19.26	10	1001	299 92	103
Striped wolmsn	1900	18.30	0	1002	255 12	80
Striped woimsn	1969	1 26	28	1002	204 65	77
Striped wolmsn	1970	E.44	20	1004	425 12	64
Striped wolmsn	1971	0.44	30	1004	152 57	95
Striped wolfnsh	1972	30.29	24	1072	1204 50	50
I norny skate	1951	1004.45	20	1074	120 51	12
Thorny skate	1952	1228.45	45	1075	\$16.20	36
1 norny skate	1953	1328.45	40	1076	1264 20	41
Thorny skate	1954	1351.48		1077	1104.90	47
Thorny skate	1955	239.15	07	1070	1104.80	80
Thorny skate	1956	177.87	27	1070	906.59	02
Thorny skate	1957	651.52	30	1090	139.62	80
L'DOFDI EXDIR	14258	921.10		* dou	*730'O	~~~

Table C.1 Total weight for each species and year surveyed on southern Grand Bank. The ordering of the species is alphabetical. The total number of tows  $({\rm N})$  is given for reference.

		Total			Total	
Species	Year	Weight (kg)	N	Year	Weight (kg)	N
Thorny skate	1050	2275 78	52	1981	403.4	.10
Thorny skate	1960	1491.5	54	1982	1233.6	71
Thorny skate	1961	1600.89	55	1984	926.86	58
Thorny skate	1962	2659.96	64	1985	3740.85	167
Thorny skate	1063	2065.85	56	1986	2631 39	115
Thorny skate	1964	2380.83	90	1987	1379.21	110
Thorny skate	1965	1709.92	66	1988	1346.97	90
Thorny skate	1966	805 15	50	1989	1261.72	112
Thorny skate	1967	655.9	19	1990	1656.61	104
Thorny skate	1968	467.45	40	1991	670.69	103
Thorny skate	1969	238.15	8	1992	786.32	89
Thorny skate	1970	940.78	28	1993	455.89	77
Thorny skate	1971	550.88	30	1994	234.05	64
Thorny skate	1972	555.88	21	1995	370.47	85
Threebeard rockling	1951	0	26	1973	0	50
Threebeard mckling	1952	0	50	1974	0	13
Threeheard rockling	1953	0	45	1975	ő	36
Threebeard rockling	1954	0	40	1976	0.05	41
Threeheard rockling	1955	0	57	1977	0.09	47
Threeheard rockling	1956	0	27	1978	0.1	89
Threeheard rockling	1957	0	30	1979	0	92
Threeheard rockling	1958	0	28	1980	0	80
Threeheard rockling	1050	0	52	1081	ő	49
Threeheard rockling	1060	0	54	1982	0	71
Threeheard rockling	1061	0.00	55	1084	0	19
Threeheard rockling	1962	0.00	64	1985	0	167
Threeheard rockling	1963	0	56	1986	0	115
Threeheard rockling	1964	0	90	1987	0	110
Threeheard rockling	1965	0	66	1988	ő	90
Threeheard rockling	1966	0	50	1989	0	112
Threeheard rockling	1967	0	19	1990	0	104
Threebeard rockling	1968	ő	40	1991	0	103
Threeheard mckling	1969	0	8	1992	0	89
Threeheard rockling	1970	0	28	1993	0	77
Threebeard rockling	1971	õ	30	1994	0	64
Threebeard rockling	1972	0	21	1995	0	85
Vahl's eelpout	1951	0	26	1973	0.23	50
Vahl's eelpout	1952	0	50	1974	0	13
Vahl's eelpout	1953	0	45	1975	0.18	36
Vahl's eelpout	1954	0	40	1976	0.45	41
Vahl's eelpout	1955	0	57	1977	0	47
Vahl's eelpout	1956	0	27	1978	25.56	89
Vahl's eelpout	1957	0	30	1979	1.13	92
Vahl's celpout	1958	0	28	1980	0.5	80
Vahl's eelpout	1959	0	52	1981	0	49
Vahl's eelpout	1960	0	54	1982	0	71
Vahl's eelpout	1961	0	55	1984	0	58
Vahl's eelpout	1962	1.29	64	1985	0.66	167
Vahl's eelpout	1963	0	56	1986	0	115
Vahl's eelpout	1964	2.26	90	1987	0	110
Vahl's eelpout	1965	0.23	66	1988	0	90
Vahl's eelpout	1966	0.42	50	1989	0	112

Table C.1 Total weight for each species and year surveyed on southern Grand Bank. The ordering of the species is alphabetical. The total number of tows (N) is given for reference.

The base of the first state		Total			Total	
Species	Year	Weight (kg)	N	Year	Weight (kg)	N
Vahl's eelpout	1967	7.71	19	1990	0.04	104
Vahi's eelpout	1968	0	40	1991	0	103
Vahi's celnout	1969	0	8	1992	0	89
Vahl's eelpout	1970	0	28	1993	0	77
Vahl's celpout	1971	0	30	1994	0	64
Vahi's eeloout	1972	0	21	1995	0	85
White hake	1951	608.73	26	1973	116.35	50
White hake	1952	1635 18	50	1974	0	13
White hake	1953	1139 15	45	1975	24.04	36
White hake	1954	328 44	40	1976	95.93	41
White hake	1955	59.39	57	1977	24 94	47
White hake	1956	88.81	27	1978	472.67	89
White hake	1957	86.81	30	1979	220.65	92
White hake	1958	51.85	28	1980	57.5	80
White hake	1959	696.8	52	1981	331.3	49
White hake	1960	1079 53	54	1982	0.1	71
White hake	1961	1741 57	55	1984	28 64	58
White hake	1062	1026 75	64	1085	61.42	167
White hake	1062	1494 20	50	1096	21.59	115
White hake	1963	7725.07	90	1987	157.66	110
White hake	1904	116.69	66	1066	126.12	00
White bake	1066	040.07	50	1080	20.2	112
White hale	1900	545.01	10	1000	20.3	104
White hale	1967	191 90	40	1001	15.97	104
White hales	1900	101.09	0	1002	10.70	205
White hale	1909	21.00	00	1002	10.75	0.9
white hate	1970	31.98	20	1993	18.00	
White hake	1971	43.37	30	1005	2.41	95
White hate	1972	144.01	21	1990	000 00	50
witch nounder	1951	144.21	20	1973	200.23	30
Witch nounder	1952	200.21	45	1974	59.06	13
Witch flounder	19054	332.13	40	1910	079.64	30
witch nounder	1904	241.99	40	1970	210.04	41
Witch flounder	1955	120.00	07	1079	74.02	
Witch Bounder	1900	132.37	20	1070	77.00	00
witch nounder	1957	91.3	30	1979	13.33	92
Witch nounder	1958	130.23	28	1980	1/3.5	80
Witch nounder	1939	014.13	54	1981	105	49
witch hounder	1960	103.9	34	1962	107	71
witch nounder	1961	329.99	55	1984	111.05	38
Witch flounder	1962	1161.2	04	1985	489.37	167
witch hounder	196.5	342.67	30	1990	229	115
witch nounder	1964	841.63	90	1987	250.7	110
Witch flounder	1965	286.03	66	1988	419.4	90
witch hounder	1966	517.23	30	1969	200.00	112
witch hounder	1967	4.04	19	1990	192.34	104
witch nounder	1968	136.76	-0	1991	99.99	103
witch nounder	1969	7.48	8	1992	154.16	69
witch flounder	1970	8.75	28	1993	50.7	
Witch flounder	1971	18.15	30	1994	85.74	04
witch nounder	1972	25.85	21	1995	29.63	60
Yellowtail flounder	1951	2.99	26	1973	1906.37	30
Yellowtail flounder	1952	75.49	51	1974	1077.28	13

Table C.1 Total weight for each species and year surveyed on southern Grand Bank. The ordering of the species is alphabetical. The total number of tows: (N) is given for reference.

Table C.1 Total weight for each species and year surveyed on southern Grand Bank. The ordering of the species is alphabetical. The total number of tows (N) is given for reference.

Species	Year	Total Weight (kg)	N	Year	Total Weight (kg)	N
Yellowtail flounder	1953	108.78	45	1975	1606.39	36
Yellowtail flounder	1954	135.37	40	1976	2094.24	41
Yellowtail flounder	1955	34.33	57	1977	2542.19	47
Yellowtail flounder	1956	49.7	27	1978	4463.43	89
Yellowtail flounder	1957	121.69	30	1979	4790.51	92
Yellowtail flounder	1958	257.3	28	1980	4651.87	80
Yellowtail flounder	1959	241.27	52	1981	2114.5	49
Yellowtail flounder	1960	144.23	54	1982	2780.4	71
Yellowtail flounder	1961	298.25	55	1984	3214.53	58
Yellowtail flounder	1962	743.44	64	1985	6815.42	167
Yellowtail flounder	1963	684.93	56	1986	4435.6	115
Yellowtail flounder	1964	1677.53	90	1987	3800.03	110
Yellowtail flounder	1965	1377.25	66	1988	1968.05	90
Yellowtail flounder	1966	1278.44	50	1989	2337.58	112
Yellowtail flounder	1967	493.73	19	1990	2991.73	104
Yellowtail flounder	1968	1104.52	40	1991	2563.03	103
Yellowtail flounder	1969	365.15	8	1992	1533.83	89
Yellowtail flounder	1970	1089.1	28	1993	2217.12	77
Yellowtail flounder	1971	728.24	30	1994	1093.41	64
Yellowtail flounder	1972	1494.15	21	1995	1894.17	85







